

HANDBOOK of CIRCULATION

Analysis and Compilation by

PHILIP L. ALTMAN

Edited by

DOROTHY S. DITTMER

RUDOLPH M. GREBE



Prepared under the direction of the Committee
on the Handbook of Biological Data

DIVISION OF BIOLOGY AND AGRICULTURE
THE NATIONAL ACADEMY OF SCIENCES
THE NATIONAL RESEARCH COUNCIL

W B SAUNDERS COMPANY

Philadelphia and London

© 1959, BY W. B. SAUNDERS COMPANY

COPYRIGHT UNDER THE INTERNATIONAL COPYRIGHT UNION

All rights reserved This book is protected by copyright No part of it may be reproduced in any manner without written permission from the publisher, except for any purpose of the United States Government.

Made in the United States of America

LIBRARY OF CONGRESS CATALOG CARD NUMBER 59-15183

Foreword

The Handbook of Circulation is the tenth in a series of publications*, each containing information, chiefly tabular, in one or more fields of the biological sciences. These handbooks have been prepared under the general direction of the Committee on the Handbook of Biological Data, Division of Biology and Agriculture, National Academy of Sciences-National Research Council.

The information for the present Handbook was prepared and contributed by leading authorities in the field of circulation. The data were assembled, tabulated, and edited by the Handbook staff, then critically reviewed and authenticated by experts in the areas covered in this volume.

On behalf of the Committee, acknowledgment is made to the numerous scientists who have been so liberal with their time and advice; to Wright Air Development Center United States Air Force, the National Institutes of Health of the Public Health Service, the Division of Biology and Medicine of the Atomic Energy Commission, the Office of Naval Research, the Office of the Surgeon General of the Army, and the Army Chemical Center, for generous support and cooperation which have made possible the production of this book. The Air Force participation in this undertaking was carried out under Contract No. AF 33(616)-3972 with the National Academy of Sciences. Dr. J. W. Heim, Aero Medical Laboratory, Wright Air Development Center, served as contract monitor.

THE COMMITTEE ON THE HANDBOOK OF BIOLOGICAL DATA

Theodore C. Byerly, Chairman

W. H. Larrimer, Executive Secretary

David B. Dill
Charles M. Goss
Frank G. Hall
J. W. Heim

Paul E. Howe
Paul J. Kramer
Milton O. Lee
Royal Shanks
H. Burr Steinbach**

Oswald Tippo
Lionel A. Walford
Paul Weiss
Raymond L. Zwemer

** ex officio

HANDBOOK STAFF

W. H. Larrimer, Director

Philip L. Altman, Analyst

Dorothy S. Dittmer, Associate Editor

Rudolph M. Grebe, Associate Editor

Judith P. Bloomer
Nellie F. Brown
Helen D. Carmack
Viola A. Crowder
S. W. Lipsman

Gilbert R. Magee
Jean A. McNeely
Mary J. Noe
Olga G. Slanczak
Marlene L. Stanley

*Handbooks published 1949-1959

Standard Values in Blood	1951
Standard Values in Nutrition and Metabolism	1953
Handbook of Biological Data	1956
Handbook of Respiration	1958
Handbook of Circulation	1959

Handbook of Toxicology, Vol. I, Acute Toxicities	1955
Handbook of Toxicology, Vol. II, Antibiotics	1957
Handbook of Toxicology, Vol. III, Insecticides	1959
Handbook of Toxicology, Vol. IV, Tranquillizers	1959
Handbook of Toxicology, Vol. V, Fungicides	1959

© 1959, BY W. B. SAUNDERS COMPANY

COPYRIGHT UNDER THE INTERNATIONAL COPYRIGHT UNION

All rights reserved. This book is protected by copyright. No part of it may be reproduced in any manner without written permission from the publisher, except for any purpose of the United States Government.

Made in the United States of America

LIBRARY OF CONGRESS CATALOG CARD NUMBER. 59-15183

Foreword

The Handbook of Circulation is the tenth in a series of publications*, each containing information, chiefly tabular, in one or more fields of the biological sciences. These handbooks have been prepared under the general direction of the Committee on the Handbook of Biological Data, Division of Biology and Agriculture, National Academy of Sciences-National Research Council.

The information for the present Handbook was prepared and contributed by leading authorities in the field of circulation. The data were assembled, tabulated, and edited by the Handbook staff, then critically reviewed and authenticated by experts in the areas covered in this volume.

On behalf of the Committee, acknowledgment is made to the numerous scientists who have been so liberal with their time and advice, to Wright Air Development Center United States Air Force, the National Institutes of Health of the Public Health Service, the Division of Biology and Medicine of the Atomic Energy Commission, the Office of Naval Research, the Office of the Surgeon General of the Army, and the Army Chemical Center, for generous support and cooperation which have made possible the production of this book. The Air Force participation in this undertaking was carried out under Contract No. AF 33(616)-3972 with the National Academy of Sciences. Dr. J. W. Heim, Aero Medical Laboratory, Wright Air Development Center, served as contract monitor.

THE COMMITTEE ON THE HANDBOOK

OF BIOLOGICAL DATA

Theodore C. Byerly, Chairman

W. H. Larrimer, Executive Secretary

David B. Dill
Charles M. Goss
Frank G. Hall
J. W. Heim

Paul E. Howe
Paul J. Kramer
Milton O. Lee
Royal Shanks
H. Burr Steinbach**

Oswald Tippo
Lionel A. Walford
Paul Weiss
Raymond L. Zwemer

** *ex officio*

HANDBOOK STAFF

W. H. Larrimer, Director

Philip L. Altman, Analyst

Dorothy S. Dattmer, Associate Editor

Rudolph M. Grebe, Associate Editor

Judith P. Bloomer
Nellie F. Brown
Helen D. Carmack
Viola A. Crowder
S. W. Lipsman

Gilbert R. Magee
Jean A. McNeely
Mary J. Noe
Olga G. Stanczak
Marlene L. Stanley

*Handbooks published 1949-1959

Standard Values in Blood	1951
Standard Values in Nutrition and Metabolism	1953
Handbook of Biological Data	1956
Handbook of Respiration	1958
Handbook of Circulation	1959

Handbook of Toxicology, Vol. I, Acute Toxicities	1955
Handbook of Toxicology, Vol. II, Antibiotics	1957
Handbook of Toxicology, Vol. III, Insecticides	1959
Handbook of Toxicology, Vol. IV, Tranquilizers	1959
Handbook of Toxicology, Vol. V, Fungicides	1959

CONTRIBUTORS AND REVIEWERS	x
INTRODUCTION	1-3
ABBREVIATIONS AND SYMBOLS	1-3
HANDBOOK OF CIRCULATION	1-3

I. CIRCULATORY ANATOMY

1. Development of the Circulatory System: Vertebrates	1
Part I. Frog	1
Part II. Chick	2
Part III. Pig	3
Part IV. Man	4
2. Heart: Man (Schematic Drawing)	6
3. Principal Arteries and Pulmonary Veins: Man (Schematic Drawing)	7
4. Principal Veins and Pulmonary Arteries: Man (Schematic Drawing)	8
5. Lymphatics: Man (Schematic Drawing)	9
6. Comparative Anatomy of the Circulatory System: Vertebrates	10
Part I. Heart	10
Part II. Blood Vessels	12
Part III. Lymphatics	16
7. Circumferential Growth of the Heart, Aorta, and Superior Vena Cava: Man (Graph)	18
8. Dimensional Growth of the Heart during First Year: Man (Graphs)	19
9. Heart, Linear Dimensions: Man	19
10. Heart, Surface Dimensions: Man	19
11. Ostial Breadths: Man	20
12. Growth of Arterial Lumina: Man	20
13. Thickness of Arterial Walls: Man	20
14. Dimensions of Main Arterial Trunks: Man	21
Part I. Upper Abdomen	21
Part II. Lower Abdomen	21
15. Atrio-Ventricular Weight Ratio during First Year: Man (Graph)	22
16. Weight of Ventricles: Man	22
17. Heart Weight during First Year: Man	23
18. Heart Weight at Various Ages: Man	23
19. Heart Weight: Vertebrates	24
Part I. Man	24
Part II. Mammals Other Than Man	25
Part III. Birds	30
Part IV. Reptiles and Amphibians	32
Part V. Fishes	33
20. Density, Surface, and Maximum Diffusion Distance of Capillaries in Various Tissues: Vertebrates	34
21. Capillary Vascularity of Muscle: Vertebrates	35
Part I. Skeletal Muscle	35
Part II. Smooth Muscle	37
Part III. Cardiac Muscle	37

II. CHEMICAL COMPOSITION AND PHYSICAL PROPERTIES

22. Chemical Composition of the Heart: Vertebrates	38
Part I. Man	38
Part II. Carnivores	39
Part III. Rodents	42
Part IV. Artiodactyla and Perissodactyla	45
Part V. Birds	46
23. Capillary Filtration Coefficients: Vertebrates	46
24. Fractional Extraction of Diffusible Substances from Capillaries: Man, Dog	47
25. Transcapillary Exchange of Large Molecules: Man, Dog, Guinea Pig	48
26. Capillary Permeability Coefficients of Various Substances in Tissue: Cat	48

27. Permeability of Cell Surfaces: Animals, Plants.	49
Part I: To Water.	49
Part II: To Inorganic Ions.	50
Part III: To Organic Substances.	52
28. Cardiovascular and Pulmonary Reflexes. Mammals.	55
29. Electrical Properties of Cardiac Tissue: Vertebrates	56
Part I: Resting and Action Potentials of Single Fibers.	56
Part II: Action Potential Characteristics of Single Fibers.	58
Part III: Electrical Constants of Single Purkinje Fibers.	59
30. Effect of pH on the Heart: Dog	59
31. Elasticity of the Arterio-venous System: Man, Dog.	60
Part I: Aorta.	60
Part II: Venous System	62

III. BLOOD VOLUMES

32. Blood Volumes.	64
Part I: Man	64
Part II: Mammals Other Than Man.	65
Part III: Vertebrates Other Than Mammals; Crustaceans and Mollusks	67
Part IV: Insects	69
33. Concentration of Blood in Tissue: Vertebrates	70
Part I: Total Blood Volume	70
Part II: Residual Blood Volume.	71

IV. CARDIAC OUTPUT

34. Cardiac Function vs Age. Man	72
35. Cardiac Index Determined under Basal Conditions: Man	72
36. Cardiac Index Determined by Direct Fick or Dye Method. Man.	73
37. Cardiac Index vs O ₂ Uptake, at Rest and during Exercise: Man	74
38. Cardiac Index vs O ₂ Uptake, during Exercise: Man (Graph)	75
39. O ₂ Uptake vs Arterio-venous O ₂ Difference, during Exercise: Man (Graph).	76
40. Cardiac Output vs O ₂ Consumption, during Exercise. Man.	77
41. Cardiac Output vs Temperature: Man, Dog	77
42. Effect of Various Conditions on Cardiac Output: Mammals	78
Part I: Man	78
Part II: Mammals Other Than Man	80

V. HEART RATE

43. Heart Rate: Vertebrates	81
Part I: Mammals	81
Part II: Birds	84
Part III: Reptiles and Amphibians.	85
Part IV: Fishes	86
44. Heart Rate: Invertebrates	87
Part I: Tunicates, Mollusks, Crustaceans, Annelids	87
Part II: Insects	91
45. Heart Rate of Embryo and Newborn: Mammals	94
46. Heart Rate, Basal and Non-basal at Various Ages: Man	95
47. Effect of Temperature on Heart Rate: Vertebrates	96
Part I: Per Degree Temperature Change.	96
Part II: Isolated Heart.	97
Part III: Intact Animals	99
48. Effect of Exercise on Heart Rate. Man.	99
49. Effect of Posture and Sleep on Heart Rate: Man.	100
50. Recovery of Heart Rate and Blood Pressure after Exercise. Man	100
Part I: Various Work Loads.	100
Part II: Step-Test Exercise.	101

VI. BLOOD PRESSURES

51. Man.	102
.	102
.	103
.	103

XI. EFFECT OF O₂ AND CO₂ CONCENTRATIONS

82. Hemodynamic Effects of Hypoxia (Low O ₂)	160
Part I: Heart Size: Mammals	160
Part II: Arterial, Venous, and Pulse Pressures: Man	161
Part III: Systemic, Pulmonary, and Right Ventricular Pressures: Man	161
Part IV: Arterial Blood Pressure during Progressive Adaptation to Altitude: Man	162
Part V: Pulse Rate Change, in Percent of Ground Level Value, during Progressive Exposure: Man	162
Part VI: Pulse Rate Change during Progressive Exposure: Man	162
Part VII: Circulation of Peripheral Blood: Man, Dog	163
Part VIII: Pulmonary Circulation: Man, Dog	164
Part IX: Intervals of the Electrocardiogram: Man	164
Part X: Amplitudes of the Electrocardiogram: Man	165
Part XI: The Electrocardiogram, Heart Rate, and Blood Pressure at Reduced Arterial O ₂ Saturation: Man	166
Part XII: Cardiac Work and Blood Pressure at Reduced Arterial O ₂ Saturation: Man	167
83. Hemodynamic Effects of Hyperoxia (High O ₂) Mammals	168
84. Hemodynamic Effects of Hypercapnia (High CO ₂)	170
Part I: Change in Pulse Rate, Cardiac Output, Blood Pressure, and Blood Flow: Mammals	170
Part II: Change in Pulse Rate and Blood Pressure: Man	172
Part III: Change in Pulse Rate, Blood Pressure, and the Electrocardiogram: Dog	173
Part IV: Mean Pressure, pH, CO ₂ Pressure, and CO ₂ Content of Arterial Blood: Man	176
Part V: Cardiac Output, Blood Pressure, and Heart Rate: Man	176
Part VI: Cardiac Output, Blood Pressure, and Heart Rate: Cat, Dog, Frog	178
Part VII: Coronary Blood Flow and Vascular Resistance: Cat, Dog	180
Part VIII: Limb Blood Flow and Vascular Resistance: Man, Dog	180
Part IX: Cerebral Blood Flow and Vascular Resistance: Man	182
Part X: Renal Blood Flow and Vascular Resistance: Man, Dog	182
Part XI: Pulmonary Blood Flow and Mean Arterial Pressure: Mammals	184
Part XII: Mesenteric Blood Flow and Vascular Resistance, and Hepatic Vascular Resistance: Dog	184

XII. EFFECT OF COMPRESSION, DECOMPRESSION, AND ACCELERATION

85. Pressure Equivalents	186
86. Effect of Compression on Pulse Rate, Systolic Blood Pressure, and Pulse Pressure: Man	186
87. Effect of Breathing O ₂ at 3-4 Atmospheres on Respiratory Rate, Pulse Rate, and Blood Pressure: Man	186
88. Effect of Breathing Air at 4 Atmospheres on Pulmonary Arterial Pressure: Rat	187
89. Effect of Recompression, with O ₂ and with Air, on Arterial Blood Pressure: Dog	187
90. Effect of Explosive Decompression on Heart Rate and Arterial Pressure: Mammals	188
91. Effect of Explosive Decompression on Arterial Pressure: Dog	189
Part I: Final Pressure Greater Than Theoretical Vapor Pressure of Body Fluids	189
Part II: Final Pressure Less Than Theoretical Vapor Pressure of Body Fluids	189
92. Effect of Acceleration on Circulatory and Respiratory Function Mammals	190

XIII. EFFECT OF RADIATION

93. Hemodynamic Effects of Ionizing Radiation: Vertebrates	196
94. Effect of Ionizing Radiation on Hematopoietic Tissue: Vertebrates	207
95. Effect of Ionizing Radiation on Peripheral Blood: Vertebrates	214
Part I: Man	214
Part II: Vertebrates Other Than Man	219
96. Effect of Ionizing Radiation on Blood Pressure and Pulse Rate: Vertebrates	228
97. Effect of Ionizing Radiation on Total Blood Volume and Plasma Volume: Vertebrates	229
98. Effect of Ionizing Radiation on Organ Blood Volume: Chicken	230

XIV. BLOOD COAGULANTS AND ANTICOAGULANTS

99. Blood Coagulation (Schematic Diagrams).	231
Part I:	231
Part II:	232
Part III:	232

131. Physiologic Alterations and Compensatory Mechanisms in Chronic Anemia: Man (Diagram)	343
132. General Symptoms of Anemia: Man and Other Animals	344
133. Cardiovascular and Pulmonary Symptoms of Anemia: Man and Other Animals	345
134. Electrocardiographic, Ballistocardiographic, and Roentgenographic Findings in Anemia: Man and Other Animals	347
135. Hemodynamic and Pulmonary Effects of Anemia: Man and Other Animals	349
APPENDICES	353-363
I. Formulas and Calculations Proposed for Clinical Use	355
II. Regression Equations.	357
III. Empirical Normal Standards for Ultra-low Frequency Ballistocardiograms	362
INDEX	365

Contributors and Reviewers

Abramson, David I.
Ahlquist, Raymond P.
Alexander, Robert S.
Altschule, Mark D.
Andres, Reubin
Andrew, Warren
Angelone, Luis
Angerer, Clifford A.
Anson, Barry J.
Asmussen, Erling
Assail, N. S.
Astrand, Per-Olof
Avlado, Domingo M., Jr.

Babers, Frank H.
Badeer, Henry
Bader, Mortimer E.
Baez, Silvio
Ballard, W. W.
Banga, I.
Barger, A. Clifford
Barnes, Glover
Barron, Donald H.
Beach, Elliot F.
Beard, Raunon L.
Behnke, A. R.
Bennett, C. W.
Berman, Herbert J.
Berne, Robert M.
Bing, Richard J.
Bishop, Jack G.
Blake, William D.
Blasius, W.
Blau, Monte
Blinks, Lawrence R.
Blumgart, Herrman L.
Bondurant, Stuart
Böving, Bent G.
Boyd, Edith
Boyden, Edward A.
Brady, Allan J.
Braunwald, Eugene
Brecher, Gerhard A.
Brody, Samuel
Brown, A. W. A.
Brown, Ellen
Buck, John B.
Burch, George E.
Burke, Jack D.
Burn, J. H.
Burton, Alan C.
Burwell, C. Sidney

Caccese, Anthony
Carter, C. L.
Carver, M. J.
Case, Robert B.
Chait, Alejandro
Chapman, Don W.
Chen, Allen
Chesley, Leon C.

Chinard, Francis P.
Clark, Elliot R.
Clarke, Norman E.
Comroe, Julius H., Jr.
Conklin, Ruth E.
Coon, William W.
Cooper, K. E.
Copenhagen, Wilfred M.
Courmand, Andre
Courtice, P. C.
Crafts, A. S.
Craneffeld, Paul F.
Cullumbine, H.
Currens, James H.

Darling, Louise
Davies, Dean F.
Dawe, A. R.
Dawes, Geoffrey S.
Dean, Robert B.
Dedichen, Jens
DeGaris, Charles F.
Delmas, A.
Dexter, Lewis
Dill, David B.
Draper, M. H.
Dresden, D.
Dudley, Horace C.
D'Zmura, Andrew P.

Ebert, Richard V.
Edelman, I. S.
Edholm, O. G.
Eisenberg, R.
Emalie, A. R. G.
Engel, Frank L.
Everett, Newton B.

Fazekas, Joseph F.
Feldman, Joseph D.
Ferencz, Charlotte
Finch, Clement A.
Fine, Jacob
Fishman, Alfred P.
Flexner, Louis B.
Foldes, Francis F.
Fowler, Noble O.
Frank, Howard A.
Freed, S. Charles
Fregly, Melvin J.
Freis, Edward D.
Friedman, J. J.
Fulton, George P.
Furth, Jacob

Ganong, William F.
Garb, Solomon
Gemmill, Chalmers L.
Gertler, Menard M.

Gibson, John G., II
Gillot, Claude
Gleckler, William J.
Glenn, William W. L.
Gordan, Gilbert S.
Gorlin, Richard
Grant, J. C. B.
Gregg, Donald E.
Griffith, John Quintin, Jr.
Grob, David
Grodzinski, Z.
Grollman, Arthur
Gross, Paul
Grossman, Jacob
Guyton, Arthur C.

Haddy, Francis J.
Hamilton, Howard L.
Hampton, John K., Jr.
Handley, Carroll A.
Hansard, Sam L.
Hardie, E. L.
Hartroft, W. Stanley
Hecht, Hans H.
Heirle, Robert W.
Heisler, Charles R.
Helm, Robert A.
Hlatt, Edwin P.
Hickam, John B.
Hoffman, Brian F.
Holtgrave, D. E.
Hull, Herbert M.
Hultgren, Herbert N.

Ingram, Marylou
Isaacs, Raphael

Jaques, Louis B.
Jensen, R.
Johnson, Richard P.
Jones, Jack Colvard

Kampmeier, Otto F.
Katz, Louis N.
Kelly, John
Kemmmerer, Arthur R.
Kemp, Norman E.
Kety, Seymour S.
Keynes, R. D.
Kibler, Hudson H.
King, Allen L.
Klein, J. Raymond
Krisely, Melvin H.
Knobil, Ernest
Knowles, Harvey C.,
Kollros, Jerry J.
Korman, Henry
Kosman, Charles E.
Krantz, John C., Jr.

Krayer, Otto
Krogman, Wilton Marion
Kruta, Vladislav
Kuo, Peter T.

Lackey, Robert W.
Lambertsen, Christian J.
Lamport, Harold
Landis, Eugene M.
Lange, Ramon L.
Larks, S.
Latimer, Homer B.
Lauson, Henry D.
Lawson, Hampden C.
Lawton, Richard W.
Leathem, James H.
Lee, Richard E.
Leonards, Jack R.
Lepeschkin, Eugene
Levy, Matthew N.
Lewis, Robert
Lichtman, S. S.
Lindsay, Hugh A.
Link, Roger P.
Lombard, Elma A.
Lowrance, E. W.
Lu, F. C.
Lucké, Baldwin
Ludwig, Daniel
Luisada, Aldo A.
Lynn, R. Beverley

Marbarger, John P.
Marks, Asher
Master, Arthur M.
Mayerson, H. S.
McGinty, Daniel A.
McGuire, Johnson
McKibbin, John M.
McKusick, Victor A.
McMichael, John
Melville, K. I.
Mendlowitz, Milton
Michaelson, Sol M.
Michels, Nicholas A.
Modell, Walter
Monie, I. W.
Monkhouse, Frank C.
Montgomery, Esther
Montgomery, P. O'B.
Moog, Florence
Morris, Lester E.
Morrison, Peter R.
Morse, Minerva
Mosher, Robert E.
Mukherjee, S. R.
Munson, Sam C.
Myers, Gordon B.

Nardone, Roland M.
Neill, Catherine A.
Newcomb, Thomas F.
Nutting, William L.

Orti, Eduardo
Osgood, Edwin E.

Page, Irvine H.
Pappenheimer, John R.
Pardee, Harold E. B.
Patterson, John L., Jr.
Penneys, Raymond
Perloff, Joseph K.
Porter, Reno R.
Pressman, David

Quénu, Louis
Quiring, Daniel P.

Ralli, Elaine P.
Redisch, Walter
Reinhardt, William O.
Remington, John W.
Renkin, Eugene M.
Reynolds, Monica
Richards, A. Glenn
Richards, Dickinson W.
Richardson, Alfred W.
Richardson, David W.
Robb, Jane Sands
Rodbard, Simon
Roeder, Kenneth D.
Rosenbaum, Francis F.
Ross, Lucille J.
Roth, Lawrence W.
Rowlands, S.
Rubin, Bernard
Rugh, Roberts
Ruskin, Arthur

Samet, Philip
Sanghvi, L. M.
Schaefer, Karl Ernst
Scheinberg, Peritz
Schmidt, Carl F.
Scott, John C.
Scott, Ralph C.
Seegers, Walter H.
Seliger, Václav
Selkurt, Ewald E.
Settle, W. R.
Severinghaus, John W.
Shepard, Roy J.
Shepler, Herbert G.
Sherrad, Theodore R.
Shipley, Reginald A.
Shock, Nathan W.
Siebens, Arthur A.
Smith, Durwood J.
Soloff, Louis A.
Sonnenschein, Ralph R.
Spencer, Joseph N.
Starr, Isaac
Stearner, S. Phyllis
Stone, Alan
Stone, Daniel J.
Sturkie, Paul D.
Sugioka, Kenneth
Swett, W. W.

Talbot, Timothy R., Jr.
Terry, Luther L.
Tocantins, L. M.

Trautwein, Wolfgang
Tritt, Edward B., Jr.

Vandam, Leroy D.
Van Harreveld, A.
Van Liere, Edward J.
Voelkel, Arno

Wade, D. L.
Wagner, Henry N., Jr.
Walcott, William W.
Walker, Sheppard M.
Walton, R. P.
Wartman, William B.
Webb, John L.
Weidmann, Silvio
Wesolowski, Sigmund A.
Wesson, Laurence G., Jr.
White, Fred N.
Whitehorn, William V.
Wiggers, Carl J.
Wigglesworth, V. B.
Wilde, Walter S.
Wilson, G. M.
Wintrobe, M. M.
Witham, A. Calhoun
Witschl, Emil
Wolpert, Arthur
Woodbury, Robert A.
Wright, H. Payling
Wright, Irving S.

Yoffey, Joseph M.
Yu, Paul N.

Zierler, Kenneth L.
Zuidema, George D.

Introduction

The Handbook of Circulation is presented as the companion volume to the Handbook of Respiration*. Its purpose is to make available in a single, comprehensive compilation useful data on circulation, organized for ready reference in the form of tables, graphs, diagrams, and drawings. Contents of this volume have been authenticated by 326 leading investigators in the field of physiology. The review process to which the tables have been subjected was designed to eliminate, insofar as possible, errors of transcription and material of questionable validity.

An explanatory headnote, designed to serve as an introduction to the subject matter or to justify the inclusion of controversial material, occasionally precedes a group of tables. Single tables, however, are usually prefaced by a short headnote containing such important information as abbreviations, definitions, units, methods, and conditions of measurement, and estimate of the range of variation. In order to interpret the data properly, it is essential that the related headnote be read with care.

Appended to the tables are the names of the contributors and a list of the literature citations. The reference abbreviations conform to the LIST OF ABBREVIATIONS FOR SERIAL PUBLICATIONS, Fourth Series, Volume X, Army Medical Library, Washington, D. C. (U. S. Government Printing Office, 1948), and the 1955 SUPPLEMENT thereto. Abbreviations for publications issued after 1955 were constructed from the "Dictionary of Abbreviated and Contracted Words" in the SYNOPSIS OF STYLE, Fourth Series, Volume II, Army Medical Library, Washington, D. C. (U. S. Government Printing Office, 1937).

Occasionally, differences in values for the same specifications, certain inconsistencies in nomenclature, and some overlapping of coverage may occur among tables. These result not from oversights or failure to choose between alternatives, but from the deliberate intention of the Handbook Staff to respect the judgment and preferences of the contributor. On the other hand, each table, with few exceptions, is internally consistent.


Values are generally presented as a mean and the lower and upper limit of the range of individual values about the mean. This range may be estimated in several ways, the method depending on the information available. Letter designations (a, b, c, d) identify types of ranges in descending order of accuracy.

(a) When the group of values is relatively large, a 95% range is derived by curve fitting. A recognized type of frequency curve is fitted to a group of measured values, and the extreme 2.5% of the area under the curve at each end is excluded (see illustration)

(b) When the group of values is too small for curve fitting, as is usually the case, a 95% range is estimated by a simple statistical calculation. Assuming a normal symmetrical distribution, the standard deviation is multiplied by a factor of 2, then subtracted from and added to the mean to give the lower and upper range limits

(c) A less dependable, but commonly applied, procedure (akes as range limits the lowest value and the highest value of the reported sample group of measurements. It underestimates the 95% range for small samples and overestimates for larger sample sizes, but may be used in preference to the preceding method where there is marked asymmetry in the position of the mean within the sample range

(d) Another estimate of the lower and upper limits of the range of variation is based on the judgment of an individual experienced in measuring the quantity in question. The trustworthiness of such limits should not be under-



* Handbook of Respiration. 1958. National Academy of Sciences-National Research Council. W. B. Saunders, Philadelphia

Abbreviations and Symbols

MEASUREMENT

yr = year
mo = month
wk = week
da = day
hr = hour
min = minute
sec = second
msec = millisecond

wt = weight
lb = pound
g = gram
kg = kilogram
mg = milligram
µg = microgram
mEq = milliequivalent
M = mole
mM = millimole
µM = micromole

ht = height
mi = mile
ft = foot
in. = inch
m = meter
cm = centimeter
mm = millimeter
µ = micron

g-m = gram-meter
g-cm = gram-centimeter
kg-m = kilogram-meter

sq in. = square inch
sq m = square meter
sq cm = square centimeter
sq mm = square millimeter
sq µ = square micron

L = liter
ml = milliliter
µL = microliter
cu cm = cubic centimeter
cu mm = cubic millimeter
cu µ = cubic micron

v = volt
kv = kilovolt
mv = millivolt
Å = Ångström unit

ppm = parts per million
rpm = revolutions per minute
av = average

vol % = volumes per cent
g % = grams per cent
mg % = milligrams per cent

°C = degree Centigrade
°F = degree Fahrenheit
> = greater than
< = less than

ROUTE OF ADMINISTRATION

IH = inhalation
IM = intramuscular
IP = intraperitoneal
IS = intraspinal
IV = intravenous

PO = oral
R = rectal
SC = subcutaneous
SL = sublingual
TOP = topical application

BIOLOGICAL SPECIFICATION

ISu = in situ
IVi = in vitro
IVv = in vivo
sp = species

RBC = red blood cell (erythrocyte)
WBC = white blood cell (leukocyte)
♂ = male
♀ = female

HANDBOOK of CIRCULATION

1. DEVELOPMENT OF THE CIRCULATORY SYSTEM. VERTEBRATES

Part I. FROG

Values are principally for *Rana pipiens* or *R. fusca*. At a given stage, age and size can be expected to vary widely with differences in geographic strains and culture conditions. Stages designated for the embryo are adapted from Shumway [3], and those for the larva from Taylor and Kollros [5].

Stage	Age	Size mm	Order of Development	Reference
(A)	(B)	(C)	(D)	(E)
			Embryo ¹	
1 15			Heart rudiment lies within heart bulge.	1
2 16	72 hr	3	Pericardial cavity indicated by split in mesoderm.	1
3			Temporary ventral mesocardium.	
4 17	84 hr	3.5	Dorsal mesocardium.	1
5			Heart a straight tube.	
6			Truncus arteriosus branched	
7			Blood island within 1st aortic arch.	
8			Dorsal aorta.	
9 18	96 hr	4	Rudiments of vitelline veins	2
10 19	118 hr	5	Heart S-shaped.	1, 3
11			Truncus arteriosus connects to 1st aortic arch, and thus to dorsal aorta.	
12			Blood islands in 2nd aortic arch, and numerous islands below posterior yolk mass.	
13			Blood in heart.	
14			Heart beat.	
15			Common cardinal veins and hepatic veins empty into sinus venosus.	
16 20	140 hr	6	Anterior lymph hearts.	2
17			Pulmonary vein rudiments.	
18 21	162 hr	7	Interatrial septum.	1, 2
19			Sino-atrial constriction.	
20			Atriobulbar constriction.	
21			Circulation through aortic arches I and II.	
22			Formation of anterior cardinal veins and arteries of the brain.	
23			Suprabranchial aortas fuse to single dorsal aorta medial to pronephros.	
24 22	192 hr	8	A few blood cells weakly ellipsoidal--most, spherical.	1-3
25			Trabeculae carneae in ventricle.	
26			Partitioning of heart complete.	
27			Circulation in tail fin.	
28 23	216 hr	8-9	Atrioventricular constriction.	1
29			Truncus arteriosus has thick endothelial cushions at level of its branching.	
30			Blood cells ellipsoidal.	
31 25	284 hr	10-11	Truncus arteriosus divided by median septum.	1, 2
32			RHC free of yolk granules.	
33			Spleen rudiment.	
			Larva ²	
34 I	3 da	13	Lymphocytes appear in spleen.	4
35 III	11 da	17	Thoracic ducts.	2
36 XXXI-XXV	76-88 da	63-253	Reorganization of aortic arches at metamorphosis.	2
1/1 At 18°C / 2/ At 20°C. / 3/ Size upon completion of metamorphosis highly variable, ranging from 16-30 mm.				2

Contributor: Kollros, Jerry J.

References [1] Kopsch, F. 1952. Die Entwicklung des braunen Grasfrosches *Rana fusca* Roessl. Georg Thieme Verlag, Stuttgart. [2] Ruth, R. 1951. The frog: its reproduction and development. Blakiston, Philadelphia. [3] Shumway, W. 1940. Anat. Rec. 78:139. [4] Fabrizio, M., and H. A. Charipper. 1941. J. Morph. 68:179. [5] Taylor, A. C., and J. J. Kollros. 1946. Anat. Rec. 94:7.

I. DEVELOPMENT OF THE CIRCULATORY SYSTEM: VERTEBRATES (Continued)

Part II: CHICK

Time of origin is expressed in days or hours of incubation, or number of pairs of somites.

Time of Origin	Structure	Site of Origin	Page In Reference
(A)	(B)	(C)	(D)
1 3-5 somites	Epimyocardium, ¹	Thickening of splanchnopleure of amniocardiac vesicles.	156, 158
2 3-5 somites and onward in later stages	Vascular system.	Initiated in blood islands at posterior margin of area opaca.	154
3 3-5 somites	Ventral aorta, ²	Hemangioblasts continuous anteriorly with primordia of heart.	157, 158
4			
5			
6			
7			
8 20 somites	cardinal vein).	and omphalomesenteric vein at level of somite 4.	
9 50-55 hr	Pulmonary vein.	Proliferation of hemangioblasts from dorsal wall of sinus venosus at level of lung buds.	255, 257
10 70 hr	Interatrial septum.	Vault of atrium between openings of sinus venosus and pulmonary vein.	432
11 35 somites	Subcardinal vein.	Series of venous islands on median surface of mesonephros.	453
12 3 da	Pulmonary artery.	Network of hemangioblasts extending caudally from ventral aorta.	253
13 3-4 da	Granulocytes.		257-260
14	Thrombocytes.		257-260
15	Endocardial cushions of atrioventricular canal.	Floor and roof of the atrioventricular canal.	433
16	Umbilical vein.	Capillary plexus in the somatopleure extending back from the duct of Cuvier.	255
17 4 da	Interventricular septum.	Along sulcus from bulboatrial angle towards apex of heart.	430
18	Subclavian artery.	Segmental artery of the 18th intersomitic space.	446
19 5 da	Neutrophils.		257-260
20	Mesenteric vein.	Dorsal mesentery, near level of dorsal pancreas.	450
21	Hepatic portal vein, definitive vessel.	Mesenteric vein in the dorsal mesentery.	450
22	Pelvic lymph sac.	Hemangioblasts from intersegmental veins of pelvic region.	462
23 5-6 da	Septum aorticopulmonale.	Truncus and bulbus arteriosus.	428, 429
24 6-7 da	Vertebral artery.	First 5 or 6 segmental arteries, and the plexus they form around the secondary sympathetic ganglia.	443
25	Thoracic duct of lymphatic system.	Intramesenchymal spaces along ventrolateral side of dorsal aorta.	463
26 7 da	Jugular lymph sac.	Fusion of lymphatic vessels near anterior cardinal vein.	463
27	Eosinophils.	Spleen (?).	257-260
28 14 da	Basophils.	Spleen (?).	257-260
29 17 da	Lymphocytes.	Spleen and other lymphatic areas.	257-260

/1/ Contractions begin at stage of 9 somites. /2/ Onset of function, with beginning of circulation at stage of 16 somites.

Contributor: Hamilton, Howard L.

Reference: Hamilton, H. L. 1952. Lillie's Development of the chick. Henry Holt, New York.

1. DEVELOPMENT OF THE CIRCULATORY SYSTEM VERTEBRATES (Continued)

Part III Pig

Stages adapted from Witschl [9]

Stage	Age da	Size mm	Order of Development	Reference
(A)	(B)	(C)	(D)	(E)
1 14	14-15	2.5	Pericardial cavity, endocardium, and epimyocardium.	1
2			Anlagen of blood vessels.	
3 15	15-16		Ventral aorta.	1
4			Fusion of paired cardiac tubes.	
5			Blood in yolk sac.	
6			Blood finally in heart and blood vessels.	
7 16	15-17		Truncus arteriosus, bulbus arteriosus, ventricle, atrium, sinus venosus.	1, 2
8			Septum primum.	1
9		4.15	1st pair of aortic arches.	3
10			Dorsal aorta.	
11		4.0	Omphalomesenteric arteries.	4
12			Allantoic arteries and veins.	
13			Subintestinal artery.	
14			Arteries of pronephros.	
15			Posterior cardinal vein.	
16			Duct of Cuvier.	
17			Umbilical veins.	
18 17	17	6.0-6.5	Longitudinal neural artery.	4
19			Lateral cardinal vein.	
20 18	16.5-18	3.8-4.5	2nd aortic arch.	1, 3
21			Endocardial cushions.	
22 19	16.5-17.5	3.6-4.57	3rd aortic arch.	1, 3
23			Interventricular septum.	
24 20	17.5	4.46-7.0	4th aortic arch.	1, 3, 4
25			Mesial cardinal vein.	
26			Venous valves.	
27 22		6.0	6th aortic arch.	3, 4
28			Primitive head vein.	
29 23	20	6.4	Foramen II in septum I	3
30 25	20	8.0-8.6	Septum II.	1, 3
31			External carotid.	
32 26	20-21	9.5-10.0	Bulbus dividing into aortic and pulmonary trunks.	1, 5, 6
33			Inferior vena cava.	
34			Spleen.	
35 27	21-22	11.0	Thymus.	7
36 28	22	12.0-14.0	Pulmonary, subclavian, basilar, and vertebral arteries.	1
37 30	28	20.7	Brachiocephalic and left common carotid arteries.	1
38 35	52	90.0	Tonsil.	8

Contributor Kemp, Norman E

References [1] Keibel, F. 1897. Normentafeln zur Entwicklungsgeschichte der Wirbelthiere. 1 Normentafel zur Entwicklungsgeschichte des Schweines (*Sus scrofa domestica*). Gustav Fischer, Jena. [2] Patten, B. M. 1948. Embryology of the pig. Blakiston, New York. [3] Heuser, C. H. 1923. Contr. Embryol. Carnegie Inst. 15: 121. [4] Sabin, F. R. 1917. Ibid. 6: 61. [5] Davis, D. M. 1910. Am. J. Anat. 10: 461. [6] Holyoke, E. A. 1936. Anat. Rec. 65: 333. [7] Badertscher, J. A. 1915. Am. J. Anat. 17: 317. [8] Levin, P. M. 1930. Anat. Rec. 45: 189. [9] Witschl, E. 1956. The development of vertebrates. W. B. Saunders, Philadelphia.

This table must be used with discretion because of certain discrepancies between embryological terminology and anatomical or functional characteristics, stages or events. The descriptions are consequently arbitrary, for many different authors. These defects have been obscured by conversion of descriptions into symbols in order to not indicate time or age relationships, but are merely space accommodations for the numerical headings. Letters primordium; B = loop formed, onset of coiling. C = completion of basic plan; D = differentiation; E = erythrocyte fetal stage; I = appearance of lymphoid cells; J = lymphocytes present in tissue; K = lengthwise subdivision, primordial; N = major channels formed; P = primitive nodules present; S = secondary follicles; T = twitch, markedly decreased; 2, 3, 4, 6 = aortic arch number.

Structure	Streeter's Horizons ¹													
	VIII	IX	X	XI	XII	XIII	XIV	XV	XVI	XVII	XVIII	XIX	XX	XXI
1 Blood forming foci														
2 Yolk sac	V	X					Y							
3 Chorion		V	(X)	Y		(X)			Z					
4 Liver									X					
5 Spleen														
6 Bone marrow														
7 Pericardial cavity	A		D										C	
Heart														
8 Atrium			M		K				L					
9 Septum primum									K					
10 Septum secundum									K					
11 Ventricle			T	M							L			
12 Bulboventricular loop			B											
13 Atrioventricular valve						A							G	
14 Semilunar valve											A			G
15 Conducting tissue								D						
Blood vessels														
16 Truncus arteriosus									K	L				
17 Aortic arches			N	2		3, 4	6							
18 Aorta			A							C	C			
19 Pulmonary artery								A						
20 Ductus arteriosus														
Lymphatic vessels														
21 Jugular region								A						
22 Retro-peritoneal region												A		
23 Cisterna chyli region												A		
24 Posterior iliac region												A		
Lymph nodes ⁷														
25 Jugular region														
26 Retro-peritoneal region														
27 Posterior iliac region														
Lymphatic tissues														
28 Palatine tonsil														A
29 Pharyngeal tonsil														
30 Lingual tonsil														
31 Peyer's patches														
32 Thymus						(A)			(A)			(F) 12	D 11 V	
33 Spleen								A						
Crown-rump length, cm	0.1	0.15	0.23	0.3	0.4	0.5	0.65	0.85	1.1	1.36	1.65	2.05	2.35	2.45

/1/ Use of Streeter's horizons as units for the first seven weeks is an attempt to minimize discrepancies arising vessels have almost reached vertex. /4/ Lymph vessels have probably reached intestinal mucosa. /5/ Thoracic lymph nodes are found in various primitive conditions well after birth. /8/ Secondary nodules present. /9/ Pit corpuscles. /13/ Malpighian bodies.

Contributors: (a) Böving, Bent G., (b) Clark, Elliot R., (c) Neill, Catherine A., (d) Kampmeier, Otto F.

References: [1] Streeter, G. L. 1951. Embryology. Reprint. Carnegie Institution, Washington. v. 2. [2] Bartelmez, [4] Kramer, T. C. 1942. Am. J. Anat. 71:342. [5] Gilmour, J. R. 1941. J. Path. Bact., Lond. 52:25. [6] Sabin, Philadelphia. v. 2, p. 709. [7] Arey, L. B. 1920. Developmental anatomy. Ed. 4. W. B. Saunders, Philadelphia. Duodecim. Ser. B, 19.1. [10] Davis, C. L. 1927. Contr. Embryol. Carnegie Inst. 19:245. [11] Odgers, P. N. B. W. G. Trumers. 1948. Am. J. Med. 5:324. [14] Walls, E. W. 1947. J. Anat., Lond. 81:93. [15] Goss, C. M. J. 1953. Proc. Nat. Acad. Sc. U. S. 39:333. [18] Congdon, E. D. 1922. Contr. Embryol. Carnegie Inst. 14:27. 1923. Traité technique d'histologie. Maloine et Fils, Paris. v. 2. [22] Grosser, O. 1912. In F. Keibel, and F. T. 1912. Ibid. v. 2, p. 390. [24] Tonkoff, W. 1900. Arch. mikr. Anat. 56:392. [25] Snook, T. 1934. Am. J. 27:193.

SYSTEM. VERTEBRATES (Concluded)

MAN


fact. The changes which constitute development are continuous but must be described in discontinuous units of time, have units which are unmeasurable, unreliable, estimated, omitted, qualified or used in different senses by emphasize essential features and reveal the quite regular sequence of events. Differences in width of columns do in parentheses and connected by dashes, are ranges. → indicates time spread. Abbreviations: A = anlage or formation, F = fusion or closure; G = growth completed, H = humerus marrow formation begins (defines onset of branching, or separation begun; L = lengthwise subdivision, branching, or separation completed; M = myocardium circulation begins; V = vascularization begins; X = hemopoiesis begins; Y = hemopoiesis maximal; Z = hemopoiesis


XXII	Weeks after Fertilization														After Birth	Reference
	8	9	10	12	14	16	18	20	24	28	32	36	38			
			(Z)-----				(Z)								1-9	
							(E)X-----Y-----Z-----		Y		Z	(E)				
H																
															1-4 1-4,10-17	
															L ²	

from uncertain age determination and individual rate of development. /2/ Six months after birth. /3/ Lymph duct complete. /6/ Lymph vessels have reached base of toes. /7/ The times given apply to the earliest nodes only. formation. /10/ Ectodermal and endodermal components fuse. /11/ Cortex-medulla differentiation. /12/ Hassall's



G. W. and H. M. Evans. 1926. Contr. Embryol. Carnegie Inst. 17.1. [3] Hensler, C. H. 1923. Ibid. 15 121. Florence R. 1912. In P. Keibel, and F. P. Mall, ed. Manual of human embryology. English ed. J. B. Lippincott. [8] Bloom, W. and G. W. Bartelmez. 1940. Am. J. Anat. 67:21. [9] Jägerskiöld, B. H. 1934. Acta Soc. med. 1937. J. Anat., Lond. 72.247 [12] Odgers, P. N. B. 1939. Ibid. 73 643 [13] Robb, J. S., C. T. Kaylor, and [19] Ehrlich, W. 1929. Am. J. Anat. 43.385 [20] Kiling, C. A. 1904. Arch. mikr. Anat. 63.575. [21] Jolly, J. F. P. Mall, ed. Manual of human embryology. English Ed. J. B. Lippincott, Philadelphia. v. 2, p. 446. [23] Lewis, Anat. 55.321. [26] Weller, G. L. 1933. Contr. Embryol. Carnegie Inst. 24.93. [27] Morris, E. H. 1938. Ibid.



2. HEART: MAN






2. HEART: MAN





4. PRINCIPAL VEINS AND PULMONARY ARTERIES- MAN
The liver and stomach are shown reflected upwards.



Reference: Dorland's illustrated medical dictionary. Ed. 23. 1957. W. B. Saunders, Philadelphia. Plate XLVIII

ULATORY SYSTEM. VERTEBRATES

17

ct, see references 11-15

Reptiles ²	Birds	Mammals			
		(Ox)	(Dog)	(Man)	
		(H)	(H)	(J)	
ly incorporated into right atrium, and a distinct only internally ⁴	Largely included in right atrium.	Incorporated into right atrium. Sino-atrial node, or "pacemaker," is a specialized tissue in atrial region (the latter was part of sinus in embryo).		Sinus venous completely incorporated in right atrium. ⁵ Sino-atrial node same as in ox and dog	1
proximately same as in teleosts	Embryonic right sino-atrial valve becomes valve of inferior vena cava.	Right valve of embryonic sinus becomes valve of inferior vena cava and of coronary sinus.		Vestige of embryonic sino-atrial junction found in adult right atrium.	2
ame as in frog, but interatrial septum contributes to valves dividing atrioventricular opening in two.	Complete interatrial septum. Extensive distribution of specialized conduction fibers	Complete interatrial septum. Histological characteristics of sino-atrial and atrioventricular nodal fibers unusually clear.	Complete interatrial septum. Sino-atrial node near opening of superior vena cava. atrioventricular node near opening of coronary sinus.	Complete interatrial septum except for small remnant of foramen ovale. Sino-atrial and atrioventricular nodes connected by typical cardiac fibers of right atrium	3
Turtle continuity of muscle as in teleosts, presence of "Purkinje-like" fibers controversial. ⁶ Alligator atrioventricular ring contains cartilage extending into base of right atrioventricular valve and similar valve of right aortic arch	Atrial and ventricular muscles separated by ring of connective tissue, except for atrioventricular bundles. Right atrioventricular valve large and muscular, left valve is bicuspid.	Connective tissue of atrioventricular ring (annulus fibrosus) contains bone. Atrioventricular bundle is unusually distinct. Right atrioventricular valve usually tricuspid, left valve usually bicuspid	Atrioventricular valves same as in ox. Ring of connective tissue separates atrial and ventricular muscle, except for atrioventricular bundle of His.	Atrioventricular ring of dense connective tissue. Right atrioventricular valve is tricuspid, left valve is bicuspid. Atrioventricular bundle of His present from atrioventricular node to ventricles ⁷	4
Turtle, snake, lizard: partially divided by incomplete septum Alligator: complete ventricular septum containing cartilage.	Complete ventricular septum.	Complete septum. Histological characteristics of Purkinje (conduction) fibers distinct.	Complete ventricular septum.	Complete interatrial septum which is membranous in its uppermost reaches.	5
Conus incorporated in ventricle and in arterial trunks	Proximal portion incorporated in right ventricle, distal into right aortic and pulmonary trunks	Proximal portion incorporated in right ventricle, distal into left aortic and pulmonary trunks.			6
Turtle: divided into right and left systemic and pulmonary trunks Alligator: right and left systemics connected by foramen of	Divided into aortic and pulmonary trunks, semilunar valves at embryonic juncture of conus and truncus.				7

(e) Grodzinski, Z. (f) DeGaris, Charles F. (g) Moog, Florence, (h) Ballard, W W

1956 Anat Rec 125 417 [3] Heinbecker, P. and G H Bishop. 1935 Am J Physiol. 114 212. [4] Hamilton J S 1953 Am J Physiol. 172 7 [6] Davies, F. and E. T. B. Francis. 1946. Biol. Rev. Cambridge Philos. Soc. anatomy. University of Chicago Press, Chicago [9] Patten, B M. 1946 Human embryology. Blakiston, Toronto Structure and development of vertebrates Macmillan, London. [12] Kingsley, J. S. 1917. Comparative anatomy Blakiston, New York

For a comprehensive review of the

Component	Cyclostomes ¹	Elasmobranchs and Teleosts	Dipnoans	Amphibians
(A)	(B)	(C)	(D)	(E)
1 Sinus venosus	Thin-walled, elongated sac, or tube, into which systemic veins open.	Smooth, thin-walled chamber into which systemic veins open. Cardiac muscle has high intrinsic contraction rate and acts as the "pacemaker." ³		Salamander: thin-walled triangular shaped chamber, shifted toward right side. Frog: separate chamber, and relatively smaller.
2 Sino-atrial junction	Sino-atrial opening guarded by pair of valves.	Sino-atrial opening guarded by pair of valves. Cardiac muscle is continuous from sinus to atrium		Approximately same as in teleosts.
3 Atrium	Single, muscular sac. Atrium lateral to ventricle.	Thin, reticulate-walled chamber. No division into right and left sides; no pulmonary veins.	Partial septation. Pulmonary vein enters to left, and sinus venosus to right, of septum.	Salamander: incomplete septum partially dividing chamber bilaterally, with sinus venosus opening into right atrium and single pulmonary vein into left atrium. Frog: complete interatrial septum.
4 Atrio-ventricular junction	Atrioventricular channel connects both vesicles; two valves present.	Atrioventricular valve composed of two cusps. Cardiac muscle of atrium continuous with that of ventricle around entire circumference of the atrioventricular junction.	Fibrocartilaginous plug regulates blood flow.	Continuity of atrial and ventricular muscles, as in teleosts. Salamander: atrioventricular valve of two or four muscular thickenings. Frog: four valve cusps.
5 Ventricle	Thick-walled, muscular sac, with smooth internal surface.	Thick-walled chamber, network of muscular trabeculae. No division into right and left sides.	Septum present.	No ventricular septum.
6 Conus arteriosus	Absent, unless two valves may be regarded as remnants.	Variable in size; relatively large in cartilaginous fish, small in teleosts. Has semilunar valves.	Divided into dorsal and ventral channels.	Semilunar valves, also a "spiral valve" coursing lengthwise.
7 Truncus arteriosus ⁸	Part bordering on ventricle enlarged to form bulbus arteriosus	Enlarged in teleosts which have reduced conus.	Divided into three paired channels.	Salamander: divided internally into right and left channels. Frog: divided both internally and externally into right and left channels.

/1/ Data from Favaro [1]. /2/ Data for alligator from White [2]. /3/ Although contraction is myogenic in origin, stimulation. /4/ Origin of beat in turtle heart is dependent on intrinsic ganglia [3]. /5/ Some authorities consider fibers are present in the turtle and in lower forms is supported by Robb [5], while the view that a specialized /7/ There are usually no muscular, atrioventricular connections other than the bundle of His in the adult [7]. arteriosus, or aortic sac [8-10].

Contributors: (a) Copenhagen, Wilfred M., (b) Andrew, Warren, (c) Monie, I. W., (d) White, Fred N.,

References [1] Favaro, G. 1901. Cyclostomi. In Bronn's Klassen und Ordnungen. Bd 6. [2] White, F. N. W. J., J. D. Boyd, and H. W. Moosman. 1952. Human embryology. Williams and Wilkins, Baltimore. [5] Robb, Z. I. 1913. [7] Kistin, A. D. 1949. Am. Heart J. 37:849. [8] Hyman, Libbie H. 1947. Comparative vertebrate [10] Arey, L. B. 1954. Developmental anatomy. W. B. Saunders, Philadelphia. [11] Goodrich, E. S. 1930. of vertebrates. Blakiston, Philadelphia [13] Nelson, O. 1953. Comparative embryology of the vertebrates.

5	Anterior cardinal veins	<p>Elasmobranchii: anterior cardinal veins unite to form common trunk and open into sinus venosus. Inferior jugular vein (ventral) present.</p> <p>Myxine: left anterior cardinal vein (internal jugular) and inferior jugular vein (ventral) join sinus venosus. Right anterior cardinal vein opens into cor portae of liver.</p>	<p>Urodele: two anterior cardinal veins (internal jugular), two inferior jugular veins.</p> <p>Anura: anterior cardinal veins (internal jugular) and inferior jugular veins fuse into one short trunk on each side and enter sinus venosus by intermedial vein (cranial).</p> <p>Dipnoi: two cardinal veins and two jugular veins.</p>	<p>Sphenodon: two anterior cardinal veins (internal jugular veins); inferior jugular veins (external) reduced and replaced by tracheal vein. These veins join subclavian vein to form anterior vena cava.</p> <p>Lacertilia: similar to conditions in Sphenodon.</p>	<p>Two anterior venae cavae, each consisting of anterior cardinal vein, vertebral vein and subclavian vein. Right cardinal vein more prominent than left because blood passes from left into right by anastomosis.</p> <p>Carnivora: Primates: Section of left vein, located between heart and anastomosis, disappears.</p>	<p>Two anterior venae cavae (unominate as in Rodentia and Insectivora. Each anterior vena cava consists of anterior cardinal, inferior jugular, and subclavian.</p> <p>Ungulata: Right vena cava more prominent than left because blood passes from left into right by anastomosis.</p>
Main Segmental Vessels						
6	Dorsal segmental vessels	<p>In all embryos, three kinds of segmental vessels are present: dorsal, lateral, and ventral segmental arteries and veins. They persist at least partially in adults and give rise to some longitudinal as well as other important stems.</p> <p>Arteries and veins alternate in successive segments. Their dorsal tips join to form the longitudinal dorsal trunk, and may persist in adults. Tributaries of dorsal segmental vessels are ventral medullary arteries and veins.</p>				
7	Dorsal rami	<p>Deep pinnae arteries and veins are segmental vessels supplying dorsal fins.</p>				
8	Ventral rami	<p>Segmental vessels run from the main longitudinal vessels (aorta, cardinal veins) in horizontal myoseptum and reach skin. Tributaries are muscular arteries and veins. Some become intercostal arteries and veins, surpassing main vessels in size. In region of paired fins and limbs, segmental vessels become subclavian and iliac arteries and veins.</p>				
9	Lateral cutaneous veins	<p>Elasmobranchii: Dipnoi, Urodele, Anura: lateral cutaneous vein in lateral line groove, below skin, from tail to region of forelimb. Originates from end tips of lateral segmental veins.</p> <p>Anura: great and small cutaneous vein.</p>		<p>Lacertilia: lateral cutaneous vein probably present in all species.</p>		
10	Abdominal vein	<p>Elasmobranchii: two lateral abdominal veins, one on each side of body wall.</p> <p>Extends from region of cloaca to shoulder girdle, where it merges into sinus venosus or into hepatic portal system.</p>			<p>Only anterior root of abdominal vein persists and is represented by umbilical vein in the embryo.</p>	
11	Epigastric arteries & veins	<p>Myxine: netlike tract.</p>				

11/ Longitudinal vessels originating from ventral rami of dorsal segmental vessels.

6. COMPARATIVE ANATOMY OF THE CIRCULATORY SYSTEM: VERTEBRATES (Continued)

Part II: BLOOD VESSELS

Vessel (A)	Cyclostomes (B)	Fishes (C)	Amphibians (D)	Reptiles (E)	Birds (F)	Mammals (G)
1 Aorta and caudal artery	Continuous vessels located ventral to the axial skeleton, extending from the region of the heart to the tip of the tail. Caudal artery enclosed in the hemal channel; aorta similarly enclosed in <u>Acipenser</u> (sturgeons) only.		Main Longitudinal Vessels			
2 Aortic arches	Six, seven, or more epi-brachial arteries.	Mainly 4-5 epi-brachial arteries. <u>Dipnoi</u> pulmonary arteries present.	One to four pair of aortic arches.	One pair of aortic arches.	Left aortic arch only.	Left aortic arch only.
3 Carotid arteries	Petromyzon internal carotid artery (dorsal), external carotid artery (ventral). Myxine internal and external carotid arteries.	Internal carotid arteries.	Internal and external carotid arteries. <u>Anura</u> carotid duct absent.	Internal and external carotid arteries originate from common carotids; many snakes possess only one common carotid. <u>Crocodylia</u> : carotid duct absent.	Internal and external carotids originate from common carotid arteries, which may be symmetrical, asymmetrical, fused, or one may be obliterated.	Various connections of internal and external carotids with aortic arch. Inominate artery, common carotid artery, carotid trunk.
4 Posterior cardinal veins	In all embryos, two posterior cardinal veins are present, being continuous with the caudal vein. Renal portal system absent as in embryo. Myxine, same as right posterior cardinal vein much thinner than left.	Elasmobranchii: caudal vein empties into renal portal veins, hepatic veins empty into posterior cardinal veins, which are sinus-like distensions. Teleostei: posterior cardinals often asymmetrical but receive renal and hepatic veins, renal portal blood derived from caudal vein and/or segmental veins, caudal vein often leads directly to posterior cardinal. Dipnoi: caudal vein, renal portal vein, asymmetric posterior or cardinal vein, and a posterior vena cava	Posterior cardinals rudimentary. Renal portal blood derived from iliac veins, and caudal vein in <u>Urodela</u> . Well-developed posterior vena cava drains kidneys.	Lacerellia: caudal vein empties into two renal portal veins. Two hepatic veins join posterior vena cava. <u>Crocodylia</u> : posterior vena cava emerges as single vessel from kidneys.	Rudimentary caudal and renal portal veins. Renal veins join femoral and internal iliac veins in kidney, which in turn join to form posterior vena cava. Posterior vena cava receives hepatic vein. No posterior cardinal veins persist.	No renal portal system. Posterior cardinals obliterated and replaced by supra-cardinal derivatives, the azygos and hemiazygos veins. Posterior vena cava is main trunk vein.

6. COMPARATIVE ANATOMY OF THE CIRCULATORY SYSTEM. VERTEBRATES (Continued)

Part II: BLOOD VESSELS (Concluded)

Vessel (A)	Cyclostomes (B)	Fishes (C)	Amphibians (D)	Reptiles (E)	Birds (F)	Mammals (G)
			Main Segmental Vessels (concluded)			
12 Lateral seg- mental vessels	Paired lateral vessels supply pronephros, mesonephros, and, when present, metanephros. Their number is much reduced in higher forms when forming renal and gonadal arteries.					
13 Ventral segmental vessels						
14 Arteries	<p>In all embryos, ventral segmental vessels originate from the main longitudinal vessels (aorta, caudal artery, caudal vein); in trunk, only arteries (mesenteric arteries) originate from main longitudinal vessels, and in tail both arteries and veins (ventral segmental arteries and veins). The paired arteries frequently unite to form a single median vessel.</p> <p><u>Myxine</u>: inter- teric arteries Dipnoi: celiac artery, two or three mesen- teric arteries <u>Teleostei</u>: mainly one celio-mesenteric artery.</p> <p>intestine as seg- mental vessels. <u>Petromyzon</u>: only one artery per- sists as celi- mesenteric artery.</p>	<p><u>Urodela</u>: celiac artery, and about 13 mesenteric arteries almost seg- mental in arrangement, distributed to intestine (Siren). Segmental arteries fuse into com- plex vessels (<u>Meno-</u> <u>branchus</u>, <u>Crypto-</u> <u>branchus</u>, <u>Salamandra</u>). <u>Anura</u>: only celio-mesen- teric artery.</p>	<p><u>Hatteria</u>: gastric artery, celiac artery, common mesenteric artery (ante- rior), posterior mesen- teric artery. <u>Lacerta</u>: gastric artery, celio-mesenteric artery, posterior mesenteric artery. <u>Crocodylia</u>: gastroso- phageal artery, celi- mesenteric artery. Many arteries (mesen- teric) reach intestine from the aorta (Boal).</p>	<p><u>Celiac artery</u>, ante- rior mesenteric artery (cranial or superior), posterior mesenteric artery (caudal or inferior).</p>	<p><u>Celiac artery</u>, ante- rior mesenteric artery (cranial or superior), posterior mesenteric artery (caudal or inferior).</p>	
15 Veins	<p>Subintestinal vein, which gets some blood directly from caudal vein, develops with invasion of intestine by ventral segmental arteries. Subintestinal vein is prominent vessel in embryonic fish and amphibians, less significant in different reptiles and birds; remnants found in left omphalomesenteric veins form the hepatic portal vein.</p> <p>Vitelline vein (omphalomesenteric) may participate in liver circulation.</p> <p>Hepatic vein opens directly into heart.</p> <p><u>Elasmobranchii</u>: subintestinal vein well developed.</p> <p><u>Myxine</u>: reduced.</p>	<p>Hepatic vein opens into posterior vena cava, subintestinal vein is incorporated into trunk of hepatic portal vein.</p>	<p><u>Vessels of the Forelimb</u></p> <p><u>Urodela</u>: subclavian artery arises from median aorta. Brachial artery, interosseal artery (main vessel), radial artery (radiomarginal), ulnar artery (ulnomarginal). Dorsal arterial arch of hand, metacarpal arteries, digital arteries.</p> <p><u>Lacertilia</u>: both subclavian arteries arise from right aortic arch. Axillary artery, interosseal artery (main vessel), ulnar artery, radial artery and median artery. Metacarpal arteries, digital arteries.</p> <p><u>Subclavian artery</u> with carotid artery originates from right aortic arch (innominate artery), left subclavian, direct from arch or trunks communia. Axillary artery, brachial artery, ulnar artery (main vessel), median artery (main).</p>			
16 Arteries	<p>Forelimbs absent</p> <p><u>Elasmobranchii</u>: subclavian artery arises from median aorta. Lateral and medial pterygial arteries, adradial arteries. <u>Teleostei</u>: subclavian artery originates from median aorta. Basal arteries, interradial arteries.</p>	<p><u>Elasmobranchii</u>: subclavian artery arises from median aorta. Lateral and medial pterygial arteries, adradial arteries. <u>Teleostei</u>: subclavian artery originates from median aorta. Basal arteries, interradial arteries.</p>	<p><u>Vessels of the Forelimb</u></p> <p><u>Urodela</u>: subclavian artery arises from median aorta. Brachial artery, interosseal artery (main vessel), radial artery (radiomarginal), ulnar artery (ulnomarginal). Dorsal arterial arch of hand, metacarpal arteries, digital arteries.</p> <p><u>Lacertilia</u>: both subclavian arteries arise from right aortic arch. Axillary artery, interosseal artery (main vessel), ulnar artery, radial artery and median artery. Metacarpal arteries, digital arteries.</p> <p><u>Subclavian artery</u> with carotid artery originates from right aortic arch (innominate artery), left subclavian, direct from arch or trunks communia. Axillary artery, brachial artery, ulnar artery (main vessel), median artery (main).</p>			


5	Cisterna chyli in lumbar area	sinus in liver region.		<u>Apoda</u> , <u>Anura</u> : absent. <u>Urodeli</u> : present; extends to thoracic region. <u>Urodela</u> : with anterior cardinal veins. <u>Anura</u> : with anterior lymph hearts.	<u>Lacertilia</u> , <u>Chelonis</u> : present. <u>Crocodylia</u> , <u>Ophidia</u> : absent.	With anterior cardinal veins.	With anterior cardinal veins.	Always present; great variation in shape and size.
6	Connections with veins	<u>Platymyxon</u> numerous connections with both posterior cardinal veins. <u>Myxine</u> : with anterior cardinal vein.	<u>Elasmobranchii</u> : with posterior cardinal vein at point where subclavian artery crosses. <u>Telostei</u> : with anterior cardinal vein.					In most mammals with left anterior cardinal vein, in some with right, in few with both.
7	Jugular lymphatic trunk	<u>Petromyzon</u> : seven peribranchial sinuses, each connected with anterior cardinal vein.	<u>Elasmobranchii</u> : two trunks connected with corresponding anterior cardinal veins. <u>Telostei</u> : two sinuses, each connected with corresponding subvertebral trunks.	<u>Urodela</u> : two trunks connected with corresponding subvertebral trunks. <u>Anura</u> : two short trunks connecting head sinuses with anterior lymph hearts.	<u>Lacertilia</u> : two trunks connected with subvertebral trunks by way of jugular sinus. <u>Chelonis</u> : two trunks enter jugular claters which connects both subvertebral trunks. <u>Crocodylia</u> : two trunks enter corresponding anterior cardinal veins.	Each of two trunks join corresponding subvertebral trunk.	Each of two trunks join corresponding subvertebral trunk.	Irregular lymph vessels join anterior cardinal vein, separately, or by way of subvertebral trunk.
8	Lateral longitudinal lymphatic trunks		<u>Elasmobranchii</u> : absent. <u>Telostei</u> , <u>Ganoidi</u> : below skin in lateral line groove from base of tail fin to head; opens into some of head sinuses or directly into duct of Cuvier.	<u>Apoda</u> , <u>Urodela</u> : in lateral line groove from base of tail to head; opens into lymph hearts. <u>Anura</u> : in tadpoles: from base of tail to anterior lymph heart, disappears during metamorphosis.	<u>Lacertilia</u> : from tip of tail to forelimbs; caudal part enters lymph heart, thoracic part enters axillary sinus. <u>Crocodylia</u> : only caudal part present. <u>Chelonis</u> : thoracic part well developed. <u>Ophidia</u> : lateral trunk reaches maxillary sinus.			In adult, present only in tail; in embryos, transitory thoracic part observed.
9	Other longitudinal lymphatic trunks		<u>Elasmobranchii</u> : absent. <u>Telostei</u> , <u>Ganoidi</u> : dorsal trunk unpaired, located in dorsal mid-line of tail and body, from base of tail fin to head; ventral trunk unpaired, located in ventral mid-line of tail and in middle of abdominal wall; spinal trunk dorsal to spinal cord.	<u>Urodela</u> : dorsal trunk unpaired, located in dorsal mid-line of tail and body; ventral trunk unpaired, located in ventral mid-line of tail; abdominal trunk paired in wall of abdomen. <u>Anura</u> : dorsal and ventral trunks only in fin of tadpoles.				

11/ Cole [7] and Favaro [8] regard these sacs as venous hearts.


Part III. LYMPHATICS

Component (A)	Cyclostomes (B)	Fishes (C)	Amphibians (D)	Reptiles (E)	Birds (F)	Mammals (G)
1 Lymph hearts	Petromyzon: absent. Myxine: one pair of pulsating sacs located in tail. ¹	Elasmobranchii: absent. Teleostei: two elongated vesicles, joined by a channel and located at base of tail. Ganoidi: similar to Teleostei, but not pulsating vesicles.	Apoda: approximately 100 spherical vesicles in trunk and tail, beneath skin in lateral line groove. Urodela: 10-20 rounded vesicles on each side of trunk, located as in Apoda. Anura: one pair of anterior, and 1-4 pairs of posterior, hearts.	Only posterior hearts. Lacertilia: two ovoid vesicles attached to both ends of transverse processes of first caudal vertebra. Ophidia: two elongated vesicles, each surrounded by bifurcated transverse processes of 4-5 caudal vertebrae. Dinosauria: volume of one heart of <i>Apatosaurus</i> , approximately 8000 ml; of <i>Diplodocus</i> , 20,000 ml. Chelonio, Crocodilia: spherical in shape.	In all embryos only one pair of posterior hearts, located in region between pelvis and femur/hearts persist in some adult birds (Struthio, Casuarina, Anser, Colymbus, Alca).	Absent even in embryos.
2 Other lymph sacs, and sinuses	Petromyzon: supra-labial sinus, orbital sinus, ocular ring sinus, deep labial sinus. Myxine: three subcutaneous sacs underlie entire skin.	Elasmobranchii: absent. Teleostei: pectoral pinnae sinus, orbital sinus, cephalic sinus, occipital sinus, lateral sinus.	Urodela: orbital sinus, lymphatic cordis, axillary sinus in larvae only. Anura: in tadpoles: mandibular sinus, circumoral sinus, pericardial sinus, temporal sinus; in adults: several subcutaneous sacs.	Lacertilia: retrocardial sinus, axillary sinus, jugular sinus, tracheal sinus, thyroïdal sinus. Ophidia: mandibular sinus Chelonio: jugular cistern. Crocodilia: absent.	Jugular and iliac lymph sacs in embryo only.	
3 Lymph nodes				Chelonio: small nodes in lower eyelid.	Microscopically discernible nodes in walls of lymph vessels. Anseriformes: two cervico-thoracic and two lumbar lymph glands, macroscopically visible.	Many lymph nodes: approximately 60 in dog, 100 in cattle, 465 in man, 6000 in horse.
4 Subvertebral lymphatic trunks (thoracic duct)	Petromyzon: unpaired sinus-like trunk beneath aorta and cardinal veins Myxine: two wide trunks located on both sides of aorta; fused into wide	Elasmobranchii, Ganoidi, Teleostei: two slender trunks located on both sides of aorta and caudal artery.	Apoda: one sinus-like extended trunk accompanies aorta. Urodela: one or two trunks. Anura: two trunks.	Lacertilia, Ophidia: sinus surrounds aorta. Chelonio: two trunks in tail, single in body cavity, and bifurcated anteriorly. Crocodilia: two slender trunks.	Two trunks located on both sides of aorta.	One or two trunks associated with aorta and caudal artery.

8. DIMENSIONAL GROWTH OF THE HEART DURING FIRST YEAR: MAN



+ - Heart length, ♂
 x - Heart length, ♀
 o - Heart breadth, ♂
 o - Heart breadth, ♀



Reference Scammon, R. E. 1927. Radiology 9 94 (Based on data of A. A. Falk. 1901. The growth of the heart in children according to age. Dissertation (in Russian). Also on data of K. Bamberg, and H. Putzig. 1919. Zchr. Kinderh. 20 195)

9. HEART, LINEAR DIMENSIONS: MAN

Values are in centimeters.

Age	Male			Female		
	Width (A)	Length (B)	Thickness (C)	Width (D)	Length (E)	Thickness (F)
1 Newborn	4.0	3.1	1.9	3.9	2.9	1.7
2 5-6 mo	4.9	3.9	2.6	4.7	3.8	2.6
3 1-2 yr	6.2	5.0	3.0	5.9	4.9	2.9
4 3-4 yr	6.6	5.5	3.3	6.2	5.4	3.1
5 5-6 yr	7.3	6.5	3.5	6.4	5.9	3.4
6 7-8 yr	7.6	7.0	3.5	7.3	6.2	3.2
7 9-10 yr	8.2	7.3	3.4	7.2	6.2	3.6
8 13-14 yr	8.2	7.8	3.7	8.3	7.8	3.8
9 Adult	10.7	9.7		9.9	9.3	

Reference Krogman, W. M. 1941. Tabulae biol., Berl. 20 664.

10. HEART, SURFACE DIMENSIONS: MAN

Values are in centimeters.

Age yr	No of Subjects	Filled Heart	
		Circumference of Base (A)	Distance from Base to Apex (B)
1 1-2.5	51	13.0	5.5
2 3-3.5	29	13.0	6.0
3 4-4.5	21	13.0	6.2
4 5-5.5	14	13.0	6.4
5 6-6.5	6	15.5	7.0
6 7-7.5	11	17.0	8.0
7 8-8.5	15	17.0	8.0
8 9-9.5	4	17.0	8.0
9 10-10.5	10	17.5	8.0
10 11-11.5	14	18.5	8.5
11 12-12.5	9	17.5	8.5
12 13-13.5	6	19.0	9.0
13 14-14.5	3	19.0	9.0

Reference. White House Conference on Child Health and Protection. 1933. Growth and development of the child. Century. New York Pt II

6. COMPARATIVE ANATOMY OF THE CIRCULATORY SYSTEM. VERTEBRATES (Concluded)


Part III. LYMPHATICS (Concluded)

Contributor: Grodzifski, Z.

References: [1] Allen, W. F. 1913-14. J. Micro. Sc., Lond. 59:part 2. [2] Allen, W. F. 1907. Proc. Washington Acad. Sc. v. 9. [3] Allen, W. F. 1908. Am. J. Anat. v. 8. [4] Baum, H. 1912. Das Lymphgefäßsystem des Rhindes. Berlin. [5] Baum, H. 1928. Das Lymphgefäßsystem des Pferdes. Berlin. [6] Clark, E. R., and E. L. Clark. 1920. Contr. Embryol. Carnegie Inst. 9:447. [7] Cole, F. J. Tr. R. Soc. Edinburgh. v. 54. [8] Favaro, G. 1905. Atti Ist. venet. sc. 65:195. [9] Glaser, G. 1933. Zschr. Anat. Entw. v. 100. [10] Grodzifski, Z. 1932. Bull. internat. Acad. sc. Cracovie. [11] Grodzifski, Z. 1929. Ibid. [12] Hoyer, H. 1934. Mém. internat. Acad. polon. sc., Cl. med. 1:205. [13] Hoyer, H. 1928. Bull. internat. Acad. sc. Cracovie p. 79. [14] Hoyer, H. 1905. Anat. Anz. 27:50. [15] Hoyer, H. 1908. Bull. internat. Acad. sc. Cracovie. p. 451. [16] Kampmeier, O. 1925-26. J. Morph. 41:95. [17] Kihara, R., and E. Naito. 1933. Fol. anat. jap 11:405. [18] Marcus, H. 1908. Morph. Jahrb. 38:590. [19] Mozejko, B. 1911. Anat. Anz. 40:469. [20] Müller, J. 1941. Abh. K. Akad. Wiss., Berlin. [21] Müller, J. 1939. Ibid. [22] Panizza, B. 1833. Sopra il sistema linfatico dei rettili. Pavia. [23] Retzius, G. 1890. Biol. Untersuch., Stockh. 1:20. [24] Sabín, F. R. 1909. Am. J. Anat. 9:43. [25] Tretjakoff, D. 1927. Morph. Jahrb. 58:209. [26] Tretjakoff, D. 1930. Ibid. v. 64. [27] Weidenreich, F. 1933. Lymphgefäßsystem. In L. Bolk, ed. Handbuch der vergleichenden Anatomie der Wirbeltiere. Bd 6.

7. CIRCUMFERENTIAL GROWTH OF THE HEART, AORTA, AND SUPERIOR VENA CAVA: MAN

Growth computed as per cent of natal value.



Reference: Scammon, R. E. 1927. Radiology 9:97.

14. DIMENSIONS OF MAIN ARTERIAL TRUNKS MAN

For information on the shapes of arterial segments, see reference 2. Values in parentheses are ranges, estimate "c" (cf. Introduction).

PART I. UPPER ABDOMEN

Measurements made in anatomic dissection rooms on male cadavers injected with preserving fluid and a colored starch. Due to shrinkage upon fixation, width measurements of vessels may be smaller than those for living bodies. Width of left gastric, splenic, and hepatic arteries measured at point of origin from the celiac artery.

	Artery (A)	No of Subjects (B)	Age yr (C)	Width mm (D)	Length mm (E)
1	Celiac trunk	47	61.4(34-81)	10.1(6-13)	24.8(11-35) ²
2	Left gastric	48	61.4(34-81)	5.5(4-9)	
3	Right gastric	48	61.4(34-81)	1.7(1-4)	
4	Splenic	48	61.4(34-81)	8.5(6-14)	
5	Common hepatic trunk	48	61.4(34-81)	8.5(6-11)	
6	Hepatic to gastroduodenal	18	62.2(43-79)		31.1(20-50)
7	Hepatic to its divisions	7	54.4(34-74)		48.1(22-80)
Terminal hepatic branches					
8	Right hepatic	43	60.8(34-81)	6.4(4-9)	
9	Left hepatic	44	61.2(34-81)	5.0(3-7)	
10	Middle hepatic	48	61.4(34-81)	3.6(2-6)	
Aberrant hepatics					
11	Accessory right hepatic from superior mesenteric	7	59.8(43-74)	5.1(2-8)	80.0(70-90)
12	Replaced right hepatic from superior mesenteric	4	66.2(57-73)	7.0(6-8)	80.0(70-100)
13	Accessory left hepatic from left gastric	5	55.7(39-69)	4.4(4-5)	
14	Replaced left hepatic from left gastric	4	50.5(43-58)	4.8(4-5)	
15	Single cystic	16	59.2(34-81)	2.7(2-4)	17.7(2-30)
Double cystic					
16	Superficial	12	66.2(58-74)	2.3(2-3)	22.2(10-33)
17	Deep	12	66.2(58-74)	2.4(2-4)	23.8(10-50)
18	Superior mesenteric	48	61.4(34-81)	11.2(8-14)	
19	Interval from celiac to superior mesenteric	48	61.4(34-81)		9.2(2-30)
20	Gastroduodenal	48	61.4(34-81)	5.3(4-7)	

1/1 External diameter measurement. 2/2 Length of celiac artery arbitrarily determined as the distance from its point of origin in the aorta to the point at which the celiac trunk branches into the hepatic and splenic arteries.

Contributor: Michels, Nicholas A.

References: [1] Michels, N. A. 1955. Blood supply and anatomy of the upper abdominal organs. J. B. Lippincott, Philadelphia. [2] Jeffords, J. V., and M. H. Knisely. 1956. Angiology, Balt. 7:105.

Part II. LOWER ABDOMEN

Measurements made on cadavers injected with latex.

	Artery (A)	No of Subjects (B)	Age yr (C)	Width mm (D)	Length mm (E)
1	Aorta	3	110.5(1-15)	7(6.5-8.0)	
2		29	(19-72)	14.4(9-18)	
3	Inferior mesenteric	3	110.5(1-15)	2.6(2.0-3.5)	58.6(50-60)
4		32	(19-72)	3.7(2.5-5.0)	120(77-170)
5	Interval from superior mesenteric to inferior mesenteric	3	110.5(1-15)		28.6(16-45)
6		24	(19-72)		53(35-70)
7	Interval from inferior mesenteric to aorta	3	110.5(1-15)		33(12-13)
8		26	(19-72)		31(15-45)
		32	110.5(1-15)		16.3(15-18)
		32	(19-72)		30.6(15-48)
		8	Adults	2.9(2-4)	5(3-12)
12	Inferior common trunk (from inferior mesenteric)	8	7 adults, 1 child	2.7(2.0-3.5)	9.1(4-22)

Contributors: Gillot, Claude, and A. Delmas

Reference: La Société Anatomique de Paris. Unpublished

11. OSTIAL BREADTHS: MAN

Values are in centimeters and are for both sexes.

Age yr		Tricuspid	Mitral	Pulmonary	Aortic
(A)		(B)	(C)	(D)	(E)
1	Newborn	3.9	3.2	2.7	2.0
2	1-5	5.3	5.0	3.6	3.3
3	6-10	6.8	6.8	4.5	4.2
4	11-15	8.3	7.6	5.9	4.9
5	16-20	11.0	9.5	6.2	5.8

Reference: Krogman, W. M. 1941. Tabulae biol., Berl. 20:664.

12. GROWTH OF ARTERIAL LUMINA: MAN

Values are in square millimeters for cross-sectional areas.

Age	No. of Subjects	Ascending Aorta	Common Carotid Artery		Subclavian Artery		Descending Thoracic Aorta	Smaller Arterial Branches	Sum of D-1 ¹	Ratio of C:J ²
			Right	Left	Right	Left				
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)	(K)
1 Newborn ³	7	27.8	5.2	4.6	7.2	5.1	23.0	2.8	47.9	0.58
2 0-3 mo	31	42.8	5.7	5.9	7.3	6.1	24.6	4.3	53.9	0.79
3 3-12 mo	23	67.6	9.1	8.7	10.8	9.3	30.7	6.8	75.4	0.90
4 1-3 yr	41	93.9	12.2	12.6	15.8	13.1	44.4	9.4	107.5	0.87
5 3-7 yr	19	133.0	16.6	15.7	20.9	17.8	62.2	13.3	146.5	0.91
6 15-18 yr	25	221.7	23.6	22.5	32.0	27.1	107.6	22.2	235.0	0.94
7 20-30 yr	89	280.7	25.5	24.5	39.6	33.1	146.5	28.1	288.0	0.99
8 30-40 yr	60	314.9	28.0	25.6	41.7	40.3	235.0	31.5	325.1	0.97

/1/ Sum of the lumina of the common carotid artery, subclavian artery, descending thoracic aorta, and smaller branches. /2/ Ratio of the lumen of the ascending aorta to the sum of the lumina of the common carotid artery, subclavian artery, descending thoracic aorta, and smaller branches. /3/ Full-term stillborn.

Reference: White House Conference on Child Health and Protection. 1933. Growth and development of the child. Century, New York. Pt. II.

13. THICKNESS OF ARTERIAL WALLS: MAN

Values are in millimeters.

Age yr		Pulmonary Artery	Aorta	Common Carotid
(A)		(B)	(C)	(D)
1	1-2	0.79	0.83	0.55
2	3-4	0.91	1.14	0.59
3	5-6	0.66	1.23	0.60
4	7-8	0.72	1.36	0.89
5	9-10	1.16	1.39	0.79
6	11-13	1.07	1.49	0.90
7	14-16	1.01	1.41	0.90
8	17-19	1.12	1.58	0.90
9	20-22	1.12	1.58	0.93

Reference: Krogman, W. M. 1941. Tabulae biol., Berl. 20:666.

17. HEART WEIGHT DURING FIRST YEAR: MAN

Values are for both sexes. Values in parentheses are ranges, estimate "c" (cf. Introduction).

Age	No. of Observations	Body Length cm	Heart Weight g
(A)	(B)	(C)	(D)
1 3-8(2-7) da	25	53.6	24.5
2 8-6(7-14) da	16	53.0	23.2
3 15.4(14-21) da	20	54.0	22.9
4 22.0(21-30) da	17	53.7	21.5
5 1.3(1-2) mo	6	58.1	30.0
6 3(2-4) mo	28	62.6	30.5
7 5(4-6) mo	14	66.4	31.6
8 7.5(6-9) mo	22	71.3	35.5
9 10.5(9-12) mo	14	79.5	45.0

Reference: White House Conference on Child Health and Protection. 1933. Growth and development of the child. Century, New York. Pt. II.


18. HEART WEIGHT AT VARIOUS AGES: MAN

Age yr	Male		Female	
	No. of Specimens	Heart Weight g	No. of Specimens	Heart Weight g
(A)	(B)	(C)	(D)	(E)
1 Birth	58	19	31	20
2 0-5-0-9	37	41	5	36
3 7-1-9	31	54	15	48
4 12-2-9	24	63	8	62
5 3-3-9	24	73	12	71
6 4-4-9	27	83	11	80
7 5-5-9	27	95	8	90
8 6-6-9	20	103	8	100
9 7-7-9	16	110	5	113
10 8-8-9	13	122	8	126
11 9-9-9	14	132	5	140
12 10-10-9	9	144	6	154
13 11-11-9	4	157	8	168
14 12-12-9	12	180	9	188
15 13-13-9	8	202	2	207
16 14-14-9	8	238	4	226
17 15-15-9	11	258	9	238
18 16-16-9	6	282	9	243
19 17-17-9	18	300	8	247
20 18-18-9	13	310	9	250
21 19-19-9	24	318	11	251
22 20-20-9	29	322	7	252
23 20-29		352		253
24 30-39		342		265
25 40-49		355		280
26 50-59		362		293
27 60-69		365		308
28 70-79		352		319

Contributor: Boyd, Edith

Reference: Boyd, E. 1952. An introduction to human biology and anatomy for first year medical students. Child Research Council, Denver

15. ATRIO-VENTRICULAR WEIGHT RATIO DURING FIRST YEAR: MAN



Reference: Scammon, R. E. 1927. Radiology 9:95.

16. WEIGHT OF VENTRICLES: MAN

Age	Right Ventricle g	Left Ventricle	
		g	% of Body Weight
(A)	(B)	(C)	(D)
Adapted from Müller			
1 Newborn	6.14	7.15	3.00
2 4-6 mo	6.55	12.35	2.97
3 7-12 mo	8.04	16.31	3.19
4 2 yr	12.42	22.00	2.92
5 3 yr	14.98	32.15	2.97
6 4-5 yr	16.24	34.20	3.04
7 6-10 yr	25.01	50.97	3.06
8 11-15 yr	34.00	67.10	2.45
9 16-20 yr	63.40	117.10	2.00
10 21-30 yr	70.80	125.10	2.44
Adapted from Falk			
11 Newborn	6.54	8.14	2.70
12 5-6 mo	9.13	18.13	2.73
13 11-12 mo	12.47	24.48	2.50
14 1-2 yr	15.92	29.29	2.59
15 5-6 yr	23.24	44.98	2.50
16 7-8 yr	25.08	52.25	2.38
17 9-10 yr	27.70	57.74	2.23
18 12-13 yr	38.00	72.87	2.16
19 14-15 yr	49.56	96.74	2.34
20 16-17 yr	66.47	136.87	2.65

Reference: Krogman, W. M. 1941. Tabulae biol., Berl. 20 663. (Adapted from W. Müller 1883. Die Massenverhältnisse des menschlichen Körpers. Hamburg-Leipzig Also from A. A. Falk. 1938. In Dragendorff's Handbuch der Anatomie des Kindes. J. F. Bergmann, Munich. v. 2).

19. HEART WEIGHT: VERTEBRATES (Continued)
Part II. MAMMALS OTHER THAN MAN

19. HEART WEIGHT: VERTEBRATES

Part II. MAMMALS OTHER THAN MAN

Species (A)	No. and Sex (B)	Body Weight kg (C)	Heart Weight (D) (E)		Habitat (F)
			g (D)	g/100 g (E)	
Primates					
1 Baboon (<i>Papio cynocephalus</i>)	1♂	19.51	79.94	0.41	Moto Umbo, Africa
2 Chimpanzee (<i>Troglodytes niger</i>)	1♂	25.75	184.6	0.72	New York
3	1♂	52.16	250.0	0.48	Uganda, Africa
4	1♀	43.99	219.0	0.50	
5 Lemur (<i>Galago senegalensis</i>)	1♀	0.20	1.38	0.69	Maji Moto, Africa
6 Lemur, ring-tailed (<i>Lemur catta</i>)	1♀	1.73	8.2	0.47	New York
7 Monkey, gray (<i>Cercopithecus mitis kibonotensis</i>)	1♀	1.22	6.56	0.54	Maji Moto, Africa
8	1♂	2.90	7.59	0.26	
9	1♂	4.55	35.32	0.78	
10 Monkey, howler, black (<i>Alouatta palliata inconnexa</i>)	10♂, 5♀	0.67	3.9	0.58	Panama
11	3♀	2.68	14.1	0.53	
12	28, 5♀	6.17	20.67	0.34	Panama
13	22	0.43	1.9	0.44	
14 Monkey, howler, brown (<i>A. palliata palliata</i>)	5♂, 5♀	1.21	8.0	0.66	
15	6, 5♀	3.12	7.8	0.25	
16	6, 5♀	5.26	48.0	0.91	South America
17 Monkey, Humboldt (<i>Lagothrix humboldti</i>)	1♀	0.212	1.5	0.71	Panama
18	18♂, 5♀	1.93	7.65	0.39	
19	19♂	1.39	5.3	0.38	Laboratory
20	7♀	3.63	12.2	0.34	
21	4♂	3.29	12.7	0.39	
22	2♀	0.41	2.75	0.67	Panama
23	18♂, 5♀	1.93	7.65	0.40	
24	19♂, 5♀	1.03	4.49	0.44	Panama
25	11♂, 5♀	2.81	14.52	0.52	
26	6♂, 5♀	7.63	32.5	0.43	
27	133♂, 5♀	0.191	1.34	0.71	Panama
28	19♂, 5♀	0.475	3.02	0.64	
29	8, 5♀	0.793	3.91	0.50	
30	2♂	4.94	29.97	0.61	Maji Moto, Africa
31	2♂	4.19	19.98	0.48	Maji Moto, Africa
32	19♂	1.23	6.56	0.53	
33	14, 5♀	3.10	18.6	0.60	Panama
34	6♂, 5♀	1.32	7.53	0.57	
35	272, 5♀	0.59	3.06	0.52	
36	60, 5♀	0.607	3.68	0.61	Panama
37	22	0.167	0.85	0.51	
38	2♂	0.24	1.25	0.52	
Carnivores					
39 Bear, grizzly (<i>Ursus horribilis</i>)	1♀	142.88	1132.5	0.79	Zoo
40	1♂	199.57	1161	0.58	Zoo
41	1♀	317.50	1220	0.38	
42	1♂	0.576	3.0	0.52	Panama
43	2, 5♀	1.54	10.0	0.65	
44	2♂	2.89	12.38	0.43	Florida
45	5♂	3.78	16.82	0.44	Ohio
46	5♂	5.03	19.61	0.39	Florida
47	2♂	1.30	7.51	0.58	Maji Moto, Africa
48	1♀	1.53	8.46	0.55	
49	1♀	2.70	5.87	0.22	Maji Moto, Africa
50	1♂	40.82	159	0.39	Zoo
51	1	0.399	2.0	0.50	Panama
52	1♂	6.25	37.97	0.61	Guatemala
53	1♀	8.51	72.71	0.85	Kansas
54	3♂	14.56	127	0.87	Maji Moto, Africa
55	2♀	12.47	120	0.96	
56	1♂	24.49	308.8	1.26	Tennessee
57	2♂	31.75	297	0.94	Canada
58	2♀	14.75	95.4	0.65	Florida
59	2♂	11.26	126.3	1.12	Ohio
60	4♀	23.71	182.6	0.77	Ohio

1/ Emaciated. 2/ Infant 3/ Juvenile 4/ Goiter. 5/ Peral.

19. HEART WEIGHT: VERTEBRATES

Values in column F, Part I, and column E, Parts II-V, are grams heart weight per 100 grams body weight.

Part I MAN

Subjects	Age yr	Sex	Body Weight kg	Heart Weight		Habitat and Ancestry
				g	g/100 g	
(A)	(B)	(C)	(D)	(E)	(F)	(G)
1 American, Negro		♂	72.57	320	0.44	New York
2	24	♂	71.2	465	0.65	
3	41	♂	86.2	365	0.43	
4	45 ¹	♂	73.0 ¹	389 ¹	0.53 ¹	Louisiana and Ohio
5 American, white		♂	58.97	280	0.48	New York; Irish
6		♂	58.97	308	0.52	New York; Jewish
7		♂ ²	59.42	265	0.45	New York, Czechoslovakian
8		♂	61.24	250	0.41	New York
9		♂	63.50	310	0.49	
10		♂	65.77	270	0.41	New York; Irish
11		♂	70.30	320	0.46	New York; Italian
12		♂	70.30	310	0.44	New York; Jewish
13		♂	72.57	260	0.36	
14	22	♂	81.6	370	0.45	German
15	25	♂	78.5	331	0.42	Czechoslovakian
16	30	♂	65.8	340	0.52	Swedish
17	45	♂	50.0	325	0.65	New York; Scottish
18	54	♂	79.4	390	0.49	Polish
19	54	♂	81.6	400	0.49	
20	41	♂ ³	74.0	382	0.52	Ohio; Polish
21 Chinese		♂	83.89	555	0.66	Honolulu
22 Danish	13	♂	36.0	220	0.61	Denmark
23	28	♂	56.0	380	0.68	
24	39	♂	61.0	410	0.67	
25	44	♀	44.0	310	0.71	
26	47	♀	61.0	300	0.49	
27	50	♀	43.0	220	0.51	
28	55	♂	51.0	380	0.75	
29	55	♂	51.0	290	0.57	
30	60	♀	45.0	410	0.91	
31	65	♀	43.0	320	0.74	
32	68	♀	46.0	260	0.57	
33	68	♂	83.0	520	0.63	
34	78	♀	63.0	450	0.71	
35 Filipino	39	♂	43.09	200	0.46	Hawaii
36 Hawaiian	36-40	♂	95.0	430	0.45	Hawaii
37 Maya Quiche	17	♂ ⁴	37.0	159	0.43	Guatemala
38 Indian	25	♂ ⁵	42.0	220	0.52	
39	28	♀ ⁴	45.8	225	0.49	
40	32	♂ ⁶	45.0	226	0.50	
41	32	♂ ⁷	49.9	240	0.48	
42	35	♂ ⁸	43.4	210	0.48	
43	45	♂ ⁴	36.3	155	0.43	

/1/ Mean value for 4 subjects. /2/ Goiter. /3/ Immigrant. /4/ Death from cachexia. /5/ Death from tuberculosis. /6/ Death from pneumonia (goiter). /7/ Death from double pneumonia. /8/ Death from septicemia (goiter).

Contributors: (a) Quiring, Daniel P., (b) Lowrance, E. W.

Reference: Quiring, D. P. 1950. Functional anatomy of the vertebrates. McGraw-Hill, New York. p. 526.

19. HEART WEIGHT: VERTEBRATES (Continued)
Part II. MAMMALS OTHER THAN MAN (Continued)

Part II. MAMMALS OTHER THAN MAN (CONTINUED)

Species (A)	No and Sex (B)	Body Weight kg (C)	Heart Weight		Habitat (F)
			g (D)	g/100 g (E)	
Rodents (concluded)					
119 Hamster, golden (<i>Cricetus cricetus</i>)	2♂	0.129	0.60	0.47	Ohio
120	2♂	0.108	0.54	0.50	
121 Hare, African (<i>Lepus capensis</i>)	1♀	2.93	30	1.02	Maji Moto, Africa
122 Hare, arctic (<i>L. arcticus arcticus</i>)	2♂	1.90	28.45	1.50	Canada
123	2♂	2.64	28.87	1.09	
124 Lemming, brown (<i>Lemmus trimacrotus</i>)	4♀	0.029	0.43	1.48	Canada
125	1♂	0.048	0.60	1.25	
126 Lemming, rock (<i>Dicrostonyx rubricatus richardsoni</i>)	4♂	0.052	0.31	0.60	Canada
127	3♂	0.055	0.38	0.69	
128 Mouse, African (<i>Mastomys coucha microdon</i>)	1♂	0.022	0.122	0.55	Maji Moto, Africa
129 Mouse, dormouse (<i>Claviglis saturatus</i>)	1♂	0.022	0.113	0.52	Maji Moto, Africa
130	1♂	0.018	0.12	0.67	
131 Mouse, Guatemala (<i>Peromyscus</i> sp) ⁹	14♂	0.012	0.098	0.81	Guatemala
132	2♀	0.019	0.11	0.61	
133	2♂	0.02	0.15	0.75	
134 Mouse, jumping (<i>Zapus hudsonicus</i>) ⁹	3♀	0.02	0.16	0.80	Ohio
135	1♂	0.015	0.13	0.87	
136 Mouse, meadow (<i>Microtus drummondii</i>) ⁹	67♂	0.024	0.16	0.67	Canada
137	42♀	0.023	0.16	0.70	
138	42♂	0.033	0.20	0.61	
139 Mouse, meadow (<i>M. pennsylvanicus pennsylvanicus</i>) ⁹	42♀	0.025	0.197	0.79	Ohio
140	53♂	0.028	0.194	0.69	
141	109♂	0.041	0.257	0.63	
142 Muskrat (<i>Ondatra zibethica alba</i>)	1♂	0.90	3.23	0.36	Canada
143 Porcupine (<i>Erethizon dorsatus</i>)	1♀	2.8	19.75	0.71	Maji Moto, Africa
144	2♀	2.7	14.4	0.53	New York
145	1♂	3.4	15.5	0.46	
146 Rabbit, Flemish giant (<i>Lepus</i> sp)	2♂	3.68	10.57	0.29	Ohio
147	22♀	2.59	9.23	0.36	
148 Rabbit, wild (<i>Silvillagus gabbi messorius</i>)	2	0.44	2.00	0.45	Panama
149 Rat, Norway (<i>Rattus norvegicus</i>)	1♀	0.26	0.73	0.37	Ohio
150	2♂	0.28	1.61	0.58	
151 Squirrel (<i>Sciurus gerrardi mortuus</i>)	1	0.29	2.00	0.69	Panama
152 Squirrel, ground (<i>Citellus paryii paryii</i>)	3♀	0.96	5.92	0.62	Canada
153	5♂	0.88	5.85	0.67	
154 Squirrel, red (<i>Sciurus hudsonicus</i>)	4♀	0.25	1.82	0.73	Canada
155	4♂	0.18	1.57	0.87	
156	1♂	0.17	2.13	1.25	Ohio
157	2♀	0.17	1.23	0.72	
158 Squirrel, red (<i>S. hudsonicus loquax</i>)	1♂	0.18	1.51	0.84	Ohio
159	1♂ ³	0.063	0.545	0.67	
Artiodactyla					
160 Bison, American (<i>Bison bison</i>)	1♂	54.88	361	0.66	Ohio
161 Buffalo (<i>Syncerus caffer caffer</i>)	1♂ ⁸	572	3050	0.53	
162	1♂	759	3628	0.48	Maji Moto, Africa
163 Bushbuck (<i>Tragelaphus scriptus massaicus</i>)	1♀	35.38	325	0.92	Maji Moto, Africa
164	1♂	53.07	350	0.66	Lake Manyara, Africa
165 Caribou, barren ground (<i>Rangifer arcticus arcticus</i>)	1♀	71.87	710	0.99	Canada
166	2♂	128.47	1066	0.85	
167	1♂	62.14	650	1.05	
168 Cattle (<i>Bos taurus</i>)	2 ²	20.75	248.7	1.20	Kentucky
169	1♀ ²	98.43	810.7	0.82	
170	1♀	378.75	2018	0.53	
171	71♀	505	1888	0.37	
172	18♀ ¹⁰	369	1533	0.42	
173 Cattle, Aberdeen angus (<i>B. taurus</i>) ¹¹	1♀	719	1950	0.27	
174 Cattle, Ayrshire (<i>B. taurus</i>) ¹¹	44♀	491	1882	0.38	
175 Cattle, Guernsey (<i>B. taurus</i>) ¹¹	62♀	450	1737	0.39	
176 Cattle, Hereford (<i>B. taurus</i>) ¹¹	7♀	371	1143	0.31	

121 Infans 131 Juvenile 184 Deceased 184

1/2 Infant 1/3 Juvenile 1/8 Pregnant 1/9 Preserved weights 1/10 Castrate 1/11 Data furnished by Swett, W. W., et al 1937 J. Agr. Res 55(4).

19. HEART WEIGHT: VERTEBRATES (Continued)
Part II. MAMMALS OTHER THAN MAN (Continued)

Species		No. and Sex	Body Weight	Heart Weight		Habitat
(A)		(B)	kg (C)	g (D)	g/100 g (E)	(F)
Carnivores (concluded)						
61	Dog, police (<i>Canis familiaris</i>)	1♂	38.42	261	0.68	Ohio
62	Fox, bat-eared (<i>Otocyon megalotis</i>)	1♀	3.34	24.65	0.74	Maji Moto, Africa
63	Fox, gray (<i>Urocyon cinereoargenteus</i> scottii)	1♂	3.76	21.97	0.58	Florida
64	Fox, red (<i>Vulpes fulva</i>)	1♀	4.63	41.75	0.90	Canada
65	Hyena, spotted (<i>Crocuta crocuta</i>)	2	62.37	447	0.72	Maji Moto, Africa
66	Jaguar (<i>Felis onca</i>)	1♀	34.47	186	0.54	Zoo
67	Jackal (<i>Thos mesomelas</i>)	2♂	2.85	21.33	0.75	Kisete Plains, Africa
68	Kinkajou (<i>Potos flavus aztecus</i>)	1♀	2.62	14.3	0.55	Guatemala
69	Leopard (<i>Felis pardus</i>)	1♂	48	200	0.42	Zoo
70		1♀	8.62	52	0.60	
71	Lion (<i>F. leo</i>)	1♂	83.91 ⁷	455	0.54	Zoo
72		1♀	90.72 ⁷	327.3	0.36	Circus
73		1♀ ^{2, 4}	50.9 ⁷	363.5	0.71	Zoo
74		1♂	117.93	810	0.69	
75		1♂	161.52	1614	1.00	
76		1♂	117.57	713	0.60	
77		1♂	94.86	743.2	0.78	
78		1♂	126.08	1078	0.86	
79		1♂	195.4	1175	0.60	Maji Moto, Africa
80		1♂	186.36	860	0.46	
81	Lion, mountain (<i>F. oregonensis</i>)	1♂	28.79	184	0.64	New Mexico
82	Mongoose (<i>Ichneumia albicauda</i>)	1♂	4.40	28.30	0.64	Maji Moto, Africa
83	Raccoon (<i>Procyon lotor pumilus</i>)	1♀	2.23	19.73	0.89	Florida
84		1♀	4.54	31.2	0.69	Zoo
85		1♂	5.22	42.5	0.81	
86	Seal (<i>Phoca richardi geronimensis</i>)	1♂	107.3	1435	1.34	California
87	Seal, bearded (<i>Erignathus barbatus</i>)	1♂	109.7	515	0.47	Canada
88		1♂	281	1245	0.44	
89	Seal, ringed (<i>Phoca hispida</i>)	3♂	39.76	281	0.71	Canada
90		2♀	39.68	302	0.76	
91	Serval (<i>Felis capensis</i>)	1♂	9.96	37.3	0.37	Maji Moto, Africa
92		2♀	5.82	28.45	0.54	
93	Tiger (<i>F. tigris</i>)	1♀	160	432	0.27	Zoo
94		1♂	209	698	0.33	
95	Walrus (<i>Odobenus rosmarus</i>)	1♂ ²	79.38	731	0.92	Canada
96		1♀ ²	55.79	650	1.17	
97		1♂	667	4536	0.68	
98	Weasel, arctic (<i>Mustela arctica</i>)	3♂	0.169	2.83	1.67	Canada
99		1♀	0.121	1.95	1.61	Ohio
100	Wolf, Russian (<i>Canis lupus lupus</i>)	1♂	22.68	246	1.08	Zoo
101	Wolf, timber (<i>C. lubilis</i>)	1♂	29.94	315	1.05	Minnesota
Rodents						
102	Agouti, brown (<i>Dasyprocta punctata</i> dariensis)	12	1.40	10	0.71	Panama
103		3, ♂♀	2.06	9.2	0.45	
104		2, ♂♀	3.17	17.54	0.55	
105	Agouti, spotted (<i>Cumulus paca</i> virgatus)	23, ♂♀	1.37	5.5	0.40	Panama
106		3, ♂♀	3.63	16.1	0.44	
107	Beaver (<i>Castor canadensis</i>)	1♂	4.18	16.57	0.40	Michigan
108		1♀	5.83	27	0.46	
109	Capybara (<i>Hydrochoerus isthmius</i>)	2, ♂♀	27.67	84.13	0.30	Panama
110		12	7.09	24.18	0.34	
111		3♂♂	14.96	55.06	0.37	
112	Chipmunk (<i>Tamias striatus fisheri</i>)	2♂	0.075	0.597	0.80	Ohio
113	Guinea pig (<i>Cavia cutleri</i>)	1♀	0.351	1.72	0.49	Ohio
114		2♂	0.361	1.27	0.35	
115		2♀	0.324	1.31	0.40	
116		46♂, 56♀	0.215	1.29	0.60	
117		10♀	0.432	1.67	0.39	
118		10♂	0.456	1.86	0.41	

/2/ Infant. /3/ Juvenile. /4/ Goiter. /6/ Thyroid enlarged in one animal. /7/ Estimated. /8/ Pregnant.

19. HEART WEIGHT: VERTEBRATES (Continued)
Part B: MAMMALS OTHER THAN MAN (Continued)

Part II - MAMMALS OTHER THAN MONOTREMES						
	Species	No. and Sex	Body Weight kg	Heart Weight		Habitat
				g	g/100 g	
	(A)	(B)	(C)	(D)	(E)	(F)
Perissodactyla (continued)						
233	Horse, polo pony (<i>Equus caballus</i>)	1♀	380.75	3370	0.92	Ohio
234	Horse, saddle (<i>E. caballus</i>)	1♂12	39.50	181.5	0.91	Kentucky
235		2♀3	150.1	1136	0.77	
236		2♂	306.5	1658	0.55	
237		2♂10	335.88	2199	0.66	
238		1♂	449	2495	0.78	Ohio
239	Horse, Shetland pony (<i>E. caballus</i>)	1♂10	242.67	2193	0.90	Kentucky
240		1♀	272.19	2508	0.92	Ohio
241		1♂	150.35	1375	0.91	
242	Horse, standardbred (<i>E. caballus</i>)	1♀12	31.75	402	1.21	Kentucky
243		1♂4	92.98	971	1.04	
244	Horse, thoroughbred (<i>E. caballus</i>)	5♂12, 5♀12	26.89	303.7	1.13	Kentucky
245		18♂2	52.45	565	1.08	
246		19♀12	54.52	406.4	1.32	
247		4♀4	116.71	1125	0.96	
248		8♂3	285.13	1959	0.70	
249		7♀	408.5	1237	0.79	
250		3♂	431.92	1488	0.80	
251		5♂10	445.76	1444	0.73	
252		10♀	441.16	1663	0.81	
253		5♂	485.31	4688	0.97	
254	Horse, western (<i>E. caballus</i>)	1♂10	426.18	1487	0.82	Ohio
255	Mule (<i>E. asinus</i>)	1♀	249.47	2144	0.84	Kentucky
256		1♀	444.52	1604	0.91	
257	Mule, Panama (<i>E. asinus</i>)	1♂1	42.64	397	0.91	Panama
258		4, ♀♀	279.2	2048	0.73	
259	Rhinoceros (<i>Rhinoceros bicornis</i>)	1♂	764	4800	0.63	Maji Moto, Africa
260	Tapir (<i>Tapirella bairdii</i>)	1♂	8.60	74	0.86	Panama
261		1♀	58.06	540	0.93	
262		1	14.26	121	0.85	
263	Zebra (<i>Equus quagga granti</i>)	1♂12	29.48	330	1.12	Zoo
264		1♀2, 4	43.09	581	1.35	
265		1♀2, 4	56.59	515	0.91	Maji Moto, Africa
266		1♂4	78.02	660	0.85	Zoo
267		2♂	254.89	1925	0.75	Maji Moto, Africa
268		18♂	297.8	1970	0.66	
269		1♂	317.5	2251	0.70	Zoo
Proboscidea, Hyracoidea, and Sirenia						
270	Elephant (<i>Loxodonta africana knochenhaueri</i>)	1♂	6554	26,080	0.39	Maji Moto, Africa
271	Hyrax (<i>Heterohyrax brucei</i>)	1♂	0.750	3.65	0.48	Lake Manyara, Africa
272	Manatee (<i>Trichechus manatus</i>)	1♂	434	3250	0.29	Florida
273		1♀	557.92	1241	0.22	
Cetaceans						
274	Porpoise (<i>Phocaena phocaena</i>)	1♂	142.43	738	0.52	Florida
275	Whale, humpbacked (<i>Megaptera nodosa</i>)	1♂	57,395	214,000	0.38	California
276		2♂	40,370	187,000	0.46	
277	Whale, sperm (<i>Physeter catodon</i>)	1♂	39,009	126,000	0.32	California
278	Whale, beluga (<i>Delphinapterus leucas</i>)	2♀	303.25	1722	0.57	Canada
279		4♀	411.6	2454	0.59	
Insectivores						
280	Mole (<i>Scalopus aquaticus</i>)	1♂	0.0396	0.272	0.69	Ohio
281	Shrew, short-tailed (<i>Blarina brevicauda</i>)	3♀♀	0.0163	0.1723	1.06	Ohio
282		2♀♀	0.0188	0.1922	1.02	
Edentates						
283	Anteater (<i>Tamandua tetradactyla chiquensis</i>)	1♂	3.409	2.0	0.58	Panama
284		2, ♀♀	3.692	18.0	0.49	
285	Armadillo (<i>Dasypus novemcinctus fenestratus</i>)	7♂, ♀♀	0.471	2.5	0.45	Panama
286		4♂, ♀♀	1.392	6.0	0.45	
287		1♂	3.701	14.2	0.40	
288		10♂	3.401	9.0	0.26	

(2) Infant (3) Juvenile (4) Gether (5) Pregnant (10) Castrate (12) Fetus.

19. HEART WEIGHT: VERTEBRATES (Continued)

Part II: MAMMALS OTHER THAN MAN (Continued)

	Species (A)	No. and Sex (B)	Body Weight kg (C)	Heart Weight g g/100 g (D) (E)		Habitat (F)
Artiodactyls (concluded)						
177	Cattle, Holstein (<i>Bos taurus</i>) ¹¹	2009	574	2245	0.39	
178		6σ ²	90	422	0.47	
179		3σ ³	241	998	0.41	
180		5σ	552	1905	0.35	
181		5σ	888	3357	0.38	
182	Cattle, Jersey (<i>B. taurus</i>) ¹¹	2189	413	1605	0.39	
183		3σ ²	51.9	304	0.59	
184		5σ ³	214	744	0.35	
185		2σ	367	1270	0.35	
186		10σ	597	2186	0.37	
187		213σ	412	1606	0.39	
188	Deer (<i>Odocoileus chiriquensis</i>)	23	2.38	5.16	0.22	Panama
189		19	13.93	15.9	0.11	
190	Deer, Indian axis (<i>Cervus axis</i>)	1σ ⁴	88.45	749	0.85	Zoo
191	Deer, white-tailed (<i>Odocoileus virginianus</i>)	1σ	65.22	632	0.97	Zoo
192	Dik-dik (<i>Rhynchotragus kirki</i>)	1σ	4.57	36.8	0.81	Maji Moto, Africa
193	Elk, American (<i>Cervus canadensis</i>)	1σ	13.61	131.1	0.96	Zoo
194	Gazelle, Thomson's (<i>Gazella thomsoni</i>)	19 ²	2.43	31.7	1.30	Maji Moto, Africa
195		2σ	24.37	245	1.01	
196	Giraffe (<i>Giraffa camelopardalis tippelskirchi</i>)	1σ	1220	4990	0.41	Maji Moto, Africa
197	Hartebeest, Coke's (<i>Bubalis cokei cokei</i>)	1σ	134	875	0.65	Moto Umbo, Africa
198	Hippopotamus (<i>Hippopotamus amphibius</i>)	19 ³	543	1610	0.30	Maji Moto, Africa
199		19	1351	4536	0.34	
200	Impala (<i>Aepyceros melampus</i>)	2σ	37.86	270.05	0.71	Maji Moto, Africa
201		2σ	57.61	378	0.66	
202	Peccary, collared (<i>Pecari angulatus bangsi</i>)	1σ ⁴	29	174	0.60	Zoo
203		12	2.25	20	0.89	Panama
204		2	13.83	70.5	0.51	
205	Reedbuck (<i>Redunca redunca tothi</i>)	2σ	31.7	242	0.76	Lake Mangara, Africa
206	Sheep (<i>Ovis aries</i>)	212	2.72	28.30	1.04	Kentucky
207		59 ²	8.55	85	0.99	
208		4σ	15.76	158.7	1.01	
209		79	52.1	276.7	0.53	
210		9σ	40.23	168	0.42	
211	Steinbok (<i>Raphicerus campestris neumanni</i>)	2σ	8.62	72.2	0.84	Maji Moto, Africa
212	Swine (<i>Sus scrofa</i>)	512, σ♀	1.42	14.25	1.00	Kentucky
213		1σ ³	4.76	52.1	1.09	
214		193	13.15	124.5	0.95	
215		19	113.2	452	0.40	
216		53σ ¹⁰	102.06	303.5	0.30	
217		369	102.06	324.39	0.32	
218	Warthog (<i>Phacochoerus aethiopicus</i>)	1σ	65.32	325	0.50	Maji Moto, Africa
219	Wildbeest (<i>Connochaetes</i> sp.)	2σ	212.07	1313	0.62	Maji Moto, Africa
Perissodactyls						
220	Burro (<i>Equus asinus</i>)	19	150.73	825	0.55	Guatemala
221		1σ ¹⁰	199.58	1184	0.59	Kentucky
222		1σ	376.48	2607	0.69	Kentucky
		1σ	412.28	3587	0.87	Ohio
		1σ	122.98	850	0.69	Guatemala
		1σ	521.64	3260	0.62	Kentucky
		1σ	70.76	546	0.77	Kentucky
		1σ	184.16	1165	0.63	Kentucky
		1σ	362.87	1427	0.39	Illinois
		1σ	402.65	3648	0.91	Ohio
		1σ	230.9	1843	0.80	Panama
		1σ	635.04	5600	0.88	Ohio
232		19	771.40	4700	0.61	

/2/ Infant. /3/ Juvenile. /4/ Götter. /10/ Castrate. /11/ Data furnished by Swett, W. W., et al. 1937. J. Agr. Res. 55(4). /12/ Fetus.

19. HEART WEIGHT: VERTEBRATES (Continued)

Part III. BIRDS (Concluded)

	Species	No. and Sex	Body Weight kg	Heart Weight		Habitat
				g	g/100 g	
	(A)	(B)	(C)	(D)	(E)	(F)
39	Hawk, red-tailed (<i>Buteo borealis</i>)	1♀	1.029	6.94	0.67	Ohio
40	Hawk, sharp-shinned (<i>Accipiter velox velox</i>)	1♀	0.320	3.70	0.71	Ohio
41	Hawk, sparrow (<i>Falco sparverius sparverius</i>)	1♂	0.112	1.12	1.00	Florida
42	Hornbill, ground (<i>Bucorvus cafer</i>)	1♂	3.25	56.61	1.74	Maji Moto, Africa
43	Hummingbird (<i>Amarilia izacati izacati</i>)	1♀	0.0048	0.1138	2.37	Guatemala
44	Loon, red-throated (<i>Gavia stellata</i>)	1♀	1.530	22.72	1.49	Canada
45		2♂	1.568	22.24	1.42	
46	Man-of-war (<i>Fregata aquila</i>)	1♂	1.405	16.59	1.18	Florida
47	Merganser, red-breasted (<i>Mergus serrator</i>)	1♀	0.770	11.01	1.43	Canada
48	Ostrich, Masai (<i>Struthio camelus massaicus</i>)	1♂	123	1205	0.98	Maji Moto, Africa
49	Owl, horned (<i>Bubo virginianus virginianus</i>)	1♂	1.177	8.60	0.73	Ohio
50	Partridge (<i>Francoelinus sephaena</i>)	1♂	0.208	1.46	0.70	Maji Moto, Africa
51	Salin (<i>Salinopsis occidentalis</i>)	20	1.30	13.13	0.29	Florida

70	Swallow, barn (<i>Hirundo erythrogastra</i>)	1♂	0.0215	0.302	1.41	Ohio
71		2♀	0.021	0.283	1.35	
72	Tesi, green-winged (<i>Nettion carolinensis</i>)	1♀	0.305	2.88	0.94	Canada
73	Vulture (<i>Pseudogyps africanus</i>)	1♀	5.27	37.85	0.72	Maji Moto, Africa

1/1 Juvenile

Contributor Quiring, Daniel P

Reference Quiring, D. P. 1950. Functional anatomy of the vertebrates McGraw-Hill, New York. p. 528.

19. HEART WEIGHT: VERTEBRATES (Continued)
Part II: MAMMALS OTHER THAN MAN (Concluded)

	Species (A)	No. and Sex (B)	Body Weight kg (C)	Heart Weight		Habitat (F)
				g (D)	g/100 g (E)	
Edentates (concluded)						
289	Sloth, three-toed (<i>Bradypus griseus</i>)	12	0.676	3.0	0.44	Panama
290	<i>griseus</i>)	43, ♂♀	2.005	6.1	0.30	
Marsupials						
291	<i>Opossum</i> [<i>Didelphis marsupialis</i>]	83, ♂♀	0.666	3.4	0.51	Panama
292	<i>etensis</i>)	4, ♂♀	1.147	5.0	0.44	
293	<i>Opossum</i> , woolly (<i>Philander laniger</i>)	13	0.158	1.5	0.95	Panama
294	<i>pallidus</i>)	1	0.222	3.0	1.35	

/2/ Infant. /3/ Juvenile.

Contributors: (a) Quiring, Daniel P., (b) Lowrance, E. W.

Reference: Quiring, D. P. 1950. Functional anatomy of the vertebrates. McGraw-Hill, New York. p. 528.

Part III: BIRDS

	Species (A)	No. and Sex (B)	Body Weight kg (C)	Heart Weight		Habitat (F)
				g (D)	g/100 g (E)	
1	Blackbird (<i>Quiscalus quiscula aeneus</i>)	1♂	0.082	1.159	1.41	Ohio
2	Bluebird (<i>Sialia sialis sialis</i>)	1♀	0.034	0.383	1.13	Ohio
3		1♂	0.029	0.493	1.70	
4	Bustard, greater (<i>Choriotis kori</i>)	1♀	5.54	60.25	1.09	Athi Plain, Africa
5	struthiunculus)	1♂	10	97	0.97	
6	Bustard, lesser (<i>Haliaeetus bociter</i>)	1♀	1.100	11.93	1.09	Maji Moto, Africa
7	Buzzard, steppe (<i>Buteo vulpinus vulpinus</i>)	1♂	0.560	4.58	0.82	Maji Moto, Africa
8	Buzzard, turkey (<i>Cathartes aura</i>)	1♂	0.495	10.24	2.07	Florida
9	Canary (<i>Serinus canarius</i>)	1♂	0.017	0.285	1.68	Ohio
10		1♀	0.015	0.133	0.89	
11	Catbird (<i>Dumetella carolinensis</i>)	1♂	0.033	0.327	0.99	Ohio
12	Cowbird (<i>Molothrus ater ater</i>)	1♀	0.066	1.06	1.61	Ohio
13	Crane, crested (<i>Balearica pavonina</i>)	1♀	4.071	41.21	1.01	Lake Manyara, Africa
14		1♂	4.825	32.37	0.68	
15	Crane, gray (<i>Grus canadensis</i>)	1♂	1.651	19.01	1.15	Florida
16	Crow (<i>Corvus brachyrhynchos</i>)	1♂	0.337	3.2	0.95	Ohio
17	Dovekie (<i>Alle alle</i>)	1♀	0.102	1.24	1.22	Florida
18		1♂	0.104	1.24	1.19	
19	Duck (<i>Nyroca affinis</i>)	1♂	1.041	13.4	1.29	Ohio
20	Duck, pintail (<i>Dafila acuta tsitzilhoa</i>)	1♀	0.670	8.34	1.24	Canada
21	Eagle, fish (<i>Haliaeetus vocifer vocifer</i>)	1♀	3.50	32.23	0.92	Maji Moto, Africa
22	Eagle, tawny (<i>Aquila rapax rapax</i>)	2♀	2.625	18.54	0.71	Maji Moto, Africa
23		2♂	2.438	14.17	0.58	
24	Egret, great white (<i>Casmerodius albus</i>)	1♀	1.03	9.25	0.90	Maji Moto, Africa
25	melanorhynchos)					
25	Egret, yellow-billed (<i>Mesophyx intermedia</i>)	1♂	0.525	5.0	0.86	Maji Moto, Africa
26	brachyrhyncha)					
26	Flamingo (<i>Phoeniconotus minor</i>)	3♀	1.483	14.68	0.99	Maji Moto, Africa
27		2♂	1.598	13.53	0.85	
27	Fowl, Leghorn (<i>Gallus gallus domesticus</i>)	10♀	0.4907	2.80	0.57	Ohio
28		10♂	0.5007	3.18	0.64	
29		16♀	0.6151	3.91	0.64	
30		8♂	0.7331	4.19	0.57	
31		1♂	2.20	8.78	0.40	Ohio
32	Fowl, white Orpington (<i>G. gallus</i>)					
					0.96	Lake Manyara, Africa
					0.88	Maji Moto, Africa
					1.68	Canada
					0.98	Florida
					1.03	Florida
					0.92	Florida

19. HEART WEIGHT: VERTEBRATES (Continued)

Part V: FISHES

Species (A)	No. and Sex (B)	Body Weight kg (C)	Heart Weight g (D) g/100 g (E)		Habitat (F)
1 Barracuda (<i>Sphyraena</i> barracuda)	1♀	4.803	7.78	0.16	Florida
	1♂	5.990	12.25	0.20	
	1♀	8.675	11.60	0.13	
	1♂	9.150	60.31	0.66	
	1♂	11.33	19.27	0.17	
	1♀	12.69	16.03	0.13	
7 Bonita (<i>Gymnosarda</i> alleterata)	1♂	6.291	29.43	0.47	Florida
8 Carp (<i>Cyprinus carpio</i>)	4♂	1.0421	1.617	0.15	Ohio
	2♀	1.061	1.554	0.13	
	2♀	4.942	49.60	1.00	
10 Catfish (<i>Catfish</i> sp)	1♂	2.900	17.65	0.61	Maji Moto, Africa
12 Cisco (<i>Argyrosomus</i> artedii)	2♂	0.1621	1.592	0.98	Canada
13 Codfish (<i>Gadus callarias</i>)	1♀	10.6	161.4	1.52	Massachusetts
	3♀	2.625	97.1	3.70	
	3♂	2.518	94.2	3.82	
16 Eel, green moray (<i>Gymnothorax funebris</i>)	1	3.510	101.5	2.89	Florida
17 Grouper, black (<i>Mycteroperca</i> <i>bonasus xanthosticta</i>)	1♂	2.712	11.66	0.43	Florida
18 Grunt, white (<i>Haemulon</i> <i>plumieri</i>)	1♂	0.30	3.55	1.18	Florida
19 Haddock (<i>Melanogrammus</i> <i>aeglefinus</i>)	6♀	5.275	132.6	4.05	North Atlantic Ocean
20 Hogfish (<i>Lachnolaimus</i> <i>maximus</i>)	1♂	0.480	0.432	0.09	Florida
21 Jack, common (<i>Caram</i> <i>hippos</i>)	1♂	2.305	4.87	0.21	Florida
22 Jack, yellow (<i>C.</i> <i>bartholomaei</i>)	1♀	4.274	11.62	0.21	Florida
23	1♂	4.912	13.96	0.29	
24 Jewfish (<i>Promicrops</i> <i>italara</i>)	1♂	32.89	49.23	0.15	Florida
25 Kingfish (<i>Scomberomorus</i> <i>regalis</i>)	1♂	2.43	5.33	0.22	Florida
26 Mackerel (<i>S. maculatus</i>)	2♀	1.457	2.93	0.20	Florida
27 Marlin, white (<i>Makaira</i> <i>albida</i>)	1♀	24.94	47.60	0.19	Florida
28 Muttonfish (<i>Lutjanus</i> <i>analis</i>)	1♀	2.49	5.12	0.21	Florida
29 Palometa (<i>Trachinotus</i> <i>glauca</i>)	1♀	8.504	24.85	0.29	Florida
30 Perch (<i>Perca flavescens</i>)	1♀	0.192	0.44	0.13	Lake Erie
31	6♂	0.167	0.3087	0.19	
32 Pike (<i>Esox lucius</i>)	4♂	0.3523	0.660	0.19	Lake Erie
33	2♀	0.3738	0.560	0.15	
34 Ray, electric (<i>Torpedo</i> <i>torpedo</i>)	1♂	0.345	0.22	0.06	Florida
35 Ray, sting (<i>Dasyatis</i> <i>sabina</i>)	2♂	17.58	47.58	0.27	Florida
36	1	2.675	6.83	0.26	
37 Sailfish (<i>Istiophorus</i> <i>americanus</i>)	1♀	25.20	49.15	0.20	Florida
38 Salmon (<i>Salmo salar</i>)	4♀	5.361	10.72	0.20	Coast of Maine
39	2♂	4.922	15.22	0.31	
40	1♂	3.93	6.42	0.16	
41	1♀	5.412	7.92	0.15	
42 Shad (<i>Alosa onkensis</i>)	1♀	0.615	2.21	0.36	Ohio
43 Shark, sand (<i>Carcharias</i> <i>littoralis</i>)	1♀	35.38	41.6	0.12	
44	1♂	35.83	40.14	0.11	
45	1♀ ¹	123	115	0.09	Florida

1/1 Pregnant.

19. HEART WEIGHT: VERTEBRATES (Continued)

Part IV: REPTILES AND AMPHIBIANS

	Species	No. and Sex	Body Weight kg	Heart Weight		Habitat
				g	g/100 g	
	(A)	(B)	(C)	(D)	(E)	(F)
1	Alligator (Alligator mississippiensis)	1♂	0.105	1.39	1.32	
2		2♂	0.351	1.62	0.46	Florida
3		1♀	52.4	137	0.26	Great Lakes Exposition
4		2♂	189	286.5	0.15	Florida
5	Crocodile (Crocodilus americanus)	1♀	134	134	1.00	Florida
6	Lizard (Lacerta viridis)	1♀	0.024	0.091	0.38	Maji Moto, Africa
7	Lizard, Gila monster (Heloderma suspectum)	1♀	0.514	4.17	0.81	Arizona
8	Lizard, horned toad (Phrynosoma cornutum)	2♂	0.025	0.125	0.50	Arizona
9	Lizard, iguana (Iguana iguana)	1♀	1.34	2.60	0.19	Guatemala
10	Lizard, iguana (Amblyrhynchus cristatus) ²	1♀	4.19	21.51	0.51	Galapagos Islands
11	Snake, black (Coluber constrictor)	1♀	0.590	6.08	1.31	Ohio
12		1♂	0.286	0.96	0.34	Laboratory
13	Snake, boa imperator (Boa imperator)	1♀	1.829	5.64	0.31	Guatemala
14	Snake, garter (Thamnophis sirtalis)	1♀ ³	0.057	1.02	1.79	Ohio
15	Snake, python (Python molurus)	1♂	6.14	18.5	0.30	Enguruka, Africa
16	Snake, water moccasin (Agkistrodon piscivorus)	1♀	0.728	4.77	0.66	Florida
17	Turtle (Clemmys guttata)	1♂	0.185	0.57	0.31	Ohio
18		1♂	2.163	7.05	0.33	
19	Turtle (Chelydra serpentina)	1♀	5.125	13.43	0.26	Maji Moto, Africa
20	Turtle, green (Chelonia mydas)	1♀	68.04	180	0.26	Florida
21		1♂	114.3	435	0.38	
22	Turtle, leathernecked (Amyda forex)	1♀	3.253	16.0	0.49	Florida
23	Turtle, snapping (Macrochelys latertina)	1♂	1.848	12.88	0.70	Ohio
24	Frog, bull (Rana catesbeiana)	1♂	0.339	1.37	0.40	North Carolina
25		6♂	0.5199	1.65	0.32	Louisiana

1/ Infant. 2/ Preserved weights. 3/ Pregnant.

Contributor: Quirling, Daniel P.

Reference: Quirling, D. P. 1950. Functional anatomy of the vertebrates. McGraw-Hill, New York. p. 528.

21. CAPILLARY VASCULARITY OF MUSCLE: VERTEBRATES

"Capillary" counts from injected cross-section may not exclude other minute vessels (e.g., arterioles, venules).

Part I SKELETAL MUSCLE

For information on capillary capacity and surface area, see references 15-24. Abbreviations: A = asphyxiated, N = nourished, R = resting, S = starved, T = trained, U = untrained, V = vasodilated, W = working, SL = sea level, HA = high altitude (13,800 ft).

Animal	Muscle	Physiologic Condition	Capillaries		Reference
			No./sq mm tissue ¹ (cross-section)	No./fiber	
(A)	(B)	(C)	(D)	(E)	(F)
Mammals					
1 Cat	Gastrocnemius	V	2343		1
2	Rectus femoris	V	2474		
3	Tibialis anterior	V	2214		
4 Dog, puppy	Gracilis	R	1690	0.7	2
5 Dog, adult	Gracilis	R	1050	0.6	2
6	Gracilis	W ²	2010	1.2	
7	Gracilis	V	2580		
8	Seminembranosus	R	3240	0.6	
9	Seminembranosus	V	5900	1.6	
10	Seminembranosus	V	2630		
11 Guinea pig	Abdominal wall	R	85		3
12	Abdominal wall	R	270		4
13	Abdominal wall	V	1400		
14	Diaphragm	W ³	2570		
15	Diaphragm	V, W ³	3000		
16	Gastrocnemius	R	1136		
17	Gastrocnemius	U	1378		5
18	Gastrocnemius	T	1924		6
19	Gastrocnemius	W ²	2000		
20	Gastrocnemius	A	2600		5
21	Gastrocnemius	V	2614		
22	Gastrocnemius, external	SL	1542		1
23	Gastrocnemius, external	HA	2183	2.5	7
24	Gastrocnemius, internal	SL	1621	1.4	
25	Gastrocnemius, internal	HA	2207	2.5	
26	Masseter	R	1927	3.5	
27	Masseter	R	2840		5
28	Masseter	U	2873		3
29	Masseter	T	3064		6
30	Masseter	W ²	3200		
31	Masseter	A	3150		5
32	Pretracheal	V	3760		
33	Rectus femoris	V	3060		3
34	Rectus femoris	SL	1702		1
35	Rectus femoris	HA	2326	2.7	7
36	Thigh muscle, anterior (white)	SL	686	3.5	
37	Tibialis anterior	V	3501	1.5	
38	Vastus lateralis (white)	SL	740		1
39	Vastus lateralis (red) ⁴	SL	1990	1.4	7
40	Vastus lateralis (red) ⁵	SL	1897	2.2	
41	Vastus lateralis (red)	HA	2407	3.0	
42 Horse	Gastrocnemius	V	1350		
43 Mouse	Gastrocnemius	R	3060	3.8	
44	Gastrocnemius	W ²	3600		3
45	Gastrocnemius	A	4650		5
46	Masseter	W ²	4593		
47	Masseter	V, W ²	5100		
48	Masseter	A	6800		
49 Rabbit	Adductor magnus (white)	V	1550		
50	Gastrocnemius	V	1344	0.7	8
51	Glimed	U		1.6	9
52	Glimed	T		0.5	10
53	Rectus femoris	V	1475	1.2	
54	Semitendinosus (red)	V	790	1.2	9
				1.2	8

1/ Differences in counts during rest, work, and vasodilatation are now attributed to constriction of the meta-arteriole and to precapillary sphincter action [13, 14]. Because of parallel capillary pattern, counts are averages.
2/ Muscle or animal stimulated before injection. 3/ Classified as working (W), on basis of involuntary use.
4/ Young animal. 5/ Older animal.

19. HEART WEIGHT: VERTEBRATES (Concluded)

Part V: FISHES (Concluded)

Species (A)	No. and Sex (B)	Body Weight kg (C)	Heart Weight		Habitat (F)
			g (D)	g/100 g (E)	
46 Shark, tiger (<i>Galeocerdo tigrinus</i>)	1♀	200	291.5	0.15	Florida
47 Sheephead (<i>Aplodinatus grunniens</i>)	1♀	0.937	1.71	0.18	Ohio
48 Smelt (<i>Osmerus mordax</i>)	1	0.04766	1.29	2.71	Lake Erie
49 Snapper, vermillion (<i>Rhomboplites aurorubens</i>)	1♂	0.202	0.33	0.16	Florida
50 Trout, brown (<i>Salmo trutta</i>)	1♂	0.292	0.385	0.13	Ohio
51 Trout, northern (<i>Cristivomer namaycush</i>)	5♀	3.24	5.17	0.16	Canada
52	1♂	2.50	4.34	0.17	
53 Trout, rainbow (<i>Salmo irideus</i>)	4♀	0.2177	0.280	0.13	Ohio
54	2♂	0.2608	0.307	0.12	
55	1♂	0.0428	0.095	0.22	
56	1♀ ³	0.1101	0.155	0.14	
57 Trout, salmon (<i>S. gairdneri</i>)	2♂	2.75	6.22	0.23	Canada
58 Trout, speckled (<i>Salvelinus fontinalis</i>)	1♀	0.255	0.99	0.39	Pennsylvania
59	1♂	0.2301	1.15	0.50	
60 Tuna (<i>Thunus secundodorsalis</i>)	1♂	5.21	31.22	0.60	Florida
61 Turbot (<i>Balistes carolinensis</i>)	1♂	0.295	0.364	0.12	Florida
62 Whitefish (<i>Coregonus clupeiformis</i>)	3♂, 3♀	0.7726	0.933	0.12	Lake Erie
63 Yellowtail (<i>Ocyurus chrysurus</i>)	1♀	0.255	0.37	0.15	Florida

/2/ Infant. /3/ Yearling

Contributor: Quiring, Daniel P.

Reference: Quiring, D. P. 1950. Functional anatomy of the vertebrates. McGraw-Hill, New York. p 528.

20. DENSITY, SURFACE, AND MAXIMUM DIFFUSION DISTANCE OF CAPILLARIES
IN VARIOUS TISSUES: VERTEBRATES

Tissue (A)	Animal (B)	Capillary Density no./sq mm (C)	Capillary Surface sq cm/cu cm tissue (D)	Maximum Diffusion Distance μ (E)
1 Adipose				
2 Fat-rich	Rat	274	52	34
3 Fat-poor	Rat	1000	222	18
4 Brain, white matter	Man	300	57	33
5 Cerebellum	Mouse	1700	330	14
6 Cerebral cortex	Man	1000	190	18
7	Mouse	1250	240	16
8 Duodenum	Mouse	2400	460	11
9 Liver	Mouse	4200	800	9
10 Muscle	Dog	2600	590	11
11	Horse	1400	240	15
12	Frog	400	190	28
13 Heart	Mouse	5300	1000	8
14 Papillary	Man	5220	990	8
15 Pancreas	Guinea pig	1900	360	13
16 Renal cortex	Mouse	4500	850	8
17 Renal medulla	Mouse	7400	1400	7
18 Ventricle, left	Man	5730	1090	8
19 Right	Man	5680	1080	8
20 Ventricular septum	Man	4450	850	8

Reference: Kety, Seymour S. 1951. Pharm. Rev., Balt. 3.5.

21. CAPILLARY VASCULARITY OF MUSCLE: VERTEBRATES

"Capillary" counts from injected cross-section may not exclude other minute vessels (e.g., arterioles, venules).

Part I SKELETAL MUSCLE

For information on capillary capacity and surface area, see references 15-24. Abbreviations: A = asphyxiated, N = nourished, R = resting, S = starved, T = trained, U = untrained, V = vasodilated, W = working, SL = sea level, HA = high altitude (13,800 ft).

Animal	Muscle	Physiologic Condition	Capillaries		Reference
			No./sq mm tissue ¹ (cross-section)	No./fiber (E)	
(A)	(B)	(C)	(D)	(E)	(F)
Mammals					
1 Cat	Gastrocnemius	V	2341		1
	Rectus femoris	V	2474		
	Tibialis anterior	R	2214		2
	Gracilis	R	1690	0.7	2
2 Dog, puppy	Gracilis	R	1030	0.6	
3 Dog, adult	Gracilis	W ²	2010	1.2	
4	Gracilis	V	2580		
5	Gracilis	V	3240	0.6	
6	Seminembranosus	R	5900	1.6	
7	Seminembranosus	V	2630		3
8	Seminembranosus	V	85		4
9	Abdominal wall	R	270		
10	Abdominal wall	V	1400		
11	Abdominal wall	W ¹	2570		
12	Diaphragm	V, W ³	3000		5
13	Diaphragm	R	1136		6
14	Gastrocnemius	U	1378		
15	Gastrocnemius	T	1924		5
16	Gastrocnemius	W ²	2000		
17	Gastrocnemius	A	2600		1
18	Gastrocnemius	V	2614		7
19	Gastrocnemius	SL	1542	2.5	
20	Gastrocnemius, external	HA	2103	3.4	
21	Gastrocnemius, external	SL	1621	2.5	
22	Gastrocnemius, internal	HA	2207	3.5	
23	Gastrocnemius, internal	R	1927		5
24	Masseter	R	2840		3
25	Masseter	U	2873		6
26	Masseter	T	3064		
27	Masseter	W ²	3200		5
28	Masseter	A	3150		
29	Masseter	V	3700		3
30	Pretracheal	V	3060		1
31	Rectus femoris	SL	1702		7
32	Rectus femoris	HA	2326	2.7	
33	Rectus femoris	SL	686	3.5	
34	Thigh muscle, anterior (white)	V	3501	1.5	
35	Tibialis anterior	V	740		1
36	Vastus lateralis (white)	SL	1990	1.4	7
37	Vastus lateralis (red) ⁴	SL	1897	2.2	
38	Vastus lateralis (red) ⁵	SL	2407	3.0	
39	Vastus lateralis (red)	HA	1350	3.8	
40	Gastrocnemius	V	3060		3
41	Gastrocnemius	R	3600		5
42	Gastrocnemius	W ²	4650		
43	Gastrocnemius	A	4393		
44	Masseter	W ²	5100		
45	Masseter	V, W ²	6800		
46	Masseter	A	1550	0.7	8
47	Adductor magnus (white)	V	1344	1.6	9
48	Gastrocnemius	U		0.5	10
49	Gluteal	T		1.2	
50	Gluteal	V	1475	1.2	9
51	Rectus femoris	V	790	1.2	8
52	Semitendinosus (red)	V			
53					
54					

vasodilatation are now attributed to constriction of the meta-
4) Because of parallel capillary pattern, counts are averages.
3) Classified as working (W), on basis of involuntary use.

21. CAPILLARY VASCULARITY OF MUSCLE: VERTEBRATES (Continued)

Part I: SKELETAL MUSCLE (Concluded)

Animal	Muscle	Physiologic Condition	Capillaries		Reference
			No./sq mm tissue ¹ (cross-section)	No./fiber	
(A)	(B)	(C)	(D)	(E)	(F)
Mammals (concluded)					
55 Rabbit (concluded)	Semitendinosus (red)	V	1198	3.1	9
56 Rat	Rectus femoris	V	3573		1
57	Tibialis anterior	V	3402		
Amphibians					
58 Frog (<i>Rana esculenta</i>)	Gastrocnemius	V	286	1.9	11
59	Obliquus oculi inferior	V	314	0.2	
60	Sartorius	V	309	1.3	
61	Submaxillaris	V	514	0.7	
62 Frog (<i>R. temporaria</i>)	Gastrocnemius	V	382	2.6	11
63	Gracilis	V	406	1.9	
64	Obliquus oculi inferior	V	393	0.2	
65	Sartorius	R	90		4
66	Sartorius	W ²	325		
67	Sartorius	V	400		3
68	Sartorius	V	426	1.8	11
69	Submaxillaris	V	652	0.7	
70 Salamander (<i>Ambystoma mexicanum</i>)	Dorsalis trunci	V	133	1.0	11
71	Extensor iliotibialis	V	264	0.4	
72 Toad (<i>Bufo calamita</i>)	Gastrocnemius	V	444	1.6	11
73	Sartorius	N, V	520	1.2	
74	Sartorius	S, V	1021	1.6	
Fishes					
75 Cod	Eye	W ³	400		3
76 Hagfish (<i>Myxine</i> sp.)	Axial		56		12
77 Trout	Eye	W ³	276		12
78	Parietal	R	123		
79	Rectus lateralis	W ³	400		

/1/ Differences in counts during rest, work, and vasodilatation are now attributed to constriction of the meta-arteriole and to precapillary sphincter action [13, 14]. Because of parallel capillary pattern, counts are averages. /2/ Muscle or animal stimulated before injection. /3/ Classified as working (W), on basis of involuntary use.

Contributors: (a) Fulton, George P., and Herbert J. Berman, (b) Shipley, Reginald A., (c) Burton, Alan C., (d) Lee, Richard E.

References: [1] Paff, G. H. 1930. *Anat. Rec.* 46:401. [2] Martin, E. G., E. C. Woolley, and M. Miller. 1931. *Am. J. Physiol.* 100:407. [3] Krogh, A. 1919. *J. Physiol., Lond.* 52:409. [4] Krogh, A. 1919. *Ibid.* 52:457. [5] Sjostrand, T. 1935. *Skand. Arch. Physiol., Berl., Suppl.* 71:1. [6] Petren, T., T. Sjostrand, and B. Sylven. 1936. *Arbeitsphysiologie* 9:376. [7] Valdivia, E. 1958. *Am. J. Physiol.* 194:585. [8] Stoel, G. 1925. *Zachr. Zellforsch.* 3:91. [9] Duff, J. W., and H. D. Bouman. 1927. *Ibid.* 5:596. [10] Vanotti, A., and M. Magiday. 1934. *Arbeitsphysiologie* 7:615. [11] Steudel, W. 1938. *Zool. Jahrb. Abt. 2 Anat. Ontog.* 65:63. [12] Gorkiewicz, G. 1947. *Bull. Internat. Acad. polon. sc., cl. med., Ser. B, II*, p. 241. [13] Fulton, G. P., and B. R. Lutz. 1940. *Science* 92:223. [14] Chambers, R., and B. W. Zweifach. 1944. *Am. J. Anat.* 75:173. [15] Scarborough, R. A. 1930-31. *Yale J. Biol.* 3:63. [16] Dukes, H. H. 1942. *The physiology of domestic animals*. Cornell University Press, Ithaca. [17] Ponder, E. 1948. *Hemolysis and related phenomena*. Grune and Stratton, New York. [18] Prosser, C. L. 1950. *Comparative animal physiology*. W. B. Saunders, Philadelphia. [19] Krogh, A. 1929. *The anatomy and physiology of capillaries*. Yale University Press, New Haven. [20] Gibson, J. G., et al. 1946. *J. Clin. Invest.* 25:848. [21] Klein, J. R. 1945. *Arch. Biochem.*, N. Y. 8:421. [22] Sharpe, L. M., G. G. Culbreth, and J. R. Klein. 1950. *Proc. Soc. Exp. Biol.*, N. Y. 74:681. [23] Pappenheimer, J. R., E. M. Renkin, and L. M. Borrero. 1951. *Am. J. Physiol.* 167:13. [24] Dunn, J. R., Jr., S. Deavers, R. A. Huggins, and E. L. Smith. 1958. *Ibid.* 195:69.

21. CAPILLARY VASCULARITY OF MUSCLE: VERTEBRATES (Concluded)

Part II. SMOOTH MUSCLE

Animal	Muscle Source	Capillaries/sq mm tissue ¹ (cross-section)	Reference
(A)	(B)	(C)	(D)
1 Dog	Circular layer of stomach	429	1
2	Longitudinal layer of stomach	432	
3	Longitudinal layer of intestine	268	2
4	Circular layer of intestine	356	

/1/ Calculated from total number of capillaries in gut smooth muscle/total volume of smooth muscle, assuming capillary length = 0.3 mm.

Contributors: (a) Fulton, George P., and Herbert J. Berman, (b) Shipley, Reginald A., (c) Burton, Alan C., (d) Lee, Richard E.

References: [1] Mall, F. P. 1889. Johns Hopkins Hosp. Rep. 1:1. [2] Mall, F. P. 1887. *Abh. K. sächs. Ges. Wiss.* 24:151.

Part III: CARDIAC MUSCLE

For information on capillary capacity and surface area, see references 9-16.

Animal	Heart Area and/or Condition	Capillaries		Reference
		No./sq mm tissue ¹ (cross-section)	No./fiber ²	
(A)	(B)	(C)	(D)	(E)
1 Man, 6 mo fetus-16 yr		3744	0.3	1
2 25-77 yr		3343	0.8	
3 18-77 yr	Hypertrophied	2483	0.8	
4 49-71 yr	Atrophied	4613	0.6	
5 Cat	Atrium	4800		2
6	Left ventricle	5718		
7	Right ventricle	5616		
8	Septum	5614		
9	Papillary muscle	4700		
10 Guinea pig		1970		3
11	Untrained	1948		
12	Trained	2819		4
13		3407		5
14 Rabbit	Hypertrophied	2950		
15	Left ventricle	3420	1.1	6
16	Left ventricle, hypertrophied	2670	1.2	
17	Right ventricle	3310	1.0	
18	Right ventricle, hypertrophied	2740	1.2	
19 Snake (<i>Tropidonotus natrix</i>)	Papillary muscle	3230		7
20	Inner	500		
21	Peripheral	1000		8
22 Snake (<i>Caretta caretta</i>)	Inner	555		
23	Peripheral	875		8

/1/ Because of parallel capillary pattern, counts are averages. /2/ Number of capillaries divided by the number of separate groups of muscle substance present in a given cross-section. /3/ High counts may be due to shrinkage of heart tissue upon fixation. [6]

Contributors: (a) Fulton, George P., and Herbert J. Berman, (b) Shipley, Reginald A., (c) Burton, Alan C., (d) Lee, Richard E.

References: [1] Roberts, J. T., and J. T. Wearn. 1941. *Am Heart J.* 21:617. [2] Wearn, J. T. 1928. *J. Exp. Med.* 47:273. [3] Petren, T., and B. Sylvén. 1937. *Morph. Jahrb.* 79:200. [4] Petren, T., T. Sjöstrand, and B. Sylvén. 1936. *Arbeitsphysiologie* 9:376. [5] Rotta, A. 1943. *Rev. argent. card.* 10:186. [6] Shipley, R. A., L. J. Shipley, and J. T. Wearn. 1937. *J. Exp. Med.* 65:29. [7] Stoel, G. 1925. *Zachr. Zellforsch.* 3:91. [8] Ludicke, M. 1939. *Zool. Jahrb. Abt. allg. Zool. Physiol. Tiere.* 59:463. [9] Ponder, E. 1948. *Hemolysis and related phenomena*. Grune and Stratton, New York. [10] Scarborough, R. A. 1930-31. *Yale J. Biol.* 3:63. [11] Dukes, H. H. 1942. *The physiology of domestic animals*. Cornell University Press, Ithaca. [12] Prosser, C. L. 1950. *Comparative animal physiology*. W. B. Saunders, Philadelphia. [13] Krogh, A. 1929. *The anatomy and physiology of capillaries*. Yale University Press, New Haven. [14] Gibson, J. G., et al. 1946. *J. Clin. Invest.* 25:848. [15] Klein, J. R. 1945. *Arch. Biochem.* 8:421. [16] Sharpe, L. M., G. O. Culbreth, and J. R. Klein. 1950. *Proc. Soc. Exp. Biol., N. Y.* 74:681.

22. CHEMICAL COMPOSITION OF THE HEART: VERTEBRATES

Values for water are given in per cent of wet tissue, values for cholesterol and phospholipid in per cent of dry weight. All other values are in mg/100 g wet weight of tissue, unless otherwise indicated. Values in parentheses are ranges, estimate "c" (cf. Introduction).

Part I: MAN

Chemical	Total Heart	Ventricle		Reference
		Left	Right	
(A)	(B)	(C)	(D)	(E)
1 Water, % ¹	79.2	79.2	80.7	B, 1; C, D, 2
2	80.1	80.1(77.2-84.9)	80.0	B, C, 3, D, 4
3	79.7	77.8	80.6	B, 5; C, D, 6
4	78.9	80.8(79.7-81.9)	82.3(80.5-83.8)	B, 7, C, D, 8
5	79.0	79.3	80.3	B, 9; C, D, 10
6		79.0(75.8-80.4)	79.9(73.8-84.7)	11
7		78.9	79.2	12
8		79.0(78.4-79.1)		13
9		78.8(78.1-81.3)		14
10 Aluminum	0.21(0.11-0.29)			15
11 Bromine	0.21(0.065-0.325)			16
12	0			17-19
13 Calcium		6.4	7.3	10
14		7.2(4.7-10.8)	8.4(4.4-15.1)	11
15		6.9	7.6	12
16		8.0(5.0-10.0)		20
17 Chlorine ²	124	120 ³	140	B, 21, C, D, 2
18		139(98-195)	182(136-222)	11
19		120	137	10
20		143(111-166) ⁴		14
21 Chromium	0.01			22
22 Copper	0.77 ⁵			9
23	0.34			23
24	1.34 ⁶			24
25	3.65 ⁷	0.33(0.21-0.49)		B, 25; C, 20
26 Fluorine	0.46 ⁶			26
27 Iron	34.6 ⁶	11.8(4.2-21.3) ⁸		B, 27, C, 3
28 Lead	0.05(0.05-0.08) ⁹			28
29	0			29
30 Magnesium		21.1	18.4	10
31		18.7(10.4-26.1)	20.3(13.4-28.3)	11
32		20.1(15.6-23.2)	19.8(16.4-22.3)	12
33		16.4(8.9-22.4) ⁸		3
34 Manganese	0.046			30
35	0.021			31, 32
36 Phosphorus, total	216(156-300) ⁴	194(180-204)	160(140-187)	B, 14; C, D, 8
37		172	141	10
38		180(132-210)	147(126-172)	11
39		203	177	12
40		161(111-215) ⁸		3
41		183.3		33
42 Potassium ¹⁰	293 ³	314 ³	196	B, 7, C, D, 2
43		290.5 ³	285.5	6
44		285(251-305)	219(189-242) ³	8
45		277	218	10
46		269(184-322)	206(146-240)	11
47		311	255	12
48		325(270-420) ⁴	261 ³	C, 14; D, 4
49	50 ⁶			34
50 Sodium ¹¹	109 ³	90.5 ³	112	B, 7, C, D, 2
51		99(83-132)	126(75-197)	11
52		92	107	12
53		114(84-142) ⁴		14
54 Zinc	10 ¹			24
55	(1.3-1.6) ⁹			35

/1/ Ventricle: apex, 79.4, base, 78.9. Right atrium, 81.2. Septum, 79.2. [2] /2/ Ventricle apex, 120, base, 120. Right atrium, 150. Septum, 120. [2] /3/ Recalculated from mg/100 g wet weight. /4/ Recalculated from mEq/100 g wet weight. /5/ Recalculated from mg/kg dry weight. /6/ mg/100 g dry weight. /7/ Recalculated from mg/kg dry weight. /8/ Recalculated from g of the metallic oxides/100 g dry weight to mg of the element/100 g wet weight. /9/ Recalculated from mg/kg wet weight. /10/ Ventricle apex, 311, base, 316. Right atrium, 139. Septum, 309. [2] /11/ Ventricle apex, 89, base, 92. Right atrium, 120. Septum, 94. [2]

22. CHEMICAL COMPOSITION OF THE HEART: VERTEBRATES (Continued)

Part I. MAN (Concluded)

Chemical	Total Heart	Ventricle		Reference (E)
		Left	Right	
(A)	(B)	(C)	(D)	
56 Cholesterol, total, %		910(600-1220) ³	1580(1220-2000) ³	36
57 Collagen		950(610-1350) ³		37
58				38
59 Creatine	202(117-262)	203(166-218)	165(154-185)	B, I, C, D, 8
60	175(145-250)	208	149	B, 5, C, D, 39
61		220	156	40
62	(170-180) ¹²	211(150-300)	148(100-250)	B, 41; C, D, 42
63		175	173	C, 43, D, 44
64		160(38-295)	114.5(38-230)	C, 45, D, 46
65		(220-285)		47
66 Phospholipid, %			7.00 ¹³	36

/3/ Recalculated from g/100 g wet weight. /12/ Recalculated from mg/g wet weight. /13/ Recalculated.

Contributors (a) Clarke, Norman E., and Robert E. Mosher, (b) Lackey, Robert W., (c) Emslie, A. R. G., (d) Beach, Elliot F., (e) McKibbin, John M.

References [1] Cowan, D. W. 1934. Am. Heart J. 9:378. [2] Clarke, N. E., and R. E. Mosher. 1952. Circulation. N. Y. 5:907. [3] Scott, L. C. 1930. J. Clin. Invest. 9:463. [4] Calhoun, J. A., G. E. Cullen, G. Clarke, and T. R. Harrison. 1930. Ibid. 9:393. [5] Herrman, G., G. Decherd, and T. Oliver. 1936. Am. Heart J. 12:689. [6] Harrison, T. R., C. Pilcher, and G. Ewing. 1930. J. Clin. Invest. 8:325. [7] Norn, M. 1929. Skand. Arch. Physiol., Berl. 55:162. [8] Mangun, G. H., and V. C. Myers. 1940. J. Biol. Chem. 135:411. [9] Chou, T., and W. H. Adolph. 1935. Biochem. J., Lond. 29:476. [10] Cullen, G. E., W. Wilkins, and T. R. Harrison. 1933. J. Biol. Chem. 102:415. [11] Muntwyler, E., R. F. Hamal, G. H. Mangun, and C. T. Way. 1936. Proc. Soc. Exp. Biol., N. Y. 35:555. [12] Wilkins, W., and G. E. Cullen. 1933. J. Clin. Invest. 12:1063. [13] Gross, H. 1940. J. Lab. Clin. M. 25:899. [14] Iseri, L. T., L. C. Alexander, R. S. McCaughy, A. J. Boyle, and G. B. Myers. 1952. Am. Heart J. 43:215. [15] Myers, V. C., and J. W. Mull. 1928. J. Biol. Chem. 78:625. [16] Neufeld, A. H. 1937. Canad. J. Res. 15(B):132. [17] Serbecu, P., and G. A. Buttu. 1934. Bull. Acad. Med., Par. 91:232. Quoted by A. H. Neufeld. 1937. Canad. J. Res. 15(B):132. [18] Lobat, A. 1913. C. rend. Acad. sc. 156:255. Quoted by A. H. Neufeld. 1937. Canad. J. Res. 15(B):132. [19] Damien, A. 1920. Bull. sc. pharm., Par. 27:609. [20] Boyle, A. J., and R. E. Mosher. Unpublished. [21] Magnus-Levy, A. 1910. Biochem. Zschr. 24:363. [22] Grushko, Y. M. 1948. Biokhimiia, Moskva 13:124. [23] White, C. P. 1921. Lancet, Lond. 2:701. [24] Eggleston, W. G. E. 1940. Biochem. J., Lond. 34:991. [25] Herkel, W. 1930. Beitr. path. Anat. 65:531. [26] Zdzarek, E. 1910. Zschr. physiol. Chem. 69:127. [27] Kooyman, J. C. 1949. Acta med. scand. 134:205. [28] Bagchi, K. N., R. D. Ganguly, and J. N. Sirdar. 1939. Ind. J. M. Res. 26:935. [29] Kehoe, R. A., F. Thammann, and J. Cholak. 1933. J. Indust. Hyg. 15:271. [30] Von Oettingen, W. F. 1935. Physiol. Rev. 15:175. [31] Reuman, C. K., and A. S. Minot. 1920. J. Biol. Chem. 42:329. [32] Richards, M. B. 1930. Biochem. J., Lond. 24:152. [33] Decherd, G., and J. E. Blum. 1938. Proc. Soc. Exp. Biol., N. Y. 38:341. [34] Gonnerman, M. 1917. Zschr. physiol. Chem. 99:255. [35] Lutz, R. E. 1924. J. Indust. Hyg. 8:177. [36] Bloor, W. R. 1936. J. Biol. Chem. 114:639. [37] Lowry, O. H., D. R. Gilligan, and E. M. Katersky. 1941. Ibid. 139:795. [38] Blumgart, H. L., D. B. Gilligan, and M. J. Schlesinger. 1940. Tr. Ass. Am. Physicians 55:113. [39] Linegar, J. Biol. Chem. 132:701. [40] Constabel, F. 1921. Biochem. Zschr. 122:152. [41] Seecof, D. P., C. R. Linegar, and V. C. Myers. 1934. Arch. Int. M. 53:574. [42] Herrman, G., G. Decherd, and T. Oliver. 1936. Am. Heart J. 12:689. [43] Volmer, H. Z. 1929. Zschr. ges. exp. Med. 65:522. [44] Bodansky, M., J. F. Pilcher, and V. B. Duff. 1937. Arch. Int. M. 59:232. [45] Bodansky, M., and J. F. Pilcher. 1937. Ibid. 59:232. [46] Bodansky, M. 1931. J. Biol. Chem. 91:147.

Part II. CARNIVORES

Chemical	Total Heart	Ventricle		Atrium		Septum	Reference (H)
		Left	Right	Left	Right		
(A)	(B)	(C)	(D)	(E)	(F)	(G)	
1 Water, %	80.8 ¹	76.9					B, I, C, 2
2	80.4 ¹	77.7 ¹	77.4 ¹				B, 3, 4, C, D, 5
3	78.7 ¹						6
4 Chlorine	145 ²	125 ²					B, 7, C, 2
5	140.93						1
6	133 ²						3, 4
7	121 ²						6

/1/ Recalculated from g/100 g dry weight to per cent of wet tissue. /2/ Recalculated from mM/kg wet weight.

22. CHEMICAL COMPOSITION OF THE HEART: VERTEBRATES (Continued)

Part II: CARNIVORES (Continued)

Chemical (A)	Total Heart (B)	Ventricle		Atrium		Septum (G)	Reference (H)
		Left (C)	Right (D)	Left (E)	Right (F)		
Cat (concluded)							
8 Copper	1.44 ⁴						8
9 Lead	0.15						9
10 Nitrogen	2765 ⁵						3,4
11 Phosphorus, total	207 ³						1
12	206.5 ³						3,4
13 Potassium	301 ³	309.12	310.62				B,1;C,D,5
14	307.4 ³	2402					B,3,4,C,2
15	330 ³						6
16	366 ³						10
17 Sodium	94 ³	942					B,6;C,2
18	110 ³						3,4
19	111.3 ³						1
20 Cholesterol, total, %		0.44(0.34-0.50)					11
21 Collagen	1193 ¹						6
22 Creatine	267						12
	(221-333)						
23 Glycogen	476						13
24 Phospholipid, %		5.72 ⁶					11
Dog							
25 Water, %	80.6	77.4	77.1	74.9	74.6		B,1;C-F,14
26	77.1	78.6	78.9			78.2	B,15,C,D,G,16
27		78.3	78.8			78.26	17
28	80.6 ¹	78.3 ¹					B,3;C,18
29		77.3	77.6				19
30		78.8					20
31		78.5					21
32		78.3					22
33		77.3					23
34 Aluminum	0.184						24
	(0-0.608)						
35	0.11						25
	(0.02-0.17)						
36	0.22						26
37 Bromine	0.19 ⁷						27
38	0.16						28
39	(0.55-0.63)						29
40 Calcium	75 ⁴	18.3	22.7				B,30;C,D,31
41		5.18	5.38	8.28	8.48		14
42		9.3					20
43 Chlorine	115 ³	70.58	70.88	72.78	70.18		B,1;C-F,14
44	119	90.22	94.42			83.82	B,15;C,D,G,16
45	136	85.22	88.42			82.72	B,28;C,D,G,17
46	112 ³	104.8 ³					B,3;C,18
47		1082					21
48		103 ²					22
49 Iron		0.279	0.279	0.199	0.199		14
50 Magnesium		21.6 ⁸	21.1 ⁸	13.98	13.88		14
51		22.7 ¹⁰					18
52 Manganese	0.021						32
53 Nitrogen, total	2852 ⁷	2960 ⁷	2900 ⁷	2480 ⁷	2360 ⁷		B,3;C-F,14
54 Phosphorus	216.5 ³	285	5.82 ⁹	4.64 ⁹	4.57 ⁹		B,1;C,20;D-F,14
55		767 ⁴	708 ⁴				31
56		223	205 ¹¹			216	33
57 Potassium	326.9 ³	2478	2488	1518	1368		B,1;C-F,14
58		345 ²	375 ²			363 ²	16

/1/ Recalculated from g/100 g dry weight to per cent of wet tissue. /2/ Recalculated from mM/kg wet weight.
 /3/ Recalculated from mM/100 g dry weight. /4/ mg/100 g dry weight. /5/ Recalculated from g/100 g dry weight.
 /6/ Recalculated. /7/ Recalculated from g/100 g wet weight. /8/ Recalculated from mEq/kg dry weight.
 /9/ mEq/100 g wet weight. /10/ Recalculated from mEq/100 g dry weight to mg/100 g dry weight. /11/ Total

22. CHEMICAL COMPOSITION OF THE HEART: VERTEBRATES (Continued)

Part II: CARNIVORES (Continued)

Part II. CARNIVORES (continued)							
Chemical	Total Heart	Ventricle		Atrium		Septum	Reference
		Left	Right	Left	Right		
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)
Dog (concluded)							
59 Potassium (concluded)	327.7 ³	333 ² 294 ³ 321 ⁷ 344 301 ² 319 ² 296 ²	363 ² 362 ⁷			354 ²	17 18 19 20 21 22 23
66 Silicon	4 ⁴						34
67 Sodium	87.43	133 ⁸ 71.5 ² 71.1 ² 89.7 ³ 75.4 ² 81.1 ²	133 ⁸ 75.4 ² 75.0 ²	69.9 ⁴	56.4 ⁸	70.2 ² 73.1 ²	B,1,C-F,14 16 17 B,3,C,18 21 22
72 Zinc	2.01					6.4	B,35,G,34
74 Cholesterol, total, %		0.61(0.49-0.90)					11
75 Collagen		415 ¹²	449 ¹² 1380 (1960-1890) ⁷			407 ¹² 1050 (1770-1320) ⁷	33 37
77 Creatine	287 (210-327) 243 (221-257)	289	268			282	B,12,C,D,G,33 38
79 Glycogen		331 603 (386-973) ¹³ 573 (270-570) ⁷ 385 ⁷	799 374-1168 ¹³ 609 (386-737) ⁷	690 (451-1004) ¹³	501 (308-609) ¹³	612 (423-941) ¹³	18 39
81							B,40,C,D,41
82							
83 Phospholipid, %			8.54 ⁶		534 ⁷		B,42,C,G,43 31

/2/ Recalculated from mM/kg wet weight. /3/ Recalculated from mM/100 g dry weight. /4/ mg/100 g dry weight.
/6/ Recalculated. /7/ Recalculated from g/100 g wet weight. /8/ Recalculated from mEq/kg dry weight. /12/ mg
collagen nitrogen/100 g wet tissue. /13/ Recalculated from mg glucose/g wet weight to mg glycogen/100 g wet
weight

Contributors. (a) Clarke, Norman E., and Robert E. Mosher. (b) Lackey, Robert W.. (c) Emalie, A. R. G.
(d) Beach, Eliot F., (e) McKibbin, John M.

References [1] Darrow, D. C., R. R. Harrison, and M. Taffel. 1939. *J. Biol. Chem.* 130:487. [2] Boyer, P. K., and C. A. Poindexter. 1940. *Am. Heart J.* 20:586. [3] Darrow, D. C., and H. C. Miller. 1942. *J. Clin. Invest.* 21:601. [4] Yarnet, H., and D. C. Darrow. 1940. *J. Biol. Chem.* 134:721. [5] Wedd, A. M. 1939. *J. Pharm. Exp. Ther.* 65:268. [6] Robertson, W. V. B., and P. Peyer. 1951. *Am. J. Physiol.* 166:217. [7] Amberson, W. R., T. P. Nash, A. G. Mulder, and D. Binns. 1938. *Ibid.* 122:224. [8] Cunningham, I. J. 1931. *Biochem. J.* 33:401. [9] Crawford, B. L., R. L. Stewart, C. E. Willoughby, and F. L. Smith. 1938. *Am. J. Cancer* 33:401. [10] Zwerner, R. L., and R. Truskowski. 1937. *Endocrinology* 21:40. [11] Bloor, W. R. 1936. *J. Biol. Chem.* 114:639. [12] Folio, O., and T. E. Buckman. 1914. *Ibid.* 17:483. [13] Evans, G., and A. T., and C. H. A. Walton. 1928. *Tr. R. Soc. Canada, Ser. III, Sect. V, 22:1.* [14] Sherrod, T. R. 1947. *Ibid.* 65:49. [15] Cameron, Blumgart, O. H. Lowry, and D. R. Gilligan. 1939. *Tr. Ass. Am. Physiologists* 54:237. [16] Hastings, A. B., H. L. Gilligan, and A. B. Hastings. 1942. *Am. J. Physiol.* 136:474. [17] Lowry, O. H., D. R. Gilligan, and A. B. Hastings. 1942. *Endocrinology* 31:597. [18] Wood, E. H. 1942. *Ibid.* 143:165. [19] Herrman, G., and P. Erhard. 1938. *Proc. Soc. Exp. Biol., N. Y.* 38:35. [20] Muntwyler, E., P. Mautz, and G. Griffin. 1938. *Proc. Soc. Exp. Biol., N. Y.* 38:35. [21] Muntwyler, E., R. C. Mellors, F. R. Mautz, and G. H. Mangun. 1940. *J. Biol. Chem.* 134:367. [22] Wood, E. H. 1942. *Ibid.* 143:165. [23] Clarke, A. P. W., and R. A. Clegghorn. 1942. *Endocrinology* 31:597. [24] Underhill, F. P., and F. L. Petermann. 1929. *Am. J. Physiol.* 90:15. [25] Myers, V. C., and D. B. Morrison. 1928. *J. Biol. Chem.* 78:615. [26] Wührer, J. 1933. *Biochem. Z.* 265:169. [27] Neufeld, A. H. 1936. *Canad. J. Res.* 14(B):160. [28] Damians, A. 1921. *Bull. ac. pharm.* 38:205. [29] Bernhardt, H., and H. Ucko. 1925. *Biochem. Z.* 155:174. [30] Neufeld, A. H. 1936. *Canad. J. Res.* 14(B):160. [31] Reed, C. L., L. M. Dillman, E. A. Thacker, and M. Schöninger. 1930. *Zschr. ges. exp. Med.* 116:423. [32] Bertrand, G., and F. Medigecare. 1913. *Ann. Inst. Pasteur, Par.* 27:1. [33] Albura, H. E., and V. C. Myers. 1939. *J. Biol. Chem.* 131:713. [34] Gonnemann, M. 1917. *Zschr. physiol. Chem.* 99:235. [35] Minard, F. R. 1944. *Tr. Inst. Nac. cienc. Méd., Madr.* 5:239. [36] Rost, E., and A. Weitzel. 1918. *Arch. Reichesgesundh.* 51:494. [37] Lowry, O. H., D. R. Gilligan, and E. M. Katersky. 1941. *J. Biol. Chem.* 139:795.

22. CHEMICAL COMPOSITION OF THE HEART: VERTEBRATES (Continued)

Part II: CARNIVORES (Concluded)

[38] Beker, J. C. 1913. Zschr. physiol. Chem. 87:21. [39] Boyle, R. W., and C. H. McDonald. 1938. Proc. S. Exp. Biol., N. Y. 39:14. [40] Fischer, N. F., and R. W. Lackey. 1925. Am. J. Physiol. 72:43. [41] Cruickshank, E. W. H., and D. L. Shrivastava. 1930. Ibid. 92:144. [42] Jensen, P. 1902. Zschr. physiol. Chem. 35:514. [43] Macleod, J. J. R., and D. J. Prendergast. 1921. Tr. R. Soc. Canada, Ser. III, Sect. V, 15 37.

Part III. RODENTS

Chemical (A)	Total Heart (B)	Ventricle		Reference (E)
		Left (C)	Right (D)	
1 Calcium		Guinea Pig		
2 Copper	6.8			
3 Magnesium	2.12 ¹			
4 Manganese	8.9			1
5 Potassium	0.021			2
6 Sodium	278			1
7 Cholesterol, total, %	133			3
8 Collagen				1
9			0.44(0.31-0.53)	1
10 Glycogen	1050 ²			4
11 Phospholipid, %	241			5
				6
12 Water, %			5.72 ³	7
13		Rabbit		4
14	78.0			
15	79.0(77.9-79.8)			8, 9
16	80.0			10
17	78.0			11
18 Boron		77.1		12
19 Bromine	3.94 ⁴	77.9		13
20 Calcium	0.065		78.0	14
21 Chlorine	67.2 ¹			15
22 Copper	1365			16
23 Iron	2.23 ⁴			17
24 Magnesium	12.4			12
25 Manganese	50.7 ¹			2
26	0.0286			18
27 Nitrogen, total	0.021			17
28 Phosphorus, total	1055(774-1326) ⁷			8
29 Potassium				3
30 Sodium	676 ¹	212(174-238) ⁸		10
31	491 ¹	238 ⁹		10
32 Zinc	122 ⁵			B, 17, C, 13
33	1.066			17
34	2.11 ¹⁰			12
35 Cholesterol, total, %				19
36 Creatine ¹³				20
37	244	0.57(0.40-0.83) ¹¹		4
38	211	0.45(0.36-0.60) ¹²		4
39 Glycogen ¹⁵	223(186-256)	176	170	B, 21, C, D, 14
40	455(350-600) ²	201(158-234)	195(160-228)	B, 22, C, D, 23
41	483(334-640) ²	150(112-225) ¹⁴		B, 24, C, 25
42	108 ²	542		B, 10, C, 26
43	380 ²			27
44 Phospholipid, %	670 ²			28
				29
			9.12 ³	30

/1/ mg/100 g dry weight /2/ Recalculated from g/100 - wet weight
 wet weight
 lated from g,
 weight. /10/
 96.5(76-144) [

g. wet weight. /15/ Right atrium 512 [26] Right atrium

22. CHEMICAL COMPOSITION OF THE HEART: VERTEBRATES (Continued)

Part III. RODENTS (Continued)				
Chemical (A)	Total Heart (B)	Ventricle		Reference (E)
		Left (C)	Right (D)	
Rat				
45 Water, %	77.4			31
46	76.6(76.0-77.2)			32
47	78.1			12
48	77.3			33
49	77.16			34
50	76.916			35
51	77.2			36
52	80.6			37
53	76.5			38
54	77.7			39
55	0.295			16
56 Bromine	70 ¹			40
57 Calcium	8.1(2.7-19.7)			41
58	53.9(51.5-61.7) ¹			42
59 Chlorine	87.517			12
60	96.5			33
61	260 ²			43
62	111			44
63	83.75			39
64 Copper	0.1918			37
65	2.8 ¹			2
66 Iron	7.2(5.0-11.1)			45
67 Magnesium	100 ¹			40
68	22.3(17.6-25.9)			41
69 Manganese	0.05718			38
70 Phosphorus, total	22119			34
71	22219			35
72	940 ¹			40
73	113(87-134)			41
74	881(714-1118) ¹			42
75	255 ⁵			39
76	124			46
77 Potassium	318			31
78	329			12
79	31420			33
80	31319			34
81	33219			35
82	1700 ¹			40
83 Sodium	86.75			31
84	91.85			33
85	90.119			34
86	88.3			35
87	300 ¹			40
88 Cholesterol, total, %		0.53(0.34-0.62)		4
89 Collagen	64021		2960 ¹	B, 39, D, 47
90 Creatine	225			48
91	174(162-195) ¹⁴			49
92	190			50
93	210(190-260) ²			51
94	207			52
95	193(179-218)			53
96	188			54
97 Glycogen	542			55
98	341			56
99	266(190-388)			57
100	582(517-662)			58
101	478			59

11/ mg/100 g dry weight 12/ Recalculated from g/100 g wet weight. 15/ Recalculated from mM/kg wet weight.
 14/ Recalculated from mg/g wet weight. 16/ Recalculated from g/100 g dry weight to per cent of wet tissue.
 17/ Recalculated from mEq/kg of tissue water. 18/ Recalculated from mg/kg dry weight. 19/ Recalculated
 from mM/100 g dry weight. 20/ Recalculated from mEq/kg dry weight. 21/ Recalculated from g/kg wet weight.

Part III: RODENTS (Concluded)

Chemical	Total Heart	Ventricle		Reference
		Left	Right	
(A)	(B)	(C)	(D)	(E)
Rat (concluded)				
102 Glycogen (concluded)	498(391-635)			60
103	659			61
104	520 ²			62
105	451			63
106	442(240-680)			64
107 Phospholipid, %		1.953		4

/2/ Recalculated from g/100 g wet weight. /3/ Recalculated.

Contributors (a) Clarke, Norman E., and Robert E. Mosher, (b) Lackey, Robert W., (c) Emslie, A. R. G., (d) Beach, Elliot F., (e) McKibbin, John M.

References: [1] Kaunetz, H. 1937. *Zschr. ges. exp. Med.* 100:121. [2] Cunningham, I. J. 1931. *Biochem. J.* Lond. 25:1267. [3] Bertrand, G., and F. Medigreane. 1913. *Ann. Inst. Pasteur, Par.* 27:1. [4] Bloor, W. R. 1936. *J. Biol. Chem.* 114:639. [5] Elster, S. K., and E. L. Lowry. 1950. *Proc. Soc. Exp. Biol., N. Y.* 75:127. [6] Elster, S. K. 1950. *J. Biol. Chem.* 186:105. [7] Cacippo, P. 1947. *Boll. Soc. Ital. Biol. Sper.* 23:147. [8] Fore, H., and R. A. Morton. 1952. *Biochem. J.* Lond. 51:600. [9] Bertrand, G., and R. Videsco. 1922. *Bull. Soc. chim. France* 31:268. [10] Wassermeyer, H., and A. Rohrbach. 1932. *Arch. exp. Path., Lpz.* 166:375. [11] Hurwitz, M., and L. Friedberg. 1942. *Arch. Path., Chic.* 34:875. [12] Manery, J. F., and A. B. Hastings. 1939. *J. Biol. Chem.* 127:657. [13] Hagan, P. S. 1939. *J. Pharm. Exp. Ther.* 67:50. [14] Decherd, G., G. Herrman, and E. H. Schwab. 1936. *Proc. Soc. Exp. Biol., N. Y.* 34:864. [15] Kondo, K., S. Mori, and M. Kajima. 1950. *Bull. Inst. Chem. Res., Kyoto Univ.* 21:97. [16] Neufeld, A. H. 1936. *Canad. J. Res.* 14(B):160. [17] Kochmann, M. 1907. *Pflügers Arch.* 119:417. [18] Urbanyi, L. 1943. *Mezo gaz. dasage Kutatasok.* 16:1. [19] Rost, E., and A. Weitzel. 1918. *Arb. Reichsgesundhmt.* 51:494. [20] Leiner, M., and G. Leiner. 1941. *Naturwissenschaften* 29:763. [21] Myers, V. C., and G. H. Mangun. 1940. *J. Biol. Chem.* 132:701. [22] Rose, W. C., O. M. Helmer, and A. Chanutin. 1927. *Ibid.* 75:543. [23] Herrman, G., G. Decherd, E. H. Schwab, P. Erhard. 1935. *Proc. Soc. Exp. Biol., N. Y.* 33:522. [24] Follin, O., and T. E. Buckman. 1914. *J. Biol. Chem.* 17:483. [25] Davies, F., E. T. B. Francis, and H. B. Stoner. 1947. *J. Physiol., Lond.* 106:154. [26] Chang, I. 1938. *Q. J. Exp. Physiol., Lond.* 28:3. [27] Macpherson, W. F., H. E. Essex, and P. C. Mann. 1931. *Am. J. Physiol.* 99:429. [28] Macleod, J. J. R., and D. J. Prendergast. 1921. *Tr. R. Soc. Canada, Ser. III Sect. V.* 15:37. [29] Jensen, P. 1902. *Zschr. physiol. Chem.* 35:514. [30] Genkin, A. M. 1946. *Blokhimia, Moskva* 2:155. [31] Pecora, L. J. 1952. *Am. J. Physiol.* 169:554. [32] Lemley, J. M., and G. R. Menely. 1952. *Ibid.* 169:61. [33] Laramore, D. C., and A. Grollman. 1950. *Ibid.* 161:278. [34] Darrow, D. C. 1944. *Proc. Soc. Exp. Biol., N. Y.* 55:13. [35] Darrow, D. C., and H. C. Miller. 1942. *J. Clin. Invest.* 21:601. [36] Cardini, C. E., and M. E. Scranes. 1942. *Ap. farm. bioquim., B. Air.* 13:102. [37] Lindow, C. W., W. H. Peterson, and H. Steenbock. 1929. *J. Biol. Chem.* 84:419. [38] Skinner, J. T., W. H. Peterson, and H. Steenbock. 1931. *Ibid.* 90:65. [39] Lowry, O. H., A. B. Hastings, T. Z. Hull, and A. N. Brown. 1942. *Ibid.* 143:271. [40] Orent-Kelles, E., and E. V. McCollum. 1941. *Ibid.* 140:337. [41] Boelter, M. D. D., and D. M. Greenberg. 1941. *J. Nutr.* 21:75. [42] Patwardhan, V. N., and R. G. Chitre. 1938. *Ind. J. M. Res.* 25:633. [43] Salvatori, A. 1938. *Atti Accad. naz. Lincei* 27:243. [44] Cameron, A. T., and C. H. A. Walton. 1928. *Tr. R. Soc. Canada, Ser. III, Sect. V.* 22:1. [45] Austoni, M. E., A. Robinovitch, and D. M. Greenberg. 1940. *J. Biol. Chem.* 134:17. [46] Casten, G. G. 1950. *Am. Heart J.* 39:353. [47] Neuman, R. E., and M. A. Logan. 1950. *J. Biol. Chem.* 186:549. [48] Alekseeva, A. M., and I. P. Kok. 1952. *Blokhimia, Moskva* 17:427. [49] Chamutin, A. 1927. *J. Biol. Chem.* 75:549. [50] Bodansky, M., and J. F. Pilcher. 1937. *Proc. Soc. Exp. Biol., N. Y.* 35:607. [51] Beard, H. H., J. K. Espenan, A. L. Koven, and P. Pirrolato. 1941. *Endocrinology* 29:762. [52] Baker, Z., and B. F. Miller. 1940. *J. Biol. Chem.* 132:233. [53] Bodansky, M. 1935. *Ibid.* 109:615. [54] Spoerdama, A. 1951. *Proc. Soc. Exp. Biol., N. Y.* 76:126. [55] Evans, G., and M. A. Bowie. 1936. *Ibid.* 35:68. [56] Long, C. N. H., and G. T. Evans. 1932. *Ibid.* 30:168. [57] Forbes, J. C., and G. M. Duncan. 1950. *Q. J. Alcohol* 11:373. [58] Schürger, G. 1947. *Klin. Wochr.* 24:593. [59] Schumann, H. 1939. *Ibid.* 18:925. [60] Shelley, W. B., C. F. Code, and M. B. Visscher. 1943. *Am. J. Physiol.* 138:652. [61] Moses, L. E. 1944. *Ibid.* 142:886. [62] Lawrence, R. D., and R. A. McCance. 1931. *Biochem. J.* Lond. 25:570. [63] Lackey, R. W., C. A. Bunde, and L. C. Harris. 1947. *Proc. Soc. Exp. Biol., N. Y.* 66:433. [64] Lackey, R. W., C. A. Bunde, A. J. Gill, and L. C. Harris. 1944. *Ibid.* 57:191.

22. CHEMICAL COMPOSITION OF THE HEART: VERTEBRATES (Continued)

Part IV. ARTIODACTYLS AND PERISSODACTYLS

Chemical	Total Heart	Ventricle		Reference
		Left	Right	
(A)	(B)	(C)	(D)	(E)
Cattle				
1 Water, %	77.4 ¹	78.8		B, 1; C, 2
2	65.8			1
3 Bromine	0			3
4 Calcium		3.6	4.3	4
5		1.94 ²		2
6 Copper	1.56 ³			5
7	0.76 ⁴ , 5			6
8 Iron		6.8 ²		2
9 Magnesium ⁶		27.2 ²	26.5 ⁷	C, 2, D, 7
10		16.1	16.1	4
11 Manganese	0.028			8
12 Phosphorus, total	211 ⁴ , 6	239	233	B, 1; C, D, 4
13	158 ⁸	214 ²		B, 1, C, 2
14 Potassium		288	266	4
15		253(247-259) ²		2
16 Sodium		78.5	90	4
17		164(130-186) ²		2
18 Tin	0.15 ⁵			9
19 Collagen		1936 ⁹	3760 ⁹	10
20 Creatine	215(207-220)	305	266	B, 11, C, D, 12
21 Fat	4396 ⁸			1
22	19.45 ¹⁰			1
Horse				
23 Water, %	63.0			13
24 Copper	1.76 ³			5
25 Tin	0.24 ⁵			9
26 Zinc	1.58 ⁵			14
27	7.1			13
28 Glycogen	330 ⁶			15
Pig				
29 Aluminum	0.55			16
30 Copper	1.49 ³			5
31 Manganese	0.020			8, 17
32 Collagen		2180 ⁹	3380 ⁹	10
Sheep				
33 Copper	1.79 ³			5
34 Manganese	0.029			8
35 Tin	0.24			9
36 Creatine	283(208-339)	342	267	B, 18, C, D, 12

1/1 Lean cattle 2/2 Recalculated from g of the metallic oxides/100 g dry weight to mg of the element/100 g wet weight 3/3 Recalculated from mg/kg dry weight to mg/100 g dry weight 4/4 Calf 5/5 Recalculated from mg/kg wet weight 6/6 Right atrium, 19 [7] 7/7 Recalculated from mg/g wet weight 8/8 Recalculated from g/100 g wet weight 9/9 mg/100 g dry weight 10/10 g/100 g wet weight.

Contributors (a) Clarke, Norman E., and Robert E. Mosher, (b) Lackey, Robert W., (c) Emalle, A. R. G., (d) Beach, Elliot F., (e) McKibbin, John M.

References [1] Francis, C. K., and P. F. Trowbridge 1910. J. Biol. Chem. 8:81. [2] Scott, L. C. 1930. J. Clin. Invest. 9:463 [3] Lobat, A. 1913. C. rend. Acad. sc 156 255. [4] Wilkins, W. 1934. Proc. Soc. Exp. Biol. N. Y. 31 1117 [5] Cunningham, I. J. 1931. Biochem. J., Lond. 25 1267. [6] McIlhargue, J. S. 1925. Am. J. Physiol. 72 583 [7] Davies, F., E. T. B. Francis, and H. B. Stoner 1947 J. Physiol., Lond. 106:154. [8] Richards, M. B. 1930. Biochem. J., Lond. 24:1572. [9] Bertrand, G., and V. Ciurea. 1931. C. rend. Acad. sc 192 780. [10] Neuman, R. E., and M. A. Logan. 1950. J. Biol. Chem. 186:549. [11] Beker, J. C. 1913. Zschr. physiol. Chem. 87:21 [12] Secor, D. P., C. R. Linegar, and V. C. Myers. 1934. Arch. Int. M. 53:574. [13] Bertrand, G., and R. Viadeco. 1920. C. rend. Acad. sc 171:744 [14] Rost, E., and A. Weissel. 1918. Arb. Reichgesundhmt. 51:494. [15] Yater, W. M., A. E. Osterberg, and H. W. Hefke. 1930. Arch. Int. M. 45 760 [16] Mackenzie, K. 1930. Biochem. J., Lond. 24:1433. [17] Bertrand, G., and F. Médigreane. 1913. Ann. Inst. Pasteur, Par 27 1 [18] Folin, O., and T. E. Buckman. 1914. J. Biol. Chem. 17 483

Part III: RODENTS (Concluded)

Chemical	Total Heart	Ventricle		Reference
		Left	Right	
(A)	(B)	(C)	(D)	(E)
Rat (concluded)				
102 Glycogen (concluded)	498(391-635)			60
103	659			61
104	520 ²			62
105	451			63
106	442(240-680)			64
107 Phospholipid, %		7.95 ³		4

/2/ Recalculated from g/100 g wet weight. /3/ Recalculated.

Contributors: (a) Clarke, Norman E., and Robert E. Mosher, (b) Lackey, Robert W., (c) Emslie, A. R. G., (d) Beach, Elliot F., (e) McKibbin, John M.

References: [1] Kaunetz, H. 1937. *Zschr. ges. exp. Med.* 100:121. [2] Cunningham, I. J. 1931. *Biochem. J.*, Lond. 25:1267. [3] Bertrand, G., and F. Medigresceane. 1913. *Ann. Inst. Pasteur, Par.* 27:1. [4] Bloor, W. R. 1936. *J. Biol. Chem.* 114:639. [5] Elster, S. K., and E. L. Lowry. 1950. *Proc. Soc. Exp. Biol.*, N. Y. 75:127. [6] Elster, S. K. 1950. *J. Biol. Chem.* 186:105. [7] Cacippo, F. 1947. *Boll. Soc. Ital. Biol. Sper.* 23:147. [8] Fore, H., and R. A. Morton. 1952. *Biochem. J.*, Lond. 51:600. [9] Bertrand, G., and R. Viadescio. 1922. *Bull. Soc. chim. France* 31:268. [10] Wassermeyer, H., and A. Rohrbach. 1932. *Arch. exp. Path., Lpz.* 166:375. [11] Hurwitz, M., and L. Friedberg. 1942. *Arch. Path., Chic.* 34:875. [12] Manery, J. F., and A. B. Hastings. 1939. *J. Biol. Chem.* 127:657. [13] Hagan, P. S. 1939. *J. Pharm. Exp. Ther.* 67:50. [14] Decherd, G. G. Herrman, and E. H. Schwab. 1936. *Proc. Soc. Exp. Biol.*, N. Y. 34:864. [15] Kondo, K., S. Mori, and M. Kajima. 1950. *Bull. Inst. Chem. Res., Kyoto Univ.* 21:97. [16] Neufeld, A. H. 1936. *Canad. J. Res.* 14(B):160. [17] Kochmann, M. 1907. *Pflügers Arch.* 119:417. [18] Urbanyi, L. 1943. *Mezo gaz. dásage Kutatasok.* 16:1. [19] Rost, E., and A. Weltzel. 1918. *Arb. Reichsgesundhamt.* 51:494. [20] Leiner, M., and G. Leiner. 1941. *Naturwissenschaften* 29:763. [21] Myers, V. C., and G. H. Mangun. 1940. *J. Biol. Chem.* 132:701. [22] Rose, W. C., O. M. Helmer, and A. Chamutin. 1927. *Ibid.* 75:543. [23] Herrman, G., G. Decherd, E. H. Schwab, and P. Erhard. 1935. *Proc. Soc. Exp. Biol.*, N. Y. 33:522. [24] Folin, O., and T. E. Buckman. 1914. *J. Biol. Chem.* 17:483. [25] Davies, F., E. T. B. Francis, and H. B. Stoner. 1947. *J. Physiol.*, Lond. 106:154. [26] Chang, L. 1938. *Q. J. Exp. Physiol.*, Lond. 28:3. [27] Macpherson, W. F., H. E. Essex, and F. C. Mann. 1931. *Am. J. Physiol.* 99:429. [28] Macleod, J. J. R., and D. J. Prendergast. 1921. *Tr. R. Soc. Canada, Ser. III Sect. V.* 15:37. [29] Jensen, P. 1902. *Zschr. physiol. Chem.* 35:514. [30] Genkin, A. M. 1946. *Biokhimiya, Moskva* 2:155. [31] Pecora, L. J. 1952. *Am. J. Physiol.* 169:554. [32] Lemley, J. M., and G. R. Meneely. 1952. *Ibid.* 169:61. [33] Laramore, D. C., and A. Grollman. 1950. *Ibid.* 161:278. [34] Darrow, D. C. 1944. *Proc. Soc. Exp. Biol.*, N. Y. 55:13. [35] Darrow, D. C., and H. C. Miller. 1942. *J. Clin. Invest.* 21:601. [36] Cardini, G. G. 1950. *Am. Heart J.* 39:353. [37] Neuman, R. E., and M. A. Logan. 1950. *J. Biol. Chem.* 186:549. [38] Aleksseva, A. M., and I. P. Kok. 1952. *Biokhimiya, Moskva* 17:427. [39] Chamutin, A. 1927. *J. Biol. Chem.* 75:549. [40] Bodansky, M., and J. F. Pilcher. 1937. *Proc. Soc. Exp. Biol.*, N. Y. 35:607. [41] Beard, H. H., J. K. Espenan, A. L. Koven, and P. Pirrolato. 1941. *Endocrinology* 29:762. [42] Baker, Z., and B. F. Miller. 1940. *J. Biol. Chem.* 132:233. [43] Bodansky, M. 1935. *Ibid.* 109:615. [44] Sjoerdama, A. 1951. *Proc. Soc. Exp. Biol.*, N. Y. 76:126. [45] Evans, G., and M. A. Bowie. 1936. *Ibid.* 35:68. [46] Long, C. N. H., and G. T. Evans. 1932. *Ibid.* 30:168. [47] Forbes, J. C., and G. M. Duncan. 1950. *Q. J. Alcohol* 11:373. [48] Schürger, G. 1947. *Klin. Wochr.* 24:593. [49] Schumann, H. 1939. *Ibid.* 18:925. [50] Shelley, W. B., C. F. Code, and M. B. Viesscher. 1943. *Am. J. Physiol.* 138:652. [51] Moses, L. E. 1944. *Ibid.* 142:686. [52] Lawrence, R. D., and R. A. McCance. 1931. *Biochem. J.*, Lond. 25:570. [53] Lackey, R. W., C. A. Bunde, and L. C. Harris. 1947. *Proc. Soc. Exp. Biol.*, N. Y. 66:433. [54] Lackey, R. W., C. A. Bunde, A. J. Gill, and L. C. Harris. 1944. *Ibid.* 57:191.

24. FRACTIONAL EXTRACTION OF DIFFUSIBLE SUBSTANCES FROM CAPILLARIES. MAN, DOG

24. FRACTIONAL EXTRACTION OF DIFFUSIBLE SUBSTANCES FROM CAPILLARIES. Diffusion through the capillary walls is affected by blood flow per amount of tissue, as well as by certain physical attributes of the diffusing molecules. Diffusion across the capillary wall is complete, or nearly complete, within a single circulation for t amipyrine {8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299, 300, 301, 302, 303, 304, 305, 306, 307, 308, 309, 310, 311, 312, 313, 314, 315, 316, 317, 318, 319, 320, 321, 322, 323, 324, 325, 326, 327, 328, 329, 330, 331, 332, 333, 334, 335, 336, 337, 338, 339, 340, 341, 342, 343, 344, 345, 346, 347, 348, 349, 350, 351, 352, 353, 354, 355, 356, 357, 358, 359, 360, 361, 362, 363, 364, 365, 366, 367, 368, 369, 370, 371, 372, 373, 374, 375, 376, 377, 378, 379, 380, 381, 382, 383, 384, 385, 386, 387, 388, 389, 390, 391, 392, 393, 394, 395, 396, 397, 398, 399, 400, 401, 402, 403, 404, 405, 406, 407, 408, 409, 410, 411, 412, 413, 414, 415, 416, 417, 418, 419, 420, 421, 422, 423, 424, 425, 426, 427, 428, 429, 430, 431, 432, 433, 434, 435, 436, 437, 438, 439, 440, 441, 442, 443, 444, 445, 446, 447, 448, 449, 450, 451, 452, 453, 454, 455, 456, 457, 458, 459, 460, 461, 462, 463, 464, 465, 466, 467, 468, 469, 470, 471, 472, 473, 474, 475, 476, 477, 478, 479, 480, 481, 482, 483, 484, 485, 486, 487, 488, 489, 490, 491, 492, 493, 494, 495, 496, 497, 498, 499, 500, 501, 502, 503, 504, 505, 506, 507, 508, 509, 510, 511, 512, 513, 514, 515, 516, 517, 518, 519, 520, 521, 522, 523, 524, 525, 526, 527, 528, 529, 530, 531, 532, 533, 534, 535, 536, 537, 538, 539, 540, 541, 542, 543, 544, 545, 546, 547, 548, 549, 550, 551, 552, 553, 554, 555, 556, 557, 558, 559, 560, 561, 562, 563, 564, 565, 566, 567, 568, 569, 570, 571, 572, 573, 574, 575, 576, 577, 578, 579, 580, 581, 582, 583, 584, 585, 586, 587, 588, 589, 590, 591, 592, 593, 594, 595, 596, 597, 598, 599, 600, 601, 602, 603, 604, 605, 606, 607, 608, 609, 610, 611, 612, 613, 614, 615, 616, 617, 618, 619, 620, 621, 622, 623, 624, 625, 626, 627, 628, 629, 630, 631, 632, 633, 634, 635, 636, 637, 638, 639, 640, 641, 642, 643, 644, 645, 646, 647, 648, 649, 650, 651, 652, 653, 654, 655, 656, 657, 658, 659, 660, 661, 662, 663, 664, 665, 666, 667, 668, 669, 670, 671, 672, 673, 674, 675, 676, 677, 678, 679, 680, 681, 682, 683, 684, 685, 686, 687, 688, 689, 690, 691, 692, 693, 694, 695, 696, 697, 698, 699, 700, 701, 702, 703, 704, 705, 706, 707, 708, 709, 710, 711, 712, 713, 714, 715, 716, 717, 718, 719, 720, 721, 722, 723, 724, 725, 726, 727, 728, 729, 730, 731, 732, 733, 734, 735, 736, 737, 738, 739, 740, 741, 742, 743, 744, 745, 746, 747, 748, 749, 750, 751, 752, 753, 754, 755, 756, 757, 758, 759, 760, 761, 762, 763, 764, 765, 766, 767, 768, 769, 770, 771, 772, 773, 774, 775, 776, 777, 778, 779, 780, 781, 782, 783, 784, 785, 786, 787, 788, 789, 790, 791, 792, 793, 794, 795, 796, 797, 798, 799, 800, 801, 802, 803, 804, 805, 806, 807, 808, 809, 810, 811, 812, 813, 814, 815, 816, 817, 818, 819, 820, 821, 822, 823, 824, 825, 826, 827, 828, 829, 8

Tissue				Blood Flow ml/min/100 g tissue	Reference
(A)	(B)	(C)	(D)	(E)	(F)
			Man		
1 Forearm	H ² OH	19	0.94(0.91-0.97)		1
2	SCN ⁻	78	0.49(0.38-0.56)		
3	H ² OH	19	0.43(0.36-0.54) ^c	4800-10,800 ¹	2
4	SCN ⁻	78	0.67(0.61-0.69) ^c		
			Dog		
5 Gracilis muscle	K ⁴²	42	0.78 ²	1.9	3
6	K ⁴²	42	0.67 ²	4.0	
7	K ⁴²	42	0.50 ²	8.6	
8	K ⁴²	42	0.43 ²	12.9	
9 Head, intact	H ² OH	20	0.89 ²		4
10	Na ²²	22	0.43 ²		
11	SCN ⁻	78	0.44 ²		
12	Creatinine	113	0.44 ²		
13	Inulin	5100	0.16(0.14-0.17)		
14 Heart, intact	K ⁴²	42	0.67(0.56-0.77)	68(75-100)	5
15 Hind leg, intact	H ² OH	19	0.76(0.62-0.77) ³		4
16	H ² OH	20	0.65 ^{2,3}		
17	Na ²²	22	0.36 ²		
18	C ¹³	36	0.49(0.39-0.59)		
19	Urea	60	0.60(0.54-0.69)		
20	SCN ⁻	78	0.54(0.32-0.77)		
21	Xylose	150	0.28(0.25-0.31)		
22	Antipyrine	188	0.62(0.59-0.65)		
23	Fe(CN) ₆	212	0.13(0.11-0.14)		
24	Inulin	5100	0.24(0.08-0.40)		
25 Kidney	H ² OH	19	0.93(0.91-0.96) ^c		6
26	SCN ⁻	78	0.77(0.64-0.85) ^c		
27	PAH ⁴	194	0.95(0.69-0.98) ^c		
28	Inulin	5100	0.79(0.49-0.94) ^c		
29 Liver, intact	H ² OH	19	0.97		4
30	Urea	60	0.90(0.88-0.91)		
31	SCN ⁻	78	0.22 ²		
32	Inulin	5100	0.15 ²		
33 Lung	H ² OH, H ² OH	19, 20	0.56(0.38-0.68)	1500-2400 ¹	7

1/ Total blood flow in lungs 2/ From single determinations. 3/ Deviations from complete equilibration ($L_c = 1.00$) may be due to a small amount of arterio-venous shunting [3, 8] 4/ p-aminobiphenylate.

Contributors (a) Renkin, Eugene M., (b) Freis, Edward D.

References [1] Freis, E. D. T. F. Higgins, and H. J. Morowitz. 1953. *J. Appl. Physiol.* 5:526. [2] Lillienfeld, L. S., E. D. Freis, E. A. Partenotte, and H. J. Morowitz. 1955. *J. Clin. Invest.* 34:1. [3] Renkin, E. M. Unpublished. [4] Chinnard, F. P., C. J. Voelberg, and T. Enns. 1955. *Am. J. Physiol.* 183:221. [5] Conn, H. L., and J. S. Robertson. 1955. *Ibid.* 181:319. [6] Freis, E. D., H. W. Schnaper, C. R. Koe, and L. S. Lillienfeld. 1958. *Circul. Res.*, N Y 6:432. [7] Chinnard, F. P., and T. Enns. 1954. *Am. J. Physiol.* 178:197. [8] Renkin, E. M. 1955. *Ibid.* 183:125. [9] Johnson, J. A., H. M. Cavert, and N. Lifson. 1952. *Ibid.* 171:687. [10] Tobias, C. A., H. B. Jones, J. H. Lawrence, and J. G. Hamilton. 1949. *J. Clin. Invest.* 28:1375. [11] Jones, H. B. 1950. In O. Glasser, ed. *Medical physics*. Year Book Publishers, Chicago. v. 2, p. 855.

Part V: BIRDS

	Chemical (A)	Total Heart (B)	Ventricle		Reference (E)
			Left (C)	Right (D)	
			Duck ¹		
1	Cholesterol, total, %		0.52		1
2	Phospholipid, %		7.80 ²		1
			Hen		
3	Cholesterol, total, %		0.54(0.41-0.70)		1
4	Phospholipid, %		7.56 ²		1
			Owl		
5	Cholesterol, total, %		0.56(0.52-0.64)		1
6	Phospholipid, %		6.16 ²		1
			Pigeon		
7	Magnesium		56(35-96) ³		2
8	Cholesterol, total, %		0.52(0.29-0.59)		1
9	Creatine		235 ³		2
10	Glycogen	155			3
11	Phospholipid, %		7.28 ²		1
			Sparrow		
12	Cholesterol, total, %		0.57		1
13	Phospholipid, %		8.55 ²		1

/1/ Wild. /2/ Recalculated. /3/ Recalculated from mg/g wet weight.

Contributors: (a) Clarke, Norman E., and Robert E. Mosher, (b) Lackey, Robert W., (c) Emslie, A. R. G., (d) Beach, Elliot F., (e) McKibbin, John M.

References [1] Bloor, W. R. 1936. *J. Biol. Chem.* 114:639. [2] Davies, F., E. T. B. Francis, and H. B. Stoner. 1947. *J. Physiol., Lond.* 106:154. [3] Sarzana, G., and G. Cascio. 1949. *Boll. Soc. Ital. biol. sper.* 25:893.

23. CAPILLARY FILTRATION COEFFICIENTS: VERTEBRATES

Kp is the proportionality constant relating the rate of net fluid transport by filtration (+F), or reabsorption (-F), to effective filtration pressure. $F = Kp[(pC - pT) - (xp - xt)]$, where pC is the mean capillary hydrostatic pressure, pT the interstitial fluid pressure, and xp and xt the colloid osmotic pressures of plasma and interstitial fluid, respectively. Values in parentheses are ranges, estimate "b" (cf. Introduction).

	Animal	Tissue	Temp °C			
	(A)	(B)	(C)			
1	Man	Forearm, intact	37	0.0049		1, 2
2	Cat	Brain	37	0.045	0.0014	3
3		Hind leg, isolated	36(34-38)	0.011(0.005-0.017)	0.00034	4, 5
4			10(8-12)	0.006(0.004-0.008)		6
5	Dog	Hind leg, isolated	37	0.014(0.010-0.018)		4
6	Rat	Hindquarters, isolated	37	0.033(0.025-0.041)		7
7	Frog	Mesenteric capillaries	24(22-26)		0.0076	8
8			0(-2 to +2)		0.0026	9

Contributor. Renkin, Eugene M.

References [1] Landis, E. M., and J. H. Gibbon, Jr. 1933. *J. Clin. Invest.* 12:105. [2] Landis, E. M., and J. C. Hortenstine. 1930. *Physiol. Rev.* 30:1. [3] Coulter, N. A., Jr. 1938. *Am. J. Physiol.* 195:459. [4] Pappenheimer, J. R., and A. Soto-Rivera. 1948. *Ibid.* 152:471. [5] Pappenheimer, J. R., E. M. Renkin, and L. M. Borrero. 1951. *Ibid.* 167:13. [6] Pappenheimer, J. R. 1953. *Physiol. Rev.* 33:387. [7] Renkin, E. M., and B. D. Zaun. 1955. *Am. J. Physiol.* 180:498. [8] Landis, E. M. 1927. *Ibid.* 82:217. [9] Brown, E., and E. M. Landis. 1947. *Ibid.* 149:302.

27. PERMEABILITY OF CELL SURFACES: ANIMALS, PLANTS

Values in parentheses are ranges, estimate "c" (cf. Introduction).

Part I. TO WATER

Values are expressed in the number of cubic microns of water that pass through 1 square micron of cell surface, per minute, per atmosphere of difference in osmotic pressure. Unless otherwise indicated, experimental temperatures are 18-22°C.

Species (A)	Cell (B)	Permeability (C)	Reference (D)
Vertebrates			
1 Man	Erythrocyte	3.6	1
2	Leukocyte	1.35	2
3 Cattle (<i>Bos taurus</i>)	Erythrocyte	2.5	3
4 Chick, mouse, rat	Fibroblasts from tissue culture of mouse and rat tumors, and of chick and rat embryos	[0.4-1.0] ¹	4
5 Fish, zebra (<i>Danio rerio</i>)	Ovarian egg	1.3	5
6	Shed egg	0.02	5
7 Frog (<i>Rana temporaria</i>)	Ovarian egg	4	5
8	Body cavity egg	0.06	5
9 Rabbit (<i>Oryctolagus cuniculus</i>)	Erythrocyte	[1.4-4.8]	6
10	Leukocyte	0.29 ²	2
11 Toad (<i>Xenopus laevis</i>)	Body cavity egg	0.07	5
Invertebrates			
12 Amoeba (<i>Amoeba chaos chaos</i>)		0.01	5
13 (<i>A. proteus</i>)		0.01	7
14 Clam (<i>Cumingia tellenoides</i>)	Unfertilized egg	0.49	3
15 Gregarina sp.		[0.20-0.25]	8
16 Oyster (<i>Ostrea virginica</i>)	Unfertilized egg	[0.6-0.7]	9
17 Polymyxa sp.		0.01	10
18 Sand dollar (<i>Dendraster</i> sp.)	Unfertilized egg	[0.1-0.4]	3
19 Sea urchin (<i>Arbacia punctulata</i>)	Fertilized egg	[0.2-0.3]	11-13
20	Unfertilized egg	0.09 ³	3, 14
21 (<i>Anthocidaris crassispina</i>)	Unfertilized egg	0.28[0.27-0.30]	15
22 (<i>Paracentrotus lividus</i>)	Fertilized egg	0.2 ⁴	16
23	Unfertilized egg	0.14	3
24 (<i>Pseudocentrotus depressus</i>)	Unfertilized egg	0.21[0.18-0.24]	15
25 (<i>Strongylocentrotus franciscanus</i>)	Unfertilized egg	[0.1-0.4]	3
26 Starfish (<i>Patiria miniata</i>)	Unfertilized egg	[0.1-0.4]	3
27 (<i>Paraster ochraceus</i>)	Unfertilized egg	[0.1-0.4]	3
28 Water flea (<i>Daphnia longispina</i>)	Ovarian egg	[0.1-0.4]	3
29 Worm (<i>Chaetopterus</i>)	Fertilized egg	0.04[0.03-0.07]	17
30 (<i>Pergamentocoeus</i>)	Unfertilized egg	0.61[0.33-1.20]	18
31 (<i>Urechis caupo</i>)	Unfertilized egg	0.49[0.23-1.09]	18
32 Zootherium sp.		0.27	19
		[0.12-0.25]	3, 9
Plants			
33 Algae (<i>Fucus vesiculosus</i>)		0.16	3, 20
34 (<i>Halimysia osterhoutii</i>)		2.5	21
35 (<i>Nitella flexilis</i>)		466	22
36 (<i>Spirogyra</i> sp.)		16	6
37 Fern (<i>Salvinia auriculata</i>)	Leaf	0.55	3, 23
38 Onion (<i>Allium</i> sp.)	Pulp	0.30	3, 24

1/1 24-28°C. 2/2 Outward flow = 1.16. 3/3 Outward flow = 0.14. 4/4 13-15°C.

Contributors (a) Angelone, Lula, and Clifford A. Angerer. (b) Lucké, Baldwin. (c) Dean, Robert B.

References [1] Jacobs, M. H. 1932. Biol. Bull. 62 178. [2] Shapiro, H., and A. K. Pappert. 1937. J. Cellul. Physiol. 10 147. [3] Lucké, B., H. K. Hartline, and R. A. Ricca. 1939. Ibid. 14 237. [4] Brues, A. M., and C. M. Masters. 1936. Am. J. Cancer 28:324. [5] Prescott, D. M., and E. Zeuthen. 1951. Acta physiol. scand. 28 77. [6] Brooks, S. C., and M. M. Brooks. 1944. The permeability of living cells. Edwards Brothers, Ann Arbor, Mich. [7] Prescott, D. M., and D. Mazia. 1954. Exp. Cell Res. 6:177. [8] Brooks, S. C. 1932. J. Cellul. Physiol. 2 223. [9] Harris, E. J. 1956. Transport and accumulation in biological systems. Academic Press, New York. [10] Kitching, J. A. 1954. Sympos. Soc. Exp. Biol., Camb. 8 63. [11] Lillie, R. S. 1916. Am. J. Physiol. 40 249. [12] Lillie, R. S. 1917. Ibid. 43 43. [13] McCutcheon, M., and B. Lucké. 1932. J. Cellul. Physiol. 2:11. [14] Harvey, E. N. 1932. Biol. Bull. 62 141. [15] Fukuda, T. R. 1936. J. Cellul. Physiol. 7:301. [16] Mazia, C. 1934. Scritti Biol. 9:31, 43. [17] Ring, J. R. 1941. The osmotic pressure and permeability to water of the parthenogenetic eggs of *Daphnia longispina*. Department of Biology, Brown University. [18] Shapiro, H. 1941. J. Cellul. Physiol. 18 143. [19] Leitch, J. L. 1934. Ibid. 4 457. [20] Resnér, B. 1935. Protoplasma, Lpz. 24:531. [21] Jacques, A. G. 1939. J. Gen. Physiol. 22 743. [22] Osterhout, W. J. V. 1950. Ibid. 33:275. [23] Huber, B., and K. Höger. 1930. Die Wasserpermeabilität des Protoplasmas. Gebrüder Borntraeger, Leipzig. [24] Levitt, J., G. W. Scarth, and R. D. Gibbs. 1936. Protoplasma, Lpz. 36:237.

25. TRANSCAPILLARY EXCHANGE OF LARGE MOLECULES: MAN, DOG, GUINEA PIG

Values in parentheses are ranges, estimate "c" (cf. Introduction).

Animal ¹	Substance	Molecular Weight	Transcapillary Exchange Rate fraction in plasma/min	Reference
(A)	(B)	(C)	(D)	(E)
1 Man	Serum albumin ²	69,000	0.00096	1
2			1/3 at 0.0038	2
3			2/3 at 0.00048	
4 Dog	Serum γ -globulin ²	156,000	0.0022(0.0012-0.0031)	3
5	Dextran	5,200 ³	0.033(0.029-0.037)	4
6		20,000 ³	0.0113(0.0097-0.0126)	
7		60,000 ³	0.0016(0.0013-0.0019)	
8		80,000 ³	0.0004(0.0003-0.0006)	
9	Serum albumin ²	69,000	0.00092(0.00035-0.00158)	5
10	Serum γ -globulin ²	156,000	0.00077	6
	Ferric β -globulin	90,000	0.0059(0.0025-0.0110)	7

/1/ Intact. /2/ Measured by use of 131 -labelled protein. /3/ Average molecular weight of fractions, separated by alcohol precipitation.

Contributor: Renkin, Eugene M.

References: [1] Sterling, K. 1951. J. Clin. Invest. 30 1228. [2] Berson, S. A., R. S. Yalow, S. S. Schreiber, and J. Post. 1953. Ibid. 32:746. [3] Myant, N. B. 1952. Clin. Sc., Lond. 11:191. [4] Grotte, G. 1956. Acta chir. scand., Suppl. 211:1. [5] Wasserman, K., and H. S. Mayerson. 1951. Am. J. Physiol. 165 15. [6] Wasserman, K., and H. S. Mayerson. 1952. Cardiologia, Basel 21:296. [7] Flexner, L. B., G. J. Vosburgh, and D. B. Cowie. 1948. Am. J. Physiol. 153:503.

26. CAPILLARY PERMEABILITY COEFFICIENTS OF VARIOUS SUBSTANCES IN TISSUE: CAT

Capillary endothelium is more highly permeable to lipid-soluble substances such as antipyrine than to lipid-insoluble substances. Presumably such substances penetrate endothelial cells as well as capillary pores. The oil:water partition coefficient of antipyrine is about 0.05, while that of the other substances listed in this table is less than 0.0002. Permeability to lipid-insoluble molecules is determined principally by molecular size. Data are means of results obtained by perfusing the substances dissolved in blood through muscle tissue in isolated hindlimbs of cat. Capillary surface area per gram of skeletal muscle is estimated to be 70 sq cm.

Substance	Molecular Weight	Capillary Permeability ¹ per g of tissue x (10 ⁻⁶)	Apparent Diffusion Area ² cm/g of tissue x (10 ³)
(A)	(B)	(C)	(D)
1 Water	18	36 ³	1.13
2 NaCl	58.4	23	1.0
3 Urea	60.1	18	0.94
4 Glucose	180	6.4	0.70
5 Sucrose	342	3.6	0.48
6 Raffinose	594	2.7	0.42
7 Inulin	5100	0.42	0.19
8 Myoglobin	16,800	0.05	0.03
9 Hemoglobin	67,800	0.00	0.00
10 Antipyrine	188	>550	

/1/ Coefficient of permeability (P) as moles diffusing per second for each mole of concentration difference per liter across membrane. /2/ Apparent diffusion area per path length, $A/\Delta x = P/D$, where D = free diffusion coefficient of solute in aqueous solution. Diminution of this quantity ($A/\Delta x$) with increasing molecular weight is attributable to restriction of passage to larger molecules through capillary pores. Estimated mean pore radius in capillaries of hindleg of cat is 30-45 Å. /3/ Obtained by extrapolating from other data to molecular weight of water.

Contributor: Renkin, Eugene M.

References: [1] Pappenheimer, J. R., E. M. Renkin, and L. M. Borrero. 1951. Am. J. Physiol. 167:13. [2] Pappenheimer, J. R. 1953. Physiol. Rev. 33:387. [3] Renkin, E. M. 1952. Am. J. Physiol. 168 538

27. PERMEABILITY OF CELL SURFACES. ANIMALS, PLANTS (Continued)

Part II: TO INORGANIC IONS (Concluded)

Part II: TO INORGANIC IONS (Concluded)							
Ion	Species	Cell	Permeability			Reference	
			Flow	Rate	Unit of Measurement		
(A)	(B)	(C)	(D)	(E)	(F)	(G)	
Invertebrates (concluded)							
56	Bromide	Squid (<i>Loligo pealei</i>)	Nerve	1	4.5	cm/hr x 10 ⁻⁶	13
57	Chloride	Cuttlefish (<i>Sepia officinalis</i>)	Nerve	1	0.6	M/sq cm/impulse x 10 ⁻¹²	18
Plants							
58	Sodium	Algae (<i>Nitella coronata</i>)		1	28.8	M/sq cm/min x 10 ⁻⁶	21
59		(<i>Valonia macrophyssa</i>)		1	0.09		
60		(<i>V. ventricosa</i>)		1	0.01		22
61		Waterplant (<i>Elodea</i> sp.)		1	28.8	M/sq cm/min x 10 ⁻⁶	21
62	Potassium	Algae (<i>Nitella clavata</i>)		1	0.01	M/sq cm/min x 10 ⁻⁶	21
63		(<i>N. coronata</i>)		1	157		
64		(<i>N. flexilis</i>)		1	0.01		
65		(<i>Valonia macrophyssa</i>)		1	0.44		
66		(<i>V. ventricosa</i>)		1	0.05		22
67		Waterplant (<i>Elodea</i> sp.)		1	157	M/sq cm/min x 10 ⁻⁶	21
68	Rubidium	Algae (<i>Nitella coronata</i>)		1	510	M/sq cm/min x 10 ⁻⁶	21
69		(<i>Valonia ventricosa</i>)		1	1.1		23
70		Waterplant (<i>Elodea</i> sp.)		1	51	cm/hr x 10 ⁻⁶	21
71	Bromide	Algae (<i>Nitella coronata</i>)		1	0.0002	M/sq cm/min x 10 ⁻⁶	21
72		Waterplant (<i>Elodea</i> sp.)		1	4.3	cm/hr x 10 ⁻⁶	21
73	Chloride	Algae (<i>Nitella coronata</i>)		1	0.0072	M/sq cm/min x 10 ⁻⁶	21
74		(<i>Valonia ventricosa</i>)		1	0.06		22
75	Phosphate	Algae (<i>Nitella coronata</i>)		1	25	M/sq cm/min x 10 ⁻⁶	21
76		(<i>N. flexilis</i>)		1	(1-6)	M/sq cm/sec x 10 ⁻¹²	8
77	Iodide	Algae (<i>Valonia macrophyssa</i>)		1	0.0018	M/sq cm/min x 10 ⁻⁶	21

Contributors: Angelone, Luis, and Clifford A. Angerer

References. [1] Harris, E. J., and M. Matzels. 1951. *J. Physiol.*, Lond. 113:506. [2] Harris, E. J., and O. P. Burn. 1949. *Tr. Faraday Soc.* 45:508. [3] Dainty, J., and K. Krnjevic. 1955. *J. Physiol.*, Lond. 128:489. [4] Matzels, M. 1954. *Sympos. Soc. Exp. Biol.*, Cambr. 8:202. [5] Krogh, A. 1946. *Proc. R. Soc.*, Lond. B133:140. [6] Keynes, R. D. 1954. *Ibid.* B142:359. [7] Hodgkin, A. L. 1951. *Biol. Rev. Cambridge Philos. Soc.* 26:339. [8] Harris, E. J. 1956. *Transport and Accumulation in biological systems*. Academic Press, New York. [9] Streeter, D. H. P., and A. K. Solomon. 1954. *J. Gen. Physiol.* 37:643. [10] Mullins, L. J., W. O. Fenn, T. R. Noonan, and L. Haege. 1941. *Am. J. Physiol.* 135:93. [11] Frazier, H. S., A. Sicular, and A. K. Solomon. 1954. *J. Gen. Physiol.* 37:631. [12] Hodgkin, A. L., and R. D. Keynes. 1954. *Sympos. Soc. Exp. Biol.*, Cambr. 8:423. [13] Rothenberg, M. A. 1950. *Biochim. biophys. acta*, Amst. 4:96. [14] Keynes, R. D. 1951. *J. Physiol.*, Lond. 114:119. [15] Lelich, J. L. 1934. *J. Cellul. Physiol.* 4:457. [16] Keynes, R. D., and P. R. Lewis. 1951. *J. Physiol.*, Lond. 113:73. [17] Keynes, R. D. 1951. *Ibid.* 113:99. [18] Keynes, R. D., and P. R. Lewis. 1951. *J. Physiol.*, Lond. 114:151. [19] Brooks, S. C. 1939. *Proc. Soc. Exp. Biol.*, N. Y. 42:557. [20] Dawson, H., and J. F. Danielli. 1943. *The permeability of natural membranes*. Cambridge University Press, London. [21] Brooks, S. C., and M. K. Brooks. 1944. *The permeability of living cells*. Edwards Brothers, Ann Arbor, Mich. [22] Cooper, W. C., Jr., M. J. Dorcas, and W. J. V. Osterhout. 1929. *J. Gen. Physiol.* 12:427. [23] Brooks, S. C. 1932. *J. Cellul. Physiol.* 2:223.

PERMEABILITY OF CELL SURFACES: ANIMALS, PLANTS (Continued)

Part II: TO INORGANIC IONS

Experimental temperatures range from 36-38°C for mammalian cells, and from 15-25°C for all other
Flow: I = inward, O = outward.

Ion	Species	Cell	Permeability		
			Flow	Rate	Unit of Measurement
(A)	(B)	(C)	(D)	(E)	(F)
1 Sodium	Man	Erythrocyte	I	(0.6-1.6)	cm/hr x 10 ⁻⁶
2			O	(6-16)	
3	Cat (<i>Felis catus</i>)	Erythrocyte	O	3.9	cm/hr x 10 ⁻⁶
4		Nerve, desheathed popliteal			M/sq cm/sec x 10 ⁻¹²
5	Chicken (<i>Gallus domesticus</i>)	Erythrocyte	I	2	cm/hr x 10 ⁻⁶
6			O	49	
7	Dog (<i>Canis familiaris</i>)	Erythrocyte	I	14	cm/hr x 10 ⁻⁶
8			O	10.2	
9	Frog (<i>Rana esculenta</i>)	Musculus extensor longus digitorum IV	I	10.9	M/sq cm/sec x 10 ⁻¹²
10	(<i>R. temporaria</i>)	Erythrocyte	I	2	cm/hr x 10 ⁻⁶
11		Sartorius muscle	I	13	M/sq cm/sec x 10 ⁻¹²
12			O	16	
13	Rabbit (<i>Oryctolagus sp.</i>)	Erythrocyte	I	100 (in vivo)	cm/hr x 10 ⁻⁶
14			O	26	
15	Rat (<i>Rattus rattus</i>)	Diaphragm	I	1.7	M/sq cm/sec x 10 ⁻¹²
16	Man	Erythrocyte	O	1.6	mM/sq cm/hr x 10 ⁻⁷
17			I	1.5	mM/sq cm/hr x 10 ⁻⁷
18	Cat (<i>Felis catus</i>)	Erythrocyte	I	0.08	mM/sq cm/hr x 10 ⁻⁷
19	Dog (<i>Canis familiaris</i>)	Erythrocyte	I	6	M/sq cm/sec x 10 ⁻¹²
20	Frog (<i>Rana esculenta</i>)	Musculus adductor longus digitorum IV	I	8	
21		Musculus extensor longus digitorum IV	I	4(3-5)	
22	(<i>R. pipiens</i>)	Erythrocyte	O	4.9(4-6)	
23	(<i>R. temporaria</i>)	Sartorius muscle	I	5.3	mM/sq cm/hr x 10 ⁻⁷
24			I	5	M/sq cm/sec x 10 ⁻¹²
25	Guinea pig (<i>Cavia porcellus</i>)	Erythrocyte	O	7	
26			I	3.2	mM/sq cm/hr x 10 ⁻⁷
27	Rabbit (<i>Oryctolagus sp.</i>)	Erythrocyte	I	2.1	mM/sq cm/hr x 10 ⁻⁷
28		Gastrocnemius muscle	I	130	cm/hr x 10 ⁻⁶
29	Rat (<i>Rattus rattus</i>)	Erythrocyte	I	5.4	M/sq cm/sec x 10 ⁻¹²
30		Diaphragm	I	38.5	M/sq cm/sec x 10 ⁻¹²
31		Gastrocnemius muscle	I	700	cm/hr x 10 ⁻⁶
32	Chloride	Frog (<i>Rana esculenta</i>)	I	11	M/sq cm/sec x 10 ⁻¹²
33	Phosphate	Frog (<i>R. temporaria</i>)	I	(15-21)	cm/hr x 10 ⁻⁶
34			O	35	
35	Sodium	Cuttlefish (<i>Sepia officinalis</i>)	I	40	M/sq cm/sec x 10 ⁻¹²
36		Nerve	I	10	
37			O	6	M/sq cm/impulse x 10 ⁻¹²
38	Squid (<i>Loligo pealei</i>)	Nerve	I	4.5	M/sq cm/impulse x 10 ⁻¹²
39	(<i>L. forbesi</i>)	Nerve	I	61(16-153)	M/sq cm/sec x 10 ⁻¹²
40			I	10(5-15)	M/sq cm/impulse x 10 ⁻¹²
41	Worm (<i>Urechis caupo</i>)	Egg	O	6.6(4.5-9.6)	
42			I	600	M/sq cm/min x 10 ⁻⁸
43	Potassium	Crab (<i>Carcinus maenas</i>)	I	19	M/sq cm/sec x 10 ⁻¹²
44		Nerve	O	22(16-33)	
45			I	0.5	M/sq cm/impulse x 10 ⁻¹²
46			O	2.9(2.7-3.6)	
47	Cuttlefish (<i>Sepia officinalis</i>)	Nerve	I	21	M/sq cm/sec x 10 ⁻¹²
48			O	28	
49			I	0.39	M/sq cm/impulse x 10 ⁻¹²
50	Squid (<i>Loligo pealei</i>)	Nerve	O	4.7	
51	(<i>L. forbesi</i>)	Nerve	I	1250	cm/hr x 10 ⁻⁶
52	Starfish (<i>Pisaster ochraceus</i>)	Egg	O	3.0(2.6-3.4)	M/sq cm/impulse x 10 ⁻¹²
53			I	600	M/sq cm/min x 10 ⁻⁸
54	Rubidium	Squid (<i>Loligo pealei</i>)	I	51	cm/hr x 10 ⁻⁶
55		Starfish (<i>Patiria miniata</i>)	I	600	M/sq cm/min x 10 ⁻⁸

27. PERMEABILITY OF CELL SURFACES: ANIMALS, PLANTS (Continued)

PART III. TO ORGANIC SUBSTANCES (Continued)

Substance (A)	Species (B)	Permeability (C)	Reference (D)
	Plants		
	Algae (<i>Ceramium diaphanum</i>)	49.8	3
	Algae (<i>Chara ceratophylla</i>)	71.6	9
	Algae (<i>Melosira</i> sp)	22.5	3
	Algae (<i>Pylaiella littoralis</i>)	5.8	3
	Algae (<i>Spirogyra</i> sp)	6.6	3
	Bacteria (<i>Beggiatoa mirabilis</i>)	83.4	3
	Moss (<i>Plagiothecium denticulatum</i>)	2	3
	Oyster plant (<i>Rhodo discolor</i>)	10.4	3
	Turmeric (<i>Curcuma rubricaulis</i>)	6.6	9
	Algae (<i>Chara ceratophylla</i>)	145	3
	Algae (<i>Ceramium diaphanum</i>)	1.3	9
	Algae (<i>Chara ceratophylla</i>)	1.2	3
	Algae (<i>Melosira</i> sp)	2	3
	Algae (<i>Pylaiella littoralis</i>)	0.1	3
	Algae (<i>Spirogyra</i> sp)	0.2	3
	Bacteria (<i>Bacterium paracoli</i>)	3.3	3
	Bacteria (<i>Beggiatoa mirabilis</i>)	63.6	3
	<i>Malanthemum</i> (<i>Malanthemum</i> sp)	9.8 ¹	10
	Moss (<i>Plagiothecium denticulatum</i>)	0.02	3
	Oyster plant (<i>Rhodo discolor</i>)	0.7	3
	Turmeric (<i>Curcuma rubricaulis</i>)	0.1	3
	Algae (<i>Ceramium diaphanum</i>)	0.1	3
	Algae (<i>Chara ceratophylla</i>)	0.08	9
	Algae (<i>Melosira</i> sp)	0.8	3
	Algae (<i>Pylaiella littoralis</i>)	0.007	3
	Algae (<i>Spirogyra</i> sp)	0.02	3
	Bacteria (<i>Bacterium paracoli</i>)	0.3	3
	Bacteria (<i>Beggiatoa mirabilis</i>)	50.3	3
	Moss (<i>Plagiothecium denticulatum</i>)	0.004	3
	Oyster plant (<i>Rhodo discolor</i>)	0.03	3
	Turmeric (<i>Curcuma rubricaulis</i>)	0.02	3
	Algae (<i>Chara ceratophylla</i>)	<0.05	9
	Algae (<i>C. ceratophylla</i>)	26.6	9
	Algae (<i>C. ceratophylla</i>)	133	9
	Algae (<i>C. ceratophylla</i>)	90	3
	Algae (<i>C. ceratophylla</i>)	28.2	3
	Algae (<i>Melosira</i> sp)	16.8	3
	Algae (<i>Pylaiella littoralis</i>)	4	3
	Moss (<i>Plagiothecium denticulatum</i>)	4.8	3
	Turmeric (<i>Curcuma rubricaulis</i>)	1.5	3
	Algae (<i>Ceramium diaphanum</i>)	0.2	3
	Algae (<i>Chara ceratophylla</i>)	1.1	3
	Algae (<i>Melosira</i> sp)	0.1	3
	Algae (<i>Pylaiella littoralis</i>)	0.1	3
	Algae (<i>Spirogyra</i> sp)	1.7	3
	Bacteria (<i>Bacterium paracoli</i>)	94.8	3
	Bacteria (<i>Beggiatoa mirabilis</i>)	5.5 ¹	10
	<i>Malanthemum</i> (<i>Malanthemum</i> sp)	0.05	3
	Moss (<i>Plagiothecium denticulatum</i>)	0.01	3
	Oyster plant (<i>Rhodo discolor</i>)	0.02	3
	Turmeric (<i>Curcuma rubricaulis</i>)	168	3
	Algae (<i>Ceramium diaphanum</i>)	216	3
	Algae (<i>Chara ceratophylla</i>)	180	3
	Algae (<i>Melosira</i> sp)	84	3
	Algae (<i>Pylaiella littoralis</i>)	27	3
	Algae (<i>Spirogyra</i> sp)	12	3
	Moss (<i>Plagiothecium denticulatum</i>)	9.6	3
	Oyster plant (<i>Rhodo discolor</i>)	13.2	3
	Turmeric (<i>Curcuma rubricaulis</i>)	0.05	3
	Algae (<i>Chara ceratophylla</i>)	0.4	3
	Algae (<i>Melosira</i> sp)	0.0004	3
	Algae (<i>Pylaiella littoralis</i>)	8.1	3
	Bacteria (<i>Beggiatoa mirabilis</i>)	0.0005	3
	Moss (<i>Plagiothecium denticulatum</i>)	0.0005	3
	Turmeric (<i>Curcuma rubricaulis</i>)	0.0005	3

/3/ Value given in moles per minute x 10⁻⁴.

27. PERMEABILITY OF CELL SURFACES: ANIMALS, PLANTS (Continued)

Part III: TO ORGANIC SUBSTANCES

Values are expressed in moles per square centimeter of surface area per minute per gram mole difference in concentration $\times 10^{-8}$, unless otherwise indicated. Temperatures range from 20-25°C.

Substance (A)	Species (B)	Permeability (C)	Reference (D)
Vertebrates ¹			
1 Ethylene glycol	Ground hog	246	1
2	Ox	9.6	2
3 Diethylene glycol	Ox	4.5	1
4 Triethylene glycol	Ox	2	1
5 Trimethylene glycol	Ox	6	1
6 Propylene glycol	Ox	19.8	1
7 Glycerol	Man	9	1
8	Ground hog	163	1
9	Ox	0.11	2
10 Erythritol	Ground hog	31.7	1
11	Ox	<0.003	1
12 Mannitol	Man	<0.33	1
13	Ground hog	2	1
14	Ox	<0.003	1
15 Propyl alcohol	Ox	636	3
16 Monoacetin	Ox	6.8	1
17 Diacetin	Ox	62.5	1
18 Acetamide	Ox	283	4
19 Butyramide	Ox	163	4
20 Formamide	Ox	293	4
21 Propionamide	Ox	114	4
22 Sucrose	Man	0	1
23 Xylose	Ground hog	0.66	1
24	Ox	<0.003	1
25 Urea	Man	1166	1
26	Ground hog	1666	1
27	Ox	1080	2
28 Thiourea	Ground hog	138	1
29	Ox	1.2	1
Invertebrates ²			
30 Ethylene glycol	Ciam (<i>Cumingia tellenoides</i>)	156(151-163)	5
31	Marine worm (<i>Chaetopterus pergamentaceus</i>)	143(140-153)	5
32	Sea urchin (<i>Arbacia punctulata</i>)	36(29-56)	5-7
33	Starfish (<i>Asterias</i> sp)	102(50-182)	7
34	Worm (<i>Phascolosoma gouldi</i>)	7.3	4
35	Protozoa (<i>Gregarina</i> sp)	40.2	3
36 Diethylene glycol	Sea urchin (<i>Arbacia punctulata</i>)	26	8
37	Worm (<i>Phascolosoma gouldi</i>)	3.7	4
38 Triethylene glycol	Worm (<i>P. gouldi</i>)	1.2	4
39 Trimethylene glycol	Sea urchin (<i>Arbacia punctulata</i>)	43	1
40	Worm (<i>Phascolosoma gouldi</i>)	7.3	4
41 Tetraethylene glycol	Worm (<i>P. gouldi</i>)	0.4	4
42 Propylene glycol	Sea urchin (<i>Arbacia punctulata</i>)	77	8
43	Worm (<i>Phascolosoma gouldi</i>)	11.7	4
44	Protozoa (<i>Gregarina</i> sp)	79.2	3
45 Glycerol	Marine worm (<i>Chaetopterus pergamentaceus</i>)	62(58-69)	5
46	Sea urchin (<i>Arbacia punctulata</i>)	5	1
47	Worm (<i>Phascolosoma gouldi</i>)	<0.004	4
48	Protozoa (<i>Gregarina</i> sp)	1.1	3
49 Monoacetin	Worm (<i>Phascolosoma gouldi</i>)	0.6	4
50 Diacetin	Worm (<i>P. gouldi</i>)	4.5	4
51 Acetamide	Sea urchin (<i>Arbacia punctulata</i>)	(55-60)	6
52	Worm (<i>Phascolosoma gouldi</i>)	15.8	4
53 Butyramide	Sea urchin (<i>Arbacia punctulata</i>)	(338-394)	6
54	Worm (<i>Phascolosoma gouldi</i>)	410	4
55 Formamide	Worm (<i>P. gouldi</i>)	9.8	4
56 Malonamide	Protozoa (<i>Gregarina</i> sp)	15	3
57 Propionamide	Sea urchin (<i>Arbacia punctulata</i>)	(134-150)	6
58	Worm (<i>Phascolosoma gouldi</i>)	62.3	4
59 Urea	Worm (<i>P. gouldi</i>)	1.6	4
60 Thiourea	Worm (<i>P. gouldi</i>)	2	4

/1/ Erythrocytes. /2/ Unfertilized eggs, except values for worm (*Phascolosoma gouldi*) which are for permeability of erythrocyte surface.

27. PERMEABILITY OF CELL SURFACES. ANIMALS, PLANTS (Continued)
 Part III: TO ORGANIC SUBSTANCES (Continued)

Substance (A)	Species (B)	Permeability (C)	Reference (D)
Ethylene glycol	Plants	49.8	3
	Algae (Ceranium diaphanum)	71.6	9
	Algae (Chara ceratophylla)	22.5	3
	Algae (Melosira sp)	5.8	3
	Algae (Pyralia littoralis)	6.6	3
	Algae (Spirogyra sp)	83.4	3
	Bacteria (Beggiatoa mirabilis)	2	3
	Moss (Plagiothecium denticulatum)	10.4	3
	Oyster plant (Rhoeo discolor)	6.6	9
	Turnerica (Curcuma rubricaulis)	145	3
Propylene glycol	Algae (Chara ceratophylla)	1.3	3
	Algae (Ceranium diaphanum)	1.2	9
Glycerol	Algae (Chara ceratophylla)	2	3
	Algae (Melosira sp)	0.1	3
	Algae (Pyralia littoralis)	0.2	3
	Algae (Spirogyra sp)	3.3	3
	Bacteria (Bacterium paracoli)	61.6	3
	Bacteria (Beggiatoa mirabilis)	9.8	10
	Maianthemum (Maianthemum sp)	0.02	3
	Moss (Plagiothecium denticulatum)	0.7	3
	Oyster plant (Rhoeo discolor)	0.1	3
	Turnerica (Curcuma rubricaulis)	0.1	3
Erythritol	Algae (Ceranium diaphanum)	0.08	9
	Algae (Chara ceratophylla)	0.8	3
	Algae (Melosira sp)	0.007	3
	Algae (Pyralia littoralis)	0.02	3
	Algae (Spirogyra sp)	0.3	3
	Bacteria (Bacterium paracoli)	50.3	3
	Bacteria (Beggiatoa mirabilis)	0.004	3
	Moss (Plagiothecium denticulatum)	0.03	3
	Oyster plant (Rhoeo discolor)	0.02	3
	Turnerica (Curcuma rubricaulis)	<0.05	9
Mannitol	Algae (Chara ceratophylla)	26.6	9
	Algae (C. ceratophylla)	133	9
	Algae (C. ceratophylla)	90	3
	Algae (C. ceratophylla)	28.2	3
	Algae (Melosira sp)	16.8	3
	Algae (Pyralia littoralis)	4	3
	Moss (Plagiothecium denticulatum)	4.8	3
	Turnerica (Curcuma rubricaulis)	1.5	3
	Algae (Ceranium diaphanum)	0.2	3
	Algae (Chara ceratophylla)	1.1	3
Malonamide	Algae (Melosira sp)	0.1	3
	Algae (Pyralia littoralis)	0.1	3
	Algae (Spirogyra sp)	1.7	3
	Bacteria (Bacterium paracoli)	94.8	3
	Bacteria (Beggiatoa mirabilis)	5.5	10
	Maianthemum (Maianthemum sp)	0.05	3
	Moss (Plagiothecium denticulatum)	0.01	3
	Oyster plant (Rhoeo discolor)	0.02	3
	Turnerica (Curcuma rubricaulis)	168	3
	Algae (Ceranium diaphanum)	216	3
Propionamide	Algae (Chara ceratophylla)	180	3
	Algae (Melosira sp)	84	3
	Algae (Pyralia littoralis)	27	3
	Algae (Spirogyra sp)	12	3
	Moss (Plagiothecium denticulatum)	9.6	3
	Oyster plant (Rhoeo discolor)	13.2	3
	Turnerica (Curcuma rubricaulis)	0.05	3
	Algae (Chara ceratophylla)	0.4	3
	Algae (Melosira sp)	0.0004	3
	Bacteria (Beggiatoa mirabilis)	8.1	3
Saccharose	Moss (Plagiothecium denticulatum)	0.0005	3
	Turnerica (Curcuma rubricaulis)	0.0005	3

/3/ Value given in moles per minute x 10⁻⁴.

27. PERMEABILITY OF CELL SURFACES: ANIMALS, PLANTS (Concluded)

PART III: TO ORGANIC SUBSTANCES (Concluded)

Substance (A)	Species (B)	Permeability (C)	Reference (D)
Plants (concluded)			
125 Urea	Algae (<i>Ceramium diaphanum</i>)	5	3
126	Algae (<i>Chara ceratophylla</i>)	6.7	9
127	Algae (<i>Melosira</i> sp)	2.5	3
128	Algae (<i>Pylaiella littoralis</i>)	0.5	3
129	Algae (<i>Spirogyra</i> sp)	1.1	3
130	Bacteria (<i>Bacterium paracoli</i>)	5	3
131	Bacteria (<i>Beggiatoa mirabilis</i>)	70.2	3
132	Maianthemum (<i>Maianthemum</i> sp)	(23.4-33.3) ³	10
133	Moss (<i>Plagiothecium denticulatum</i>)	0.2	3
134	Oyster plant (<i>Rhoeo discolor</i>)	0.2	3
135	Turmeric (<i>Curcuma rubricaulis</i>)	0.09	3
136 Methyl urea	Algae (<i>Ceramium diaphanum</i>)	5.8	3
137	Algae (<i>Chara ceratophylla</i>)	11.4	3
138	Algae (<i>Melosira</i> sp)	7.2	3
139	Algae (<i>Pylaiella littoralis</i>)	1.9	3
140	Algae (<i>Spirogyra</i> sp)	1.7	3
141	Maianthemum (<i>Maianthemum</i> sp)	75 ³	10
142	Moss (<i>Plagiothecium denticulatum</i>)	0.7	3
143	Oyster plant (<i>Rhoeo discolor</i>)	0.5	3
144	Turmeric (<i>Curcuma rubricaulis</i>)	0.5	3
145 Thiourea	Algae (<i>Chara ceratophylla</i>)	12.8	9
146	Maianthemum (<i>Maianthemum</i> sp)	90 ³	10
147	Oyster plant (<i>Rhoeo discolor</i>)	21 ³	10

/3/ Value given in moles per minute $\times 10^{-4}$.

Contributors: Angerer, Clifford A., and Luis Angelone

References: [1] Jacobs, M. H. 1952. In E. S. G. Barron, ed. *Modern trends in physiology and biochemistry*. Academic Press, New York. [2] Jacobs, M. H. 1934. *J. Cellul. Physiol.* 4:161. [3] Davson, H., and J. F. Danielli. 1952. *The permeability of natural membranes*. Cambridge University Press, London. [4] Love, W. E. 1953. *Biol. Bull.* 105:128. [5] Lucké, B., H. K. Hartline, and R. A. Ricca. 1939. *J. Cellul. Physiol.* 14:237. [6] Jacobs, M. H., and D. R. Stewart. 1932. *Ibid.* 1:71. [7] Stewart, D. R., and M. H. Jacobs. 1932. *Ibid.* 1:83. [8] Stewart, D. R., and M. H. Jacobs. 1936. *Ibid.* 7:333. [9] Collander, R., and H. Bärlund. 1934. *Acta bot. fenn.* 11:1. [10] Brooks, S. C., and M. M. Brooks. 1944. *The permeability of living cells*. Edwards Brothers, Ann Arbor, Mich.

28. CARDIOVASCULAR AND PULMONARY REFLEXES. MAMMALS

28. CARDIOVASCULAR AND PULMONARY REFLEXES						
Receptors		Afferent Nerves (C)	Effective Stimulus (D)	Response		
Type (A)	Location (B)			Cardiovascular (E)	Pulmonary (F)	
Blood Vessel and Heart Reflexes						
1	Systemic arterial pressure receptors	Carotid sinns	Branch of glossopharyngeal	Increase of transmural pressure.	Fall in blood pressure; decrease in heart rate and cardiac output, peripheral vasodilatation due to decrease in vasoconstrictor tone.	Slowing or arrest of breathing; bronchoconstriction.
2		Common carotid	Branches of vagi and aortic nerves	Increase of transmural pressure.		
3		Aortic arch	Aortic depressor	Increase of transmural pressure.		
4		Mesenteric arteries (Pacinian corpuscles)	Splanchnic	Increase of transmural pressure.	Uncertain.	
5		Thoracic aorta	Dorsal roots of thoracic nerves 1-12	Increase of transmural pressure.	Peripheral vasodilatation.	
6	Systemic arterial chemoreceptors	Carotid bodies	Branch of glossopharyngeal	Asphyxia, anoxia, or hypercapnia.	Peripheral vasoconstriction; primary bradycardia, secondary reflex increase in heart rate due to hyperpnea.	Hyperpnea.
7		Aortic bodies (4 main groups)	Aortic nerves and/or vagi	Asphyxia, anoxia, or hypercapnia.		
8	Venous pressure receptors	Right and left atria	Vagi	Increase of transmural pressure or volume.	Fall in blood pressure and decrease in heart rate; change in blood volume (?).	
9	Pulmonary arterial pressure receptors	Pulmonary codus	Vagi	Increase of transmural pressure.	Fall in blood pressure and decrease in heart rate.	
10	Ventricular receptors	Wall of left ventricle	Vagi	Increase of transmural pressure.	Fall in blood pressure and decrease in heart rate.	
Lung Reflexes						
11	Pulmonary deflation receptors	Small pulmonary vessels (?)	Vagi	Strong deflation; multiple minute embolism, or pulmonary congestion.	Fall in blood pressure and decrease in heart rate.	Rapid, shallow breathing.
12	Pulmonary stretch receptors	Smaller air passages	Vagi	Inflation of lungs.		Inhibition of breathing.
13	Slowly adapting bronchial receptors	Trachea and bronchi	Vagi	Inflation or deflation of lungs.		Inhibition of breathing; bronchoconstriction.
14	Rapidly adapting tracheal and bronchial receptors	Trachea and bronchi	Vagi	Mechanical or chemical.		Cough; bronchoconstriction.
15	Cough receptors	Bronchial tree	Sympathetic	Inhalation of irritant vapors.		Cough.
16	Deflation receptors	Muscles of chest wall (?)	Dorsal roots of thoracic nerves 1-4	Suction to trachea.		Inspiratory effort.

Contributor: Dawes, Geoffrey S.

Contributor: Dawes, Geoffrey S.

References: [1] Dawes, G. S., and Comroe, J. H., Jr. 1954. *Physiol. Rev.* 34 167. [2] Aviado, D. M., Jr., and C. F. Schmidt 1955. *Ibid.* 35 247. [3] Heymans, C., and E. Neil. 1958. *Reflexogenic areas of the cardiovascular system.* J and A. Churchill, London.

29. ELECTRICAL PROPERTIES OF CARDIAC TISSUE: VERTEBRATES

Part E. RESTING AND ACTION POTENTIALS OF SINGLE FIBERS

Magnitude, in millivolts (mv), determined by direct measurement with an intracellular microelectrode filled with 3M KCl. Technique: ISu = In situ, IVt = In vitro. Values in parentheses are ranges, estimate "c" unless otherwise indicated (cf. Introduction).

Animal	Tissue	Fiber Diameter ¹ μ (C)	No. of Observations	Technique	Membrane Potential ²			Reference
					Resting mv (P)	Action mv (G)	Overshoot mv (H)	
1 Man	Ventricle	16 (10)	5	ISu	65	95	30	(1)
2 Cat	Ventricle	16 (10)		ISu	80(65-95)	100(80-120)	16(4-28) ^b	1, 2
3	Atrium	10 (30)	625	IVt	60(36-91)	55(30-95)	4.7	3
4	Papillary muscle	16 (10)	160	IVt	80(66.8-89.2) ^b	116(115-117) ^b	28	4
5	Purkinje fiber	(13)		IVt	96	121	25	5
6 Dog	Ventricle	16 (11)	200	ISu	80(65-95)	100(80-120)	16(4-28) ^b	6
7	Ventricle	16 (11)	33	ISu	82(76-96)	102(96-114)		3
8	Atrium	10 (10)	100	ISu	85(68-94)	100(82-118)	15(1-29) ^b	6
9	Atrium	10 (10)	100	IVt	85(68-95)	105(82-120)	20	3
10	Atrium	10 (10)	50	IVt	83(70-94)	100(86-119)	17	7
11	Papillary muscle	16 (11)	100	IVt	85(70-95)	105(82-125)	20	8
12	Purkinje fiber	30 (22)	400	IVt	89(86-92) ^b	121(117-125) ^b	32	7
13	Purkinje fiber	30 (22)	312	IVt	90(78-102) ^b	121	31(21-41) ^b	9
14	Signus			IVt	75(65-80)	84(70-98)	11	10
15 Goat	Purkinje fiber	75 (40)	57	IVt	94(78-110) ^b	135(123-147) ^b	41	8
16 Guinea pig	Ventricle	(13)	241	IVt	52.8	67.2	14.4	11
17	Atrium	(8)	64	ISu	57	68	11	12, 13
18 Mouse	Ventricle	(11)	50	IVt	86(73-99) ^b	98(86-110) ^b	12	14
19 Rabbit	Atrium	(11)	30	IVt	70	108	28	15
20	Atrium	(11)	27	IVt	79	97	19	16
21	Atrium	(11)	35	IVt	73	98	25	17
22	Atrium	(11)	35	IVt	73	89	16	18
23	Atrium	(11)		IVt	78(70.2-85.8) ^b	92(83.6-100.4) ^b	14	19
24	Sino-atrial node			IVt	56	57	1	19
25	Sino-atrial node		25	IVt	61	66	5	18
26	Sino-atrial node		30	IVt	66(51-74)	79(63-94)	13	8
27 Rat	Ventricle	(9)	128	IVt	83(65-101) ^b	101	18(2-30) ^b	20
28	Atrium	(8)	857	IVt	62(51-73) ^b	75.2(59.2-90.8) ^b	13.2(6.0-22.4) ^b	21
29 Sheep and calf	Purkinje fiber	60		IVt	98(91-110)	132	34(24-40)	22

		195	150	39.5 (10-70)	53.5 (13-100)	14	23
30 Chick embryo	Ventricle	44	150	29.2 (10-41)	39.2 (11-61)	10	23
31	Atrium	10	150	85	110	25	24
32	Ventricle	10	150	56 (55-90)	65 (55-90)	16	25
33	Atrium	150	150	74.2 (50-110)	74.2 (50-110)	13	23
34	Ventricle	239	150	58.4 (40-80)	77.2 (30-132)	18.9 (13-26)	28
35	Ventricle	10	150	64.5 (27-112)	102.5 (95-112)	15-15	29
36	Ventricle	485	150	84.5 (77-89)	102.5 (95-112)	15-15	12, 13
37	Ventricle	10	150	45-60	125	30	
38	Truncus arteriosus	95	150				

/1/ Values in parentheses are for fiber diameters determined from formalin-fixed paraffin sections [30]. /2/ Technical difficulties encountered in measuring small fibers make it advisable to accept low values with caution.

Contributors: (a) Hoffman, Brian F., (b) Brady, Allan J., (c) Weidmann, Silvio, (d) Draper, M. H., (e) Trautwein, Wolfgang, (f) Webb, John L., (g) Cranefield, Paul F.

- References: [1] Truex, R. C., and W. M. Copenhaver. 1947. *Am. J. Anat.* 80:173. [2] Woodbury, J. W., J. Lee, K. A. Merendino, and A. J. Brady. 1953. *Circul. Res.* N. Y. 5:179. [3] Hoffman, B. F., and E. E. Suckling. 1952. *Am. J. Physiol.* 170:357. [4] Burgen, A. S. V., and K. G. Terroze. 1952. *J. Physiol.* Lond. 119:139. [5] Trautwein, W., U. Gottstein, and J. Dudel. 1954. *Flügers Arch.* 260:40. [6] Trautwein, W., and K. Zink. 1952. *Flügers Arch.* 260:46. [7] Hoffman, B. F., and E. E. Suckling. 1953. *Am. J. Physiol.* 173:312. [8] Trautwein, W., and J. Dudel. 1958. *Flügers Arch.* 266:653. [9] Trautwein, W., U. Gottstein, and K. Federschimdt. 1953. *Ibid.* 258:243. [10] Draper, M. H., and S. Weidmann. 1951. *J. Physiol.* Lond. 115:74. [11] Johnson, E. A. 1956. *J. Pharm. Exp. Ther.* 117:237. [12] Woodbury, J. W., and A. J. Brady. 1956. *Science* 123:100. [13] Woodbury, J. W., and A. J. Brady. 1956. *Unpublished*. [14] Draper, M. H., and J. F. Lamb. 1956. *Unpublished*. [15] Cranefield, P. F., and B. F. Hoffman. 1958. *Brit. J. Pharm.* Lond. 115:74. [16] Vaughan Williams, E. M. 1956. *Circul. Res.* N. Y. 5:664. [17] West, T. C. 1955. *J. Pharm. Exp. Ther.* 115:283. [18] Cranka, M., and J. L. Webb. 1955. *Circul. Res.* N. Y. 5:604. [19] Hollander, P. B., and J. L. Webb. 1955. *J. Pharm. Exp. Ther.* 104:103. [20] F. Curtner. 1957. *Helvet. physiol. pharm. acta* 15:73. [21] Fingel, E., L. A. Woodbury, and H. Hecht. 1952. *J. Pharm. Exp. Ther.* N. Y. 4:444. [22] Ware, H. P. 1957. *Am. J. Physiol.* 164:307. [23] Sano, T., M. Ono, and T. Shimamoto. 1956. *Circul. Res.* N. Y. 4:444. [24] Weidmann, S. 1956. *J. Physiol.* Lond. 132:157. [25] Sano, T., M. Ono, and T. Shimamoto. 1956. *Circul. Res.* N. Y. 4:444. [26] Draper, M., E. Stein, and S. Myers. 1954. *Ibid.* 2:488. [27] Woodbury, J. W., and W. Trautwein. 1956. *J. Gen. Physiol.* 39:715. [28] Draper, M., E. Stein, and S. Myers. 1957. *Ibid.* 3:190. [29] Q. J. J. Exp. Physiol., Lond. 44:91. [30] F. J. A. L. Bennett, and A. R. McIntyre. 1959. *Q. J. J. Exp. Physiol.*, Lond. 44:91. [31] M. H., and M. Mya-Tu.

29. ELECTRICAL PROPERTIES OF CARDIAC TISSUE: VERTEBRATES (Concluded)

Part II: ACTION POTENTIAL CHARACTERISTICS OF SINGLE FIBERS

For a summary of conduction velocity, see reference 21. Determined with two intra- or extracellular recording electrodes. Values in parentheses are ranges, estimate "c" unless otherwise indicated (cf. Introduction).

Animal (A)	Tissue Fiber (B)	Total Duration ¹ msec (C)	Conduction Velocity m/sec (D)	Upstroke Duration msec (E)	Depolarization Maximum Rate v/sec (F)	Reference (G)
1 Man	Ventricle	200				1
2 Cat	Ventricle	(200-400)				2
3	Atrium ²	250	0.3		11.5	3
4	Papillary muscle ²					
5 Dog	Ventricle ²	(300-500)	0.95(0.9-1.0) ^b	1.1		4
6	Atrium	(100-200)	0.88(0.34-1.05)	2.0	150	2, 5
7	Atrium ² , papillary muscle	250	1.0		>150	7
8	Purkinje fiber ²	2.5				8
9	Purkinje fiber ²	(250-400)	2.0(1.3-3.2)	0.5	610	9
10	Sinus	300				6
11 Goat	Purkinje fiber ³	300	2.2(1.7-2.7)	0.5	800	10
12 Guinea pig	Ventricle	(70-100)				11, 12
13	Atrium	(100-150)				13
14 Mouse	Atrium	30				14
15 Rabbit	Atrium	135				15
16	Atrium	150				16
17	Sino-atrial node	(200-300)				17
18	Sino-atrial node	189				18
19 Sheep and calf	Purkinje fiber	400				19
20 Chick embryo	Ventricle ⁴	120(100-400) ^b				20
21	Atrium ⁴	50(30-70) ^b				21
22 Turtle	Ventricle	(600-1400)				22
23	Atrium	(400-700)				23
24 Frog	Ventricle ⁵		0.174	7.4	10	24
25	Sinus venosus	(400-600)				25
26	Transverse aortic arch	560				26

^{1/1} Conveys order of magnitude only, since this measurement varies markedly with changes in frequency of contraction. ^{2/2} At 37-38°C. ^{3/3} At 33.5°C. ^{4/4} From 7-day-old embryo. ^{5/5} At 18-19°C.

Contributors: (a) Hoffman, Brian F., (b) Brady, Allan J., (c) Weidmann, Silvio, (d) Draper, M. H., (e) Trautwein, Wolfgang, (f) Webb, John L.

References: [1] Woodbury, J. W., J. Lee, K. A. Merendino, and A. J. Brady. 1957. *Circul. Res.*, N. Y. 5:179. [2] Hoffman, B. F., and E. E. Suckling. 1952. *Am. J. Physiol.* 170:357. [3] Burgen, A. S. V., and K. G. Terroux. 1953. *J. Physiol.*, Lond. 119:139. [4] Trautwein, W., U. Gottstein, and J. Dudel. 1954. *Pflügers Arch.* 260 40. [5] Schaefer, H., and W. Trautwein. 1949. *Ibid.* 231:417. [6] Trautwein, W., U. Gottstein, and J. Dudel. 1958. *Ibid.* 266:653. [7] Hoffman, B. F., and E. E. Suckling. 1954. *Am. J. Physiol.* 179:123. [8] Trautwein, W., U. Gottstein, and K. Federschmidt. 1953. *Pflügers Arch.*

258 243. [9] Draper, M. H., and S. Weidmann. 1951. *J. Physiol.*, Lond. 115:74. [10] Johnson, E. A. 1956. *J. Pharm. Exp. Ther.* 117:237. [11] Woodbury, J. W., and A. J. Brady. 1956. *Science* 123:100. [12] Woodbury, J. W., and A. J. Brady. Unpublished. [13] Draper, M. H., and J. F. Lamb. Unpublished. [14] Marshall, J. M. 1957. *Circul. Res.* N. Y. 5: 664. [15] West, T. C. 1955. *J. Pharm. Exp. Ther.* 115:283. [16] Weidmann, S. 1953. *J. Physiol.*, Lond. 129:568. [17] Flügge, E., L. A. Woodbury, and H. H. Hecht. 1952. *J. Pharm. Exp. Ther.* 104:103. [18] Sano, T., M. Ono, and T. Shimamoto. 1956. *Circul. Res.* N. Y. 4: 444. [19] Medes, E. 1955. *Arch. Biol.*, Fir. 55:298. [20] Hutter, O. F., and W. Trautwein. 1956. *J. Gen. Physiol.* 39:715. [21] Myer, Tu. M. 1957. *J. Exp. Med.* 106:199.

Part III ELECTRICAL CONSTANTS OF SINGLE PURKINJE FIBERS

Determined with two intracellular electrodes. One electrode provided a polarising pulse which was recorded by the other at various distances. A long (100 msec) square pulse of anodal current was used. Abbreviations: λ = fiber space constant, R_m = membrane resistance, C_m = membrane capacity, T_m = membrane time constant, R_i = specific resistance of myoplasm, R_0 = extracellular resistance. Values in parentheses are ranges, estimate "c" (cf. Introduction).

Animal	No. of Determinations	Fiber Diameter μ	Fiber Length mm	λ mm	R_m Ω -sq cm	C_m μ F/eq cm	T_m msec	R_i Ω -cm	R_0 Ω -cm	Reference
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)	(K)
1 Goat	8	75	4.3	1.9	1940	12.4	19.5	105	51	1
		(53-104)	(3.0-6.4)	(1.2-2.5)	(750-3380)	(9-29)	(18-39)	(52-190)	51	2
2 Sheep and calf	4	75		1.62	1220	11.3	13.82	154		
					(480-2400)	(10.7-12.2)		(81-240)		

/1/ Falls to 20 sq m during action potential upstroke. /2/ C isolated.

Contributors (a) Hoffman, Brian F., (b) Brady, Allan J., (c) Weidmann, Silvio

References [1] Weidmann, S. 1952. *J. Physiol.*, Lond. 118:348. [2] Coraboeuf, E., and S. Weidmann. 1954. *Helvet. physiol. pharm. acta* 12:32.

30. EFFECT OF pH ON THE HEART: DOG

pH	Heart Rate beats/min	PR Interval sec	Depolarisation QRS, sec	Repolarisation Process			QTc ¹	Reference
				ST Segment	Wave			
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)
1 7.42	150	0.09	0.04	-0.3	+1.3	0.35	1	
2 6.23	120	0.13	0.05	-3.5	+3.0	0.35		
3 7.42	160	0.09	0.04	-0.5	+1.2	0.35	2	
4 8.04	100	0.11	0.04	-3.5	-4.0	0.35		

/1/ QT interval divided by the square root of the cycle length (Bazett's K). /2/ Normal /3/ Causes decrease in contractility and automaticity; death from cardiac arrest. /4/ Produces monophasic currents in animals and predisposes to cardiac arrhythmias; death usually from ventricular fibrillation.

Contributor: Gertler, Menard M.

References. [1] Gertler, M. M., H. E. Hoff, and D. G. Humm. 1946. *Am. J. Physiol.* 146:478. [2] Terroux, K. G., M. M. Gertler, and H. E. Hoff. 1947. *Ibid.* 148:1.

31. ELASTICITY OF THE ARTERIO-VEINOUS SYSTEM: MAN, DOG

Intra-individual comparisons of elasticity at various levels of pressure (P) and volume (V) can be obtained by applying the following formula: elasticity in mm Hg/ml = $\frac{\Delta P}{\Delta V} \text{ mm Hg/ml}$. The resulting values when large indicate a high degree of stiffness (low extensibility).

Part II: AORTA

Aorta		Specification	Pressure mm Hg	Elasticity	Reference	Aorta		Specification	Pressure mm Hg	Elasticity	Reference
(A)		(B)	(C)	(D)	(E)	(A)		(B)	(C)	(D)	(E)
1 Thoracic, ml/cm length		20-24 yr	Man		1	71 Abdominal, ml/cm length (concluded)		18 yr, 50 kg (concluded)	150	1.99	2
2			0	1.00		74 Total, ml/cm length		18 yr, 50 kg	0	1.25	
3			25	1.24		75					
4			50	1.55		76					
5			75	1.90		77					
6			100	2.40		78					
7			125	2.90		79					
8			150	3.35		80					
9			175	3.70		81					
10			200	3.95		82					
11			225	4.10		83 Arch, ml/cm length		14.5 kg	0	0.83	2
12		29-31 yr	0	1.15		84					
13			25	1.40		85					
14			50	1.80		86					
15			75	2.25		87					
16			100	2.65		88					
17			125	2.90		89					
18			150	3.15		90					
19			175	3.30		91					
20			200	3.45		92					
21			225	3.55		93					
22		36-42 yr	0	1.25		94					
23			25	1.50		95					
24			50	1.95		96					
25			75	2.35		97					
26			100	2.65		98					
27			125	2.85		99					
28			150	3.00		100					
29			175	3.15		Thoracic, ml/cm length		14.5 kg	0	0.50	
30			200	3.25		101					
31			225	3.30		102					
32		47-52 yr	0	1.30		103					
33			25	1.65		104					
34			50	2.05		105					
35			75	2.35		106					
36			100	2.60		107					
37			125	2.80		108					
38			150	2.95		109					
39			175	3.05		Abdominal, ml/cm length		14.5 kg	0	0.36	
			200	3.15		101					
						102					
						103					
						104					
						105					
						106					
						107					
						108					
						109					

	71-78 yr	225	3.20	110	Total, ml/cm length	14.5 kg	0	0.50
40		0	1.75	111			25	0.61
41		25	2.15	112			50	0.76
42		50	2.35	113			75	0.93
43		75	3.00	114			100	1.12
44		100	3.15	115			125	1.27
45		125	3.30	116			150	1.40
46		150	3.35	117			175	1.48
47		175	3.40	118			200	1.53
48		200	3.45	119			225	1.58
49		225	3.50	120			250	1.63
50	Thoracic, ml/cm length	0	1.02	121		Aorta filled at rate of 100 ml/min	25	0.22
51		25	1.23	122			50	0.75
52		50	1.55	123			75	1.33
53		75	1.98	124			100	1.98
54		100	2.49	125			125	2.35
55		125	2.91	126			150	3.10
56		150	3.19	127			175	3.35
57		175	3.27	128			200	3.87
58		200	3.27	129			225	4.12
59	Arch, ml/cm length	0	2.11	130		Aorta filled at rate of 10 ml/min	25	0.25
60		25	2.96	131			50	0.80
61		50	4.14	132			75	1.40
62		75	5.78	133			100	2.05
63		100	6.99	134			125	2.65
64		125	7.50	135			150	3.20
65		150	7.94	136			175	3.65
66		175	8.25	137			200	4.00
67		200	8.25	138			225	4.30
68	Abdominal, ml/cm length	0	0.67	139			25	20.5
69		25	0.88	140			50	24.0
70		50	1.14	141			75	30.5
71		75	1.43	142			100	35.5
72		100	1.73				125	40.0
		125	1.87				150	42.5

Contributors: (a) Haddy, Francis J., (b) Alexander, Robert S.

References: [1] Hallock, F., and L. C. Benson. 1937. *J. Clin. Invest.* 16:595. [2] Remington, J. W., and W. F. Hamilton. 1945. *Am. J. Physiol.* 144:546. [3] Alexander, R. S. 1954. *Circul. Res.*, N. Y. 2:140. [4] Remington, J. W., W. F. Hamilton, and P. Dow. 1945. *Am. J. Physiol.* 144:536.

31. ELASTICITY OF THE ARTERIO-VEIN SYSTEM: MAN, DOG (Concluded)

Part II: VEIN SYSTEM

Component		Pressure mm Hg	Elasticity	Reference	Component		Pressure mm Hg	Elasticity	Reference
(A)		(B)	(C)	(D)	(A)		(B)	(C)	(D)
Man					Dog (concluded)				
1	Common iliac vein, ml	0	0.60	1	78	Mesenteric venous system	20	2.70	5
2		1	2.25		79	filled at 100 ml/min, Δ ml	22	2.90	
3		2	2.50		80	(concluded)	24	3.10	
4		3	2.70		81		26	3.30	
5		4	2.90		82		28	3.48	
6		5	3.10		83		30	3.65	
7		6	3.25		84		32	3.83	
8		7	3.40		85		34	4.03	
9		8	3.55		86		36	4.25	
10		9	3.70		87		38	4.50	
11		10	3.80		88		40	4.75	
12		11	3.90		89	Mesenteric venous system	4	0.00	
13	Internal mammary vein, ml	0	0.020	1	90	filled at 10 ml/min, Δ ml	6	0.60	
14		1	0.500		91		8	1.16	
15		2	0.540		92		10	2.00	
16		3	0.560		93		12	3.00	
17		4	0.570		94		14	4.08	
18		5	0.580		95		16	5.15	
19		6	0.590		96	Pulmonary vascular bed	5	20	6
20		7	0.595		97	immediately upon filling,	10	40	
21		8	0.600		98	Δ ml	15	60	
22		9	0.605		99		20	75	
23		10	0.610		100		25	88	
24		11	0.615		101		30	104	
25		12	0.620		102		35	117	
26		13	0.625		103		40	131	
Dog					104		45	142	
27	Right atrium, ml	0	0.0	2	105		50	153	
28		5	7.0		106		55	163	
29		10	12.0		107	Pulmonary vascular bed	5	20	
30		15	16.5		108	5 sec after filling, Δ ml	10	43	
31		20	20.0		109		15	63	
32		25	23.0		110		20	80	
33		30	26.0		111		25	94	
34	Left atrium, ml	0	0.0		112		30	111	
35		5	3.0		113		35	126	
36		10	5.0		114		40	139	
37		15	8.0		115		45	154	
38		20	9.0		116		50	167	
39		25	10.0		117		55	180	
40		30	10.5		118	Pulmonary vascular bed	5	60	
41	Inferior vena cava, ml	0	2.5	3	119	immediately upon	10	87	
42		1	6.9		120	withdrawal, Δ ml	15	109	

43	2	8.9	121	20	127
44	3	9.7	122	25	138
45	4	10.4	123	30	148
46	5	10.8	124	35	156
47	6	11.3	125	40	162
48	7	11.6	126	45	169
49	8	11.9	127	50	176
50	9	12.1	128	55	180
51	10	12.3	129	5	4.0
52	11	12.5	130	10	10.5
53	12	12.8	131	15	16.0
54	13	13.0	132	20	20.0
55	14	13.2	133	25	23.0
56	15	13.4	134	30	25.0
57	16	13.6	135	35	26.5
58	0	0.00	136	40	28.0
59	2	0.95	137	45	29.0
60	4	1.85	138	50	29.5
61	6	2.70	139	5	4.5
62	8	3.50	140	10	11.5
63	10	4.25	141	15	17.0
64	12	4.90	142	20	21.5
65	14	5.40	143	25	24.0
66	16	5.85	144	30	26.0
67	18	6.20	145	35	27.5
68	20	6.40	146	40	29.0
69	22	6.50	147	45	29.5
70	4	0.00	148	5	7.5
71	6	0.45	149	10	15.5
72	8	0.95	150	15	21.5
73	10	1.40	151	20	24.5
74	12	1.70	152	25	26.5
75	14	1.95	153	30	28.0
76	16	2.20	154	35	29.0
77	18	2.45	155	40	29.5

Contributor Haddy, Francis J.

References: [1] Ryder, H. W., W. E. Motte, and E. B. Ferris, Jr. 1944. *J. Clin. Invest.* 23:333. [2] Little, R. C. 1949. *Am. J. Physiol.* 158:337. [3] Green, H. D. 1944. In O. Glasser, ed. *Medical Physics*. Year Book Publishers, Chicago. P. 213. [4] Alexander, H. S. 1948. *Proc. Soc. Exp. Biol.* N. Y. 67:410. [5] Alexander, R. S. 1954. *Circul. Res.* N. Y. 2:140. [6] Sarnoff, S. J., and E. Berglund. 1952. *Am. J. Physiol.* 171:238.

32. BLOOD VOLUMES

For a summary of blood volume methods and interpretations, see reference 24. Part I. Values in parentheses are ranges, estimate "c" (cf. Introduction), Part II: MAN

Values for plasma and RBC volumes were obtained by various dilution methods. Unless otherwise indicated, Evans' blue dye (T-1824) was used to measure plasma volume, and red cells tagged with radioactive phosphorus (P^{32}) were used to measure RBC volume. Venous hematocrit values were obtained by centrifuging venous samples for 30 minutes at 3000 rpm and at a radius of 15 cm; values were not corrected for trapped plasma. Total blood volume was calculated from the two other values listed on the same line. Where blood volumes have been estimated directly from dilution with tagged red cells, it has been assumed that the venous RBC concentration represents the total body RBC concentration. Subjects were normal, unanesthetized adults of the white race, under average environmental temperature (22-28°C) and sea level barometric pressure conditions.

Subjects	Sex	No.	Venous Hematocrit		-volume	RBC Volume	Total Blood Volume	Reference
			(A)	(B)	(C)	(D)	(E)	(F)
1	Male	40				29.8(21.5-36.3)	77.7(63.8-97.0)	1
2		20	44.1(39.3-49.4)		52.5		81.6(65.4-95.2)	2
3		49	44.7(36.0-51.8)		58.2		77.7(62.7-97.7)	3
4		11	47.2(42.0-53.0)		48.2		76.7(60.5-92.6)	4
5		31	44.1(34.9-49.0)		61.85		74.2(60.0-96.0)	5
6		53	44.76		56.5		82.3(59.8-101.7)	6
7		41	45.06		44.4		85.1	7
8		32	42.6(37.9-49.2)		55.9		82.1	8
9		30	46.3		48.57			9
10		21						10
11		59						11
12		25						12
13		88						13
14		30				31.8(25.4-38.7)		14
15		11						15
16	Female	20	39.8(37.1-41.9)		41.1		65.5(45.8-81.0)	16
17		41	40.0(33.7-44.4)		47.4		69.1	17
18		16	42(37-46)		41.5(28.9-52.3)		82.8	18
19		35	39.2(34.5-43.8)		48.2		74.3(63.0-97.5)	19
20		30			41.5		66.1(46.3-85.4)	20
21		4			44.8		64.1(49.6-77.5)	21
22	Unspecified	13			47.1(39.7-52.2)		63.1	22
23		33	43.8(34.5-50.0)		44.8		72.3	23
24		25	44.1(40.3-48.5)		44.8(36.9-54.7)		76.4(66.5-83.1)	24
25		41	44.1(40.3-48.5)		43.8(32.6-53.7)		81.3(59.2-103.7)	25
26		45	44.1(40.3-48.5)		45.4(32.3-54.2)		78.1(58.4-97.1)	26
27		10	44.1(40.3-48.5)		41.1(31.6-48.5)		81.3(56.9-100.0)	27
28		21			30.3(23.7-42.1)		68.3(49.3-91.8)	28
29		45			36.0(30.2-42.2)		71.4	29
					39.65		75.710	30

1/1) RBC labelled with radioactive iron (Fe^{55} or Fe^{59}). 12/ Calculated from: reading of packed cells. 13/ Tubes centrifuged until cells completely packed. 14/ Obtained from specific gravity of blood in copper sulfate solution. 15/ Plasma diluted with radioactive, iodinated, human serum albumin. 16/ Hematocrit corrected for trapped plasma by a factor of 0.96 or 0.95. 17/ Plasma diluted with radioactive chromium (Cr^{51}) in the form of chromic chloride or 0.315. 18/ RBC labelled with radioactive chromium (Cr^{51}) in the form of sodium chromate. 19/ g RBC/kg. 10/ Hematocrit corrected for trapped plasma by a factor

Contributors: (a) Reynolds, Monica, (b) Brown, Ellen, (c) Burke, Jack D., (d) Hansard, Sam L., (e) Walcott, William W.

References: [1] Gibson, J. D., W. C. Peacock, A. M. Seligman, and T. Sack. 1946. *J. Clin. Invest.* 25: 818. [2] Von Forest, R. 1951. *Acta med. scand.* Suppl. 234. [3] Gibson, J. G., II, and W. A. Evans, Jr. 1957. *J. Clin. Invest.* 36: 317. [4] Davis, L. J. 1942. *Edinburgh M. J.* 49: 445. [5] Storaasli, J. P., H. Krieger, H. L. Freidelt, and W. R. Holden. 1950. *Surg. Gyn. Obst.* 91: 459. [6] Gregersen, M. L., and J. L. Nickerson. 1950-51. *J. Appl. Physiol.* 3: 129. [7] Noble, R. P., and M. I. Gregersen. 1946. *J. Clin. Invest.* 25: 172. [8] Steinbeck, A. W. 1950. *Austral. J. Exp. Biol.* 28: 437. [9] Henrich, A. O., Michelson, H. L., Taylor, E. Monowyer, F. R. Maiz, and L. Griffith. 1945. *Ann. Surg.* 121: 152. [10] Frank, H., and S. J. Gray. 1953. *J. Clin. Invest.* 32: 591. [11] Griffin, G. E., W. E. Abbott, M. P. Pryde, E. Monowyer, F. R. Maiz, and L. Griffith. 1945. *Ann. Surg.* 121: 152. [12] Sterling, K., and S. J. Gray. 1950. *J. Clin. Invest.* 29: 1614. [13] Reddy, W. A., R. M. French, F. Y. K. Lau, K. G. Scott, and W. E. White. 1954. *Circulation*, N. Y. 9: 571. [14] Samet, P., H. W. Fritts, Jr., A. P. Fishman, and A. Courmand. 1957. *Medicine*, Balt. 36: 211. [15] Harrington, C. R., E. Poehlin, and J. R. Smith. 1940. *Clin. Sc.* 1, Lond. 4: 311. [16] Berlin, N. I., G. M. Hyde, R. J. Parsons, J. H. Lawrence, and S. Port. 1951. *Proc. Soc. Exp. Biol.* N. Y. 76: 831. [17] Steinbeck, A. W. 1956. *Austral. J. Exp. Biol.* 34: 95. [18] Reeve, E. B., and N. Veall. 1949. *J. Physiol.*, Lond. 108: 12. [19] Stewart, J. D., and G. M. Bourne. 1940-41. *J. Laborat. Clin. M.* 26: 1383. [20] Brady, L. W., D. Y. Cooper, M. Chodzin, J. E. McClenahan, E. R. King, and R. Williams. 1953. *Surg. Gyn. Obst.* 97: 28. [21] Gray, S. J., and H. Frank. 1953. *J. Clin. Invest.* 32: 1000. [22] Heversy, G. K. H. Koster, G. Strueman, E. Warburg, and K. Zerkow. 1944. *Acta med. scand.* 116: 561. [23] Likoff, W., D. Bettkowitz, S. Geizer, H. Strauss, and A. Reale. 1955. *Am. Heart J.* 49: 1. [24] Gregersen, M. L., and R. Rawson. 1959. *Physiol. Rev.* 39: 307.

Part II: MAMMALS OTHER THAN MAN

Values for plasma and RBC volumes were obtained by various dilution methods. Unless otherwise indicated, Evans' blue dye (T-1824) was used to measure plasma volume, and red cells tagged with radioactive phosphorus (^{32}P) were used to measure RBC volume. Venous hematocrit values were obtained by centrifuging venous samples for 30 minutes at 1000 rpm and at a radius of 15 cm; values were not corrected for trapped plasma. Total blood volume was calculated from the two other values listed on the same line. Where blood volumes have been estimated directly from dilution with tagged red cells, it has been assumed that the venous RBC concentration represents the total body RBC concentration. Subjects were normal, unanesthetized, adult animals of unspecified sex, unless otherwise indicated. Female animals were not pregnant or lactating, unless otherwise indicated.

Animal	No. of Subjects (B)	No. of Observations (C)	Venous Hematocrit % cells (D)	Plasma Volume ml/kg (E)	RBC Volume ml/kg (F)	Total Blood Volume ml/kg (G)	Reference (H)
1 Monkey	23	23	40.4 (29-49)	46.7 (31-63)		73.1 (52-109)	1
2						65.9 (49-71)	2
3 Cat	52	73		47.7 (34-56)			
4	13	13		46.5 (34.3-65.8)	18.8 (22.5-44.4)	31.5 (31.9-114.6)	3
5 Dog	39	39		53.1 (31.7-83.6)	47.9 (15.8-64.3)	100.9 (91.2-121.4)	5
6	103	10		58.0 (31.9-88.0)	42.9 (15.8-64.8)	94.7 (86.3-114.8)	6
7				51.8 (42.2-56.8)	39.0 (24.0-55.0)	94.7 (76.3-107.3)	7
8	11	22		45.2 (43.7-53.0)	37.2 (20.4-48.7)	73.6 (65.8-79.8)	8
9	13	13		46.5 (40.2-59.2)	37.2 (18.6-66.3)	83.8 (58.9-115.7)	9
10	14	14		46.3 (36.7-67.4)	37.2 (18.6-66.3)	83.8 (58.9-115.7)	10
11	21	21			66.1 (42.8-83.6)	95.7 (71.1-127.8)	11
12	15	15		42.0		81.4 (61.0-101.0)	12
13	50	50		43.7 (37.6-60.0)	19.4	92.7 (84.0-97.3)	13
14	15	72		54.7 (35-65)		83.2 (60.0-107.5)	14
15	106	106		43.1 (28-52)		100.0 (79.6-121.0)	15
16	23	21		57.0 (43.0-83.4)		94.0	16
17				52.0			17

1/1/ Calculated from hematocrit value and plasma volume measured with radioactive, iodinated, human serum albumin. 12/ RBC labelled with radioactive iron (^{59}Fe or ^{51}Cr). 13/ Anesthetized. 14/ Plasma diluted with radioactive, iodinated, human plasma. 15/ RBC labelled with radioactive chromium (^{51}Cr) in the form of sodium chromate. 16/ RBC labelled with methemoglobin. 17/ Calculated from dilution with radioactive phosphorus (^{32}P). 18/ Calculated from dilution with RBC labelled with radioactive phosphorus (^{32}P). 19/ Calculated from RBC volume and hematocrit value. 10/ Calculated RBC volume and calculated plasma volume. 11/ Hematocrit corrected for trapped plasma by a factor of 0.96 or 0.95.

32. BLOOD VOLUMES (Continued)

Part II: MAMMALS OTHER THAN MAN (Concluded)

Animal (A)	No. of Subjects (B)	No. of Observations (C)	Venous Hematocrit % cells (D)	Plasma Volume ml/kg (E)	RBC Volume ml/kg (F)	Total Blood Volume ml/kg (G)	Reference (H)
Carnivores (concluded)							
Dog (concluded)		10 ³		47.0		89.2	
19	18	36	46.9(32.0-59.4) ¹¹	53.6(41.1-74.2)		97.5(78.2-123.0)	18
20	12	12	45.7(38.6-50.0) ¹¹	54.7(45.2-71.2)		102.0(79.0-127.5)	19
21	16 ²	16		51.6(37.1-70.8)			20
22				52.7(39.6-70.6) ¹³			21
Rodents							
Guinea pig	10	10		38.6		72.0 ¹⁴	22
23	13	13		39.4(35.1-48.4) ¹⁵		75.3(67.0-92.4) ¹⁶	23
Mouse	3	11		48.8	29.0	77.7	24
25	5	5		57.4(55-63)			
26				49.2(41-54) ¹⁷			
Rabbit	29	29		38.8(27.8-51.4)	16.8(13.7-25.5)	55.6(44.0-70.0)	25
27	71	71	35.2(28.6-41.0) ¹¹		17.2(13.4-22.8)	57.3(47.8-69.5)	
28	60	60		50		70.8	26
29	39	67	38.8(27-48)	42.3(29-58)		69.8(50-93)	27
30	20	20	37.2(25.8-43.2)	43.5(35.1-49.8)		69.4(57.6-78.3)	28
31	15 ³	15		45		72.4	18
32	33	35		31.3	23.6	57.5	29
Rat	34	34				63(46-74) ¹⁹	30
35	22 ³	22	48.1(44-52)		23.2(18.9-25.5) ²	49.6(41.9-54.0)	31
36	42 ³ , 20	42	45.0(43-54) ²¹		21.2(17.2-25.8)	45.2(34.6-59.5)	32
37	10 ³	100	49.2(44.0-55.4) ²¹	27.4(24.6-31.1)		54.3(50.4-58.1)	33
Artiodactyla							
Cattle, beef	4	4	41 ¹¹			57	34
40	10	10	32.4(30.3-34.9) ¹¹	38.8(36.3-40.6)		57.4(52.4-60.6)	35
Dairy cows	3	3	40.9	36.6		62.0	36
41	20	20		55.9(42.6-75.1)	14.1(9.1-19.3) ⁵	70.5(56.8-89.4)	37
Goat	30	30		53		70(57-82)	26
42	6	6				58(50-73) ²²	38
Sheep	4	4	34 ¹¹			58	34
3-yr-old ewes	4	4	38 ¹¹			59	34
43	14	14				69.4	39
Swine, av wt = 52 kg	4	4		41.9	27.5 ²	69.4	39
Av wt = 50 kg	4	4			25.9(20.2-29.0)	55(61-68)	40
Av wt = 45 kg	4	4	39.1(30.3-43.1) ¹¹				
Perissodactyla							
50 Burro	16	16				652 ³	34
51 Horse	17	17			35	63	
52	2	2				72.0	
Light weight	6	6	43.3(37-56)	61.9(45.5-79.1) ²⁴	47.1(39.6-57.3)	109.6(94.3-136.6)	26
Heavy weight	4	4	40.3(37-46)	43.2(30.6-64.1) ²⁴	28.5(23.1-37.6)	71.7(56.7-101.7)	41
Marsupials							
55 Opossum	10	10		37.8(29.6-52.2)	19.2(14.2-29.2) ²⁵	57.0(44.5-69.8) ²⁵	42

12/ RBC labelled with radioactive iron (⁵⁹Fe or ⁵⁵Fe). 13/ Anesthetized. 15/ RBC labelled with radioactive chromium (⁵¹Cr) in the form of sodium chromate. 11/ Hematocrit corrected for trapped plasma by a factor of 0.96 or 0.95. 12/ Calculated from plasma volume and arterial hematocrit value. 13/ Plasma diluted with Rose Bengal. 14/ Calculated from dye measured in whole hemolyzed cardiac blood. 15/ Calculated from blood volume and an

32. BLOOD VOLUMES (Continued)
Part III: VERTEBRATES OTHER THAN MAMMALS; CRUSTACEANS AND MOLLUSKS (Concluded)

Animal	No. and Sex	Method	Hematocrit % cells	Plasma Volume ml/100 g	RBC Volume ml/100 g	Total Blood Volume ml/100 g	Reference
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)
12 Duck, diving, redhead, and canvasback	6	T-1824	37	7.1		11.1	3
13 Duck, white Peking	42	151 human serum albumin					
14 Hawk, red-tailed	1	T-1824	43	6.55		10.2	5
15 Lark, red-throated	3	T-1824	54	3.5		6.2	3
16 Owl, great-horned	1	T-1824	43	6.1		13.2	
17 Pheasant, ring-necked	4♂	T-1824	30	4.5		6.4	
18 Pigeon	2♂	T-1824	33	3.2		4.8	
19 Pigeon	2♂	Na ₂ Cr ₂ O ₄			3.2		1
20 Pigeon	2♀	Na ₂ Cr ₂ O ₄			3.4		
21 Frog (Rana pipiens)	10	T-1824	52	4.4		9.2	3
22 Frog (Rana pipiens)	10	T-1824	13.94	7.0		8.2	6
23 Frog, bull (H. catesbeiana)	2	T-1824	15.54	8.0		9.5	6
24 Bullhead, yellow (Amelurus natalis)	6	T-1824		1.25		1.6	6
25 Cod, ling (Ophiodon elongatus)	9	T-1824	30.14			2.5	7
26 Ratfish (Chimaera colliei)	2♀	T-1824	184			2.6	7
27 Crayfish (Cambarus virilis)	11	T-1824					
28 Mussel, fresh-water (Lamprellia ventricosa, Ambilema costalis)	25	T-1824				25.1	6
/3/ Immature. /4/ Cardiac blood. /5/ Per cent by body volume.							9.55

Contributors: (a) Reynolds, Monica, (b) Brown, Ellen, (c) Burke, Jack D.

References: [1] Rodman, G. P., E. G. Ebaugh, Jr., and M. R. S. Fox. 1957. Blood, N. Y. 12:355. [2] Newell, G. W., and C. S. Shaffner. 1950. Poultry Sci. 29:78. [3] Bond, C. F., and P. W. Gilbert. 1958. Am. J. Physiol. 194:519. [4] Hegsted, D. M., D. Wilson, J. P. Milner, and P. H. Glona. 1952. Proc. Soc. Exp. Biol., N. Y. 78:114. [5] Portman, O. W., K. F. McConnell, and R. H. Rigdon. 1952. Ibid. 81:599. [6] Prosser, C. L., and S. J. F. Weinstein. 1950. Physiol. Zool. 23:113. [7] Martin, A. W. 1950. In M. H. Hatch, ed. Studies honoring Trevor Kincaid. University of Washington Press, Seattle.

Part IV: INSECTS

Blood volume varies according to sex, stage of development, age, nutrition, rearing status, method of blood extraction, coagulability, and method of volume determination. Method: A = assumed or calculated; C = chloride, D = dye, E = exsanguination; H = cell dilution, M = manganese dilution, R = C14 insulin. Unit of measurement. BW = body weight, live or fresh unless otherwise specified. Values in parentheses are ranges, estimate "b" or "c" as indicated by superscript (cf. Introduction)

Order	Species	Stage	Method	Blood Volume		Reference
				Value	Unit of Measurement	
1 Neuroptera	<i>Stasis lutea</i> (elder fly)	Larva	D	25	% BW	1
			E	25	% BW	1
			A	20	% BW	2
			E	(5-20)c	mg/insect	3
2 Blattaria	<i>Blattella germanica</i> (German roach)	Adult	E	17	% BW	4
				17	% BW	4
				17	% BW	4
				17	% BW	4
3	<i>Periplaneta americana</i> (American roach)	Nymph, ♂	D	19.4(18.8-20.4)b	% BW (individual)	5
				20(16.3-23.7)b	% BW (pooled)	
				16.8(14.8-18.8)b	% BW (pooled)	
				15.7(13.2-18.2)b	% BW	
4		Nymph, ♀	D	19.8(19.1-20.5)b	% BW	5
				18.4(14.5-22.7)b	% BW (individual)	
				19.5(17.2-21.8)b	% BW (pooled)	
				15.8(12.3-21.3)b	% BW	
5		Adult, ♂	D	27.9(23.8-31.2)b	% BW	5
				27.9(23.8-31.2)b	% BW	
				20.9(18.8-23.0)b	% BW	
				16.9(11.9-21.9)b	% BW	
6		Adult, ♀	D	16.9(11.9-21.9)b	% BW	5
				16.9(11.9-21.9)b	% BW	
				16.9(11.9-21.9)b	% BW	
				16.9(11.9-21.9)b	% BW	
7	<i>P. fuliginosa</i>	Nymph to adult	H	14.7-6.9)c	ml/insect	6
				14.7-6.9)c	ml/insect	
				14.7-6.9)c	ml/insect	
				14.7-6.9)c	ml/insect	
8	<i>Anabrus simplex</i> (Mormon cricket)	Adult	E	10.5-6.10)c	ml/insect	7
				10.5-6.10)c	ml/insect	
				10.5-6.10)c	ml/insect	
				10.5-6.10)c	ml/insect	
9	<i>Locusta migratoria migratorioides</i> (African migratory locust)	Nymph	E	< 2	cu mm/insect	8
				< 2	cu mm/insect	
				< 2	cu mm/insect	
				< 2	cu mm/insect	
10	"Termite queen"	Adult, ♀	E	0.1	ml/insect	9
				0.1	ml/insect	
				0.1	ml/insect	
				0.1	ml/insect	
11	<i>Dytiscus sp.</i>	Larva	E	10.9-25.4)c	% BW	10
				10.9-25.4)c	% BW	
				10.9-25.4)c	% BW	
				10.9-25.4)c	% BW	
12	<i>Hydroporus sp.</i>	Larva	E	10.9-25.4)c	% BW	11
				10.9-25.4)c	% BW	
				10.9-25.4)c	% BW	
				10.9-25.4)c	% BW	
13	<i>Popillia japonica</i> (Japanese beetle)	Larva	E	0.03	ml/insect	12
				0.03	ml/insect	
				0.03	ml/insect	
				0.03	ml/insect	
14	<i>Tenebrio molitor</i> (meal worm)	Larva	E	0.22	ml/g BW	13
				0.22	ml/g BW	
				0.22	ml/g BW	
				0.22	ml/g BW	
15	<i>Agapeta galbina</i>	Pupa, ♂ and ♀	E	3.3	% BW (mean)	14
				3.3	% BW (mean)	
				3.3	% BW (mean)	
				3.3	% BW (mean)	
16	<i>Bombyx mori</i> (silkworm)	Larva	E	0.35	ml/insect	15
				0.35	ml/insect	
				0.35	ml/insect	
				0.35	ml/insect	
17		Pupa	E	31.2(27.6-34.8)b	% BW (dry)	16
				28.6(25.6-31.6)b	% BW (dry)	
				10.1(-0.11)c	ml/g BW	
				10.0(-0.35)c	ml/insect	
18		Adult	E	0.05	ml/insect	17
				0.05	ml/insect	
				0.05	ml/insect	
				0.05	ml/insect	
19	<i>Celerio euphorbiae</i> (hawk moth)	Larva, feeding	E	18.6(17.2-21.0)b	% BW	18
				18.6(17.2-21.0)b	% BW	
				18.6(17.2-21.0)b	% BW	
				18.6(17.2-21.0)b	% BW	
20		Pupa (prepupa)	E	18.0-18.6)c	% BW	18
				18.0-18.6)c	% BW	
				18.0-18.6)c	% BW	
				18.0-18.6)c	% BW	
21		Pupa	E	35	% BW (dry)	19
				35	% BW (dry)	
				35	% BW (dry)	
				35	% BW (dry)	
22				10.4-0.7)c	ml/insect	9
				10.4-0.7)c	ml/insect	
				10.4-0.7)c	ml/insect	
				10.4-0.7)c	ml/insect	

34. CARDIAC FUNCTION VS AGE: MAN

Cardiac output determined by the dye-dilution method, unless otherwise indicated. Values in parentheses are ranges, estimate "b" unless otherwise indicated (cf. Introduction).

Age yr	No. and Sex	Body Surface Area, sq m	Heart Rate beats/min	Stroke Volume ml/beat	Cardiac Output L/min	Cardiac Index L/min/sq m	Reference
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)
1 2-29 hr	10					2.5(0.9-3.7) ^c	1
2 12.4	5♂, 5♀	1.29			5.55 ¹	4.30	2
3 23.6	9♂	1.75(1.67-1.83)	76.9(67.7-97.8)	85.6(73.4-97.8)	6.49(5.47-7.51)	3.72(3.16-4.28)	3
4 34.1	10♂	1.86(1.74-1.98)	71.1(64.9-78.5)	91.8(77.6-106.0)			3
5 43.3	11♂	1.81(1.73-1.89)	69.1(63.1-75.1)	78.3(68.9-87.7)			3
6 54.8	11♂	1.67(1.57-1.77)	69.8(64.2-75.4)	67.2(61.0-73.4)			3
7 65.4	10♂	1.67(1.55-1.79)	63.0(56.6-69.4)	69.5(59.7-79.3)			3
8 71.0	10♂						3
9 73.3	9♂	1.61(1.51-1.71)	65.8(58.6-73.0)	63.0(53.6-72.4)	4.05(3.55-4.55)	2.54(2.18-2.90)	4
10 82.0	7♂	1.64(1.50-1.78)	67.0(52.0-82.0)	60.1(49.9-70.3)	3.87(3.09-4.65)	2.36(1.90-2.80)	3

/1/ Determined by the Fick method.

Contributors: (a) Wade, O. L., (b) Shock, Nathan W.

References: [1] Prec, K. T., and D. E. Cassels. 1955. *Circulation*, N. Y. 11:789. [2] Brotnacher, L., and F. Fleming. 1957. *Guy's Hosp. Rep.*, Lond. 106:268. [3] Brandfonbrener, M., M. Landowne, and N. W. Shock. 1955. *Circulation*, N. Y. 12:557. [4] Smith, W. W., N. S. Wikler, and A. C. Fox. 1954. *Ibid.* 9:352.

35. CARDIAC INDEX DETERMINED UNDER BASAL CONDITIONS: MAN

Values in parentheses are ranges, estimate "c" unless otherwise indicated (cf. Introduction).

Method	No. of Observations	Cardiac Index L/min/sq m	Reference
(A)	(B)	(C)	(D)
1 Acetylene	51	2.21(1.85-2.77)	1
2	100	2.26(1.70-2.90)	2
3	15	3.44(2.65-5.80) ¹	3
4 Ballistocardiogram	20	3.35	4
5	16	3.48	5
6	50	6.21 ¹	6
7	48	6.79(5.13-10.97)	7
8 Dye (Hamilton)	6	4.10(3.16-5.10)	8
9	5	3.69(2.29-5.25)	9
10	8	4.00(2.80-5.30)	10
11 Ethyl iodide	24	2.40(1.45-3.72)	11
12 Fick	30	3.31	12
13	10	3.16(2.00-4.32) ^b	13
14	6	3.87(2.59-4.80)	8
15	13	3.74(3.16-4.40)	14
16	6	3.80(2.00-5.60) ^b	15
17	5	3.32(2.23-4.41)	9
18	7	3.80(2.80-4.70)	16

/1/ Cardiac output, in liters per minute.

Contributor: Ferencz, Charlotte

References: [1] Grollman, A. 1929. *Am. J. Physiol.* 90:210. [2] Lewis, W. H., Jr. 1938. *Ibid.* 121:517. [3] McMichael, J. 1937. *Q. J. Exp. Physiol.*, Lond. 27:55. [4] Paine, R. M., and N. W. Shock. 1950. *Circulation*, N. Y. 1:1026. [5] Scarborough, W. R., R. Penneys, C. B. Thomas, B. M. Baker, and R. E. Mason. 1951. *Ibid.* 4:190. [6] Tanner, J. M. 1949. *J. Clin. Invest.* 28:567. [7] Taylor, H. L., and K. Tiede. 1952. *Ibid.* 31:209. [8] Werkö, L., H. Lagerlöf, H. Bucht, B. Wehle, and A. Holmgren. 1949. *Scand. J. Clin. & Invest.*, Oslo 1:109. [9] Hamilton, W. F., R. L. Riley, A. M. Attyah, A. Courmand, D. M. Fowell, A. Himmelstein, R. P. Noble, J. W. Remington, D. W. Richards, Jr., N. C. Wheeler, and A. C. Whittham. 1948. *Am. J. Physiol.* 153:309. [10] Freis, E. D., H. W. Schnaper, R. L. Johnson, and G. E. Schreiner. 1952. *J. Clin. Invest.* 31:131. [11] Starr, I., J. S. Donald, A. Margolis, R. Shaw, L. H. Collins, and C. J. Gamble. 1934. *Ibid.* 13:561. [12] Courmand, A., R. L. Riley, E. S. Breed, E. F. Baldwin, and D. W. Richards, Jr. 1945. *Ibid.* 24:106. [13] Stead, E. A., Jr., J. V. Warren, A. J. Merrill, and E. S. Brannon. 1945. *Ibid.* 24:326. [14] Lagerlöf, H., and L. Werkö. 1948. *Acta physiol. scand.* 16:75. [15] Hickam, J. B., and W. H. Cargill. 1948. *J. Clin. Invest.* 27:10. [16] Dexter, L., J. L. Whittenberger, F. W. Haynes, W. T. Goodale, R. Gorlin, and C. G. Sawyer. 1951. *J. Appl. Physiol.* 3:439.

36. CARDIAC INDEX DETERMINED BY DIRECT PICK OR DYE METHOD: MAN

All subjects resting (supine), and free of cardiovascular disease. Values in parentheses are ranges, estimate "b" unless otherwise indicated (cf. Introduction).

Age yr (A)	No. and Sex (B)	Condition (C)	Cardiac Index L/min/sq m (D)	Reference (E)
Direct Pick Method				
1 13-780	53, 9d	Hospital patients	3.5(2.9-4.1)	1
2 15-35	6	Hospital patients	3.8(2.0-5.6)	2
3 17-49	2d, 4f	Normal	3.9(2.1-5.7)	3
4 17-64	33	Hospital patients	3.8(1.7-5.9)	4
5 18-57	11d, 7f	Hospital patients	3.2(1.5-4.9)	5
6 19-26	7d	Normal	3.5(2.8-4.2)	6
7 19-59	3d, 7f	Hospital patients	3.8(2.5-5.1)	7
8 20-30	7d	Normal	3.3(1.7-4.9) ¹	8
9 21-52	13d	Normal	3.1(2.3-3.9) ²	9
10 21-56	6f	Hospital patients	3.0(1.7-4.3) ²	9
11 21-57	11d	Hospital patients	3.7(2.8-4.6) ²	9
12 21-62	18, 4f	Hospital convalescents	4.0(1.8-6.2)	10
13 22-35	9d	Normal	3.1(1.9-4.3) ²	11
14 23-38	13d	Normal	3.7(2.9-4.5)	12
15 24-42	9d	Normal	3.3(1.4-5.2)	13
16 39-63	3d, 2f	Normal	4.1(3.2-4.8) ²	14
17 Unspecified	5	Normal	3.6(2.0-5.2)	15
18 Unspecified	8d	4 normal, 4 with syphilis	4.2(2.9-5.5)	16
Dye Method				
19 13-780	53, 9d	Hospital patients	3.2(1.8-4.6)	1
20 14-68	17	Normal	3.3(2.1-4.5)	17
21 16-41	23 d, 1f	Normal	3.6(2.4-4.8)	18
22 17-49	2d, 4f	Normal	4.1(2.7-5.5)	3
23 17-64	10d	Normal	2.9(2.1-3.7)	4
24 18-39	22d	Normal	3.6(2.4-4.8)	19
25 18-57	11d, 7f	Hospital patients	3.6(2.0-5.6)	5
26 19-86	67d	Normal	2.9(1.3-4.5)	20
27 21-36	7d	Normal	4.0(2.4-5.6)	21
28 21-42	8d, 1f	Normal	3.5(2.3-4.7)	18
29 21-55	2d, 8f	Normal	3.5(1.9-5.1)	19
30 22-26	10f	Normal	3.7(1.5-5.9)	23
31 26-69	5d	Normal	3.5(2.2-4.8)	24
32 26-36	7d, 2f	Normal	3.6(2.2-5.0)	25
33 30-40	12d	Normal	4.1(2.4-5.8)	26
34 39-63	6d, 4f	Normal	3.3(1.6-5.0)	27
35 40-57	12d	Normal	3.3(2.3-4.3)	28
36 Unspecified	9	Normal	3.9(3.4-4.4)	29
37 Unspecified	8	Normal		

/1/ Oxygen uptake not measured at time blood samples taken. /2/ Right atrial blood samples.

Contributor: Wade, O. L.

- References. [1] Doyle, J. T., J. S. Wilson, C. Lepine, and J. V. Warren. 1953. *J. Laborat. Clin. M.* 41:29. [2] Hickam, J. B., and W. H. Cargill. 1948. *J. Clin. Invest.* 27:10. [3] Werkö, L., H. Lagerlöf, H. Buchi, B. Wehle, and A. Holmgren. 1949. *Scand. J. Clin. & Invest.*, Oslo 1:109. [4] Doyle, J. T., J. S. Wilson, E. H. Estes, and J. V. Warren. 1951. *J. Clin. Invest.* 30:345. [5] Neely, W. A., F. C. Wilson, J. P. Milnor, J. D. Hardy, and H. Wilson. 1954. *Surgery* 35:22. [6] Rotta, A., A. Canepa, A. Hurtado, T. Velasquez, and R. Chaves. 1956. *J. Appl. Physiol.* 9:328. [7] Donald, K. W., J. M. Bishop, G. Cumming, and O. L. Wade. 1953. *Clin. Sc.*, *Clin. Invest.* 29:651. [8] Chapman, C. B., H. L. Taylor, C. Borden, R. V. Ebert, A. Keys, and W. S. Carlson. 1950. *J. Biol. Med.* 24:106. [9] Bolomey, A. A., A. J. Michie, E. S. Breed, E. de F. Baldwin, and D. W. Richards, Jr. 1945. 28:10. [10] Stead, E. A., Jr., J. V. Warren, A. J. Merrill, and E. S. Brannon. 1945. *Ibid.* 24:326. [11] Lagerlöf, H., and L. Werkö. 1949. *Acta physiol. scand.* 16:73. [12] Lewis, C. S., A. J. Sampela, M. C. Daines, and H. H. Hecht. 1952. *Circulation*, N. Y. 6:874. [13] Kopehman, H., and G. de J. Lee. 1953. *Clin. Sc.*, *Lond.* 10:383. [14] Dexter, L., J. L. Whittenberger, F. W. Haynes, W. T. Goodale, R. Gorlin, and C. G. Sawyer. 1951. *J. Appl. Physiol.* 3:439. [15] Dexter, L., J. W. Dow, F. W. Haynes, J. L. Whittenberger, B. C. Ferris, W. T. Goodale, and H. K. Hellem. 1950. *J. Clin. Invest.* 29:602. [16] Etsten, B. E., and T. H. Li. 1954. *Anesthesiology* 15:217. [17] Kowalski, H. J., W. H. Abelmann, W. F. McKeely, N. R. Frank, and L. B. Ellis. 1954. *Am. J. M. Sc.* 228:622. [18] Chapman, C. B., and R. S. Fraser. 1954. *Circulation*, N. Y. 9:57. [19] Brandfonbrener, M.,

36. CARDIAC INDEX DETERMINED BY DIRECT FICK OR DYE METHOD: MAN (Concluded)

J. Landowne, and N. W. Shock. 1955. *Ibid.* 12:557. [21] Freis, E. D., H. W. Schnaper, R. L. Johnson, and G. E. Schreiner. 1952. *J. Clin. Invest.* 31:131. [22] Smith, W. W., N. S. Wikler, and A. C. Fox. 1954. *Circulation*, 1, Y. 9:352. [23] Hamilton, W. E., R. L. Riley, A. M. Attiyah, A. Courmand, D. M. Fowell, A. Himmelstein, R. P. Noble, J. W. Remington, D. W. Richards, Jr., N. C. Wheeler, and A. C. Witham. 1948. *Am. J. Physiol.* 153:309. [24] Kattus, A. A., A. V. Rivin, A. Cohen, and G. S. Soffo. 1955. *Circulation*, N. Y. 11:447. [25] Ebert, R. V., J. W. Borden, H. S. Wells, and R. H. Wilson. 1949. *J. Clin. Invest.* 28:1134. [26] Kopelman, H., and G. de J. Lee. 1951. *Clin. Sc.*, Lond. 10:383. [27] Fraser, R. S., and C. B. Chapman. 1954. *Circulation*, N. Y. 9:193. [28] Ball, D., H. Kopelman, and A. C. Witham. 1952. *Brit. Heart J.* 14:363. [29] McIntosh, H. D., J. F. Burnum, J. B. Hickam, and J. V. Warren. 1954. *Circulation*, N. Y. 9:511.

37. CARDIAC INDEX VS O₂ UPTAKE, AT REST AND DURING EXERCISE: MAN

Values in parentheses are ranges, estimate "c" (cf. Introduction).

	Age yr	No. and Sex	Body Surface Area sq m	Method	Condition	O ₂ Uptake ml/min/sq m	Cardiac Index L/min/sq m	Reference
	(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)
1	Young adults	3♂	1.82(1.73-1.93)	Fick	Rest	155(132-164)	3.3(2.9-3.7)	1
2					Exercise	695(373-1153)	7.8(5.1-10.4)	
3	23(15-35)	8 ¹		Fick	Rest	157(115-193)	4.2(2.6-5.7)	2
4					Exercise	315(254-398)	6.1(4.2-9.7)	
5	24.5(20-29)	3♂ ² , 1♀ ²	1.84(1.54-1.99)	Fick	Rest	132(120-150)	4.32(3.37-5.20)	3
6					Exercise	295(253-339) ³	5.25(4.06-6.14) ³	
7	25(18-39)	23♂, 10♀	1.81 ⁴	Dye	Rest	359(216-511) ⁵	3.59(2.16-5.11) ⁵	4
8					Exercise	670 ⁴	6.41(4.03-14.90) ⁶	
9	29(24-39)	9♂	1.87(1.76-2.10)	Fick	Rest	135(103-169) ⁷	3.3(2.2-4.4) ⁷	5
10					Exercise	247(208-302) ⁸	5.0(4.4-5.8) ⁸	
11	31(26-36)	3♂	1.83(1.78-1.87)	Fick	Rest	126(111-147)	3.27(2.62-3.83)	6
12					Exercise	609(269-912)	6.67(4.36-9.12)	
13	32(21-42)	8♂, 1♀	1.86(1.45-2.16)	Dye	Rest	154(124-203)	3.45(2.72-4.40)	7
14					Exercise	473(301-582)	5.31(3.14-8.05)	
15	32(25-49)	3♂ ² , 1♀ ²	1.90(1.52-2.13)	Fick	Rest	144(142-146)	4.20(3.83-4.82)	3
16					Exercise	398(362-432) ⁹	6.00(5.49-6.47) ⁹	
17	33(27-42)	3♂ ² , 1♀ ²	1.82(1.54-2.06)	Fick	Rest	145(129-160)	4.64(3.51-5.77)	3
18					Exercise	564(517-610) ¹⁰	6.70(6.39-7.06) ¹⁰	
19	34(26-52)	3♂ ² , 1♀ ²	1.93(1.57-2.23)	Fick	Rest	137(124-155)	4.51(3.75-5.09)	3
20					Exercise	1047(935-1163) ¹¹	9.19(8.31-10.47) ¹¹	
21	Unspecified	71 ²	1.74(1.46-1.96)	Fick	Rest	137(106-169)	3.8(2.8-4.7)	8
22					Exercise	417(231-725)	5.5(4.4-6.9)	


1/ Hospital patients with normal cardiovascular system. /2/ Subjects during rest apprehensive about impending exercise tests. /3/ Work level = 98 kg-m/min. /4/ Approximate value. /5/ 32 observations. /6/ 30 observations. /7/ 9 observations. /8/ 8 observations. /9/ Work level = 199 kg-m/min. /10/ Work level = 358 kg-m/min. /11/ Work level = 743 kg-m/min. /12/ 5 normal subjects, 1 with asymptomatic syphilis, and 1 with asymptomatic aortic systolic murmur.

Contributors (a) Wade, O. L., (b) Shock, Nathan W.

References: [1] Freedman, M. E., G. L. Snider, P. Brostoff, S. Kimmelblot, and L. N. Katz. 1955. *J. Appl. Physiol.* 8:37. [2] Hickam, J. B., and W. H. Cargill. 1948. *J. Clin. Invest.* 27:10. [3] Donald, K. W., J. M. Bishop, G. Cumming, and O. L. Wade. 1955. *Clin. Sc.*, Lond. 14:37. [4] Chapman, C. B., and R. S. Fraser. 1954. *Circulation*, N. Y. 9:57. [5] Lewis, C. S., A. J. Samuels, M. C. Daines, and H. H. Hecht. 1952. *Ibid.* 6:874. [6] Riley, R. L., A. Himmelstein, H. L. Motley, H. M. Weiner, and A. Courmand. 1948. *Am. J. Physiol.* 152:372. [7] Kowalski, H. J., W. H. Abelmann, W. F. McNeely, N. R. Frank, and L. B. Ellis. 1954. *Am. J. M. Sc.* 28:622. [8] Dexter, L., J. L. Whittenberger, F. W. Haynes, W. I. Goodale, R. Gorlin, and C. G. Sawyer. 1951. *J. Appl. Physiol.* 3:439.

38. CARDIAC INDEX VS O₂ UPTAKE, DURING EXERCISE: MAN

Values for 16 normal subjects, plotted for the 3rd, 4th, and 5th minutes of exercise (data of Donald, et al [1]), and for 3 subjects at different levels of exercise (data of Freedman, et al [2]). Solid line represents the regression line of the data, and broken lines the 95% range. Cardiac index = $3.70 + 0.00553$ oxygen uptake.



Contributor: Wade, O. L.

- References [1] Donald, K W, J. M. Bishop, G. Cumming, and O. L. Wade. 1955. Clin. Sc., Lond 14 37.
[2] Freedman, M. E., G. L. Snider, P. Brostoff, S. Kimmelblot, and L. N. Katz. 1955. J. Appl. Physiol. 8 37.

M. Landowne, and N. W. Shock. 1955. *Ibid.* 12:557. [21] Freis, E. D., H. W. Schnaper, R. L. Johnson, and G. E. Schreiner. 1952. *J. Clin. Invest.* 31:131. [22] Smith, W. W., N. S. Wikler, and A. C. Fox. 1954. *Circulation*, N. Y. 9:352. [23] Hamilton, W. E., R. L. Riley, A. M. Attiyah, A. Courmand, D. M. Fowell, A. Himmelstein, R. P. Noble, J. W. Remington, D. W. Richards, Jr., N. C. Wheeler, and A. C. Witham. 1948. *Am. J. Physiol.* 153:309. [24] Kattus, A. A., A. V. Rivlin, A. Cohen, and G. S. Sollo. 1955. *Circulation*, N. Y. 11:447. [25] Ebert, R. V., C. W. Borden, H. S. Wells, and R. H. Wilson. 1949. *J. Clin. Invest.* 28:1134. [26] Kopelman, H., and G. deJ. Lee. 1951. *Clin. Sc.*, Lond. 10:383. [27] Fraser, R. S., and C. B. Chapman. 1954. *Circulation*, N. Y. 9:193. [28] Ball, J. D., H. Kopelman, and A. C. Witham. 1952. *Brit. Heart J.* 14:363. [29] McIntosh, H. D., J. F. Burnum, J. B. Hickam, and J. V. Warren. 1954. *Circulation*, N. Y. 9:511.

37. CARDIAC INDEX VS O₂ UPTAKE, AT REST AND DURING EXERCISE: MAN

Values in parentheses are ranges, estimate "c" (cf. Introduction).

	Age yr (A)	No. and Sex (B)	Body Surface Area sq m (C)	Method (D)	Condition (E)	O ₂ Uptake ml/min/sq m (F)	Cardiac Index L/min/sq m (G)	Reference (H)
1	Young adults	3♂	1.82(1.73-1.93)	Fick	Rest	155(132-164)	3.3(2.9-3.7)	1
2					Exercise	695(373-1153)	7.8(5.1-10.4)	
3	23(15-35)	8 ¹		Fick	Rest	157(115-193)	4.2(2.6-5.7)	2
4					Exercise	325(254-398)	6.1(4.2-9.7)	
5	24.5(20-29)	3♂ ² , 1♀ ²	1.84(1.54-1.99)	Fick	Rest	132(120-150)	4.32(3.37-5.20)	3
6					Exercise	255(253-339) ³	5.25(4.06-6.14) ³	
7	25(18-39)	23♂, 10♀	1.81 ⁴	Dye	Rest		3.59(2.16-5.11) ⁵	4
8					Exercise	670 ⁴	6.41(4.03-14.90) ⁶	
9	29(24-39)	9♂	1.87(1.76-2.10)	Fick	Rest	135(103-169) ⁷	3.3(2.2-4.4) ⁷	5
10					Exercise	247(208-302) ⁸	5.0(4.4-5.8) ⁸	
11	31(26-36)	3♂	1.83(1.78-1.87)	Fick	Rest	126(111-147)	3.27(2.62-3.83)	6
12					Exercise	609(269-912)	6.67(4.36-9.12)	
13	32(21-42)	8♂, 1♀	1.86(1.45-2.16)	Dye	Rest	154(124-203)	3.45(2.72-4.40)	7
14					Exercise	473(301-582)	5.31(3.14-8.05)	
15	32(25-49)	3♂ ² , 1♀ ²	1.90(1.52-2.13)	Fick	Rest	144(142-146)	4.20(3.83-4.82)	3
16					Exercise	398(362-432) ⁹	6.00(5.49-6.47) ⁹	
17	33(27-42)	3♂ ² , 1♀ ²	1.82(1.54-2.06)	Fick	Rest	145(129-160)	4.64(3.51-5.77)	3
18					Exercise	564(517-610) ¹⁰	6.70(5.39-7.06) ¹⁰	
19	34(26-52)	3♂ ² , 1♀ ²	1.93(1.57-2.23)	Fick	Rest	137(124-155)	4.51(3.75-5.09)	3
20					Exercise	1047(935-1163) ¹¹	9.19(8.31-10.47) ¹¹	
21	Unspecified	7 ¹²	1.74(1.46-1.96)	Fick	Rest	137(106-169)	3.8(2.8-4.7)	8
22					Exercise	417(231-725)	5.5(4.4-6.9)	


/1/ Hospital patients with normal cardiovascular system. /2/ Subjects during rest apprehensive about impending exercise tests. /3/ Work level = 98 kg-m/min. /4/ Approximate value. /5/ 32 observations. /6/ 30 observations. /7/ 9 observations. /8/ 8 observations. /9/ Work level = 199 kg-m/min. /10/ Work level = 358 kg-m/min. /11/ Work level = 743 kg-m/min. /12/ 5 normal subjects, 1 with asymptomatic syphilis, and 1 with asymptomatic apical systolic murmur.

Contributors: (a) Wade, O. L., (b) Shock, Nathan W.

References: [1] Freedman, M. E., G. L. Snider, P. Brostoff, S. Kimelblot, and L. N. Katz. 1955. *J. Appl. Physiol.* 8:37. [2] Hickam, J. B., and W. H. Cargill. 1948. *J. Clin. Invest.* 27:10. [3] Donald, K. W., J. M. Bishop, G. Cumming, and O. L. Wade. 1955. *Clin. Sc.*, Lond. 14:37. [4] Chapman, C. B., and R. S. Fraser. 1954. *Circulation*, N. Y. 9:57. [5] Lewis, C. S., A. J. Samuels, M. C. Daines, and H. H. Hecht. 1952. *Ibid.* 6:874. [6] Riley, R. L., A. Himmelstein, H. L. Motley, H. M. Weiner, and A. Courmand. 1948. *Am. J. Physiol.* 152:372. [7] Kowalski, H. J., W. H. Abelmann, W. F. McNeely, N. R. Frank, and L. B. Ellis. 1954. *Am. J. M. Sc.* 228:622. [8] Dexter, L., J. L. Whittenberger, F. W. Haynes, W. I. Goodale, R. Gorlin, and C. G. Sawyer. 1951. *J. Appl. Physiol.* 3:439.

38. CARDIAC INDEX VS O₂ UPTAKE, DURING EXERCISE: MAN

Values for 16 normal subjects, plotted for the 3rd, 4th, and 5th minutes of exercise (data of Donald, et al [1]), and for 3 subjects at different levels of exercise (data of Freedman, et al [2]). Solid line represents the regression line of the data, and broken lines the 95% range. Cardiac index = $3.70 + 0.00553$ oxygen uptake.




Contributor: Wade, O L.

- References: [1] Donald, K W, J M. Bishop, G. Cumming, and O L. Wade. 1955. Clin. Sci., Lond. 14 37.
[2] Freedman, M E, G. L. Snider, P. Brostoff, S. Kimselblot, and L. N. Katz. 1955. J. Appl. Physiol. 8 37.

39. O₂ UPTAKE VS ARTERIO-VEINOUS O₂ DIFFERENCE, DURING EXERCISE: MAN

Values for 16 normal subjects plotted for the 3rd, 4th, and 5th minutes of exercise. Solid line represents the regression line of the data, and broken lines the 95% range. Regression line formula: $A-V \text{ O}_2 \text{ content difference, ml/100 ml (normal temperature and pressure)} = \frac{18.77 \times \text{O}_2 \text{ uptake, ml/min}}{\text{O}_2 \text{ uptake, ml/min} + 694.44}$



Contributor: Wade, O. L.

Reference: Donald, K. W., J. M. Bishop, G. Cumming, and O. L. Wade. 1955 Clin. Sc., Lond. 14.37.

40. CARDIAC OUTPUT VS O₂ CONSUMPTION, DURING EXERCISE: MAN

Values in parentheses are ranges, estimate "c" (cf. Introduction).

Exercise ¹ kg-m/min	Weight kg	Body Surface Area, sq m	Method	O ₂ Consumption L/min	Cardiac Output L/min	Reference
(A)	(B)	(C)	(D)	(E)	(F)	(G)
1 0 (Basal)	65	1.76	Acetylene	0.230(0.158-0.300)	3.87(2.96-4.61)	1, 2
2 0 (Basal)	65	1.77	Direct Fick	0.244(0.167-0.277)	5.51(3.80-7.30)	3
3 0 (Sitting)	65	1.73	Direct Fick	0.236(0.184-0.298)	6.6(4.7-8.3)	4
4 102	65	1.78 ²	Indirect Fick-CO ₂	0.630(0.621-0.638)	8.8	5
5 264	65	1.78 ²	Indirect Fick-CO ₂	0.865(0.839-0.891)	11.3(10.9-11.7)	5
6 525	65	1.78 ²	Indirect Fick-CO ₂	1.369(1.348-1.390)	14.1(13.8-14.4)	5
7 725	65	1.78 ²	Indirect Fick-CO ₂	1.814(1.653-1.974)	17.1(15.5-18.7)	5
8 840	64	1.75	Acetylene	2.08(2.05-2.10)	19.2(19.0-19.4)	6
9 960	72	1.92	Acetylene	2.72(2.77-2.45)	22.1(20.6-23.6)	6
10 1200	75	1.98	Acetylene			6
11 1440	75	1.98	Acetylene			6
12 1680	75	1.98	Acetylene			6

1/1 Bicycle ergometer. 1/2 Surface area = 0.11 x Wt^{2/3}.

Contributor: Barger, A. Clifford

References [1] Grollman, A. 1929. Am. J. Physiol. 90:210. [2] Grollman, A. 1932. Cardiac output of man in health and disease. C. C. Thomas, Springfield, Ill. [3] Cournaud, A., R. L. Riley, E. S. Breed, E. deF. Baldwin, and D. W. Richards, Jr. 1945. J. Clin. Invest. 24:106. [4] Dexter, L., J. L. Whittenberger, F. W. Haynes, W. T. Goodale, R. Gorlin, and C. G. Sawyer. 1951. J. Appl. Physiol. 3:439. [5] Douglas, C. G., and J. S. Haldane. 1922. J. Physiol., Lond. 56:69. [6] Christensen, E. H. 1931. Arbeitsphysiologie 4:470.

41. CARDIAC OUTPUT VS TEMPERATURE: MAN, DOG

Stroke volume varies according to the alteration in total circulation effected by a particular method of raising body temperature. The hot bath technique produces large peripheral dilatation, and, if blood pressure is maintained, an increase in cardiac output will occur. However, in the early stages of a fever induced by intravenous pyrogens, the heart rate increases but there is intense peripheral vasoconstriction, with no change or even diminution in cardiac output. Values in parentheses are ranges, estimate "c" (cf. Introduction).

Animal	Environment	Body Temperature °C	Stroke Volume ml/beat	Cardiac Output L/min	Reference
(A)	(B)	(C)	(D)	(E)	(F)
1 Man ¹	Cold room, 6°C		72	4.2 ²	1
2	Cool room, 20°C		72	4.2 ²	
3	Hot room, 36°C		70	4.7 ²	
4	Hot room, 45°C		65	5.5 ²	
5 Man	24°C	36	133	8.5 ³	2
6	Hot bath, 47°C	40	151	15.9 ³	
7 Man ⁴	Comfortable	33.1(32.9-33.8)		5.0(2.2-9.6) ⁵	3
8	Hot, humid room	36.4(35.2-37.5)		20.2(3.5-36.6) ⁵	
9 Man ⁶	Comfortable	32.6(29.6-34.2)		5.1(2.3-7.1) ⁵	3
10	Hot, humid room	35.7(34.4-36.2)		10.0(4.1-15.0) ⁵	
11 Man ⁷	Cool room, 15°C	36.7	39.6	2.38	4
12	Cool room, 22°C	36.7	42.2	2.80	
13	Hot room, 30°C	36.7	81.1	5.10	
14	Hot room, 40°C	36.7	108.1	7.85	
15	Hot room, 50°C	38.0	95.0	11.50	
16 Dog ⁸	Cold room, 11°C	20	20.3	0.60	5
17	Cold room, 11°C	25	16.8	0.95	
18	Cold room, 11°C	30	16.6	1.52	
19	Cold room, 11°C	35	16.5	2.33	
20	Cold room, 11°C	38	20.5	2.99	
21 Dog ⁹	Ice water	20	16.7	0.47	5
22	Ice water	25	13.5	0.84	
23	Ice water	30	12.8	1.26	
24	Ice water	35	12.1	1.55	
25	Ice water	38	9.9	1.34	

1/1 1 subject only. 1/2 Determined by acetylene method. 1/3 Determined by ethyl-iodide method. 1/4 3 normal subjects. 1/5 Determined by Fick method. 1/6 4 patients with congestive cardiac failure. 1/7 Single observations on 1 subject; age, 19 yr, weight, 71.5 kg, height, 180 cm. 1/8 Anesthetized animals; average weight, 24.6 kg. 5 observations for each entry. 1/9 Anesthetized animals, average weight, 10.5 kg; 13 observations for each entry. Contributors. (a) Cooper, K. E., (b) Blasius, W., (c) Wade, O. L. References [1] Grollman, A. 1930. Am. J. Physiol. 95:263. [2] Henderson, Y., and H. W. Haggard. 1925. Ibid. 73:193. [3] Burch, G. E., and A. Hyman. 1957. Am. Heart J. 53:665. [4] Westler, K., and R. Thauer. 1943. Zschr. ges. exp. Med. 112:345. [5] Brendel, W., C. Albers, and W. Usinger. 1958. Pflügers Arch. 266:341.

Part I: MAN

Abbreviations: A = acetylene, B = ballistocardiogram, C = catheterization (Fick), D = dye (Hamilton), E = ethyl iodide, Hb = hemoglobin. Values in parentheses are ranges, estimate "c" unless otherwise indicated (cf. Introduction).

	Condition	Subjects, No. and Sex	Age yr	Method	Cardiac Output L/min	Cardiac Index L/min/sq m	Reference
	(A)	(B)	(C)	(D)	(E)	(F)	(G)
1	Anemia; Hb = 4.87(1.9-10.5) g	8♂, 109 ¹	57(23-84)	C	9.22(6.9-14.3)		1
2	Anemia; Hb = 7.03(2.4-13.0) g	7♂, 109 ²	32.5(17-67)	C	8.08(3.9-13.6)	4.98(2.5-8.3)	2
3	Anemia; Hb = 7.87(5.8-10.8) g	11♂, 19 ³	22.5(12-39)	C	9.34(6.5-14.5)	6.06(5.1-7.5)	3
4	Anemia; Hb = 5.56(4.4-7.8) g	1♂, 109 ⁴	46.5(22-73)	C	8.45(5.9-13.4)	5.37(4.0-8.8)	4
5	Anemia; Hb = >7 g	10 ⁵	38(17-67)	C		3.60(2.5-5.1)	2
6	Anemia; Hb = <7 g	14 ⁵	34(19-67)	C		5.70(2.9-8.3)	
7	Anemia, rest	10♂, 19	22(12-39)	C		6.17(5.1-7.5)	3
8	Anemia, exercise; O ₂ uptake = 250(194-304) ml/sq m	10♂, 19	22(12-39)	C		8.25(6.1-10.0)	
9	Anemia, rest	1♂, 109	46.5(22-73)	C		5.40(4.0-8.8)	4
10	Anemia, exercise; O ₂ uptake = 410(210-579) ml/sq m	1♂, 109	46.5(22-73)	C		8.20(5.2-13.1)	
11	Anemia	5 ⁵		E		2.64(1.55-3.80)	5
12	Anoxemia; arterial O ₂ saturation = 70%	16 ⁵		B		4.97	6
13	Anoxia (acute), 10% O ₂ in N ₂	5♂	(23-47)	C	5.20 ⁶		7
14	Anxiety	5♂		C		5.5	8
15		12♂, 48 ⁸	31(21-52)	C		4.4(3.37-5.77)	9
16	Arrhythmia	8 ⁵		E		1.78(1.12-2.62)	5
17	Arteriosclerosis	10♂	71(62-77)	D		2.67(2.2-3.2)	10
18	Arterio-venous aneurysms	6♂	Young soldiers	C	7.4(4.8-9.2)	4.16(2.70-5.15)	11
19		2	28(20-36)	C	14.6(13.8-15.4)	7.56(7.17-8.14)	4
20		11	29(19-51)	C	11.1(6.4-16.0)		12
21		14		D		5.69(4.43-8.44)	13
22	Beriberi, alcoholic	1♂	29	C		7.89; 4.2 ¹⁰	14
23		1♀	28	C		6.09; 4.6 ¹⁰	15
24	Beriberi, alcoholic edematous	1♂	35	C	11.8 ⁹ ; 5.4 ¹⁰		16
25	Beriberi, occidental	1		C		4.39; 2.5 ¹⁰	17
26		1		C		7.2 ⁹ ; 2.7 ¹⁰	
27		8♂, 39	44(34-57)	D		4.27(2.76-6.08)	18
28		4♂, 29	46(39-57)	D		4.46(3.53-7.72)	19
29		6♂, 28	50(38-64)	D		5.33(3.24-11.20)	18
30		3♂, 18	38(16-51)	D		4.89(4.33-5.72)	19
31		8 ⁵		C		2.87(2.1-3.7)	20
32		6 ⁵		C		2.96(1.31-4.17)	21
33		6 ⁵		D		2.93(1.31-4.70)	22
34	Coronary arterial disease	11		B		2.89	6
35		8		E		2.76(1.29-5.20)	5
36	Fatty liver	3♂	43(41-45)	D		4.76(3.57-5.88)	18
37	Hypertension	12 ⁵		E		2.35(1.27-3.95)	5
38		3 ⁵		C		4.95(4.03-6.70)	21
39		3 ⁵		D		5.31(3.34-8.25)	
40		18 ⁵		C		5.23(1.85-5.91)	22
41		18 ⁵		D		3.31(2.00-5.50)	
42		5 ⁵		C		3.2(2.5-3.7)	23
43	Hyperthyroidism	3♂, 119	34(17-55)	D		5.24(3.3-6.3)	24
44		5♂, 59	34(12-59)	C		5.87(3.9-7.3)	
45	Hyperthyroidism, rest	2♂, 39	39(21-62)	C		6.36(5.44-8.39)	4
46	Hyperthyroidism, exercise; O ₂ uptake = 672(474-1012) ml/sq m	2♂, 39	39(21-62)	C		9.76(6.8-12.6)	
47	Mitral insufficiency	11 ⁵		C		1.91(1.85-2.50)	25
48	Mitral lesion	12 ⁵		C		3.4(2.11-4.64)	22
49		12 ⁵		D		3.45(2.36-4.69)	
50	Mitral stenosis	7 ⁵		C		3.44(2.4-5.2)	20
51		18 ⁵		C		2.33(1.49-3.48)	25
52	Myocardial infarction	13 ⁵		D		2.7(1.5-4.7)	23

/1/ 6 acute, 5 chronic, 7 chronic with cardiac failure. /2/ 10 with iron deficiency, 4 with pernicious anemia, 2 with sickle cell anemia, 1 with aplastic anemia. /3/ 10 with sickle cell anemia, 1 with sickle cell anemia and cor pulmonale, 1 with sickle cell anemia and rheumatic fever. /4/ 7 with iron deficiency, 2 with pernicious anemia, 2 macrocytic. /5/ Number of observations. /6/ Breathing air: cardiac output, 5.74 L/min. /7/ Pulse rate exceeding 82 beats/min and metabolic rate elevated 10% above basal level. /8/ Apprehensive at prospect of exercising with catheters in heart. /9/ Before treatment. /10/ After treatment.

42. EFFECT OF VARIOUS CONDITIONS ON CARDIAC OUTPUT: MAMMALS (Continued)

Part I: MAN (Concluded)

	Condition	Subjects, No. and Sex	Age yr	Method	Cardiac Output L/min	Cardiac Index L/min/sq m	Reference
	(A)	(B)	(C)	(D)	(E)	(F)	(G)
53	Myocardial infarction (acute), without shock	8♂, 1♀	57(51-66)	D		2.4(1.3-2.9)	10
54	Myocardial infarction (acute), with shock	5♂, 2♀	62(48-80)	D		1.6(1.3-2.2)	
55	Oxygen, 100%	16♂		B		3.34	6
56	Paget's disease	6♂, 7♀	67(56-74)	C	6.74(3.7-13.3)		26
57		2	(54-55)	C	5.45(5.18-5.72)	3.47(3.27-3.66)	27
						4.64(3.11-4.87)b	28
62	142(116-168) ^b L/min/sq m Pregnancy, 36-40 wk, O ₂ uptake = 150(130-170) ^b L/min/sq m	11♀	22.3(17-29)	C	5.53(4.09-6.97) ^b	3.44(2.56-4.32) ^b	
63	Pulmonary disease	5♂		C		3.86(3.4-5.5)	20
64		10♂		C		3.92(1.98-6.50)	21
65		10♂		D		4.21(1.59-6.60)	21
66		2♂		C		3.96(1.20-4.73)	22
67		2♂		D		3.98(1.78-4.19)	22
68	Shock, mild	4♂		C		2.46(2.18-2.71)	29
69	Sitting, legs outstretched	17♂ ¹¹ , 19♀ ¹¹	37(19-59)	C	5.57(2.45-11.19)		30
70	Supine	17♂ ¹¹ , 19♀ ¹¹	37(19-59)	C	5.93(2.45-11.38)		
71	Standing	15♂		A	3.03(2.86-4.68)		31
72	Tilting, 70°	6♂		C		2.7(2.0-3.9)	8
73	Trauma (skeletal), moderate	4♂		C		2.66(1.30-3.01)	29

/5/ Number of observations //1/ Normal subjects and hospital patients.

Contributors: (a) Wade, O. L., (b) Ferencz, Charlotte

- References: [1] Sharpey-Schafer, E. P. 1944. *Clin. Sc.*, Lond 5:125. [2] Brannon, E. S., A. J. Merrill, J. V. Warren, and E. A. Stead, Jr. 1945. *J. Clin. Invest.* 24:332. [3] Leight, L., T. H. Snider, G. O. Clifford, and H. K. Helms, 1954. *Circulation*, N. Y. 10:652. [4] Bishop, J. M., K. W. Donald, and O. L. Wade. 1955. *Clin. Sc.*, Lond. 14:329. [5] Starr, I., J. S. Donald, A. Margolis, R. Shaw, L. H. Collins, and C. J. Gamble. 1934. *J. Clin. Invest.* 13:561. [6] Scarborough, W. R., R. Penneys, C. B. Thomas, B. M. Baker, and R. E. Mason. 1951. *Circulation*, N. Y. 4:190. [7] Motley, H. L., A. Courmand, L. Werks, A. Himmelstein, and D. Dredale. 1947. *Am. J. Physiol.* 150:315. [8] Stead, E. A., Jr., J. V. Warren, A. J. Merrill, and E. S. Brannon. 1945. *J. Clin. Invest.* 24:326. [9] Donald, K. W., J. M. Bishop, G. Cumming, and O. L. Wade. 1955. *Clin. Sc.*, Lond. 14:37. [10] Smith, W. W., N. S. Wikler, and A. C. Fox. 1954. *Circulation*, N. Y. 9:352. [11] Epstein, F. H., O. W. Shadle, F. R. Ferguson, and M. E. McDowell. 1953. *J. Clin. Invest.* 32:543. [12] Cohen, S. M., O. G. Edholm, S. Howarth, J. Michael, and E. P. Sharpey-Schafer. 1948. *Clin. Sc.*, Lond. 7:35. [13] Schreiner, G. E., N. Freinkel, J. W. Athens, and A. Stone. 1953. *Circulation*, N. Y. 7:718. [14] Lehey, W. J., D. B. Arst, M. Silver, C. R. Kleeman, and P. Kunkel. 1953. *Am. J. Med.* 14:248. [15] Fowler, N. O. 1953. *Ann. Int. M.* 38:478. [16] Burwell, C. S., and L. Dexter. 1947. *Tr. Ass. Am. Physicians* 60:59. [17] Campbell, J. A., L. A. Selvestone, and D. L. Donovan. 1951. *J. Clin. Invest.* 30:632. [18] Kowalski, H. J., and W. H. Abelmann. 1953. *Ibid.* 32:1025. [19] Kowalski, H. J., W. H. Abelmann, and W. F. McNeely. 1954. *Ibid.* 33:768. [20] Hickam, J. B., and W. H. Cargill. 1948. *Ibid.* 27:10. [21] Hamilton, W. F., R. L. Riley, A. M. Attvay, A. Coward, M. Fowell, and A. Himmelstein. R. P. m. 1948. *Am. J. Physiol.* 153:309. and J. Clin. Sc. Invest., Oslo 1952. *J. Clin. Invest.* 31:131. [22] W. Noble, M. Fowell, and A. Himmelstein. R. P. m. 1948. *Am. J. Physiol.* 153:309. and J. Clin. Sc. Invest., Oslo 1952. *J. Clin. Invest.* 31:131. [23] Daley, J. Draper, A. R. Heinbecker, R. [24] Lequime, J., and H. Denolin. 1955. *Ibid.* 12:215. [25] Howarth, S. 1953. *Clin. Sc.*, Lond 12:271. [26] Bader, R. A., M. E. Bader, D. J. Rose, and E. Braunwald. 1955. *J. Clin. Invest.* 34:1524. [27] Courmand, A., R. L. Riley, S. E. Bradley, E. S. Breed, R. P. Noble, H. D. Lauson, M. I. Gregersen, and D. W. Richards, Jr. 1943. *Surgery* 13:964. [30] Donald, K. W., J. M. Bishop, G. Cumming, and O. L. Wade. 1955. *Clin. Sc.*, Lond. 12:199. [31] McMichael, J. 1937. *Q. J. Exp. Physiol.*, Lond 27:55.

42. EFFECT OF VARIOUS CONDITIONS ON CARDIAC OUTPUT: MAMMALS (Concluded)

Part II: MAMMALS OTHER THAN MAN

Condition or Activity: U = urethane, M = morphine; B = butallylonal; Bt = barbiturate, P = sodium pentobarbital. Method. F = Direct Fick; W = Wettterer; R = Rotameter, NaCl = Stewart's NaCl; B-R = Broemser-Ranke. Values in parentheses are ranges, estimate "c" unless otherwise indicated (cf. Introduction).

Animal	No. of Subjects	Weight ¹ kg	Body Surface Area, sq. m	Condition or Activity	Method	O ₂ Consumption L/min	Cardiac Output L/min	Reference
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)
1 Cat	10			Anesthetized (U)	F		0.332(0.127-0.537) ^b	1
2	5			Anesthetized (U)	F		(0.225-0.431)	2
3		4.1		Basal, anesthetized (U)	W		0.587	3-5
4 Dog	6	6.4	0.39 ²	Basal, anesthetized (M)	F	0.057(0.042-0.068)	1.12(0.65-1.57)	6-8
5	3	11.8		Basal, anesthetized (B)	W		1.13(0.81-1.58)	3, 4, 9
6	10			Anesthetized (M + Bt)	F		1.01(0.672-1.44)	10
7	10			Anesthetized (M + Bt)	R		0.994(0.668-1.46)	10
8		14.4	0.68 ²	Anesthetized (M + Bt)	NaCl		1.82(1.14-2.50) ^b	11
9		18.5	0.78 ²	Anesthetized (M + Bt)	NaCl		2.24(1.26-3.22) ^b	11
10		23.9	0.93 ²	Anesthetized (M + Bt)	NaCl		2.66(2.00-3.32) ^b	11
11	53	16.1	0.71	Basal	F	0.106(0.063-0.184)	2.21(1.20-3.84)	12
12	2	15.0	0.68	Shivering due to cold water immersion	F	0.234(0.232-0.236)	4.49(3.66-5.31)	12
13		21.6	0.87 ²	Sitting or standing	F	0.206(0.150-0.245)	3.30(1.8-4.3)	13
14								
15								
16								
17								
18								
19				100 Treadmill, 5 mi/hr, 100	F	1.402(1.380-1.420)	12.15(12.1-12.2) ⁴	
20 Ferret	10	0.91		Anesthetized (Bt)	F		0.139(0.082-0.200)	14
21 Goat	21	23.7	0.91 ⁵	Basal	F	0.176(0.078-0.329)	3.11(1.37-5.60)	15
22 Horse	1	283.0	4.30 ⁶	Standing	F	1.364	18.8	
23				Treadmill, 47.6 m/min, 0°	F	2.965	31.4	16
24				Treadmill, 42.3 m/min, 80°	F	3.084	37.2	
25	1	342.0	4.90 ⁶	Standing	F	2.480	24.0	
26				Treadmill, 56.9 m/min, 6.5°	F	4.414(4.315-4.513)	53.1(46.6-59.5)	
27 Rabbit	8	3.2	0.21 ⁷	Anesthetized	F	0.021(0.014-0.028)	0.35(0.26-0.48)	17
28	38	2.1		Unanesthetized or lightly anesthetized	F		0.25(0.12-0.38)	18
29	3	4.6		Anesthetized (U)	W		0.527(0.284-0.763)	3, 4, 9
30	4	4.6		Anesthetized (U)	B-R		0.232(0.162-0.398)	19-21
31 Rat ⁸	52	0.18	0.03 ⁹	Anesthetized (P)	F		0.047(0.015-0.079)	22

1/1 Average. 1/2 Body surface area = $0.112 \times W^{2/3}$. 1/3 3 dogs pregnant during part of study. 1/4 Average for 2-4 experiments on same dog. 1/5 Body surface area = $0.11 \times W^{2/3}$. 1/6 Body surface area = $0.10 \times W^{2/3}$. 1/7 Body surface area = $0.098 \times W^{2/3}$. 1/8 At altitude of 1 mile. 1/9 Body surface area = $0.091 \times W^{2/3}$.

Contributors (a) Barger, A. Clifford, (b) Grossman, Jacob, (c) Blasius, W., (d) Lawson, Hampden C., and D. E. Holtgrave

References: [1] Huggett, A. St G 1924 J. Physiol., Lond. 59:373. [2] Doi, Y 1921. Ibid. 55:43. [3] Wettterer, E. 1937. Zschr. Biol. 98:26. [4] Wettterer, E. 1938. Ibid. 99:158. [5] Wettterer, E., and B. Deppe. 1939. Ibid. 99:320. [6] Harrison, T. R., C. P. Wilson, and A. Blalock 1925. J. Clin. Invest. 1:547. [7] Harrison, T. R., and A. Blalock 1926. Ibid. 2:435. [8] Harrison, T. R., and B. W. Leonard 1926. Ibid. 3:1. [9] Deppe, B., and E. Wettterer. 1940. Zschr. Biol. 100:105. [10] Seely, R. J., W. E. Nerlick, and D. E. Gregg 1950. Circulation, N. Y. 1:1261. [11] Wiggers, H. C. 1944. Am. J. Physiol. 140:519. [12] Marshall, E. K., Jr. 1926. Am. J. Physiol. 77:459. [13] Barger, A. C., V. Richards, J. Metcalfe, and B. Günther. Unpublished. [14] Kempf, J. E., and H. T. Chang. 1949. Proc. Soc. Exp. Biol., N. Y. 72:711. [15] Barcroft, J., A. E. Boycott, J. S. Dunn, and R. A. Peters 1919. Q. J. Med., Oxf. 13:35. [16] Zuntz, N., and O. Hagemann. 1898. Untersuchungen über den Stoffwechsel des Pferdes bei Ruhe und Arbeit. Parey, Berlin. [17] Leegard, F. 1926. Acta med scand., Suppl. 17:1. [18] Odaira, T. 1925. Tohoku J. Exp. Med. 6:325. [19] Deppe, B., and E. Wettterer. 1939. Zschr. Biol. 99:307. [20] Broemser, P., and O. F. Ranke. 1930. Ibid. 90:467. [21] Broemser, P., and O. F. Ranke. 1933. Zschr. Kreislaufforsch. 25:11. [22] Blood, F. R., D. L. Smith, and F. E. D'Amour. 1950. Am. J. Physiol. 163:268.

43. HEART RATE. VERTEBRATES

Heart rate varies with species, sex, age, size, environment, and temperature. Values in parentheses are ranges. estimate "c" (cf. Introduction).

Part I: MAMMALS

	Species	Weight g (B)	Heart Rate beats/min (C)	Reference (D)
			(40-56) (45-50)	1 1
1	Ass (<i>Equus asinus</i>)	400,000	138(128-144)	e
2			660	2
3	Badger (<i>Taxidea taxus</i>)	4.3	30 ¹	2
4	Bat (<i>Pipistrellus pipistrellus</i>)	4.3	750(600-900)	1, 2
5		9.4	76 ¹	2
6	Bat (<i>Plecotus auritus</i>)	9.4	140	3
7			102	3
8	Beaver (<i>Castor canadensis</i>)		(25-32)	1
9			(120-140)	1
10	Camel (<i>Camelus bactrianus</i>)		(110-130)	4, 5
11	Cat (<i>Felis domesticus</i>)		(116-128)	4, 6
12		2500	245(161-290) ³	7
13		754	240	8
14		1312	300 ⁴	4
15		117	(35-40)	1
16	Cattle (<i>Bos taurus</i>)		(45-50)	1
17			106(100-215) ⁴	3
18			45	9, 10
19	Cattle (<i>B. taurus</i>)	500,000	43	11
20			(60-70)	5
21	Cattle, dairy, ♀		71.6	12-14
22			684(660-702)	e
23	Chipmunk, least (<i>Eutamias minimus</i>)		(70-120)	1, 5
24	Dog (<i>Canis familiaris</i>)		(100-200)	1
25			(50-56) ⁵	5
26			(80-130)	5
27		20,000	85	4, 8
28		15,000	(72-82)	4, 6
29		9600	96	4, 8
30		6500	120	15
31		5000	(105-125)	4, 6
32		1750	180 ⁴	4
33		1040	208(145-275) ^{3, 4}	7
34	Dog, dachshund (<i>C. familiaris</i>)		115(96-129)	16
35	Dog, spitz (<i>C. familiaris</i>)		106	16
36	Dog, St. Bernard (<i>C. familiaris</i>)		(74-80)	16
37	Dolphin (<i>Delphinus delphis</i>)		150	1
38	Dormouse (<i>Muscardinus avellanarius</i>)	17	(590-670)	4, 17
39		23	(580-780)	4, 17
40		14	610	4, 18
41		14	104 ¹	18
42	Elephant (<i>Elephas indicus</i>)	2,000,000	41	19
43		2,000,000	(25-28)	1
44		3,000,000	(46-50)	4, 20
45		<4,090,000	30(22-39)	21
46	Ferret, European (<i>Putorius sp</i>)		231(216-242)	e
47	Giraffe (<i>Camelopardalis giraffa</i>)		66	1
48	Goat (<i>Capra hircus</i>)		(60-80)	1
49			(70-90)	4, 5
50			(80-100)	6
51		33,000	135	4, 21
52	Guinea pig (<i>Cavia porcellus</i>)		280	5
53			300	4, 15
54		300	290	4, 22, 23
55		400	267	4, 24, 25
56		593	260 ³	26
57		584	200(132-288) ³	1, 27
58		437	269(225-312) ³	7

1/1 Hibernating 1/2 Diving 1/3 Anesthetized 1/4 Young animal. 1/5 Basal rate.

HEART RATE: VERTEBRATES (Continued)

Part I. MAMMALS (Continued)

	Species	Weight g (B)	Heart Rate beats/min (C)	Reference (D)
61	Hamster (<i>Mesocricetus auratus</i>)	103	347(276-420) ³	7
62		90	400 ³	26
63	Hedgehog (<i>Erinaceus</i> sp)	485	263(200-325) ³	7
64	Hedgehog (<i>E. europaeus</i>)	520	300(280-320)	1, 2
65		520	(48-70) ⁶	2
66		911	189	8
67	Horse (<i>Equus caballus</i>)		37(32-44)	1, 5
68			(23-46)	16
69			(25-40)	28
70			63(60-71) ⁴	1
71		450,000	(36-40)	4, 29
72		400,000	(34-46)	1
73		380,000	55	8
74	Horse, thoroughbred (<i>E. caballus</i>)		(38-45)	5
75	Hyaena (<i>Hyaena hyaena</i>)		55	1
76	Lemming, collared (<i>Dicrostonyx rubicatus</i>)		416(348-465)	e
77	Lion (<i>Felis leo</i>)		40	1
78	Manatee (<i>Trichechus</i> sp)	170,000-330,000	50	30
79		170,000-330,000	30 ²	30
80	Marmot (<i>Marmota marmota</i>)		80	1
81		3600	186(160-206)	4, 31
82	Monkey (<i>Macaca irus</i>)		215	32
83	Monkey (<i>M. mulatta</i>)		165	a
84			215	f
85	Mouse (<i>Mus musculus</i>)		300	1
86			600	4, 5
87			655	4
88		29	376 ³	26
89		17.4	500(450-550) ³	7
90		12	670 ⁴	4, 33
91	Mouse, white-footed (<i>Peromyscus</i> sp)	22	420 ³	26
92	Mouse, wild	22	480 ³	26
93			(46-50)	1
			120	1, 34
			60	1
		140,000-180,000	110	3
		140,000-180,000	50 ²	3
98	Rabbit (<i>Lepus</i> sp)	1344	251(167-330) ³	7
99	Rabbit (<i>L. cuniculus</i>)		(120-160)	1
100			235	1
101			187	14
102			240	4
103		1434	220	4, 8
104		2000	205	4, 35
105	Rabbit (<i>L. europaeus</i>)	2500	64(60-70)	4, 36
106	Rat (<i>Mus rattus</i>)		400	1, 5
107			420	4
108			(300-320)	4, 14
109			(300-500)	28
110			(350-550)	37
111		200	(360-520)	4, 17
112	Rat (<i>Rattus norvegicus</i>)		362	38
113			520 ⁴	f
114		252	352(260-450) ³	7
115	Rat, white	6.3-8.0	301(279-317) ⁷	39
116		10	309(292-331) ⁷	39
117		16-20	488(407-624) ⁷	39
118		92-210	305(270-350) ⁵	39
119		92-210	538 ⁷	39
120		92-210	656 ⁸	39
121		237	347 ³	26
122	Seal (<i>Phoca vitulina</i>)	20,000-25,000	100	40
123		20,000-25,000	10 ²	40

/2/ Diving. /3/ Anesthetized /4/ Young animal. /5/ Basal rate. /6/ Cold. /7/ Held. /8/ Fettered

43. HEART RATE VERTEBRATES (Continued)

Part I: MAMMALS (Continued)

	Species (A)	Weight g (B)	Heart Rate beats/min (C)	Reference (D)
124	Sheep (<i>Ovis aries</i>)		180	14
125			(60-80)	1,5
126		50,000	(70-80)	24

134			(1300-4000)++	44
135			371	42
136	Swine (<i>Sus domesticus</i>)	100,000	(60-80)	1,5
137		130,000	(70-86)	4,6
138	Tapir (<i>Tapirus indicus</i>)		44	3
139	Tiger (<i>Felis tigris</i>)		64	3
140	Voie, field (<i>Microtus arvalis</i>)	21.3	5223	7
141	Wallaby (<i>Macropus</i> sp)		125	14
142	Weasel, long-tailed (<i>Mustela frenata</i>)		182(172-192)	e
143	Weasel, short-tailed (<i>M. erminea</i>)		357(300-420)	e
144	Whale, beluga (<i>Delphinapterus</i> sp)		(15-16)	43

/1/ Hibernating. /3/ Anesthetized /9/ Awakened from sleep. /10/ Excited, struggling

Contributors. (a) Robb, Jane Sands, (b) Garb, Solomon, (c) Kruta, Vladislav, and Václav Seliger, (d) Woodbury, Robert A., (e) Morrison, Peter R., and A. R. Dawe, (f) Walker, Sheppard M., (g) Lombard, Elira A., (h) Nardone, Roland M

References. [1] Lehmann, G. 1925. *Tabulae biol.*, Berl. Bd 1, p. 138. [2] Buchanan, F. 1911. *J. Physiol.*, Lond. 42 21. [3] Irving, L., P. F. Scholander, and S. W. Grinnell. 1941. *J. Cellul. Physiol.* 17:145. [4] Clark, A. J. 1927. *Comparative physiology of the heart*, Macmillan, New York. [5] Dukes, H. H. 1947. *The physiology of domestic animals*. Cornell University Press, Ithaca. [6] Reichert, A. 1910. *Zbl. Biochem.* 10:170. [7] Kruta, V. 1958. *Some aspects of the comparative physiology of the heart*. (In Czech.) *Babáková sbírka*, Prague. [8] Vierordt, K. 1877. *Grundriss der Physiologie der Menschen*. Ed. 5. p. 162. [9] Ralston, N. P. 1940. *Res. Bull. Univ. Missouri Agr. Exp. Sta.* 317. [10] Blaxter, K. L. 1943. *Vet. J.*, Lond. 99 2. [11] Benedict, F. G., and E. G. Retzman. 1923. *Pub. Carnegie Inst. Wash.* 324. [12] Alfredson, B. V., and J. F. Sykes. 1943. *J. Agr. Res.* 65 61. [13] Alfredson, B. V., and J. F. Sykes. 1940. *Proc. Soc. Exp. Biol.*, N. Y. 43:580. [14] Davies, F., and E. T. B. Francis. 1950. *J. Anat.*, Lond. 86 302. [15] Buchanan, F. 1910. *Science Progr.*, Lond. 5 60. [16] Ellinger, R. 1894. *Vergleichende physiol. Untersuchungen über die normale Pulsfrequenz der Haustiere*. Thesis. [17] Oppenheimer, E. 1922. *Zschr. ges. exp. Med.* 28:96. [18] Buchanan, F. 1910. *J. Physiol.*, Lond. 40 43. [19] Forbes, A. 1921. *Am. J. Physiol.* 55 385. [20] Evans, G. H. 1910. *Elephants and their diseases*. Rangoon. [21] Gley, E., and A. Quinquand. 1922. *Arch. nferi. physiol.* 7 392. [22] Koenigsfeld, H., and E. Oppenheimer. 1922. *Zschr. ges. exp. Med.* 28 106. [23] Schott, E. 1920. *Arch. exp. Path.*, Lpz. 37:309. [24] Benedict, F. G. 1925. *J. Biol. Chem.* 20 301. [25] Süßel, H. 1910. *Pflügers Arch.* 135 249. [26] Lombard, E. A. 1952. *Am. J. Physiol.* 171 189. [27] Harrington, D. W. 1898. *Ibid.* 1:383. [28] Prosser, C. L., et al. 1950. *Animal physiology* W. B. Saunders, Philadelphia p. 552. [29] Sisson, S. 1914. *The anatomy of the domestic animals*. Ed. 2. W. B. Saunders, Philadelphia. [30] Scholander, P. F., and L. Irving. 1941. *J. Cellul. Physiol.* 17 169. [31] Hecht, S. 1916. *Zschr. ges. exp. Med.* 4 259. [32] De Waart, A., and C. J. Storm. 1934. *Acta brevia neerl.* 4 130. [33] Buchanan, F. 1908. *J. Physiol.*, Lond. 37 69. [34] Hunt, R., and D. W. Harrington. 1897. *J. Exp. M.* 2 712. [35] Hering, H. E. 1895. *Pflügers Arch.* 60 425. [36] MacWilliams, J. A. 1893. *Proc. R. Soc.*, Lond. B53 464. [37] Sprague, H., and M. Rappaport. 1941. *Am. Heart J.* 26 662. [38] Grad, B. 1952. *Endocrinology* 50 94. [39] Hoskins, R. G., M. O. Lee, and E. P. Durrant. 1927. *Am. J. Physiol.* 82 621.

43. HEART RATE: VERTEBRATES (Continued)

Part I: MAMMALS (Concluded)

- [40] Irving, L., P. F. Scholander, and S. W. Grinnell. 1941. J. Cellul. Physiol. 18:283. [41] Morrison, P. R., A. R. Dawe, and F. A. Ryser. 1953. Fed. Proc., Balt. 12:100. [42] Johnson, G. E. 1928. J. Exp. Zool. 50:15. [43] White, P. D., R. L. King, and J. L. Jenks. 1953. N. England J. M. 248:69.

Part II: BIRDS

	Species	Weight g	Heart Rate beats/min	Reference
	(A)	(D)	(C)	(D)
1	Blackbird (<i>Turdus merula</i>)	58	390-560	1, 2
2	Bramblefinch (<i>Fringilla montifringilla</i>)	22	900-920	1, 2
3	Buzzard (<i>Buteo buteo</i>)	658	301(206-351)	3, 4
4	Buzzard (<i>B. vulgaris</i>)		282	1, 5
5			300	1
6	Canary (<i>Serinus canarius</i>)		690(570-840)	6
7		16	514 ¹	7
8		16	1000 ²	7
9	Cardinal (<i>Richmondia cardinalis</i>)	40	375 ¹	7
10		40	800 ²	7
11	Cassowary (<i>Casuarus casuarus</i>)	60,000	70	1
12	Catbird (<i>Dumetella</i> sp.)		330(318-354)	8
13	Chaffinch (<i>Fringilla coelebs</i>)	15	700	1, 2
14	Chickadee, black-capped (<i>Parus atricapillus</i>)	12	480 ¹	7
15		12	1000 ²	7
16	Chicken (<i>Gallus bankiva</i>)	1000	354	5
17		1980	312(178-458)	3, 4
18		1920-3120	346(304-390)	3, 9
19	Chicken (<i>Gallus gallinaceus</i>)		(150-180)	10-12
20			(200-400)	10, 13
21	Crane, Stanley (<i>Arthropoides paradisea</i>)	11,000	120	1
22	Crow, hooded (<i>Corvus cornix</i>)	360	378(312-492)	3, 4
23	Dove, mourning (<i>Zenaidura macroura</i>)	130	135 ¹	7
24		130	570 ²	7
25	Duck, domestic (<i>Anas platyrhynchos</i>)	2060	240	1, 14
26		2304	212(133-268)	1, 4
27	Duck, wild (<i>Anas boschas</i>)		(120-200)	15
28			40 ³	15
29		770-1000	190(185-195)	16
30		785	317(229-420)	3, 4
31	Falcon (<i>Falco peregrinus</i>)		347	1, 4
32	Falcon (<i>F. tinnunculus</i>)	159	367	1, 4
33	Goldfinch (<i>Carduelis elegans</i>)	16	920(914-925)	1, 14
34	Goose (<i>Anser</i> sp.)	2800	144	1, 5
35		4000	80	1, 17
36	Goshawk (<i>Astur palmarius</i>)	960	347	3, 4
37	Greenfinch (<i>Chloris chloris</i>)		710	1, 2
38		25-27	773(703-848)	1, 9, 14
39	Gull (<i>Larus canus</i>)	388	401(360-483)	3, 4
40	Hummingbird (<i>Archiocheilus colubris</i>)	4	615 ¹	7
41	Jackdaw (<i>Corvus monedula</i>)	140	342(326-358)	3, 4
42	Kingfisher (<i>Alcedo isipda</i>)	42	440	3, 9, 14
43	Kite (<i>Milvus milvus</i>)	950	258	3, 4
44	Ostrich (<i>Struthio camelus</i>)	60,000	65(60-70)	1
45	Parrot (<i>Psittacus erithacus</i>)	430	320	1, 3, 14
46	Pigeon (<i>Columba</i> sp.)		(150-300)	3
47			130(120-141)	1, 9
48			300(220-360)	3
49			200	9
50		237	244	3, 4
51		240-370	185(141-225)	3, 9, 14
		20	890	1, 2
			570(520-620)	5
		341	380(352-440)	1, 3, 4, 16

/1/ Basal rate. /2/ Maximum rate on nest. /3/ Apneic.

43. HEART RATE. VERTEBRATES (Continued)

Part II. BIRDS (Concluded)

	Species	Weight	Heart Rate	Reference
	(A)	(B)	(C)	(D)
55	Sparrow, chipping (<i>Spizella passerina</i>)	12	440 ¹	7
56		12	1060 ²	7
57	Sparrow, English (<i>Passer domesticus</i>)	20	460	3
58		21-27	(640-910)	1, 2
59		28	804(745-850)	3, 9, 14
60		28	350 ¹	7
61		28	902 ²	7
62	Sparrow, song (<i>Melospiza melodia</i>)	20	450 ¹	7
63		20	1020 ²	7
64	Starling (<i>Sturnus vulgaris</i>)		388(375-400)	6
65	Swan (<i>Cygnus alar</i>)		257	18
66	Thrasher, brown (<i>Toxostoma</i> sp.)		278(270-294)	8
67	Titmouse, great (<i>Parus major</i>)	14	870	1, 2
68	Towhee (<i>Pipilo erythrophthalmus</i>)	40	445 ¹	7
69		40	810 ²	7
70	Turkey, domestic (<i>Meleagris gallopavo</i>)	8750	93	3, 4
71	Vulture (<i>Gyps fulvus</i>)	8310	199	3, 4
72	Warbler, yellow (<i>Dendroica aestiva</i>)		480 ¹	8
73			832 ²	8
74	Wren, house (<i>Troglodytes aedon</i>)	11	450 ¹	7
75		11	950 ²	7

/1/ Basal rate. /2/ Maximum rate on test.

Contributors: (a) Robb, Jane Sands, (b) Garb, Solomon, (c) Kruta, Vladislav, and Václav Seliger, (d) Woodbury, Robert A., (e) Walker, Sheppard M., (f) Nardone, Roland M.

References [1] Clark, A. J. 1927. Comparative physiology of the heart. Macmillan, New York. [2] Oppenheimer, E. 1922. Zschr. ges. exp. Med. 28:96. [3] Lehmann, G. 1925. Tabulae biol., Berl. Bd. 1, p. 157. [4] Sübel, H. 1910. Pflügers Arch. 135:249. [5] Vierordt, K. 1877. Grundriss der Physiologie der Menschen. Ed. 5 p. 162 [6] Woodbury, R. A., and W. T. Hamilton. 1937. Am. J. Physiol. 119:663. [7] Odum, E. P. 1945. Science 101:153. [8] Ohm, E. P. 1941. Ecol. Monogr. 11(3):299. [9] Buchanan, F. 1909. J. Physiol., Lond. 38:62. [10] Dukes, H. H. 1947. The physiology of domestic animals. Cornell University Press, Ithaca. [11] Prosser, C. L., et al. 1950. Animal physiology. W. B. Saunders, Philadelphia. p. 552. [12] McNally, E. H. 1941. Poultry Sc. 20:266 [13] Henderson, E. W., and E. W. Hathaway. 1943. Ibid. 22:44. [14] Buchanan, F. 1910. Science Progr., Lond. 5:60 [15] Koppányi, T., and N. Kleitman. 1927. Am. J. Physiol. 82:672. [16] Buchanan, F. 1913. J. Physiol., Lond. 47:4. [17] Mosso, A. 1901. Arch. Ital. biol. 35:21. [18] Davies, P., and E. T. B. Francis. 1950. J. Anat., Lond. 86:302.

Part III: REPTILES AND AMPHIBIANS

	Species	Specification	Heart Rate	Reference
	(A)	(B)	(C)	(D)
1	Blindworm (<i>Anguis fragilis</i>)		64	1
2	Crocodile (<i>Crocodylus</i> sp.)		(10-60)	2, 3
3		72 g	(22-47)	2
4		150°C	30	4
5		23 50°C	70	1
6	Lizard (<i>Lacerta viridis</i>)	15-19 g	(60-66)	2
7	Lizard, sand (<i>L. agilis</i>)		50	5
8	Ringsnake (<i>Coluber natrix</i>)		68	1
9		169 g	60	1, 6
10	Tortoise (<i>Emys lotaria</i>)		(23-43)	1, 2
11			(9-60)	1
12			(16-36)	5

43. HEART RATE: VERTEBRATES (Continued)

Part III: REPTILES AND AMPHIBIANS (Concluded)

Species	Specification	Heart Rate beats/min	Reference
(A)	(B)	(C)	(D)
13 Tortoise (<i>Pseudomys rugosa</i>)		(21-44)	1
14 Tortoise (<i>Testudo denticulata</i>)	5000 g; 28-30°C	17(11-37)	7
15 Tortoise (<i>T. graeca</i>)	135 g	(10-20)	1, 3
16 Tortoise (<i>Thalassochelys</i> sp)		11	1
17 Viper (<i>Vipera berus</i>)		40	1
18 Frog (<i>Rana pipiens</i>)		(40-50)	1
19 Frog (<i>R. pipiens</i> , <i>R. temporaria</i>)	20°C	(37.5-60)	1
20 Frog (<i>R. pipiens</i> , <i>R. temporaria</i>)	22°C	(5-9)	8
21 Salamander (<i>Salamandra</i> sp)		(34-39)	8
22 Salamander (<i>Salamandra</i> sp)		(30-40)	9
23 Toad (<i>Bufo</i> sp)		(40-50)	1
24		48	10

Contributors: (a) Robb, Jane Sands, (b) Kruta, Vladislav, and Václav Seliger, (c) Garb, Solomon, (d) Walker, Sheppard M., (e) Nardone, Roland M., (f) Woodbury, Robert A.

References: [1] Lehmann, G. 1925. *Tabulae biol.*, Berl. Bd 1, p. 136. [2] Buchanan, F. 1909. *J. Physiol.*, Lond. 39:25. [3] Gaskell, W. H. 1883. *Ibid.* 4:43. [4] Rubner, M. 1924. *Biochem. Zschr.* 148:222, 268. [5] Laurens, H. 1920. *Pflügers Arch.* 182:50. [6] Hoffmeister, F. 1889. *Ibid.* 44:360. [7] Benedict, F. G. 1932. *Physiology of large reptiles*. Carnegie Institution, Washington. [8] Barcroft, J., and J. J. Izquierdo. 1931. *J. Physiol.*, Lond. 71:145. [9] Riemüller, J. 1932. *Pflügers Arch.* 230:782. [10] Hoffmeister, F. 1888. *Arch. ges. Physiol* 44:366.

Part IV. FISHES

Species	Heart Rate beats/min	Reference
(A)	(B)	(C)
1 Barbel (<i>Barbus fluviatilis</i>)	(35-58)	1
2 Bass, largemouthed (<i>Micropterus salmoides</i>)	(36-90) ¹	2
3 Blenny (<i>Zoarces viviparus</i>)	20(5-50)	3
4 Bullhead (<i>Ameiurus</i> sp)	(71-86)	1
5 Bullhead (<i>Ameiurus</i> sp)	22(5-50)	3
6 Carp (<i>Cyprinus carpio</i>)	(40-60)	1
7 Chub (<i>Leuciscus dobula</i>)	75(72-78)	4
8 Chub (<i>Leuciscus dobula</i>)	18 ²	5, 6
9 Cod (<i>Gadus morrhua</i>)	(24-40) ³	7
10 Cod (<i>Gadus morrhua</i>)	(48-60)	1
11 Dogfish, spiny (<i>Acanthias vulgaris</i>)	(40-50)	1
12 Dogfish, spotted (<i>Scylium canicula</i>)	65 ⁴	5, 8
13 Dogfish, spotted (<i>S. catulus</i>)	(39-48)	1
14 Dogfish, spotted (<i>S. catulus</i>)	(16-50)	5
15 Dragonet (<i>Callionymus lyra</i>)	44 ⁵	9
16 Dragonet (<i>Callionymus lyra</i>)	(60-84)	1
17 Eel, common (<i>Anguilla</i> sp)	(39-68)	5
18 Eel, common (<i>Anguilla</i> sp)	(20-30)	5
19 Eel, common (<i>Anguilla</i> sp)	(1-2) ⁶	5
20 Eel, common (<i>Anguilla</i> sp)	(48-56)	1
21 Eel, common (<i>Anguilla</i> sp)	(46-68) ⁷	2
22 Eel, common (<i>Anguilla</i> sp)	(33-50)	1
23 Eel, marine (<i>Conger conger</i>)	(36-40)	5
24 Goldfish (<i>Carassius</i> sp)	(62-86)	1
25 Gurnard (<i>Trigla hirundo</i>)	(36-60)	5
26 Herring (<i>Telestes multicellus</i>)	(52-66)	1
27 Perch (<i>Perca fluviatilis</i>)	(38-54)	1
28 Pike (<i>Esox lucius</i>)	(30-42)	5
29 Platice (<i>Pleuronectes platessa</i>)	(54-76)	1

1/ 120 g, 20°C. 2/ 24 cm. 3/ 30 cm, 18°C. 4/ 26°C. 5/ 1000 g; 16°C. 6/ Hibernating. 7/ 36 cm

43. HEART RATE: VERTEBRATES (Concluded)

Part IV. FISHES (Concluded)

Species	Heart Rate beats/min	Reference
(A)	(B)	(C)
30 Pogge (<i>Agonus cataphractus</i>)	(81-90)	1
31 Ray, electric (<i>Torpedo</i> sp)	(16-50)	5
32 Ray, thornback (<i>Raja clavata</i>)	(38-40)	1
33 Roach (<i>Rutilus rutilus</i>)	(50-71)	1
34 Rockling (<i>Motella mustela</i>)	(64-82)	1
35 Scorpion fish (<i>Scorpaena scrofa</i>)	(11-24)	1
36 Sculpin (<i>Cottus scorpius</i>)	(55-74)	1
37 Shark, gray (<i>Carcharias</i> sp)	30 ⁸	9, 10
38	18 ⁹	9, 10
39 Skate (<i>Raja</i> sp)	(16-50)	5
40 Skate (<i>R. batis</i>)	(21-33)	1
41 Stickleback (<i>Gasterosteus aculeatus</i>)	(60-100)	9, 11
42 Tench (<i>Tinca tinca</i>)	(31-42)	1
43 Tope (<i>Galeus canis</i>)	(34-40)	1
44 Trout (<i>Salmo trutta</i>)	(30-46)	1
45 Wrasse (<i>Labrus mixtus</i>)	(40-81)	1

18/ 100 cm 19/ 200 cm.

Contributors: (a) Kruta, Vladislav, and Václav Seliger, (b) Woodbury, Robert A., (c) Robb, Jane Sands, (d) Walker, Sheppard M., (e) Nardone, Roland M.

References [1] Biellig, W. 1931. *Zachr. vergl. Physiol.* 15: 488. [2] Koff, W. M. 1908. *Physiologische Arch.* 122: 37. [3] Hart, I. J. 1944. *Proc. Florida Acad. Sc.* 7: 221. [4] Woodbury, R. A., and W. T. Hamilton. 1937. *Am. J. Physiol.* 119: 661. [5] Lehmann, G. 1925. *Tabulae biol.*, Berl., Bd 1, p. 156. [6] Brünings, W. 1899. *Physiologische Arch.* 75: 599. [7] Thesen, J. E. 1896. *Arch. zool. exp., Par.* 3(3): 122. [8] Bottazzi, F. 1901. *Zbl. Physiol.*, Wien 14: 665. [9] Clark, A. J. 1927. *Comparative physiology of the heart.* Macmillan, New York. [10] Lyon, E. P. 1926. *J. Gen. Physiol.* 8: 279. [11] Ruhmer, M. 1924. *Biochem. Zachr.* 148: 222, 268.

44. HEART RATE: INVERTEBRATES

Heart rate varies with species, sex, age, size, environment, and temperature. Values in parentheses are ranges, estimate "c" (cf. Introduction)

Part I. TUNICATES, MOLLUSKS, CRUSTACEANS, ANNELIDS

Species	Specification	Temp. °C	Heart Rate beats/min	Reference
(A)	(B)	(C)	(D)	(E)
Tunicates				
1 Appendicularia sp	0.25 cm		250	1
2 Ascidia depressa	8 cm		(31.3-33.6)	2
3 Clona intestinalis			(17-32)	3
4 Clavellina lepadiformis			(30-50)	4
5 Cyclosalpa pinnata	8 cm		(26-30)	5
6 Molgula manhattanensis		21-22	(30-50)	5
7 Perophora annectens			45.1	6
8 Phallusia mammillata	20 cm		(8-9)	7
9 Salpa africana			(26-37)	5
10 S. africana maxima		15-18	(21-34)	3
11 S. bicaudata		19	(13-32)	8
12 S. democratica	6 cm		107	5
13 S. fusiformis	1 cm		(61-69)	8
14 S. pinnata	4 cm		108	3
Mollusks				
15 Acroloxus lacustris (fresh-water snail)	0.0004 g	20	68.6	9
16	0.0015 g	20	57.1	9
17 Anodonta cygnea (fresh-water mussel)			(12-15)	10
18			(26-29)	11

Part I: TUNICATES, MOLLUSKS, CRUSTACEANS, ANNELIDS (Continued)

Species		Specification	Temp. °C	Heart Rate beats/min	Reference
(A)		(B)	(C)	(D)	(E)
Mollusks (continued)					
19	<i>Anodonta cygnea</i> (fresh-water mussel)			(4-6)	12
20	(concluded)			(1.6-3.5)	13
21				(3-6)	14
22		1.96 g	20	28.1	9
23		12.1 g	20	19.7	9
24		25.5 g	20	17.4	9
25		42 g	20	14.6	9
26		64 g	20	13.0	9
27		87 g	20	12.7	9
28	<i>A. fluviatilis</i>			4(2-6)	12
29	<i>A. mutabilis</i>	Empty		(6-7)	15
30	<i>Anomia simplex</i>			52	16
31	<i>Aplysia</i> sp	Empty		(12-20)	4
32		Exposed		(8-15)	4
33		Perfused		(15-35)	4
34	<i>A. deplians</i>		23-24	(33-34)	17
35		Exposed		(7-8)	17
36	<i>A. limacina</i>			(20-35)	18
37		347 g	20	32.3	9
38		870 g	20	28.0	9
39	<i>Ariolimax</i> sp			(35-40)	4
40	<i>Arion</i> sp			(20-36)	19
41	<i>A. empiricorum</i> (land snail)	16.6 g	20	48.8	9
42	<i>Bulla</i> sp			(10-18)	4
43	<i>Cardium echinatum</i>	Empty		(6-7)	15
44	<i>Carinaria meditor</i>			54	4
45	<i>Cepaea memerosa</i>	0.24 g	20	65.4	9
46		0.92 g	20	63.0	9
47		2.05 g	20	52.9	9
48	<i>Cryptochiton</i> sp			(5-7)	4
49	<i>Cyclos cornea</i>			(40-50)	20
50	<i>Dreissensia polymorpha</i>	0.1-0.2 cm		(54-60)	21
51		2 cm		(30-35)	21
52		0.00065 g	20	60	9
53		0.67 g	20	35	9
54	<i>Eledone</i> sp			(35-38)	22
55				(25-30)	23
56		Perfused in situ		(20-25)	24
57	<i>Fulgar carica</i>			(0.8-1.0)	25
58	<i>Haliotis tuberculata</i>			(40-50)	26-28
59	<i>Helix hortensis</i>			40	23
60	<i>H. pomatia</i> (land snail)			(40-50)	4
61				(30-35)	5
62				(20-40)	10
63				(53-55)	4, 5
64		In situ, empty		(10-15)	4
65		Suspended, empty		(35-40)	4
66		Suspended, filled		(50-60)	4
67		0.123 g	20	60.3	9
68		0.37 g	20	68.0	9
69		0.63 g	20	47.6	9
70		8.2 g	20	42.4	9
71		14.3 g	20	37.9	9
72		24.8 g	20	35.2	9
73		36.7 g	20	37.1	9
74		At same arterial and venous pressure	20	25.3	29
75		At peak efficiency	20	22.4	29
76	<i>Ischnochiton</i> sp			(15-25)	4, 30
77	<i>Limax agrestis</i> (land snail)	0.213 g	20	90.6	9
78	<i>Limnaea auricularia</i> (fresh-water snail)	0.1-0.2 cm		100	21, 31
79		2 cm		(42.5-46.0)	21, 31
80		0.0008 g	20	112	9
81		0.66 g	20	46	9

44. HEART RATE INVERTEBRATES (Continued)

Part I: TUNICATES, MOLLUSKS, CRUSTACEANS, ANNELIDS (Continued)

Species		Temp. °C	Heart Rate beats/min	Reference
(A)	(B)	(C)	(D)	(E)
Mollusks (continued)				
82	<i>Limnaea stagnalis</i> (fresh-water snail)	20	55.4	9
83		20	21.0	9
84	<i>Loligo</i> sp		60	4
85			(70-80)	4
86	<i>Lucapina</i> sp	20	53.6	9
87	<i>Melanoides tuberculata</i>	20	39.4	9
88			(20-30)	4
89	<i>Monterelina</i> sp		(2-61)	32
90	<i>Murex trunculus</i>	20	32.9	9
91	<i>Musculium</i> sp	20	35.5	9
92		20	27.5	9
93			14	4
94	<i>Mya arenaria</i>		(5-10)	10
95			(15-25)	27
96	<i>Mytilus edulis</i>		10	4
97		20	49.2	9
98			(5-7)	4
99	<i>Natica</i> sp		(35-40)	33
100	<i>Octopus vulgaris</i>		(8-18)	4
101			(12-20)	4
102			(49-53)	33
103			(53-59)	33
104			(25-30)	27
105	<i>Ostraea edulis</i>		50	34
106	<i>Pecten jacobaeus</i>		22	35
107			36	36
108	<i>Phyllurhoe</i> sp	20	65.2	9
109	<i>Physa acuta</i>	20	65.7	9
110		20	58.2	9
111		20	54.8	9
112		20	34.4	9
113			(60-75)	21
114	<i>Planidium</i> sp	20	74	9
115		20	58	9
116			(40-45)	23
117	<i>Planorbis corneus</i> (fresh-water snail)	20	56.7	9
118		20	43.3	9
119		20	45.0	9
120		20	39.0	9
121		20	35.9	9
122		20	32.1	9
123		20	26.0	9
124		20	25.2	9
125		20	29.6	9
126		20	21.3	9
127		20	20.3	9
128		20	13.6	9
129		20	15.9	9
130		20	25.7	9
131	<i>P. marginatus</i> (fresh-water snail)	20	(20-30)	4
132	<i>Pleurobranchia</i> sp		67(50-80)	3
133	<i>Pterotrachea</i> sp		50	37
134			48(36-57)	38
135		13-14	40	39
136	<i>Sepia officinalis</i>		(18-30)	40, 41
137			26	5
138	<i>Succinea</i> sp			
139	<i>S. putris</i>	20	74.2	9
140		20	71.7	9
141		20	69.1	9
142	<i>Sycotypus</i> sp		(12-20)	30
143	<i>Theodoxus fluviatilis</i>	20	80.2	9
144		20	65.6	9
145		20	63.8	9
146		20	49.8	9

44. HEART RATE: INVERTEBRATES (Continued)

Part I: TUNICATES, MOLLUSKS, CRUSTACEANS, ANNELIDS (Continued)

Species		Specification	Temp. °C	Heart Rate beats/min	Reference
(A)		(B)	(C)	(D)	(E)
Mollusks (concluded)					
147	<i>Triopa</i> sp			(30-40)	4
148	<i>Unio pictorum</i>	5.3 g	20	25.7	9
149	<i>Valvata piscinalis</i>			100	21, 31
Crustaceans					
150	<i>Alpheus dentipes</i>	0.16 g	20	181	42
151	<i>Apherusa bispinosa</i>	0.02 g	20	335	42
152	<i>Asellus aquaticus</i>	0.0008 g	20	310	42
153		0.0061 g	20	242	42
154		0.035 g	20	(177-182)	43
155	<i>Astacus fluviatilis</i> (fresh-water crab)			(30-60)	5, 44, 45
156		Empty		(10-20)	46
157		Filled		60	46
158	<i>A. marinus</i> (crayfish)			50(30-87)	47
159	<i>Calinassa subterranea</i>	1.8 g	20	136	42
160	<i>Callinectes sapidus</i>		22-23	(25-84)	25
161	<i>Cancer pagurus</i> (hermit crab)			(90-120)	4
162		Empty		(15-30)	4
163		Perfused		(15-50)	4
164		200 g	20	105	42
165	<i>Caprella</i> sp	0.04 g	20	140	42
166	<i>Daphnia</i> sp (water flea)			140	48
167				(200-250)	49
168				(200-380)	50
169			20-22	228(188-289)	3
170	<i>D. magna</i>		20	240	51
171	<i>D. pulex</i>	0.000025 g	20	486	42
172		0.00024 g	20	470	42
173		0.0013 g	20	444	42
174		0.0011 g	20	490.6	43
175		0.0008 g	20	(381-418)	42
176	<i>Dromia vulgaris</i>	150 g	20	67	42
177		250 g	20	64	42
178	<i>Eriphia spinifrons</i>	75 g	20	90	42
179	<i>Gammarus pulex</i>	0.0011 g	20	339	42
180		0.035 g	20	258	42
181	<i>Hippolyte</i> sp	0.028 g	20	216	42
182	<i>Homarus vulgaris</i>	450 g	20	50	42
183	<i>Hyale camptonyx</i>	0.02 g	20	285	42
184	<i>Leander serratus</i>	0.77 g	20	146	42
185	<i>Limnadia</i> sp			(130-170)	52
186	<i>Limulus polyphemus</i> (king crab)			20(18-28)	4
187				15	25
188	<i>Lysmata seticaudata</i>	0.56 g	20	156	42
189	<i>Maia</i> sp			25	53
190				46	54
191	<i>M. verrucosa</i>	79 g	20	100	42
192	<i>Mysis</i> sp		19-20	260(140-320)	3
193	<i>M. lamorn</i>	0.02 g	20	285	42
194	<i>Palinurus</i> sp	Empty		35(30-50)	4
195	<i>P. vulgaris</i>	220 g	20	107	42
196	<i>Pontonia custos</i>	0.81 g	20	167	42
197	<i>Porcellana</i> sp	Larva	20	170	3
198	<i>Potamobius astacus</i>	30 g	20	100	42
199	<i>Sparassus virescens</i>	Moderate activity		120	55
200		Intense activity		(200-230)	55
201	<i>Squilla eusebia</i>		26	240	3
202	<i>S. mantis</i>	27 g	20	90	42
Annelids					
				7(6-8)	56
				6	57
				17(15-20)	35
				8	35
			7	4.6	58
			35	50	58

44. HEART RATE: INVERTEBRATES (Continued)

Part I. TUNICATES, MOLLUSKS, CRUSTACEANS, ANNELIDS (Concluded)

Contributors (a) Kruta, Vladislav, and Václav Seliger, (b) Robb, Jane Sands, (c) Nardone, Roland M., (d) Jones, Jack Colvard

- References: [1] Moss, E. L. 1871. Tr. Linn. Soc., Lond. 27:299. [2] Von Skramlik, E. 1938. Pubb. Staz. zool. Napoli 17. [3] Knoll, P. 1893. Sitzber. Akad. Wiss. Wien 102:387. [4] Carlson, A. J. 1906. Am. J. Physiol. 16:47. [5] Lehmann, G. 1925. Tabulae biol., Berl., Bd 1. [6] Ritter, W. E. 1893. Proc. California Acad. Sc. 2:37. [7] Koehnlein, H. 1933. Pubb. Staz. zool. Napoli 13:114. [8] Hykes, O. V. 1925. Pub. biol. Ec. vét., Brno 4:113. [9] Schwartzkopff, J. 1956. Verhandlungen der deutschen zoologischen Gesellschaft. Hamburg, p. 463. [10] Yung, E. 1881. Arch. zool. exp., Par. 9:421. [11] Baker, F. 1897. J. Cincinnati Soc. Nat. Hist. 19:2. [12] Willem, V., and A. Minne. 1898. Mém. couron. Acad. méd. Belgique 57:4. [13] Koch, W. 1917. Pflügers Arch. 166:281. [14] Ten Cate, J. 1923. Arch. néerl. physiol. 8:43. [15] Nevinkoff. 1907. Physiol. russe 5:15. [16] Navez, A. E. 1936. Mém. Mus. Hist. nat. Belgique 3:701. [17] Bottazzi, F., and P. Enriques. 1900. Arch. Ital. biol. 34:111. [18] Straub, W. 1901. Pflügers Arch. 86:504. [19] Niertrauss, V. E. 1907. Rauwolfine als Harigut Dissertation Utrecht. [20] Garkewitz, S. 1926. Arch. internat. physiol., Liège 26:229. [21] Ludwig, W. 1937. Zachr. vergl. Physiol. 24:319. [22] Fuchs, S. 1895. Pflügers Arch. 60:188. [23] Von Skramlik, E. 1941. Erg. Biol. 18:88. [24] Frédéricq, H. 1939. Bull. Acad. Belgique 25:611. [25] Dubuisson, M., and A. M. Monner. 1931. Arch. internat. physiol., Liège 34:180. [26] Von Skramlik, E. 1932. Herzmuskel und Extrazelle. Jena [27] Von Skramlik, E. 1929. Pflügers Arch. 221:503. [28] Von Skramlik, E. 1929. Zachr. vergl. Physiol. 9:553. [29] Schwartzkopff, J. 1953. Experientia, Basel 9:428. [30] Carlson, A. J. 1905. Am. J. Physiol. 13:396. [31] Ludwig, W. 1938. Arb. biol. Forsch. Inst. 10:93. [32] Julien, A., and G. Morin. 1932. J. physiol. path. gén. 30:900. [33] Frédéricq, H. 1914. Arch. internat. physiol., Liège 14:126, 132. [34] Dogiel, J. 1877. Arch. mikr. Anat. Entw. 4:59. [35] Clark, A. J. 1927. Comparative physiology of the heart. Macmillan, New York. [36] Snyder, C. D. 1906-07. Am. J. Physiol. 17:350. [37] Glasser, O. 1925. J. Gen. Physiol. 9:269. [38] Rywosch, D. 1905. Pflügers Arch. 109:355. [39] Bert, P. 1867. Mém. Soc. sc. Bordeaux 5:115. [40] Kruta, V. 1936. C. rend. Soc. biol. 122:582. [41] Kruta, V. 1937. Cas. lek. Cesk. 76:1328, 1345. [42] Schwartzkopff, J. 1955. Experientia, Basel 11:323. [43] Schwartzkopff, J. 1955. Biol. Zbl. 74:480. [44] Hoffman, P. 1911. Arch. Anat. Physiol., Lpz. p. 135. [45] Prosser, C. L., et al. 1950. Comparative physiology of the heart. W. B. Saunders, Philadelphia. p. 552. [46] Hoffman, P. 1912. Zachr. Biol. 59:300. [47] Von Brücke, E. T., and Y. Satake. 1912. Zachr. allg. Physiol. 14:312. [48] Pickering, J. W. 1895. J. Physiol., Lond. 17:356. [49] Von Leyding, F. 1860. Naturgeschichte der Daphniden. Tübingen. [50] Hykes, O. V. 1925. C. rend. Soc. biol. 95:58. [51] Proksova, M. 1950. Acta Acad. Sc. nat. Moravo-Silesiacae 22:31. [52] Lereboullet. 1850. Mém. Soc. Mus. Hist. nat. Strasbourg 4. [53] Von Brücke, E. T. 1925. In H. Winterstein, ed. Handbuch der vergleichenden Physiologie. Jena v. 1, p. 827. [54] Pollmann, O. 1915. Biol. Zbl. 65:398. [55] Wagner, W. 1893. Ann. Sc. nat. 15:311. [56] Carlson, A. J. 1908. Am. J. Physiol. 22:353. [57] Gaskell, W. H. 1914. Proc. R. Soc., Lond. B205:163. [58] Rogers, C. G. 1911. Am. J. Physiol. 28:81.

Part II INSECTS

Insect	Specification	Temp. °C	Heart Rate beats/min	Reference
(A)	(B)	(C)	(D)	(E)
1 <i>Campodea</i> sp	Thysanura			
2 <i>Machilis</i> sp		16-18	(79-81)	1
		20	(150-160)	2
3 <i>Clofen dipterum</i>	Nymph			
4 <i>C. rhodani</i>	Nymph	15-16	(31-34)	3
		15-16	(77-112)	3
5 <i>Anax junius</i>	Odonata			
		28	60(28-100)	4
6 <i>Blatta orientalis</i>	Orthoptera			
7	Nymph to adult			
8	Nymph to adult			
9		18	(75-105) ¹	5
			67.2(50-107) ¹	6
10 <i>Blattella germanica</i>	Nymph		(60-80)	6
11 <i>Dermapteron andersoni</i>	Nymph		(35-65) ¹	7
12 Grasshopper		25-27	120	8
13 <i>Locusta migratoria danica</i>	Nymph to adult		(20-28)	9
14	Male	25	60 ¹	10
15	Male	26	(25-75) ¹	11
16	Female	29	(80-120) ¹	11
17 <i>Melanoplus differentialis</i>	Nymph to adult	29	(80-130) ¹	11
18	Nymph to adult	25	63 ¹	12
19 <i>Periplaneta americana</i>	Nymph, male	21-26	(40-70)	13
20	Nymph, female		94	14
	Nymph to adult		100	14
			(80-110) ¹	15

¹ Heart exposed by dissection and examined under physiological saline

44. HEART RATE: INVERTEBRATES (Continued)

Part II: INSECTS (Continued)

Insect	Specification	Temp. °C	Heart Rate beats/min	Reference
(A)	(B)	(C)	(D)	(E)
Orthoptera (concluded)				
21 <i>Periplaneta americana</i>	Nymph to adult		(90-130) ¹	16
22 (concluded)	Nymph to adult	20-26	117.3(99-147)	17
23	Nymph to adult	26-27	(90-100) ¹	18
24	Nymph to adult	29	49(18-70) ¹	19
25	Adult	18	72.9(22-150) ¹	20
26	Adult, male		99	14
27	Adult, female		99	14
28 <i>Stenopelmatus longispina</i>	Nymph	25	41(30-52) ¹	21
Mallophaga				
29 <i>Menopon pallidum</i>			(112-120)	22
Anoplura				
30 <i>Pediculus</i> sp			(30-48)	23
Hemiptera				
31 <i>Belostoma</i> sp			(34-35)	24
32 <i>Nezara viridula</i>		27	100	25
33 <i>Notonecta</i> sp			70	24
34 <i>Rhodnius prolixus</i>	Nymph	25-27	30	8
Homoptera				
35 <i>Aphis</i> sp		18	(68-80)	6
36 <i>Macrostiphum tulipae</i>		23	83	26
Neuroptera				
37 <i>Chrysopa</i> sp		18	(50-60)	6
Trichoptera				
38 <i>Phryganid</i>	Larva	25	26	26
Lepidoptera				
39 <i>Bombyx mori</i>			58	27
40	Larva		(20-67)	28
41	Larva		(66-76.8)	29
42	Larva	25	54	30
43	Larva, stage 2		70	31
44	Larva, stage 3		65.7	31
45	Larva, stage 4		54.8	31
46	Larva, stage 5		45.7	31
47	Larva, stage 5	27	55.9	32
48	Larva, calm		(44-66)	31
49	Larva, excited		94	31
50	Larva, spinning		55	31
51	Adult, resting		(40-50)	33
52	Adult, active		(110-140)	33
53 <i>Celerio euphorbiae</i>			39	34
54 <i>Cerura virula</i>		21-27	(49-88)	33
55 <i>Colias eurytheme</i>	Larva		(60-120)	6
56 <i>C. hyale</i>			(54-66)	35
57 <i>Dendrolimus pini</i>	Pupa		(50-60)	6
58 <i>Ephestia kuehniella</i>	Pupa		(6-11)	36
59 <i>Galleria mellonella</i>		25.0-29.5	(60-69)	37
60	Larva, spinning	25	(50-60)	37
61 <i>Parage maera</i>		20-30	(40-80)	6
62 <i>Pieris brassicae</i>	Larva		29	38
63	Larva, parasitized with <i>Apanteles</i>		6	38
64 <i>Prodenia eridania</i>	Larva	29	58(40-90) ¹	19
65 <i>Saturnia pavonia major</i>	Pupa		(0-39.8)	39
66 <i>Sphinx ligustri</i>	Larva	22	(38-54)	33
67	Pupa		27	34
Coleoptera				
68 <i>Dytiscus marginalis</i>			60 ¹	40
69	Intact		(30-70)	40
70	Isolated fragment		6	40
71	Isolated heart		15	40
72 <i>Lucanus cervus</i>		18	15.7	41
73	Larva	13	14.1	41
74 <i>Oryctes nasicornis</i>		25	18	30
75		24-33	(17-31)	6

/1/ Heart exposed by dissection and examined under physiological saline.

44. HEART RATE. INVERTEBRATES (Continued)

Part II. INSECTS (Continued)

Insect	Specification	Temp. °C	Heart Rate beats/min	Reference
(A)	(B)	(C)	(D)	(E)
Coleoptera (concluded)				
76 <i>Prionus laticollis</i>		29	118	27
77 <i>Tenebrio molitor</i>	Larva		(15-17)	42
78	Larva, postmolt		20	42
79	Pupa		10	42
Hymenoptera				
80 <i>Anthophora retusa</i>	Adult	28	142	6
81 <i>Sphex</i> sp.			(300-354)	27
Diptera				
82 <i>Aedes diversus</i>	Larva	18	50(47.5-53.6)	43
83 <i>A. togoi</i>	Larva	25	48(41.7-52.5)	44
84 <i>Anopheles maculipennis</i>	Adult	20	(79-110)	27
85	Adult, after blood meal	20	109.9	27
86	Adult, digesting	20	91.7	27
87 <i>A. quadrimaculatus</i>	Larva, stage 1	25-27	131.7	45
88	Larva, stage 2	25-27	134.3	45
89	Larva, stage 3	25-27	118.6	45
90	Larva, stage 4	25-27	106.6	45
91	Pupa	25-27	109.1	45
92	Adult, female	25-27	151.2	45
93	Adult, female	25-27	80.7 ^{1, 2}	45
94 <i>Calliphora</i>	Larva	18	120.8	46
95	Larva	15-19	60 ¹	47
96 <i>Chironomus</i> sp.	Larva		(61.5-133.2)	48
97 <i>C. dorsalis</i>	Larva		162	49
98 <i>C. plumosus</i>	Larva	17	90	49
99 <i>Corethra albipes</i>	Larva	27	(19-29)	50
100 <i>C. plumicornis</i>	Larva		(21-24)	51
101	Larva	17-18	(15-16)	52
102	Larva		16.8(12-41) ¹	53
103	Larva, calm		(12-14)	54
104	Larva, active		22	54
105 <i>Culex</i> sp.	Larva		(50-100)	55
106 <i>Culex pipiens</i>	Larva, stage 2		100	56
107	Larva, stage 4		90	56
108	Larva, stage 4		60(56.5-65.5)	43
109	Larva, stage 4	25-27	(110-140)	8
110 <i>Culicoides nubeculosus</i>	Larva		(30-85)	57
111 <i>Drosophila funebris</i>		30	231	58
112	Adult	28-29	235	58
113 <i>Hippoboscra equina</i>		23.5	120	59
114 <i>Mochlonysa velutina</i>	Larva	18	(20-28)	43
115	Larva	21	(27-31.5)	43
116 <i>Oestrus</i>			(40-44)	6
117 <i>Pachyrhina ferruginea</i>	Adult, male	22	128	27
118 <i>Phormia</i>	Larva	18	60 ¹	47
119 <i>Tipula maxima</i>	Larva		24	60
120 <i>T. selene</i>	Larva	18	48	61
121	Pupa	18	(40-60)	61

// Heart exposed by dissection and examined under physiological saline //2/ Standard error on 100 insects, ± 1.8 .

Contributor Jones, Jack Colvard

- References. [1] Grassé, P. P. 1949. *Traité de zoologie*. T. 9. [2] Marten, W. 1919. *Zachr. Morph. Okol. Tiere* 36: 41. [3] Fox, H. M., and B. G. Simmonds. 1933. *J. Exp. Biol.*, Lond. 10: 67. [4] Malud, N. S. R. 1935. *Ann. Entom. Soc. America* 28: 332. [5] Yeager, J. F., A. Hager, and J. M. Straley. 1935. *Ibid.* 28: 256. [6] Kozhanchikov, I. V. 1932. *Bull. Leningrad Inst. Control Farm and Forest Pests* 2: 149. [7] Yeager, J. F., and A. Hager. 1934. *Iowa State Coll. J. Sc.* 8: 391. [8] Jones, J. C. 1953. *Ann. Entom. Soc. America* 46: 366. [9] Douglas, J. R. 1943. *Univ. California Pub. Entom.* 7(10) 207. [10] Walling, E. V. 1906. *J. Exp. Zool.* 3: 621. [11] Yamashiki, T., and T. Ishii. 1950. *Oyo-Kontyu* 5: 155. [12] Jahn, T. L., and B. S. Koel. 1948. *Ann. Entom. Soc. America* 41: 258. [13] Jahn, T. L., F. Crescitelli, and A. B. Taylor. 1937. *J. Cellul. Physiol.* 10: 439. [14] Coon, B. F. 1944. *J. Econom. Entom.* 37: 785. [15] Krigsman, B. J., D. Dresden, and N. E. Berger. 1950. *Bull. Entom. Res.*, Lond. 41: 141. [16] Naidu, M. B. 1955. *Ibid.* 46: 205. [17] Bellemare, E. R., and J. Belcourt. 1955. *Canad. J. Zool.* 33: 175. [18] Orser, W. B., and A. W. A. Brown. 1951. *Ibid.* 29: 54. [19] Yeager, J. F., and J. B. Gahan. 1937.

44. HEART RATE: INVERTEBRATES (Continued) Part II. INSECTS (Continued)

Insect (A)	Specification (B)	Temp. °C (C)	Heart Rate beats/min (D)	Reference (E)
21 <i>Periplaneta americana</i> (concluded)	Orthoptera (concluded) Nymph to adult Nymph to adult Nymph to adult Adult Adult, male Adult, female	20-26 26-27 29 18	(90-130) ¹ 117.3(99-147) (90-100) ¹ 49(18-70) ¹ 72.9(22-150) ¹ 99	16 17 18 19 20 14
22				
23				
24				
25				
26				
27				
28 <i>Stenopelmatus longispina</i>	Nymph		99	14
29 <i>Menopon pallidum</i>	Mallophaga	25	41(30-52) ¹	21
30 <i>Pediculus</i> sp	Anoplura		(112-120)	22
31 <i>Belostoma</i> sp	Hemiptera		(30-48)	23
32 <i>Nezara viridula</i>				
33 <i>Notonecta</i> sp				
34 <i>Rhodnius prolixus</i>	Nymph	27	(34-35)	24
35 <i>Aphis</i> sp	Homoptera		100	25
36 <i>Macrosiphum tulipae</i>		25-27	70	24
			30	8
37 <i>Chrysopa</i> sp	Neuroptera	18	(68-80)	6
		23	83	
38 <i>Phryganid</i>	Trichoptera	18	(50-60)	6
39 <i>Bombyx mori</i>	Lepidoptera	25	26	26
40				
41				
42	Larva		58	27
43	Larva		(20-67)	
44	Larva		(66-76.8)	28
45	Larva, stage 2	25	54	29
46	Larva, stage 3		70	30
47	Larva, stage 4		65.7	31
48	Larva, stage 5		54.8	31
49	Larva, stage 5		45.7	31
50	Larva, calm	27	55.9	31
51	Larva, excited		(44-66)	32
52	Larva, spinning		94	31
53	Adult, resting		55	31
54	Adult, active		(40-50)	31
55			(110-140)	33
56 <i>Celerio euphorbiae</i>			39	33
57 <i>Cerura virula</i>	Larva	21-27	(49-88)	34
58 <i>Colias eurytheme</i>			(60-120)	33
59 <i>Dendrolimus pini</i>	Pupa		(54-66)	6
60 <i>Ephestia kuehniella</i>	Pupa		(50-60)	35
61 <i>Galleria mellonella</i>			(6-11)	6
62 <i>Parage maera</i>	Larva, spinning	25.0-29.5	(60-69)	36
63 <i>Pieris brassicae</i>		25	(50-80)	37
64	Larva	20-30	(40-80)	37
65 <i>Prodenia eridania</i>	Larva, parasitized with <i>Apanteles</i>		29	6
66 <i>Saturnia pavonia major</i>	Larva		6	38
67 <i>Sphinx ligustri</i>	Pupa	29	58(40-90) ¹	38
	Larva		(0-39.8)	19
	Pupa	22	(38-54)	39
8 <i>Dytiscus marginalis</i>	Coleoptera		27	33
9				34
10	Intact		60 ¹	
11	Isolated fragment		(30-70)	40
12 <i>Lucanus cervus</i>	Isolated heart		6	40
			15	40
13 <i>Oryctes nasicornis</i>	Larva	18	15.7	41
		13	14.1	41
		25	18	30
		24-33	(17-31)	6

/ Heart exposed by dissection and examined under physiological saline.

46. HEART RATE, BASAL AND NON-BASAL AT VARIOUS AGES MAN

Heart rate varies with sex, age, environment, and temperature. Values in parentheses are ranges, estimate "b" for columns B and C, and estimate "c" for column D (cf. Introduction).

Age yr	Heart Rate, beats/min			Reference
	Basal		Non-Basal	
	Male (B)	Female (C)		
(A)	(B)	(C)	(D)	(E)
1 <1			134(101-160)	1
2 1	116	122	115(84-136)	B, C, 2, D, 1
3 2	104	103	108(84-134)	
4 3	92	86	108(80-124)	
5 4		87	103(80-133)	
6 5		91	98(70-128)	
7 6	87	80	93(72-128)	
8 7	85	73	94(72-112)	
9 8	80	77	89(72-114)	
10 9	81	85	91(68-120)	
11 10	79	80	87(56-106)	
12 11			89(60-120)	1
13 12	69 5(53.3-85.7)	71.4(55.8-87.0)	88(67-112)	B, C, 3, D, 1
14 13	65 9(53.9-77.9)	67.8(51.8-83.8)	87(66-114)	
15 14	67.1(53.1-81.1)	68.3(54.5-82.1)	82(60-112)	
16 15	65.2(54.2-76.2)	67.3(50.1-84.5)	83(66-112)	
17 16	61 6(48.6-74.6)	65.8(50.6-81.0)	80(54-108)	
18 17	58.6(49.0-68.2)	64.1(50.3-77.9)	76(54-104)	
19 18		73	77(60-108)	
20 19	65	71	74(52-104)	
21 20	64	69	71(59-99)	
22 20-40	62	68		
23 21			71(41-94)	1
24 22			70(56-100)	
25 23			71(50-96)	
26 24	61 0(47-75)	69 2(52.4-86.0)	72(50-96)	
27 25-30			72(52-102)	
28 25-44			71(50-104)	
29 30	59.3(47.1-71.5)	64 8(47.8-81.8)		
30 30-35			70(58-104)	
31 34	71 7(64.9-78.5)			
32 35-40			72(56-100)	
33 40-45			72(50-104)	1
34 40-60	57	70		2
35 43	69 1(63.1-75.1)			
36 45-49				
37 50-55			72(49-100)	
38 50-59			72(52-94)	
39 55	69 8(64.2-75.4)		73(48-108)	
40 55-60				
41 60-65			75(48-108)	
42 60-70	66	71	73(54-100)	
43 65	63 0(56.6-69.4)		74(52-100)	
44 65-70				B, C, 2, D, 4
45 70-75			75(52-96)	
46 70-79			75(54-104)	
47 >70	65	73	75(50-104)	
48 73	65.8(58.6-73.0)			
49 75-80				
50 >80			72(50-94)	
51 82	67 0(52-82)		77(63-98)	

Contributors (a) Johnson, Richard P. (b) Shock, Nathan W.

References [1] Tigerstedt, R. A. 1902 Lehrbuch der physiologie des Kreislaufes. In J. Mackenzie, ed. The study of the pulse. Young J. Pentland, Edinburgh and London. [2] Sullivan, W. D., and E. Holt. 1925. Arch. Int. M. 35:224 [3] Shock, N. W. 1944 Am. J. Dis. Child 68:17. [4] Bowerman, W. G., and J. H. Brett. 1941. Q. Rev. Biol. 16:90 [5] Brandfonbrener, M., M. Landowne, and N. W. Shock. 1955 Circulation, N. Y. 12:557.

- J. Agr. Res. 55:1. [20] Battini, S., G. Natalizi, and M. Boccacci. 1956. Riv. parassit. 17:179. [21] Davenport, D. 1949. Physiol. Zool. 22:35. [22] Wedl, C. 1855. Akad. der Wiss. 17:173. [23] Müller, J. 1915. Das österreichische Sanitätswesen 27:1. [24] Locy, W. A. 1884. Am. Natur. 18:13. [25] Beard, R. 1953. In K. D. Roeder, ed. Insect physiology. J. Wiley and Sons, New York. pp. 232-272. [26] Kirachner, R. 1932. Zschr. angew. Entom. 19:544. [27] Isaguzhinskaya, L. V. 1954. Biull. Mosk. obshch. ispyt. priro. otel Biol. 59:41. [28] Yokoyama, T. 1939. Proc. Imper. Acad. Japan 15:94. [29] Tischler, N. 1935. J. Econom. Entom. 28 215. [30] Polimanti, O. 1915. Biol. Zbl. 35:143. [31] Masera, E. 1933. Riv. biol. 15:225. [32] Campbell, F. L. 1926. J. Pharm. Exp. Ther. 28:277. [33] Newport, C. 1837. Proc. R. Soc., Lond. 127:259. [34] Fischer, E. 1931. Mitt. schweiz. ent. Ges. 15:46. [35] Fischer, E. 1918. Entom. Rdsch., Stuttgart 35:9. [36] Zeller, H. 1938. Zschr. Morph. Okol. Tiere 34:663. [37] Millman, N. 1938. Some aspects of the physiology of the heart of *Galleria mellonella* (bee moth). M. S. thesis, Brown University, Providence. p. 52. [38] Tarasova, K. L. 1936. Bull. Inst. Zool. appl., Leningr. 6:15. [39] Tirelli, M. 1935. Arch. zool. Ital. 22:279. [40] Duwez, Y. 1936. C. rend. Soc. biol. 122:84. [41] Lasch, W. 1913. Zschr. allg. Physiol. 14:312. [42] Rengel, C. 1896. Zschr. wiss. Zool. 62:1. [43] Wixforth, O. 1927. Arch. Naturgesch. 90A(5):193. [44] Koldsumi, K. 1931. J. Soc. trop. Agr., Taiwan 3:354. [45] Jones, J. C. 1954. J. Morph. 94:71. [46] Myers, J. G. 1927. Bull. Entom. Res., Lond. 17:219. [47] Levy, R. 1928. C. rend. Soc. biol. 99:1482. [48] Matsuki, T. 1928. Sc. Rep. Tohoku Univ., Ser. 4, 3:55. [49] Dubulsson, M. 1929. Arch. biol., Par. 39:247. [50] Davis, C. C. 1956. J. Cellul. Physiol. 47:449. [51] Kruger, F. 1931. Zschr. angew. Entom. 18:344. [52] Tzonis, V. K. 1936. Zool. Anz. 116:81. [53] Florey, E. 1951. Pflschber. 7:81. [54] Dogiel, J. 1877. Mém. Acad. Imp. Sci., S. Petersb., 7 Ser., 24:1. [55] Watson, G. I. 1937. Ann. Trop. M. Parasit., Liverp. 31:417. [56] Sautet, J., and Y. Audibert. 1946. Bull. Soc. path. exot., Par. 39:137. [57] Lawson, J. W. H. 1951. Tr. R. Entom. Soc., Lond. 102:511. [58] Pertunnen, V. 1955. Ann. Entom. Fenn. 21:78. [59] Hase, A. 1927. Zschr. Morph. Okol. Tiere 8:187. [60] Brown, J. M. 1910. Tr. Linn. Soc., Lond. 11:125. [61] Wetinger, O. 1927. Zschr. wiss. Zool. 129:453.

45. HEART RATE OF EMBRYO AND NEWBORN: MAMMALS

Heart rate varies with species, sex, age, size, environment, and temperature. Values in parentheses are ranges, estimate "c" (cf. Introduction).

Animal	Heart Rate beats/min	Reference
(A)	(B)	(C)
Embryo		
1 Man, 5th mo	156(150-160)	1
2 6th mo	154(141-155)	
3 7th mo	149(118-156)	
4 8th mo	142(129-152)	
5 9th mo	146(131-173)	
6 Dog	(120-170)	2
7 Goat	(120-246)	1
8 Monkey	(100-200)	1
9 Ox	161	2
10 Rat	(95-256)	1
Newborn		
11 Man, premature	145(110-185)	3
12 Man, full-term	(130-140)	
13 Cat	300	2, 4
14 Dog	(160-180)	2
15 Goat	(145-240)	1
16 Horse	(100-120)	5
17 Ox	(141-160)	2
18 Pig	227	4
19 Rabbit	220	6
20 Rat	161(121-201)	4

Contributors: (a) Johnson, Richard P., (b) Woodbury, Robert A.

References: [1] Barcroft, J. 1936. Physiol. Rev. 16:103 [2] Clark, A. J. 1927. Comparative physiology of the heart. Macmillan, New York. [3] Suttiff, W. D., and E. Hall. 1925. Arch. Int. M. 35:224. [4] Marcuse, F. L., and A. N. Moore. 1943. Am. J. Physiol. 139:49. [5] Reichert, A. 1909. Klinische Untersuchungen über die normale Pulsfrequenz unserer Haustiere. Inaugural Dissertation. Giessen, Germany. [6] Hamilton, W. E., R. A. Woodbury, and B. Woods. 1937. Am. J. Physiol. 119:206

47. EFFECT OF TEMPERATURE ON HEART RATE: VERTEBRATES (Continued)

Part III INTACT ANIMALS

Animal	Experimental Condition	Site of Temperature Recording	Heart Rate		Reference
			At Temp., °C	beats/min	
(A)	(B)	(C)	(D)	(E)	(F)
1 Man	Anesthetized or unanesthetized in water at 2-12°C ¹	Rectum	30	Atrial fibrillation	1
			34	50	
			37	80-90	
	Sedated with barbiturate, and cooled with brine-circulated blanket	Deep rectum	29	60	2
			32	83	
7 Cat	Deeply anesthetized with chloralose	Deep rectum	35	95	3
			37	100	
			30	100	
			32	150	
			34	175	
14 Dog	Anesthetized with sodium pentothal, artificial respiration when necessary	Right heart blood	36	190	4,5
			38	225	
			40	210	
			17	17	
			19	24	
			21	32	
			23	44	
			25	63	
	Anesthetized with sodium pentobarbital, O ₂ plus inhalation anesthetics with respirator	Rectum	27	84	
			29	108	
			31	132	
			33	147	
33 Guinea pig	Deep urethane narcosis	Deep rectum	35	161	3
			37	174	
			22	38	
			24	51	
			26	70	
			28	89	
			30	108	
			32	123	
	Unanesthetized and restrained ²	Colon	34	147	
			36	175	
			30	212	
			32	216	
45 Hamster, golden	Unanesthetized, during arousal from hibernation	Cheek pouch	34	227	7
			36	240	
			38	256	
			40	287	
			42	283	
			23	75	
			25	120	
			30	210	
	Unanesthetized, during arousal from hibernation	Cheek pouch	35	295	
			38	350	
			5	14	
			10	50	
52 Monkey ³	Unanesthetized, during arousal from hibernation	Cheek pouch	15	105	8
			20	185	
			25	270	
			30	380	
			35	500	
	Anesthetized with sodium pentobarbital, artificial respiration with 100% O ₂	Rectum	23	50	
			30	100	
			33	140	
			37	167	
56 Rabbit, albino	Unanesthetized ⁴	Deep rectum	20	20-40	10
			25	40	
			28	70	
			39	200	

/1/ Immediately after immersion, heart rate = 120-140 beats/min. Lethal temperature variable and uncertain.
 /2/ At body temperature of less than 23°C, heart rate irregular. /3/ Age, 12-18 months. /4/ After immersion, heart rate = 270 beats/min. At body temperature of less than 20°C, heart beat irregular and weak.

47. EFFECT OF TEMPERATURE ON HEART RATE: VERTEBRATES

Part I: PER DEGREE TEMPERATURE CHANGE

Animal	Temp. range °C	Condition or Method Effecting Temp. Change	Heart Rate Change per °C Temp. Change beats/min	Heart Rate Change per °F Temp. Change beats/min	Reference
(A)	(B)	(C)	(D)	(E)	(F)
1 Man	36.0-41.0	Fever induced by short radio-waves	7.4-30.6	4.1-17.0	1
2	36.9-38.9	Lobar pneumonia	10.4-27.2	5.8-15.1	2
3	35.6-39.4	Peptone injections	17.6	9.8	2
4	36.9-38.9	Hot baths	18	10	3
5	36.1-37.3	Regression of resting heart rate	19.8	11.01	4

15	0-30	heart Perfused isolated heart	1.7-5.0	0.9-2.8	14
----	------	----------------------------------	---------	---------	----

/1/ Of body, or of isolated heart. /2/ Urethane anesthetized. /3/ Anesthetized

Contributor: Cooper, K. E.

References [1] Bierman, W. 1933. Brit. J. Phys. M. 7:155. [2] Lyon, D. M. 1927. Q. J. Med., Oxf. 20:205. [3] Bazett, H. C. 1924. Am. J. Physiol. 70:412. [4] Tanner, J. M. 1951. J. Physiol., Lond. 115:391. [5] Hamilton, J. B., M. Dresbach, and R. S. Hamilton. 1937. Am. J. Physiol. 118:71. [6] Starling, E. H., and F. P. Knowlton. 1912. J. Physiol., Lond. 44:207. [7] Brenning, R. 1938. Über die Wirkung von Erhöhung der Körpertemperatur auf dem Kreislauf. Almqvist and Wiksell, Uppsala. [8] Kisch, B. 1923. Pflügers Arch. 198:105. [9] Lutz, W. 1948. Zschr. Kreislaufforsch. 37:266. [10] Ruhe, C. W. E., and R. H. Horn. 1955. Am. J. Physiol. 182:325. [11] Fairfield, J. 1948. Ibid. 155:355. [12] Crismon, J. M. 1944. Arch. Int. M. 74:235. [13] Barcroft, J., and J. J. Izquierdo. 1931. J. Physiol., Lond. 71:145. [14] Clark, A. J. 1920. Ibid. 54:275.

Part II: ISOLATED HEART

Animal	Experimental Condition	Site of Temperature	Heart Rate, beats/min At Temp., °C								Reference
			10	15	20	25	30	35	40	(M)	
2 Cat	oxygenated Ringer's Denervated heart-lung preparation	Blood in venous cannula							75	102 153 165	2
3 Dog	Denervated heart-lung preparation	Sino-atrial node tissue							45	88 132 160	3
4 Rabbit	Atria immersed in oxygenated Ringer's	Ambient fluid					30	65	100	147 230	4
5 Frog	Whole heart immersed in oxygenated Ringer's, during summer	Ambient fluid	92	13	23	35	44	52	60		5
6	Whole heart immersed in oxygenated Ringer's, during winter	Ambient fluid	5	8	20	30	42	57	72		4

/1/ 13 weeks. /2/ Only in South African frog

Contributor: Badeer, Henry

References [1] Garrey, W. E., and S. E. Townsend. 1948. Am. J. Physiol. 152:219. [2] Knowlton, F. P., and E. H. Starling. 1912. J. Physiol., Lond. 44:206. [3] Badeer, H. 1951. Am. J. Physiol. 167:76. [4] Clark, A. J. 1920-21. J. Physiol., Lond. 54:275. [5] Taylor, N. B. 1931. Ibid. 71:156

48. EFFECT OF EXERCISE ON HEART RATE: MAN

Values in parentheses are ranges, estimate "c" (cf. Introduction).

Exercise	Age yr	No of Subjects	Sex	Body Weight kg	Body Height cm	Heart Rate beats/min	Reference
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)
1 6 min on treadmill or bicycle ergometer	4-6	10	♂	20.8(16.0-27.8)	113.5(107-126)	203(188-214)	1
	7	7	♀	18.4(17.4-21.9)	111.6(108-114)	204(176-214)	
	7-9	12	♂	30.7(25.1-36.5)	135.0(125-143)	208(191-220)	
	14	14	♀	27.2(20.6-33.0)	132.0(121-142)	212(194-233)	
	10-11	13	♂	36.5(31.1-44.7)	145.4(132-157)	211(200-227)	
	13	13	♀	32.5(27.0-37.4)	140.6(129-148)	209(192-220)	
	12-13	19			141.4(130-169)	205(175-237)	
	13	13					
	14-15	10					
	11	11					
	16-18	9					
	16-17	10					
	20-23	42					
	20-25	44					
5 min on treadmill	20-29	11					2
	31-38	11					
	40-48	10					
	48-55	8					
	59-66	7					
Bicycle ergometer ¹							1
600 kg-m/min		31	♀			138(120-150)	
900 kg-m/min		21	♂			128(102-148)	
		31	♀			168(146-192)	
1200 kg-m/min		21	♂			148(130-169)	
1500 kg-m/min		21	♂			167(148-188)	
Running until nearly exhausted, then up and down flight of stairs ²	9	1	♂	32	134.6	189.3	3
	10	2		21.1(16.1-31.1)	132.1(117-147.3)	180(163-197)	
	11	4					
	12	1					
	13	15					
	14	3					
	15	1					

1/ Well-trained subjects 2/ Duration of exercise, 2-4 minutes. 3/ One-minute count.

Contributors (a) Åstrand, Per-Olof, (b) Asmussen, Erling, (c) Johnson, Richard P

References [1] Åstrand, P.-O 1952. Experimental studies of physical working capacity in relation to sex and age. Ejnar Munksgaard, Copenhagen. [2] Robinson, S. 1938 Arbeitsphysiologie 10:251. [3] Boas, E. P., and E F Goldschmidt 1932 The heart rate. C. C Thomas, Springfield, Ill. p 90.

49. EFFECT OF POSTURE AND SLEEP ON HEART RATE: MAN

Values in parentheses are ranges, estimate "c" (cf. Introduction)

Specification	No of Subjects	Sex	Age yr	Heart Rate beats/min	Reference
(A)	(B)	(C)	(D)	(E)	(F)
1 Basal	182	♂	College	65(45-105)	1
2 Recumbent	252	♂	College	66(40-100)	
3 Sitting	252	♂	College	73(48-103)	
4 Standing	194	♂	College	82(54-124)	
5 Sleeping	51	♂	28.1 ¹	59.5(52.8-67.1)	2
	52	♀	24.6 ¹	65.3(57.7-75.4)	
6 Awake	51	♂	28.1 ¹	77.8(61.2-111.8)	
	52	♀	24.6 ¹	83.9(67.1-120.6)	

1/ Average

Contributor Johnson, Richard P

References [1] Brouha, L., and C. W Heath. 1943 N England J. M. 228 473 [2] Boas, E. P., and E. F. Goldschmidt 1932 The heart rate C. C Thomas, Springfield, Ill. p 23.

47. EFFECT OF TEMPERATURE ON HEART RATE: VERTEBRATES (Concluded)

Part III: INTACT ANIMALS (Concluded)

Animal	Experimental Condition	Site of Temperature Recording	Heart Rate		Reference
			At Temp., °C	beats/min	
(A)	(B)	(C)	(D)	(E)	(F)
60 Rabbit, albino	Anesthetized with sodium pentobarbital	Colon	10	13	11
61 (concluded)			12	24	
62			14	34	
63			16	45	
64			18	59	
65			20	70	
66			22	84	
67			24	96	
68			26	115	
69			28	132	
70			30	158	
71	Unanesthetized and restrained	Colon	32	185	13
72			34	211	
73			36	234	
74			38	256	
75			40	276	
76 Rat, albino	Anesthetized with sodium pentobarbital	Deep rectum	15	45	12
77			20	120	
78			25	195	
79			30	270	
80			35	345	
81	Unanesthetized and restrained	Colon	15	45	13
82			20	110	
83			25	210	
84			30	300	
85			35	405	
86	Squirrel, ground	Food pouch	37	460	14
87			10	38	
88			15	102	
89			20	218	
90	Frog, South African	Intraventricular blood	25	357	15
91			2	4	
92			5	11	
93			10	20	
94			15	18	
95			20	27	
96			25	34	
97			30	31	
98	Fish (cod, eel, flounder, pollack, sculpin, skate)	Ambient fluid	36	25	16
99			0	6	
100			5	12	
101			12	18	
102			15	25	
103			20	32	
104			25	38	
105			30	45	

Contributor: Badeer, Henry

References [1] Gagge, A. P., and L. P. Herrington 1947 Annual Rev. Physiol. 9 409. [2] Talbot, J. H. 1951. Hypothermia. In Transactions of the first conference on cold injury, 1951 Josiah Macy, Jr., Foundation, New York. p. 145. [3] Barcroft, J., and J. J. Izquierdo 1931 J. Physiol., Lond 71:346. [4] Hegnauer, A. H., W. J. Shriber, and H. O. Haterius. 1950. Am. J. Physiol. 161 455. [5] Hegnauer, A. H., J. Flynn, and H. D'Amato. 1951. Ibid 167-69. [6] Deterling, R. A., Jr., E. Nelson, S. Bhonslay, and W. Howland 1959 Arch Surg 70 87. [7] Gosselin, R. E. 1949 Am. J. Physiol. 157 103 [8] Chatfield, P. O., and C. P. Lyman. 1950. Ibid. 163 566. [9] Bering, E. A., Jr., J. A. Taren, J. D. McMurrey, and W. F. Bernhard. 1956. Surg. Gyn Obst 102:134 [10] Ariel, I., F. W. Bishop, and S. L. Warren. 1943. Cancer Res 3 448 [11] Bartlett, R. G., Jr 1957 J Appl. Physiol. 10:143 [12] Crismon, J. M. 1944. Arch Int M. 74:235. [13] Hamilton, J. B., M. Dresbach, and R. S. Hamilton 1937. Am J. Physiol. 118:71 [14] Johnson, G. E. 1929 Biol. Bull 57 107. [15] Taylor, N. B. 1931. J. Physiol., Lond 71 156. [16] Britton, S. W. 1923-24 Am J. Physiol. 67 411.

41	Diastolic blood pressure, mm. Hg	Basal	74(59-93)	77(49-103)	80(62-98)	79(62-92)	77(66-88)	83(63-111)	75(64-86)
42	0.5	76(61-91)	81(52-110)	82(60-104)	80(70-90)	80(60-96)	78(60-96)	81(54-108)	76(55-123)
43	2.0	77(49-109)	81(57-105)	81(68-94)	80(61-93)	77(61-89)	79(63-105)	81(54-108)	76(55-123)
44	5.0	74(60-88)	77(48-108)	80(60-100)	80(61-93)	77(61-89)	79(63-105)	81(54-108)	76(55-123)
45	10.0								

Contributor: Shock, Nathan W.

Reference: Shock, N. W., and A. H. Norris. Unpublished.

Part II. STEP-TEST EXERCISE
Exercise walking over two 9-inch steps the number of times required to perform 222 kgm of work in 1.5 min. Subjects seated during recovery. Blood pressure determined by auscultatory method. Values in parentheses are ranges, estimate "0" (cf. Introduction).

Condition	Recovery Time, min	20-29 yr (17 subjects) (C)	30-49 yr (123 subjects) (D)	50-59 yr (119 subjects) (E)	60-69 yr (128 subjects) (F)	70-79 yr (28 subjects) (G)	80-89 yr (19 subjects) (H)
(A)	(B)						
1 After standing	0.25	60.1(37-103)	80.4(48-106)	81.1(54-108)	78.4(53-103)	81.1(54-107)	80.1(56-104)
2	1.25	75.0(54-96)	76.4(49-101)	78.3(52-104)	74.8(53-97)	76.3(52-100)	73.3(49-97)
3	5.00	77.1(53-101)	76.4(52-102)	77.8(51-105)	75.0(56-94)	75.6(51-101)	73.7(53-95)
4 After exercise	0.25	83.2(63-103)	87.7(67-107)	89.3(44-142)	89.3(64-114)	95.4(68-122)	92.9(64-122)
5	1.25	77.8(57-99)	77.3(57-107)	77.2(49-105)	77.2(53-101)	81.6(59-105)	79.5(54-106)
6	5.00	80.6(50-102)	78.3(50-106)	79.3(44-114)	76.5(56-98)	78.1(52-104)	73.3(51-97)
7 After standing	0.25	120.3(93-147)	126.4(81-171)	135.1(92-176)	131.1(91-171)	148.0(83-213)	145.5(101-191)
8	1.25	139.6(91-149)	126.9(77-177)	134.1(100-170)	128.6(99-159)	147.4(85-209)	142.1(98-186)
9	5.00	115.4(92-138)	122.7(83-163)	128.8(93-165)	124.8(92-158)	140.7(79-203)	136.3(93-179)
10 After exercise	0.25	129.1(107-153)	136.1(92-180)	146.0(108-170)	141.3(101-181)	162.3(88-236)	155.1(101-209)
11	1.25	122.8(98-151)	140.6(106-176)	140.6(106-176)	141.0(104-178)	162.7(87-239)	156.3(108-208)
12	5.00	118.6(93-145)	125.4(85-165)	132.3(98-168)	131.5(97-167)	149.5(84-216)	143.6(101-187)
13 After standing	0.25	76.4(58-96)	85.9(60-112)	86.2(62-110)	80.9(52-110)	81.1(47-115)	77.8(52-104)
14	1.25	76.3(60-92)	86.4(60-112)	86.9(65-109)	80.5(56-106)	81.1(50-112)	77.7(47-107)
15	5.00	75.3(60-92)	81.5(69-111)	86.1(64-108)	78.5(55-103)	79.3(48-110)	77.0(50-104)
16 After exercise	0.25	75.2(59-91)	87.4(69-109)	84.2(58-110)	78.0(54-102)	80.4(45-115)	76.1(53-99)
17	1.25	79.5(65-93)	87.6(67-112)	89.3(66-113)	82.2(58-106)	85.0(50-120)	81.2(53-109)
18	5.00	80.8(65-93)	89.3(66-112)	89.4(65-113)	84.2(58-110)	83.9(51-117)	80.5(56-106)

Contributor: Shock, Nathan W.

Reference: Norris, A. H., N. W. Shock, and M. J. Ylengst. 1933. Circulation, N. Y. 8:321.


50. RECOVERY OF HEART RATE AND BLOOD PRESSURE AFTER EXERCISE; MAN

Part I: VARIOUS WORK LOADS

Exercise: cranking movement of arms. Subjects recumbent during exercise and recovery. Blood pressure determined by auscultatory method. Values in parentheses are ranges, estimate "B" (cf. Introduction).


Variable	Recovery Time, min	20-29 yr (10 subjects) (C)	30-39 yr (11 subjects) (D)	40-49 yr (10 subjects) (E)	50-59 yr (9 subjects) (F)	60-69 yr (10 subjects) (G)	70-79 yr (10 subjects) (H)	80-89 yr (9 subjects) (I)
Work Load = 500 kg-m at 135 kg-m/min								
1 Heart rate, beats/min	Basal	59(41-77)	63(45-81)	70(54-86)	69(43-95)	64(42-86)	65(40-90)	59(47-71)
2	0.5	74(58-90)	77(56-98)	82(60-104)	79(59-99)	83(57-109)	91(70-112)	92(54-130)
3	2.0	63(40-86)	67(49-85)	71(57-85)	69(48-90)	69(43-95)	71(47-95)	74(49-99)
4	5.0	62(53-71)	66(50-82)	73(52-84)	68(47-89)	66(38-94)	67(41-93)	66(45-87)
5	10.0	61(45-77)	66(49-83)	71(55-87)	68(44-92)	67(44-90)	67(42-92)	64(47-87)
6 Systolic blood pressure, mm Hg	Basal	116(99-137)	119(87-151)	118(101-135)	120(89-151)	120(87-153)	141(74-208)	163(95-231)
7	0.5	130(96-164)	133(102-192)	143(95-230)	138(110-166)	152(104-200)	178(106-250)	177(101-235)
8	2.0	123(95-151)	128(84-172)	130(101-159)	135(108-162)	142(102-182)	158(76-240)	178(101-235)
9	5.0	118(88-150)	119(84-154)	122(101-143)	125(101-144)	124(101-147)	150(74-226)	156(99-213)
10	10.0	115(90-140)	119(85-153)	119(103-135)	121(98-144)	124(101-147)	150(74-226)	156(99-213)
11 Diastolic blood pressure, mm Hg	Basal	76(64-88)	82(55-109)	86(65-95)	77(64-90)	73(60-86)	84(60-108)	78(61-95)
12	0.5	79(64-94)	83(49-117)	84(68-105)	80(72-88)	79(60-98)	88(63-113)	85(56-114)
13	2.0	76(59-93)	79(46-112)	84(63-105)	79(66-92)	77(61-95)	86(65-114)	86(65-107)
14	5.0	76(59-93)	76(47-107)	82(64-100)	78(65-91)	76(62-90)	83(57-113)	80(64-96)
15	10.0	73(59-91)	74(46-105)	81(67-99)	78(65-91)	75(64-86)	83(55-111)	79(64-94)
Work Load = 500 kg-m at 267 kg-m/min								
16 Heart rate, beats/min	Basal	61(41-81)	63(49-77)	65(53-77)	65(41-114)	62(43-81)	63(36-90)	60(50-70)
17	0.5	81(60-102)	82(64-100)	89(62-116)	90(63-117)	95(65-125)	101(69-133)	101(62-140)
18	2.0	67(58-76)	67(48-86)	72(55-88)	72(49-95)	70(40-100)	76(52-100)	80(50-110)
19	5.0	62(53-71)	65(46-84)	66(52-80)	68(52-84)	70(48-92)	70(48-92)	67(49-85)
20	10.0	64(50-78)	55(45-85)	58(53-83)	70(46-94)	67(42-92)	70(46-94)	65(50-80)
21 Systolic blood pressure, mm Hg	Basal	115(104-126)	117(95-138)	116(93-138)	118(91-145)	127(85-169)	141(91-191)	163(102-222)
22	0.5	130(106-154)	136(104-168)	143(93-193)	147(93-197)	166(106-226)	178(116-236)	177(96-262)
23	2.0	123(102-144)	130(103-157)	138(96-180)	140(83-191)	149(105-193)	169(108-234)	181(108-266)
24	5.0	116(100-138)	122(97-145)	125(92-158)	129(105-153)	130(99-161)	149(87-211)	172(118-236)
25	10.0	111(102-130)	120(91-149)	118(95-140)	125(93-157)	126(94-158)	141(84-198)	166(111-221)
26 Diastolic blood pressure, mm Hg	Basal	74(62-86)	79(59-99)	82(63-101)	79(62-96)	77(65-99)	82(56-108)	77(64-90)
27	0.5	76(63-89)	82(60-104)	84(60-108)	82(68-96)	84(61-107)	88(62-114)	87(57-117)
28	2.0	73(65-85)	79(59-99)	83(62-104)	82(66-98)	80(58-102)	84(62-106)	88(67-109)
29	5.0	73(65-85)	77(58-96)	81(63-99)	80(64-96)	78(64-92)	83(58-108)	83(69-97)
30	10.0	75(63-87)	78(58-98)	80(63-97)	81(68-94)	78(60-96)	82(56-108)	81(66-96)
Work Load = 350 kg-m at 450 kg-m/min								
31 Heart rate, beats/min	Basal	61(45-77)	60(49-71)	65(53-81)	65(41-89)	62(42-82)	65(37-93)	56(42-75)
32	0.5	76(61-93)	77(60-98)	89(68-110)	81(48-114)	87(61-123)	96(56-136)	92(59-125)
33	2.0	70(55-105)	73(50-84)	73(57-89)	71(48-94)	70(41-99)	75(46-104)	78(54-102)
34	5.0	62(40-84)	68(48-76)	67(52-82)	66(45-87)	66(38-94)	69(42-96)	66(50-82)
35	10.0	61(46-74)	69(47-79)	68(54-82)	66(42-90)	65(39-91)	69(41-97)	63(46-80)
36 Systolic blood pressure, mm Hg	Basal	116(92-140)	113(87-139)	114(98-130)	121(91-151)	125(88-164)	144(76-229)	157(99-215)
37	0.5	123(93-153)	127(91-163)	127(98-156)	138(119-157)	148(109-187)	171(105-212)	177(96-247)
38	2.0	118(91-145)	125(93-157)	125(99-147)	137(119-180)	141(100-182)	167(103-231)	180(96-264)
39	5.0	116(94-138)	120(88-152)	121(91-151)	130(100-160)	128(94-162)	152(80-224)	169(106-232)
40	10.0	116(92-133)	117(88-146)	115(93-137)	126(99-153)	124(96-152)	143(66-220)	156(98-218)

51. ARTERIAL BLOOD PRESSURE VS AGE AND SEX: MAN (Concluded)
Part III. SYSTOLIC: FEMALE



Contributor: Hartrott, W. Stanley
Reference: Master, A. M., C. I. Garfield, and M. B. Walters. 1952. Normal blood pressure and hypertension. Lea and Febiger, Philadelphia p 100


Part IV: DIASTOLIC: FEMALE



Contributor: Hartrott, W. Stanley
Reference: Master, A. M., C. I. Garfield, and M. B. Walters. 1952. Normal blood pressure and hypertension. Lea and Febiger, Philadelphia p 102

51. ARTERIAL BLOOD PRESSURE VS AGE AND SEX: MAN


Part I: SYSTOLIC: MALE



Contributor: Hartroft, W. Stanley

Reference: Master, A. M., C. I. Garfield, and M. B. Walters. 1952. Normal blood pressure and hypertension. Lea and Febiger, Philadelphia. p. 99.

Part II: DIASTOLIC: MALE



Contributor: Hartroft, W. Stanley

Reference: Master, A. M., C. I. Garfield, and M. B. Walters. 1952. Normal blood pressure and hypertension. Lea and Febiger, Philadelphia. p. 101

53. ARTERIAL BLOOD PRESSURE

Part I. MAN


Values are in mm Hg. Values in parentheses are ranges, estimate "b" (cf. Introduction).

No and Sex (A)	Age yr (B)	Male		Female		Reference (G)
		Systolic (C)	Diastolic (D)	Systolic (E)	Diastolic (F)	
1	Birth	77	42			1
2	1 da	82	40			
3	2 da	85	43			
4	4 da	91	50			2
5	6 mo-1 yr	89(60-118)	60(50-70) ¹			
6	1	96(66-126)	66(41-92) ¹			
7	2	99(74-124)	64(39-89) ¹			
8	3	100(75-125)	67(44-90) ¹			
9	4	89(78-100)	60(46-74)	89(74-102)	60(47-73)	3
10	5	92(80-104)	62(47-77)	92(79-105)	62(49-73)	
11	6	94(81-107)	64(49-79)	94(80-108)	64(49-80)	
12	7	97(84-110)	65(50-80)	97(83-111)	66(51-81)	
13	8	100(87-113)	67(53-81)	100(86-114)	64(50-78)	
14	9	101(88-114)	68(55-81)	101(87-115)	69(55-83)	
15	10	103(90-116)	69(57-81)	103(89-117)	70(57-83)	
16	11	104(91-117)	70(59-81)	104(90-118)	71(58-84)	
17	12	106(93-119)	71(61-81)	106(92-120)	72(58-84)	
18	13	108(95-121)	72(62-82)	108(95-121)	73(58-88)	
19	14	110(97-123)	73(63-83)	110(97-123)	74(57-91)	
20	15	112(98-126)	75(64-86)	112(98-126)	76(57-95)	
21	16	118(95-142)	73(53-93)	116(92-140)	72(54-91)	4
22	17	121(96-146)	74(56-93)	116(93-139)	72(54-90)	
23	18	120(96-143)	74(55-94)	116(94-139)	72(55-89)	
24	19	122(92-151)	75(54-95)	115(92-138)	71(54-89)	
25	20-24	123(96-150)	76(57-96)	116(93-139)	72(53-91)	
26	25-29	125(100-150)	76(60-95)	117(94-139)	74(56-92)	
27	30-34	126(99-153)	79(60-98)	120(92-147)	75(54-96)	
28	35-39	127(99-155)	80(60-101)	124(97-151)	76(56-98)	
29	40-44	129(100-159)	81(63-100)	127(94-161)	80(59-100)	
30	45-49	130(97-163)	82(61-103)	131(92-169)	82(59-104)	
31	50-54	135(97-172)	83(61-106)	137(96-179)	84(59-106)	
32	55-59	138(101-175)	84(62-106)	139(97-180)	84(61-106)	
33	60-64	142(100-183)	85(60-109)	144(100-188)	85(60-110)	
34	65-69	143(92-194)	83(64-102)	154(97-211)	85(58-112)	5
35	70-74	145(93-197)	82(52-112)	159(108-210)	85(55-115)	
36	75-79	146(104-188)	81(56-106)	158(106-210)	84(58-110)	
37	80-84	145(95-195)	82(63-101)	157(102-212)	83(57-109)	
38	85-89	145(88-192)	79(50-108)	154(99-209)	82(48-116)	
39	90-94	145(99-191)	78(54-102)	150(104-196)	79(55-101)	
40	95-106	146(92-200)	78(53-103)	149(103-195)	81(57-106)	

1/1 Point of muffling taken as the diastolic pressure

Contributors (a) Master, Arthur M., (b) Van Liere, Edward J., and Hugh A. Lindsay, (c) Hartroft, W. Stanley

References [1] Smith, C. A. 1951. The physiology of the newborn infant. Ed 2. C. C. Thomas, Springfield, Ill.
 [2] Allen-Williams, G. M. 1945. Arch Dis Childh., Lond. 20 125. [3] Faber, H. K., and C. A. James. 1921. Am. J. Dis Child 22 7. [4] Master, A. M., C. I. Garfield, and M. B. Walters. 1952. Normal blood pressure and hypertension. Lea and Febiger, Philadelphia [5] Master, A. M., R. P. Lasser, and H. L. Jaffe. 1957. Proc. Soc. Exp Biol. N Y 94 463.



Contributor: Hartroft, W. Stanley

References [1] Master, A. M., C. I. Garfield, and M. H. Walters. 1952. Normal blood pressure and hypertension. Lea and Febiger, Philadelphia. p. 104. [2] Master, A. M., L. I. Dublin, and H. H. Marks. 1950. *J. Am. M. Ass.* 143:1464. [3] Gover, Mary. 1948. *Pub. Health Rep., Wash.* 63:1083. [4] Russek, H. I., and B. L. Zorman. 1946. *Geriatrics* 1:113. [5] Robinson, S. C., and M. Brucer. 1939. *Arch. Int. M.* 64:409.

35		173♂	None	Cuff on tail, systolic by palpation, diastolic by auscultation	98(90-104)	64(45-86)	13
36		43♀	None	Cuff on tail, systolic by palpation, diastolic by auscultation	90(86-93)	59(43-84)	
37	Jackass	4	Nembutal or luminal	Cuff, mercury manometer	90-104	54-60	1
38	Mouse	15	Urethane or ether	Optical manometer	60-136		14
39		12	Local		124(95-125)	81(67-90)	15
40		3	None	Plethysmograph	141(115-158)	106(102-110)	16
41		19	2.5-5 mo	Plethysmograph	111(95-131)		
42		14	13-14 mo	Plethysmograph	136(114-158)		
43		13	31-32 mo	Plethysmograph	151(130-184)	(120-135)	1
44	Opussum	32	None	Hamulation manometer	110(95-130)	80(60-90)	17
45	Rabbit	9	None	Pressure cup on central artery	82(70-90)		19
46		35	None	Cuff on carotid loop mercury manometer	120(80-150)		20
47		12	Newborn	Hypodermic manometer	35	1	21
48	Flat	6	Nembutal	Optical manometer	150(75-150)	90(60-100)	22
49		70	Young adult	Plethysmograph	116(88-130)		23
50		100	Either and/or amylal	Cuff, foot microcopy	106(78-132)		24
51		18♂	Either and/or amylal	Optical manometer	(60-140)		116
52		23♀	Diast.	Mercury manometer	105(87-122)		25
53		18	Young adult	Plethysmograph	120(40-180)		26
54		68♂	None	Strain gauge foot pulse	90(60-120)		27
55		68♂	Nembutal	Cuff, microcopy	110(80-140)		28
56		68♂	Either	Cuff, microcopy	120(45-180)		29
57		68♂	Nembutal	Optical manometer	125(102-140)		
58		40♀	135-200 g	Plethysmograph	164(94-172)		
59		40♀	Amylal, 4 mg/100 g	Plethysmograph	117(110-126)		
60		61	Either, shallow	Plethysmograph	104(95-110)		
61		62	Either, deep	Plethysmograph	64(45-70)		
62		63	Morphine, 10 mg/100 g	Plethysmograph	104(66-115)		
63		64	Urethane, 4 mg/100 g	Plethysmograph	78(72-84)		
64		65	Nembutal, 20 mg/100 g	Plethysmograph	129(88-184)	91(58-145)	28
65		66	Pentobarbital	Optical manometer	130(108-165)		29
66	Rat, Norway	124♂	Adult	Aortic cuff; mercury manometer			136
67	Rat, Sprague-Dawley	20♂	105-106 g	Mercury manometer			114(90-140)
68	Seal (Phoca vitulina)	1	Young		169(144-185)	108(98-120)	157(94-206)
69	Sheep	13	Local				1
70		4					1
71							1
72	Swine						1

1/1/ Fasted 7-30 days. 12/ 50 determinations. 13/ Number of determinations. 14/ Monthly male. 15/ 540 determinations. 16/ Approximately.

Contributors. (a) Van Liere, Edward J., and Hugh A. Lindsay, (b) Conklin, Ruth E., (c) Freed, S. Charles, (d) Freis, Edward D., (e) Heister, Charles R., and Jack R. Leonards, (f) Link, Roger P., (g) Roodhard, Simon, (h) Woodbury, Robert A.

53. ARTERIAL BLOOD PRESSURE (Continued)

Part II MAMMALS OTHER THAN MAN

Values in parentheses are ranges, estimate "c" (cf. Introduction).

Animal	No. and Sex (B)	Age or Weight (C)	Anesthetic (D)	Method (E)	Systolic (F)	Diastolic (G)	Mean (H)	Reference (I)
1 Bat	1							
2 Cat	4		None	Cuff on carotid loop; mercury manometer			50 (140-170)	1
3	2		Barbital	Optical manometer	120	75		2
4	3		Ether	Optical manometer	120	15		3
5	191♂	1.4-4.9 kg	Dial-urethane	Mercury manometer			129(67-216)	4
6	208♀	1.4-4.9 kg	Dial-urethane	Mercury manometer			121(62-200)	4
7 Cattle								
8	♀		Local		134(124-166)	88(80-120)		1
9	♀						125	5
10	4	Young	Chloral hydrate				156	6
11							257(133-277)	1
12							130(110-140)	c
13 Dog	67♂	Young	Nembutal	Mercury manometer			134(85-190)	7
14	80♀		Sodium barbital	Mercury manometer			125(60-170)	c
15			Sodium barbital	Mercury manometer			108(92-120)	8
16	35		None	Hypodermic, Tyco's sphygmomanometer	120			9
17	13		None	Cuff, auscultation	112(95-136)	56(43-66)		10
18	22		Nembutal	Capacitance manometer	156(108-198)	100(75-121)		11
19	22		Nembutal	Oscillometer	149(108-189)	100(75-122)		12
20	22		Nembutal	Auscultation	149(100-187)	104(79-127)		13
21	99♂		Morphine	Hypodermic, optical manometer	180(100-275)	89(30-140)		14
22	116♀		Morphine	Hypodermic, optical manometer	180(100-275)	89(30-140)		15
23	4							16
24	41							17
25	2			Cuff, auscultation	82(74-86)	41(35-51)	159(143-172)	18
26	4		Urethane				124(118-135)	19
27							120	20
28					120(112-126)	84(76-90)		21
29							75(7-111)	22
30							196-200	23
31	Adult		Nembutal	Mercury or condenser manometer			120(120-170)	24
32	6	13-18 yr	Local					25
33	5♂	4.5 mo-1.5 yr	None	Cuff on tail, systolic by palpation, diastolic by oscillation or auscultation	80	50	159(152-194)	26
34	3♀	4.5 mo-1.5 yr	None	Cuff on tail, systolic by palpation, diastolic by palpation or auscultation	80	50		27
35	120				85(70-98)	48(40-58)		28

18	Vile (<i>Milvius migrans</i>)	4	Local	133(120-140)	103(100-115)	194	1
19	Pigeon	2	Local	118(110-125)	80		2
20	Robin	3	Local	108(80-135)			2
21	Sparrow, pin-feather	2	Local	123(115-130)	140		1
22	Fiddling	1	Local	180	130(100-160)	161	1
23	Adult	2	Local	160(150-210)		193	1
24	Starling					171	
25	Stork						
26	Turkey						
27	Vulture						

1/1 3 breeds

Contributors: (a) Van Liere, Edward J., and Hugh A. Lindsay, (b) Conklin, Ruth E., (c) Freed, S. Charles, (d) Frels, Edward D., (e) Heisler, Charles R., and Jack R. Leonards, (f) Rodbard, Simon, (g) Woodbury, Robert A.

References: [1] Lehmann, G. 1925. *Tabulae biol.*, Berl. Bd 1, p. 143. [2] Woodbury, R. A., and W. F. Hamilton. 1937. *Am. J. Physiol.* 119 663.

[3] Woodbury, R. A., and B. E. Abreu. 1944. *Ibid.* 142:114. [4] Dukes, H. H. 1947. *The physiology of domestic animals.* Cornell University Press, Ithaca.

Part IV REPTILES AND AMPHIBIANS
Values in parentheses are ranges, estimate "c" (cf. Introduction).

Animal (A)	No. of Subjects (b)	Method (c)	Systolic (d)	Diastolic (e)	Mean (f)	Reference (g)
1 Crocodile					130-55	1
2 Snake, ring	11	Optical manometer	44(38-55)	37(28-47)	89	2
3 Turtle (<i>Pseudemys elegans</i>) ¹		Hamilton manometer	31	25		3
4 Turtle (<i>P. elegans</i>) ²		Hamilton manometer	27	21	(18-17)	1
5 Turtle (<i>P. elegans</i>) ²					(18-35)	1
6 Turtle (<i>P. rugosa</i>)			(30-35)			1
7 Frog			(20-40)		(5-30)	2
8 Frog			43(35-56)	31(24-44)		4
9 Frog (<i>Rana catesbeiana</i>)	6 ¹	Optical manometer	32(28-36)	21(18-24)		4
10 Frog (<i>Rana catesbeiana</i>)	6 ²	Cuff, web microscopy	31(27-36)	21(16-26)		1
11 Frog (<i>R. pipiens</i>)	6 ³	Cuff, web microscopy			48	1
12 Frog (<i>R. pipiens</i>)	6 ⁴					1
13 Toad, earth						

1/1 At 22°C. 1/2 At 16°C. 1/3 Local anesthesia. 1/4 No anesthesia.

Contributors: (a) Van Liere, Edward J., and Hugh A. Lindsay, (b) Conklin, Ruth E., (c) Freed, S. Charles, (d) Frels, Edward D., (e) Heisler, Charles R., and Jack R. Leonards, (f) Rodbard, Simon, (g) Woodbury, Robert A.

References: [1] Lehmann, G. 1925. *Tabulae biol.*, Berl. Bd 1, p. 142. [2] Woodbury, R. A., and W. F. Hamilton. 1937. *Am. J. Physiol.* 119 663.

[3] Rodbard, S., and D. Feldman. 1946. *Proc. Soc. Exp. Biol.*, N. Y. 63:43. [4] Dieter, R. N., and F. H. Scott. 1929. *Am. J. Physiol.* 91:263.

53. ARTERIAL BLOOD PRESSURE (Continued)
Part II. MAMMALS OTHER THAN MAN (Concluded)

- References: [1] Lehmann, G. 1925. *Tabulae biol.*, Berl. Bd 1, p. 143. [2] Liddell, E. G. T., and H. M. Carleton, 1936-37. *Q. J. Exp. Physiol.*, Lond. 26:155. [3] Woodbury, R. A., and B. E. Abreu. 1944. *Am. J. Physiol.* 142:114. [4] Root, M. A. 1950. *Ibid.* 162:308. [5] Dukes, H. H. 1947. The physiology of domestic animals. Cornell University Press, Ithaca. [6] Dukes, H. H., and L. H. Schwarte, 1931. *J. Am. Vet. M. Ass.* 79:37. [7] Van Liere, E. J., J. C. Suckney, and D. F. Marsh. 1949. *Science* 109:489. [8] Parkies, W. M. 1934. *Am. J. Physiol.* 107:518. [9] Wilhelmj, C. M., E. B. Waldman, and T. F. McCuire. 1951. *Ibid.* 166:296. [10] Romagnoli, A. 1953. *Cornell Vet.* 43:161. [11] Hamilton, W. F., et al. 1940. *Am. J. Physiol.* 128:231. [12] Chatfield, P. O., and C. P. Lyman. 1950. *Ibid.* 163:566. [13] Covington, N. G., and G. W. McNutt. 1931. *J. Am. Vet. M. Ass.* 79:603. [14] McMaster, P. D. 1941. *J. Exp. M.* 74:29. [15] Woodbury, R. A., and W. F. Hamilton. 1937. *Am. J. Physiol.* 119:663. [16] Wu, C. H., and M. B. Vlascher. 1947. *Fed. Proc.*, Balt. 6:231. [17] Rodbard, S. 1940. *Am. J. Physiol.* 129:448. [18] Anderson, H. C. 1923. *Proc. Soc. Exp. Biol.*, N. Y. 20:295. [19] Dominguez, R. 1924. *J. Metab. Res.* 6:123. [20] Hamilton, W. F., R. A. Woodbury, and E. B. Woods. 1937. *Am. J. Physiol.* 119:206. [21] Snider, R. H., H. S. Kupperman, and W. F. Hamilton. 1944. *Ibid.* 141:625. [22] Byrom, F. B., and C. Wilson. 1918. *J. Physiol.*, Lond. 93:301. [23] Griffith, J. Q., Jr. 1934. *Proc. Soc. Exp. Biol.*, N. Y. 35:394. [24] Durant, R. R. 1927. *Am. J. Physiol.* 81:679. [25] Olmsted, F. A. C. Corcoran, and I. V. Page. 1951. *Circulation*, N. Y. 3:722. [26] Farties, E. J., and J. Q. Griffith, Jr. 1949. The rat in laboratory investigations. Ed. 2. J. B. Lippincott, Philadelphia. [27] Sulkin, N. M., and K. R. Brizze. 1947. *Proc. Soc. Exp. Biol.*, N. Y. 64:125. [28] Schroeder, H. A. 1942. *J. Exp. M.* 75:513. [29] Farrell, G. L., and E. Anderson. 1949. *Proc. Soc. Exp. Biol.*, N. Y. 72:461. [30] Irving, L., F. F. Scholander, and S. W. Grinnell. 1942. *Am. J. Physiol.* 135:557.

Part III: BIRDS

Values in parentheses are ranges, estimate "c" (cf. Introduction).

Animal (A)	No. of Subjects (B)	Anesthetic (C)	Method (D)	Blood Pressure, mm Hg Systolic (E)	Diastolic (F)	Mean (G)	Reference (H)
1 Buzzard (<i>Buteo buteo</i>)							
2 Canary	6			175(110-250)	154(150-160)	171	1
3 Chicken ¹	5	Barbital	Optical manometer	130	85		2
	1	Morphine	Optical manometer	130	85		3
4 Chickens, ♂	13	Ether	Optical manometer	130	85		3
5 Chickens, ♀						135	4
6 Crow (<i>Corvus cornix</i>)						168-171	1
7 Crow (<i>C. frugilegus</i>)						147	1
8 Duck, domestic						151	1
9 Duck, wild						162	1
10 Falcon (<i>Falco tinnunculus</i>)						179	1
11 Fowl						103	1
12 Goose				150	120		1
13 Gull (<i>Larus canus</i>)						129-176	1
14 Hawk (<i>Accipiter</i>)						179	1
15 Pigeon (<i>Columba</i>)						178	1
16 Jackdaw (<i>Corvus monedula</i>)						119	1

54. CAPILLARY BLOOD PRESSURE: VERTEBRATES

All measurements made directly by microcannulation [8] Capillary: A = arterial, V = venous. Values in parentheses are ranges, estimate "c" (cf. Introduction).

	Tissue	Animal	Condition	Capillary	Blood Pressure		Reference
					mm Hg	cm H ₂ O	
	(A)	(B)	(C)	(D)	(E)	(F)	(G)
1	Eponychium (finger)	Man	Normal	A	30.6(7.5-60.0)		1
2				V	21.9(7.5-47.0)		
3		Man	Normal	A	29.3(13.9-44.4)	43.5(28.6-65.0) ^{1,2}	E, 2, F, 2, 3
4				V	14.1(8.3-19.9) ¹	16.5(8.0-24.5) ^{1,2}	
5		Man	Hypertension	A	35.9(7.5-70.5)		1
6				V	22.8(9.5-43.0)		
7		Man	Hyperemia	A		86(71.0-93.0)	3
8				V		154.5-66.5)	
9	Mesentery	Guinea pig		A	28.0(23.0-36.0)	38.5(31.0-49.0) ^{3,4}	E, 4, F, 3, 5
10				V	12.0(9.5-14.5)	17.0(13.0-19.5) ^{3,4}	
11		Rat		A	22.0(16.0-25.0)	30.0(22.0-34.0) ³	E, 6, F, 3, 5
12				V	12.7(11.0-14.5)	17.0(15.0-20.0) ³	
13		Frog		A		14.4(5.0-22.0) ⁵	3, 4
14				V		10.1(6.7-18.0) ⁵	
15	Muscle	Frog	Normal	A	10.7(8.1-13.2)	14.9(11.0-18.0) ⁶	E, 5, 7, F, 3, 7
16				V	7.1(5.1-9.2)	9.5(7.0-12.7) ⁶	
17		Frog	Hyperemia	A		20.1(17.0-26.0)	3
18				V		16.0(12.0-17.5)	
19	Web	Frog	Normal	A	10.2(8.1-13.2)	13.9(10.0-19.0) ⁶	E, 7, F, 3, 7
20				V	8.1(5.9-10.3)	9.6(8.5-13.0) ⁶	
21		Frog	Normal	A		14.5(10.0-20.5) ⁷	3
22				V		10.0(8.5-15.5) ⁷	
23		Frog	Hyperemia	A		19.5(14.0-26.5)	3
24				V		16.5(15.0-17.5)	

[1] Varies directly with arteriolar vasodilatation produced by emotion, heat, or trauma. Varies inversely with arteriolar vasoconstriction produced by emotion or cold. Varies minimally in a single capillary with time, and also from capillary to capillary. [2] Varies directly with venous pressure as affected by hydronautic pressure or venous obstruction. [3] Decerebrate. [4] Anesthetized (veronal ether). [5] Flithed. [6] Anesthetized (urethane). [7] Curarized.

Contributors (a) Griffith, John Quintin, Jr., (b) Nendlowitz, Milton, (c) Redisch, Walter

References [1] Eichna, L. W., and J. Bordley, III 1942 J. Clin. Invest. 21 711. [2] Landis, E. M. 1930. Heart, Lond 15 209. [3] Landis, E. M. 1934. Physiol. Rev. 14 404. [4] Landis, E. M. 1926 Am. J. Physiol. 75 548 [5] Landis, E. M. 1930 Ibid 93 353. [6] Landis, E. M. 1928. Ibid. 83 528. [7] Landis, E. M. 1931. Ibid 98 704 [8] Carrier, E. B., and P. B. Rehberg 1923. Skand Arch. Physiol., Berl. 44 20.

55. VENOUS BLOOD PRESSURE. MAN

Reference level is the phlebostatic axis. Subjects in supine position, breathing quietly. Forced respiration (e.g., Valsalva's maneuver) profoundly influences venous pressure. Values in parentheses are ranges, estimate "c" (cf. Introduction)

	Vein	Blood Pressure
		mm H ₂ O
	(A)	(B)
1	Median basilic vein at elbow	
2	3-5 yr	46(30-63)
3	5-10 yr	58(33-74)
4	Adult ♀	94(60-126)
5	Adult ♂	100(50-140)
6	Femoral	111(98-128)
7	Abdominal	115(70-160)
8	Dorsal metacarpal	130(70-170)
9	Great saphenous at ankle	150(110-190)
10	Dorsal pedal	175(124-210)

Contributor Terry, Luther L.

Reference Burch, G. E. 1950 A primer of venous pressure. Lea and Febiger, Philadelphia

53. ARTERIAL BLOOD PRESSURE (Concluded)

Part V: FISHES

Values in parentheses are ranges, estimate "c" (cf. Introduction).

	Animal (A)	Blood Pressure, mm Hg			Reference (E)
		Systolic (B)	Diastolic (C)	Mean (D)	
1	Barbel			42	1
2	Carp ¹	43(40-45)			2
3	Eel			(65-70)	1
4	Pike			(35-64)	1
5	Ray (Raja sp)	20			1
6		21			1
7	Ray (R. punctulata)	16	7		3
8	Ray, electric			(16-18)	1
9	Salmon			75(47-120)	1
10	Shark, cat	(8-9)			3
11		33	29		3
12				(30-37)	1

/1/ 3 fish under local anesthesia; values derived by optical manometer method

Contributors: (a) Van Liere, Edward J., and Hugh A. Lindsay, (b) Freis, Edward D., (c) Heisler, Charles R., and Jack R. Leonards, (d) Rodbard, Simon, (e) Woodbury, Robert A.

References: [1] Lehmann, G. 1925. *Tabulae biol.*, Berl. Bd 1, p. 142. [2] Woodbury, R. A., and W. F. Hamilton. 1937. *Am. J. Physiol.* 119:663. [3] Prosser, C. L., et al. 1950. *Comparative animal physiology*. W. B. Saunders, Philadelphia.

Part VI: INVERTEBRATES

Values in parentheses are ranges, estimate "c" (cf. Introduction).

	Animal (A)	Blood Pressure, mm Hg			Reference (D)
		Systolic (B)		Mean (C)	
1	Crab	4			1
2	Lobster			9	1
3	Lobster (Homarus sp)	10			1
4	Mussel (Anodonta sp)	3			2
5	Octopus (Octopus sp)			(25-80)	1
6	Octopus (O. vulgaris)	60			1
7	Sea hare (Aplysia sp)	(20-40)			3
8	Squid	(25-80)			1

Contributors: (a) Van Liere, Edward J., and Hugh A. Lindsay, (b) Rodbard, Simon, (c) Heisler, Charles R., and Jack R. Leonards, (d) Woodbury, Robert A.

References: [1] Lehmann, G. 1925. *Tabulae biol.*, Berl. Bd 1, p. 142. [2] Willem, V., and V. A. Minne. 1898. *Mém. Acad. Belgique* 57:8. [3] Straub, W. 1904. *Pflügers Arch.* 103:429.

56. PULSE CHARACTERISTICS: MAN (Concluded)

Part II: PULSE WAVES

Chamber or Vessel	Method	Maximal Velocity m/sec	No. of Waves	Wave Configuration	Reference
(A)	(B)	(C)	(D)	(E)	(F)
1 Right atrium	Right heart catheterization.		2	Small, positive wave in presystole; collapse during ventricular systole; positive wave at the opening of tricuspid valve; collapse in early diastole.	1-5
2 Left atrium	Left heart catheterization.		2	Small, positive wave in presystole; collapse during ventricular systole; positive wave at opening of mitral valve; collapse in early diastole.	5, 6
3 Left ventricle	or retrograde arterial catheterization.			lar systole, plus smaller vibrations.	
5 Pulmonary arteries	Right heart catheterization, or by electrokymography.	2	1	1 positive wave during ventricular systole.	4, 8
6 Pulmonary arterioles	Right heart catheterization, or by electrokymography.	2.75	2	1 in early systole and 1 in early diastole.	4, 8
7 Pulmonary veins	Left heart catheterization, or deduced from animal experimentation and from collateral data in man.		2	Small, positive wave in presystole; collapse during ventricular systole; positive wave at opening of mitral valve; collapse in early diastole.	1-5
8 Aortic arch	Retrograde arterial catheterization, or indirect pulse tracing.	4(3-5)	2	1 positive wave in systole, plus small wave in early diastole (dicrotic wave).	4, 9-12
9 Thoracic aorta	Retrograde arterial catheterization, or indirect pulse tracing.	5(4-6)	2	1 positive wave in systole, plus small wave in early diastole (dicrotic wave).	4, 9-12
10 Abdominal aorta	Retrograde arterial catheterization, or indirect pulse tracing.	6(5-7)	2	1 positive wave in systole, plus small wave in early diastole (dicrotic wave).	4, 9-12
11 Carotid arteries	Indirect pulse tracing, or deduced from animal experimentation and from collateral data in man.	5(4-6)	2	1 positive wave in systole, plus small wave in early diastole (dicrotic wave).	4, 11-13
12 Femoral arteries	Direct measurement, or indirect pulse tracing.	6(5-8)	2	1 positive wave in systole, plus small wave in early diastole (dicrotic wave).	4, 11-14
13 Bronchial arteries	Direct measurement, or indirect pulse tracing.	4(5-8)	2	1 positive wave in systole, plus small wave in early diastole (dicrotic wave).	4, 11-14
14 Tibial arteries	Direct measurement, or indirect pulse tracing.	8(6-10)	2	1 positive wave in systole, plus small wave in early diastole (dicrotic wave).	4, 11-14
15 Jugular veins	Indirect pulse tracing.	1.5	3	Small, positive wave in presystole, plus small wave in systole; collapse during ventricular systole; positive wave at opening of tricuspid valve; collapse in early diastole.	4, 12, 13
16 Venae cavae	Right heart catheterization.		2	Small positive wave in presystole; collapse during ventricular systole; positive wave at opening of tricuspid valve; collapse in early diastole.	4, 12, 13

11/ In cases of atrial septal defect

Contributor Luisada, Aldo A.

- References [1] Courmand, A., and H. A. Ranges 1941 Proc Soc Exp Biol, N. Y. 46:462. [2] Dexter, L., et al. 1950 J Clin Invest 29:602 [3] Lagerlöf, H., and L. Werkö, 1949 Scand J Clin. & Invest., Oslo 1:147. [4] Luisada, A. A. 1953 The heart beat. P. B. Hoeber, New York. [5] Luisada, A. A., and C. K. Liu. 1958. Intracardiac phenomena Grune and Stratton, New York. [6] Björk, V. O. 1954 Acta chir scand. 107:406. [7] Salans, A. H., et al. 1951. Circulation, N. Y. 4:510. [8] Fleischner, F. G., P. J. Romano, and A. A. Luisada. 1948 Proc. Soc. Exp Biol, N. Y. 67:535 [9] Baggett, H. C., and N. B. Dreyer. 1922. Am. J. Physiol. 63:94. [10] Dow, P., and W. F. Hamilton. 1939 Ibid. 125:60 [11] Hamilton, W. F. 1949. Textbook of human physiology. Philadelphia [12] Wiggers, C. J. 1950. Physiology in health and disease. Ed 5 Lea and Febiger, Philadelphia [13] Fonder, E. 1946. In W. H. Howell, ed. Textbook of physiology. W. B. Saunders, Philadelphia [14] Bramwell, J. C., and A. V. Hill. 1923. Heart, Lond 10:233

56. PULSE CHARACTERISTICS: MAN

Values in parentheses are ranges, estimate "c" (cf. Introduction).

Part I. BLOOD PRESSURES

Chamber or Vessel	Method	Ventricular Pressure		Mean Pressure mm Hg	Pulse Pressure mm Hg	Reference
		Systolic mm Hg	Diastolic mm Hg			
(A)	(B)	(C)	(D)	(E)	(F)	(G)
	retrograde arterial catheterization, or indirect measurement. ¹					
5 Pulmonary artery	Right heart catheterization.	20(11-29)	9(4-13)	(8-19)	(7-16)	3
6 Pulmonary arterioles ²	Right heart catheterization.	15	5	(5-13)	10	3-7
7 Pulmonary veins	Left or right heart catheterization ¹ , or deduced from animal experimentation and from collateral data in man.	4	8	5(3-8)		9, 14
8 Aorta	Retrograde arterial catheterization, or left heart catheterization.	120(100-150)	70(60-90)	95	50	11
9 Carotid arteries	Deduced from animal experimentation and from collateral data in man.	120	70	95	50	12, 14
10 Femoral arteries	Direct or indirect measurement.	150	80	115	70	15-17
11 Bronchial arteries	Direct or indirect measurement.	120	80	100	40	13, 15
12 Tibial arteries	Direct or indirect measurement.	140	80	110	60	12, 14, 15
13 Small veins	Deduced from animal experimentation and from collateral data in man			18(15-20)		18
14 Medium veins	Direct measurement.			5(3.5-7.0)		12, 14, 18
15 Venae cavae	Right heart catheterization.			-1(-5 to +4)		12, 17

/1/ In cases of atrial septal defect. /2/ "Pulmonary capillary" pressure or "wedge pressure."

Contributor Luisada, Aldo A.

References [1] Cournand, A., and H. A. Ranges 1941. *Proc. Soc. Exp. Biol.*, N. Y. 46:462 [2] Bloomfield, R. A., et al. 1946. *J. Clin. Invest.* 25:639. [3] Dexter, L., et al. 1950. *Ibid.* 29:602. [4] Werkö, L., et al. 1948. *Acta physiol. scand.* 16:75. [5] Werkö, L., et al. 1949. *Scand. J. Clin. & Invest.*, Oslo 1:147. [6] Werkö, L., et al. 1953. *Circul. Res.*, N. Y. 1:337. [7] Fowler, N. O., et al. 1953. *Am. Heart J.* 46:264. [8] Calazel, P., et al. 1951. *Bull. Johns Hopkins Hosp.* 88:20. [9] Luisada, A. A., and C. K. Liu. 1958. *Intracardiac phenomena*. Grune and Stratton, New York. [10] Björk, V. O. 1954. *Acta chir. scand.* 107:406. [11] Salans, A. H., et al. 1951. *Circulation*, N. Y. 4:510. [12] Ponder, E. 1946. In W. H. Howell, ed. *Textbook of physiology* W. B. Saunders, Philadelphia. [13] Weiss, A. 1951. *Arch. Kreislaufforsch.* 17:176. [14] Wiggers, C. J. 1950. *Physiology in health and disease* Ed. 5 Lea and Febiger, Philadelphia [15] Luisada, A. A. 1948. *Heart* Williams and Wilkins, Baltimore. [16] Luisada, A. A. 1953. *The heart beat* P. B. Hoeber, New York. [17] Hamilton, W. F. 1949. *Textbook of human physiology*. Ed 2. F. A. Davis, Philadelphia. [18] Landis, E. M. 1934. *Physiol. Rev.* 14:416

58. CIRCULATION TIME: VERTEBRATES

Circulation time = time required for an indicator substance introduced into the blood stream to traverse a particular vascular circuit. Values in parentheses are ranges, estimate "c" unless otherwise indicated (cf. Introduction).

Circuit		Indicator	Registration Time sec	Reference
(A)	(B)	(C)	(D)	
Man (Newborn)				
1 Arm to conjunctiva (first 5 da of life)	Fluorescein	10.2(5-14)	1	
2 Ductus venosus to lips	Fluorescein	4.4(3.3-6.0)	2	
3 Umbilical vein to lips	Fluorescein	4.8(3.1-7.0)	2	
4 Wrist to eye	Fluorescein	7.8(5-10)	3	
Man (Child)				
5 Arm to arm	Radioactive sodium	11(5-17)	4	
6 Arm to arm (14 da-6 yr)	Riboflavin	8.9(6.2-15.1)	5	
7 Arm to arm (6-13 yr)	Riboflavin	10.4(9.2-15.3)	5	
8 Arm to carotid sinus	Sodium cyanide	(9.0-13.5)	6	
9 Arm to foot	Riboflavin	28.1(25-35)	5	
10 Arm to lips (1-24 mo)	Fluorescein	7.0(5.0-9.1)	7	
11 Arm to lips (4-24 mo)	Fluorescein	6.5(5.0-8.5)	8	
12 Arm to lips (2-13 yr)	Fluorescein	11.5(7-16)	7	
13 Arm to lips (3-13 yr)	Fluorescein	8.5(5.0-12.5)	8	
14 Arm to lips (5-10 yr)	Fluorescein	10	9	
15 Arm to mouth (4-24 mo)	Fluorescein	6(4.5-7.5)	10	
16 Arm to mouth (2-4 yr)	Fluorescein	6.5(5.5-7.5)	10	
17 Arm to mouth (4-7 yr)	Fluorescein	8(6.5-9.5)	10	
18 Arm to mouth (7-12 yr)	Fluorescein	10.7(8.5-13.0)	10	
19 Arm to mouth (12-15 yr)	Fluorescein	12(10.1-14.0)	10	
20 Arm to opposite brachial artery (birth-2 yr)	Radioactive sodium	7(3-12)	4	
21 Arm to tongue or nose	Thiamine HCl	(4-8)	11	
22 Wrist to eye (2-12 mo)	Fluorescein	10(7.6-13.0)	3	
Man (Adult)				
23 Antecubital vein to radial artery	Evans' blue (T-1824)	17.5(14.5-23.0)	12	
24 Arm to arm	Brilliant vital red	(30-45)	13	
25 Arm to arm	Congo red	(8-12)	14	
26 Arm to arm (♂, rest)	Evans' blue (T-1824)	18.2(14.3-24.3)	15	
27 Arm to arm (♂, exercise)	Evans' blue (T-1824)	12.1(9.5-17.9)	15	
28 Arm to arm (♀, rest)	Evans' blue (T-1824)	15.4(10.2-20.2)	15	
29 Arm to arm (♀, exercise)	Evans' blue (T-1824)	9.9(7.3-11.5)	15	
30 Arm to arm (rest)	Evans' blue (T-1824)	22.5(17-29)	16	
31 Arm to arm (exercise)	Evans' blue (T-1824)	20.2(14-26)	16	
32 Arm to arm (42-58 yr, rest)	Evans' blue (T-1824)	18.4(10.3-24.8)	17	
33 Arm to arm (42-58 yr, exercise)	Evans' blue (T-1824)	12.3(9.8-16.1)	17	
34 Arm to arm	Evans' blue (T-1824)	30.1(22.4-37.8)	18	
35 Arm to arm	Evans' blue (T-1824)	19	19	
36 Arm to arm	Fluorescein	16.6(9-25)	20	
37 Arm to arm (15-79 yr)	Fluorescein	(12-26)	21	
38 Arm to arm	Fluorescein	18.6(13.5-25.6)	5	
39 Arm to arm	Radium-C	18(12-23)	22	
40 Arm to arm (13-65 yr)	Riboflavin	19.1(14.5-26.2)	5	
41 Arm to arm	Thorium-X	(10-20)	23	
42 Arm to brain (semirecumbent subject)	Thiopental	13(10-16)	24	
43 Arm to carotid sinus	Lobeline	10.6	25	
44 Arm to carotid sinus	Lobeline	8.5(5.0-12.5)	26	
45 Arm to carotid sinus	Lobeline	8.3(5-12)	27	
46 Arm to carotid sinus	Lobeline	9.5(7-12)	28	
47 Arm to carotid sinus	Lobeline	10.3(7-15)	29	
48 Arm to carotid sinus	Lobeline	9.9(6.4-13.5)	30, 31	
49 Arm to carotid sinus	Sodium cyanide	20.1(12-33)	32	
50 Arm to carotid sinus	Sodium cyanide	15.6(12.5-18.0)	33	
51 Arm to carotid sinus	Sodium cyanide	15.6(9-21)	34	
52 Arm to carotid sinus	Sodium cyanide	17(7-92)	35	
53 Arm to carotid sinus	Sodium cyanide	20.1(12.4-33.2)	36	
54 Arm to carotid sinus (warm subject)	Sodium cyanide	18.5(13-27)	37	
55 Arm to carotid sinus (cold subject)	Sodium cyanide	32.3(25-47)	37	
56 Arm to carotid sinus (pregnant)	Sodium cyanide	14.5(10-24)	38	

57. HEART PRESSURES: MAN

Part I: BIRTH TO TWO WEEKS

Additional values for some of the variables are available in the literature [2, 4]. Values in parentheses are ranges, estimate "c" (cf. Introduction).

Pressure	Right Heart		Left Heart		Reference
	No. of Subjects	Value mm Hg	No. of Subjects	Value mm Hg	
(A)	(B)	(C)	(D)	(E)	(F)
Atrium					
1 Systolic	8	0(-6 to +4)	5	1.6(-2 to +10)	1
2 Diastolic	8	-2(-8 to +2)	5	-0.4(-4 to +9)	1
3 Mean		(0-3)		(1-4)	2
Ventricle					
4 Systolic	8	35(20-48)			1
5 Diastolic	8	0(-10 to +8)			1
Pulmonary artery					
6 Systolic	13	41(16-80)			1,3
7 Diastolic	13	20(0-50)			1,3
8 Mean ¹	6	36(17-45)			3
Femoral artery ⁴					
9 Systolic			6	72(55-90)	3
10 Diastolic			6	46(20-65)	3
11 Mean ¹			5	54(40-80)	3

/1/ Electrically derived. /2/ Pressures obtained by direct puncture.

Contributors. Hultgren, Herbert N., and R. Eisenberg

References [1] Adams, F. H., and J. Lind. 1957. *Pediatrics*, Springfield, 19:431. [2] Rudolph, A. M., and G. G. Cayler. 1958. *Pediat. Clin. N. America* 5:907. [3] Rowe, R. D., and L. S. James. 1957. *J. Pediat.*, S. Louis 51:1. [4] Keith, J. D., R. D. Rowe, and P. Viad. 1958. *Heart disease in infancy and childhood*. Macmillan, New York.

Part II: ADULTS

Additional values for some of the variables are available in the literature [8-11]. Values in parentheses are ranges, estimate "c" (cf. Introduction).

Pressure	Right Heart		Left Heart ¹		Reference
	No. of Subjects	Value mm Hg	No. of Subjects ²	Value mm Hg	
(A)	(B)	(C)	(D)	(E)	(F)
Atrium ³					
1 Mean	24	2.8(1-5)	5	7.4(4-10)	B, C, 1, 2, D, E, 3, 4
2 A wave	9	5.6(2.5-7)	3	8.7(6-12)	B, C, 1; D, E, 3, 4
3 X wave	7	2.9(1-5.5)			1
4 C wave	5	3.8(1.5-6)	3	8.7(6-12)	B, C, 1, D, E, 3, 4
5 X' wave	8	1.7(0-5)			1
6 V wave	7	4.6(2-7.5)	4	10.3(7-14)	B, C, 1, D, E, 3, 4
7 Y wave	7	2.4(0-6)			1
Ventricle					
8 Peak systolic	33	24.7(17-32)	6	90.2(79-121)	B, C, 1, 2, 5, D, E, 3, 4
9 End diastolic	13	4(1-7)	6	8.8(4-14)	B, C, 2, D, E, 3, 4
10 Mean diastolic			6	6.2(3-10)	3, 4
Pulmonary artery					
11 Mean	31	14.7(9-19)			1, 2
12 Peak systolic	51	22.3(11-29)			1, 2, 6, 7
13 End diastolic	51	6.7(4-13)			1, 2, 6, 7
14 Pulmonary wedge ("capillary")	24	8.6(4.5-13)			1, 7

/1/ Zero point, mid-atrial level. /2/ Studies performed on anesthetized, open-chest patients. /3/ For atrial configurations, see p. 156.

Contributor Fowler, Noble O.

References [1] Fowler, N. O., R. N. Westcott, and R. C. Scott. 1953. *Am Heart J* 46:264. [2] Lagerlöf, H., and L. Werkö. 1948. *Acta physiol. scand.* 16:75. [3] Braunwald, E., H. L. Moscovitz, S. S. Amram, R. P. Lasser, and S. O. Sapin. A. Himmelstein, M. M. Ravitch, and A. J. Gordon. 1955. *Circulation*, N. Y. 12:69. [4] Gordon, A. J., E. Braunwald, and M. M. Ravitch. 1954. *Circul. Res.*, N. Y. 2:432. [5] Bloomfield, R. A., H. D. Laseen, A. Courand, E. S. Breed, and D. W. Richards, Jr. 1946. *J. Clin. Invest.* 25:639. [6] Borden, C. 1948. *Minnesota Med J.* 31:1216. [7] Dexter, L., J. W. Dow, F. W. Haynes, J. L. Whittenberger, B. G. Ferris, W. T. Goodale, and H. K. Hellens. 1950. *J. Clin. Invest.* 29:602. [8] Braunwald, E., A. P. Fishman, and A. Courand. 1956. *Circul. Res.*, N. Y. 4:100. [9] Wynn, A., M. B. Mathews, I. K. R. McMillan, and R. Daley. 1952. *Lancet*, Lond. 2:216. [10] Facquet, J., J. J. Welti, P. Ahomme, J. M. Lemoine, and P. Solignac. 1955. *Acta card.*, Brux 10:139. [11] Wood, P. 1958. *Brit Heart J*, 20:557.

58. CIRCULATION TIME: VERTEBRATES (Continued)

Circuit		Indicator	Registration Time sec	Reference
(A)		(B)	(C)	(D)
Man (Adult) (continued)				
118	Arm to medulla	Papaverine	20.8(15.4-27.0)	69
119	Arm to mouth	Calcium chloride	(9-16)	70
120	Arm to palpebral muscle	Succinylcholine	22.7(15-32)	71
121	Arm to perineum	Macasol	26(9.2-42.8) ^b	51
122	Arm to perineum	Macasol	21.5(12-32)	52
123	Arm to perineum	Macasol	21.2(10-34)	54
124	Arm to rectum	Fluorescein	20(19-21)	41
125	Arm to respiratory center (♂, 60-95 yr)	Lobeline	12.8	72
126	Arm to respiratory center (♀, 60-95 yr)	Lobeline	16.6	72
127	Arm to respiratory center	Sodium succinate	15.8(12.0-19.5)	73
128	Arm to respiratory center (student)	Sodium succinate	20(16.1-26.1)	74
129	Arm to respiratory center (trained athlete)	Sodium succinate	27.3(20.5-43.2)	74
130	Arm to respiratory center	Theophylline ethylenediamine	12.1(6.8-22.0)	75
131	Arm to respiratory center	Theophylline ethylenediamine	12.5(10-15)	76
132	Arm to throat	Calcium gluconate	18(12-29)	43
133	Arm to throat	Magnesium sulfate	(11-17)	77
134	Arm to throat	Macasol	15.8(7-22)	78
135	Arm to tongue	Calcium bromide	(10-17)	79
136	Arm to tongue	Calcium gluconate	(10.0-18.6)	80
137	Arm to tongue	Calcium gluconate	(7.1-24.0)	81
138	Arm to tongue	Calcium gluconate	12.7(8-16)	84
139	Arm to tongue	Calcium gluconate	12.1(10.3-15.3)	82
140	Arm to tongue (pregnant, 1st trimester)	Calcium gluconate	12.4(10.3-15.0)	83
141	Arm to tongue (pregnant, 2nd trimester)	Calcium gluconate	11.3(9.2-15.4)	83
142	Arm to tongue (pregnant, 3rd trimester)	Calcium gluconate	10.2(8.0-16.2)	83
143	Arm to tongue	Decholin	12.3(10-14)	84
144	Arm to tongue (pregnant)	Decholin	14.7(12.5-17.5)	84
145	Arm to tongue (recumbent subject)	Decholin	14(10-17)	24
146	Arm to tongue (semirecumbent subject)	Decholin	9(7-11)	24
147	Arm to tongue	Decholin	27.3(9-63)	39
148	Arm to tongue	Decholin	13.6(8.0-25.4)	85
149	Arm to tongue	Decholin	13.4(10-18)	86
150	Arm to tongue	Decholin	22.9(13-31)	87
151	Arm to tongue	Decholin	(7.4-19.0)	81
152	Arm to tongue	Decholin	(10-15)	88
153	Arm to tongue	Decholin	(8-14)	89
154	Arm to tongue	Decholin	18(15-20)	33
155	Arm to tongue	Decholin	13(10-16)	90
156	Arm to tongue	Fluorescein	15.6(10.8-21.0)	80
157	Arm to tongue	Intra-sul (sulfur and salts)	11.6(10-17)	65
158	Arm to tongue	Macasol	17.2(11-27)	58
159	Arm to tongue	Macasol	14.6(7-22)	52
160	Arm to tongue	Macasol	13.7(5-24)	54
161	Arm to tongue	Macasol	16.7(7.1-26.3) ^b	53
162	Arm to tongue (♂)	Magnesium sulfate	19.1(8.7-29.5) ^b	44
163	Arm to tongue (♀)	Magnesium sulfate	16.1(5.5-26.7) ^b	44
164	Arm to tongue	Magnesium sulfate	12.9(7.0-17.8)	91
165	Arm to tongue	Magnesium sulfate	(10.0-20.8)	80
166	Arm to tongue	Saccharin	12.6(7.5-16.0)	84
167	Arm to tongue	Saccharin	26.1(15-60)	39
168	Arm to tongue	Saccharin	11.9(9.0-15.8)	92
169	Arm to tongue	Saccharin	10.7(6-18)	62
170	Arm to tongue	Saccharin	(11.4-31.8)	93
171	Arm to tongue	Thiamine HCl	(5-23)	11
172	Foot to carotid sinus	Sodium cyanide	27.7(19-40)	34
173	Foot to carotid sinus	Sodium cyanide	36(15-95)	35
174	Foot to carotid sinus	Sodium cyanide	38.7(22-67)	32, 36
175	Foot to foot	Fluorescein	27.2(21-35)	20
176	Foot to groin	Radioactive sodium	18(16.2-19.8) ^b	94
177	Foot to groin	Radioactive sodium	21	95
178	Foot to groin (pregnant, 36 wk)	Radioactive sodium	36.8	96
179	Foot to groin (in labor)	Radioactive sodium	50.3	96
180	Foot to groin (puerperium)	Radioactive sodium	15.8	96

58. CIRCULATION TIME: VERTEBRATES (Continued)

Circuit		Indicator	Registration Time sec	Refer- ence
(A)		(B)	(C)	(D)
Man (Adult) (continued)				
57	Arm to cheek	Fluorescein	20.7(9-47)	39
58	Arm to conjunctiva	Fluorescein	10(7.0-15.6)	40
59	Arm to conjunctiva	Fluorescein	9(8-11)	41
60	Arm to conjunctiva	Fluorescein	11	42
61	Arm to ear	Evans' blue (T-1824)	14.2(10-21)	43
62	Arm to ear (♂)	Evans' blue (T-1824)	15.8(6.4-25.2) ^b	44
63	Arm to ear (♀)	Evans' blue (T-1824)	13.2(3.4-23.0) ^b	44
64	Arm to ear	Methylene blue ¹	(9-16)	45
65	Arm to face	Evans' blue (T-1824)	10.5(8.9-12.1)	46
66	Arm to face	Histamine	23(13-30)	47
67	Arm to face	Indigo carmine	9.1(6.6-11.6)	46
68	Arm to face	Methylene blue	12.9(10.8-15.0)	46
69	Arm to femoral artery (♂, sea level)	Evans' blue (T-1824)	16.5(11.2-21.5)	48
70	Arm to femoral artery (♀, sea level)	Evans' blue (T-1824)	15.3(11.1-23.0)	48
71	Arm to femoral artery (♂, high altitude)	Evans' blue (T-1824)	20.7(15.1-27.5)	48
72	Arm to femoral artery	Sodium ¹³¹ I	11.2(5.5-18.5)	49
73	Arm to foot	Fluorescein	27.2(21-35)	20
74	Arm to foot	Fluorescein	23	41
75	Arm to foot	Fluorescein	24.9(21-33)	30
76	Arm to foot	Fluorescein	(34-36)	42
77	Arm to foot	¹³¹ I albumin	20	51
78	Arm to foot	Macasol	28(10-48)	52
79	Arm to foot	Macasol	37.4(18.4-56.4) ^b	53
80	Arm to foot	Macasol	34(15-59)	54
81	Arm to foot	Radioactive sodium	45(15-90)	55
82	Arm to foot	Riboflavin	28.1(25-35)	5
83	Arm to hand	Fluorescein	(23-28)	42
84	Arm to hand	Fluorescein	24.3(21-30)	50
85	Arm to hand	Macasol	26	52
86	Arm to hand	Macasol	28.1(14.9-41.3) ^b	53
87	Arm to hand	Macasol	23.6(10-39)	54
88	Arm to left ventricle	Diodrast	8(6-9)	56
89	Arm to left ventricle	Diodrast and fluorodensograph	5.6(4.4-6.9)	57
90	Arm to right atrium (17-39 yr)	Radium-C	6.4(3-14)	22
91	Arm to right atrium (40-65 yr)	Radium-C	7.3(4-13)	22
92	Arm to right ventricle	Diodrast and fluorodensograph	1.6(1.3-2.1)	57
93	Arm to heart	Radium-C	6.7(2-14)	22
94	Arm to heart	Sodium ¹³¹ I	1.8(0.8-5.0)	49
95	Arm to lips (10-25 yr)	Fluorescein	20	9
96	Arm to lips (25-60 yr)	Fluorescein	25	9
97	Arm to lips	Fluorescein	(13.0-20.9)	41
98	Arm to lips (20-30 yr)	Fluorescein	15.4	41
99	Arm to lips (30-40 yr)	Fluorescein	15.8	41
100	Arm to lips (40-50 yr)	Fluorescein	16.3	41
101	Arm to lips (50-70 yr)	Fluorescein	17.0	41
102	Arm to lips (70 yr and over)	Fluorescein	18.2	41
103	Arm to lips (exercise)	Fluorescein	6	41
104	Arm to lips	Fluorescein	9	42
105	Arm to lung (recumbent or semirecumbent subject)	Ether	6(4-8)	24
106	Arm to lung	Ether	8(6-12)	58
107	Arm to lung	Ether	5.5(3-8) ^b	59
108	Arm to lung	Ether	(4-8)	60
109	Arm to lung	Ether	(6-9)	61
110	Arm to lung	Ether	6.5(4-8)	28
111	Arm to lung	Ether	6.3(2-11)	62
112	Arm to lung	Ether	7.2(3-12)	63
113	Arm to lung	Ether	5.7(4.1-9.0)	64
114	Arm to lung	Intra-sui (sulfur and salts)	7.1(4-10)	65
115	Arm to lung	Paraldehyde	6(2.8-9.3) ^b	66
116	Arm to medulla	Aminophyllin	12.4(7.1-20.4)	67
117	Arm to medulla	Aminophyllin	14.6(12-19)	68

/1/ Photoelectric method.

58. CIRCULATION TIME: VERTEBRATES (Continued)

Circuit	Indicator	Registration Time		Reference
		(B)	(C)	
(A)				
Dog (concluded)				
238 Right external jugular to left carotid			3.7(3.0-4.4)	123
239 Right external jugular to left external jugular vein	Lithium acetate ³		9.3	123
240 Right external jugular to left external jugular vein	Shadacol		7.8(4.9-10.2)	114
241 Right heart to left heart	Acetylcholine		5.5	119
242 Right ventricle to sino-atrial node	Acetylcholine		6(4.4-6.8)	121
243 Root of aorta to sino-atrial node	Acetylcholine		1.3(1.1-1.7)	121
244 Superior vena cava to sino-atrial node	Acetylcholine		7.8(6.4-9.9)	121
245 2-3 cm above root of aorta to sino-atrial node	Acetylcholine		3.4(2.6-4.3)	121
246 Total circulation	Thiocyanate		(8.9-12.8)	122
247 Total circulation			(10-11)	120
Rabbit				
248 Ear vein to carotid sinus	Sodium cyanide		3.9	114
249 Ear vein to ear vein	Methylene blue		6	114
250 Ear vein to ear vein	Lithium chloride		4.7	124
251 Ear vein to eye	Fluorescein		(5-6)	124
252 Ear vein to medulla	Sodium cyanide		1.9(1.3-5.2)	114
253 Femoral artery to femoral vein			2.8	123
254 Femoral vein to right atrium	Bismuth oil		6	118
255 Jugular vein to cranial vein			8.4	123
256 Left jugular vein to right femoral artery	Sodium chloride		(3.3-4.5)	123
257 Left jugular vein to right jugular vein	Sodium chloride		(4.1-6.1)	123
258 Left jugular vein to right renal artery	Sodium chloride		(3.8-5.4)	123
259 Left jugular vein to right renal vein	Sodium chloride		(10.8-12.7)	123
260 Portal circulation			(3.9-7.0)	123
261 Pulmonary circulation			2.9	123
262 Renal circulation			8	123
263 Right jugular vein to right carotid artery	Sodium chloride		(2.4-3.4)	123
264 Total circulation			10.5	123
Sheep				
265 Brachiocephalic artery to anterior caval channel	Iodine		(5-8)	125
266 Pulmonary circulation, fetal (ductus open)	Iodine		2.7(2-6)	126
267 Pulmonary circulation, at birth (ductus closed)	Iodine		1.4(1.0-3.3)	126
Chicken				
268 Sciatic vein to arterioles in systemic circuit (normal body temp., 41.5°C)	Acetylcholine		2.8(1.3-5.0)	127
269 Sciatic vein to arterioles in systemic circuit (body temp., 45°C)	Acetylcholine		1.2	127
270 Sciatic vein to arterioles in systemic circuit (body temp., 35°C)	Acetylcholine		4.0	127
271 Sciatic vein to arterioles in systemic circuit (body temp., 30°C)	Acetylcholine		8.2	127
272 Sciatic vein to arterioles in systemic circuit (body temp., 25°C)	Acetylcholine		12.2	127

[3] Also hexamethylene tetramine

Contributors (a) Rowlands, S., and S. R. Mukherjee. (b) Morris, Lester E., and Herrman L. Blumgart. (c) Wright, H. Payling. (d) Davies, Dean F., (e) Edholm, O. G., (f) Ruskin, Arthur

References: [1] Lesser, R. E., R. Meyer, and A. T. Henderson. 1952. *Am. J. Dis. Child.* 83:645. [2] Slobody, L. B., G. D. Rook, M. Levbar, and M. Morcy. 1950. *Pediatrics*, Springf. 6:254. [3] Oberst, B. B., and F. LaRoche. 1954. *J. Pediat.* S. Louis 54:580. [4] Hubbard, J. P., W. N. Preston, and R. A. Ross. 1942. *J. Clin. Invest.* 21:613. [5] Winsor, T., W. Adolph, W. Ralston, and G. M. Leiby. 1947. *Am. Heart J.* 34:80. [6] McGuire, J., and F. Goldman. 1937. *Ibid.* 14:230. [7] Witsberger, C. M., and H. G. Cohen. 1943. *J. Pediat.* S. Louis 22:726. [8] Gasul, B. M., J. J. Marino, and J. R. Christian. 1949. *Ibid.* 34:460. [9] Strömbeck, J. P., and O. Björk. 1945. *Nord med.* 26:878. [10] Bulgarelli, R., and F. Garioni. 1952. *Minerva pediat.*, Tor. 4:576. [11] Ruskin, A., and G. Dechard. 1947. *Am. J. M. Sc.* 213:337. [12] Kopelman, H., and G. de J. Lee. 1951. *Clin. Sc.*, Lond. 10:383. [13] Thompson, W. O., J. M. Alper, and P. K. Thompson. 1928. *J. Clin. Invest.* 5:605.

58. CIRCULATION TIME: VERTEBRATES (Continued)

Circuit		Indicator	Registration Time sec	Reference
(A)	(B)	(C)	(D)	
Man (Adult) (concluded)				
181 Foot to lung	Ether	(16-34)	59	
182 Foot to lung	Ether	32.5(22-40)	97	
183 Foot to tongue	Decholin	47.3(26-100)	97	
184 Foot to tongue (prone position)	Histamine	37.2(18-72)	98	
185 Hand to axilla (puerperium)	Radioactive sodium	11.7	96	
186 Lung to ear	Helium	4.6(3-6)	58	
187 Lung to ear	Oxyhemoglobin ²	5.2(4.8-7.1)	99	
188 Lung to ear	100% nitrogen	5.2(4.1-7.0)	100	
189 Lung to face	Amyl nitrite	19.5(14-25)	79	
190 Lung to face (10-20 yr)	Amyl nitrite	17.4	79	
191 Lung to face (20-30 yr)	Amyl nitrite	18.4	79	
192 Lung to face (30-40 yr)	Amyl nitrite	18.0	79	
193 Lung to face (40-50 yr)	Amyl nitrite	20.0	79	
194 Lung to face (50-60 yr)	Amyl nitrite	21.7	79	
195 Lung to face (60-70 yr)	Amyl nitrite	21.7	79	
196 Lung to face (70 yr and over)	Amyl nitrite	24.0	79	
197 Lung to medulla	Carbon dioxide	(11.5-16.5)	101	
198 Lung to medulla	Carbon dioxide	(5-10)	102	
199 Pulmonary artery to brachial artery	Evans' blue (T-1824)	(11-10-15)	12	
200 Pulmonary artery to brachial artery	Evans' blue (T-1824)	11.5(6.1-16.9) ^b	103	
201 Pulmonary artery to right ventricle	P32-labelled cells	(7.0-8.5)	104	
202 Pulmonary circulation, right atrium to brachial artery (17-39 yr)	Radium-C	10.7(6.5-17.0)	22	
203 Pulmonary circulation, right atrium to brachial artery (40-65 yr)	Radium-C	12.9(7.5-17.5)	22	
204 Pulmonary circulation, right ventricle to left ventricle	Thorotrast	1.7(1.4-2.5)	105	
205 Portal circulation, rectum to lung	Ether	(11-25)	106	
206 Portal circulation, rectum to lung	Ether	(20-25)	107	
207 Right heart to abdominal aorta	Evans' blue (T-1824)	7.8(5.6-9.0)	108	
208 Right heart to dorsalis pedis	Evans' blue (T-1824)	17.6(12.8-22.2)	108	
209 Right heart to ear	Evans' blue (T-1824)	8(6.3-9.3)	108	
210 Right heart to femoral artery	Evans' blue (T-1824)	9.7(7.5-10.7)	108	
211 Right heart to radial artery	Evans' blue (T-1824)	10.7(8.6-13.6)	108	
212 Thoracic aorta to abdominal aorta	Evans' blue (T-1824)	1.5(0.9-2.1)	108	
213 Thoracic aorta to dorsalis pedis	Evans' blue (T-1824)	11.8(8.1-22.3)	108	
214 Thoracic aorta to ear	Evans' blue (T-1824)	21.5(18.3-35.2)	108	
215 Thoracic aorta to femoral artery	Evans' blue (T-1824)	2.9(1.8-3.8)	108	
216 Thoracic aorta to radial artery	Evans' blue (T-1824)	24.3(20.3-36.5)	108	
217 Total circulation	P32-labelled cells	(7-9)	104	
Cat				
218 Femoral vein to carotid artery	p32-labelled cells	(3.0-3.5)	109	
219 Femoral vein to carotid artery	p32-labelled cells	(3-5)	110	
220 Femoral artery to carotid artery	p32-labelled cells	(9-11)	110	
221 Femoral artery to femoral vein	p32-labelled cells	(4-8)	110	
222 Left common carotid to right atrium	Radium-C	6(3.0-9.5)	111	
223 Subclavian vein to carotid artery	p31-albumin-labelled plasma	11.2	112	
224 Subclavian vein to carotid artery	p32-labelled cells	9.6	112	
Cow				
225 Mammary vein to opposite mammary vein	Fluorescein	52	113	
Dog				
226 External jugular vein to carotid sinus	Sodium cyanide	8.7(5.4-13.8)	114	
227 Femoral vein to carotid sinus	Sodium cyanide	11.7(8-16)	115	
228 Femoral vein to carotid sinus	Sodium cyanide	(6-15)	116	
229 Femoral vein to conjunctiva	Fluorescein	12.6(9-16)	117	
230 Femoral vein to right atrium	Bismuth oil	(16-18)	118	
231 Jugular vein to right heart	Shadacol	(1.0-2.5)	119	
232 Left heart to right heart	Shadacol	6	119	
233 Left ventricle or aorta to femoral artery	Thiocyanate	4	120	
234 Left ventricle (apex) to sino-atrial node	Acetylcholine	1.8(1.7-2.1)	121	
235 Leg vein to sino-atrial node	Acetylcholine	6.7(4.0-9.5)	121	
236 Main pulmonary artery to sino-atrial node	Acetylcholine	5.2(3.9-7.6)	121	
237 Pulmonary circulation (lesser circuit)		(4-6-2)	222	

/2/ Measured with Millikan oximeter.

58. CIRCULATION TIME: VERTEBRATES (Concluded)

- J. S. Cassella, and D. D. Smith. 1958. *J. Appl. Physiol.* 13:303. [88] Cottrell, J. D., and D. C. Cuddie. 1942. *Brit. M. J.* 1:170. [89] Wintermiller, M., J. Deutsch, and Z. Brüll. 1931. *Med. Klin., Berl.* 27:986. [90] Tarr, L., B. S. Oppenheimer, and R. V. Sager. 1933. *Am. Heart J.* 8:766. [91] Bernstein, M., and S. Simkins. 1939. *Ibid.* 17:218. [92] Fishberg, A. M., W. M. Hitzig, and F. H. King. 1933. *Proc. Soc. Exp. Biol., N. Y.* 30:651. [93] Esser, K. H., and K. Berliner. 1943. *Ann. Int. M.* 19:64. [94] Wright, H. P., S. B. Osborn, and D. G. Edmonds. 1948. *Lancet, Lond.* 2:767. [95] Levi, J. E., and E. F. Lewison. 1950. *Bull. Johns Hopkins Hosp.* 86:370. [96] Wright, H. P., S. B. Osborn, and D. G. Edmonds. 1950. *Surg. Gyn. Obst.* 90:481. [97] Oliveira, J. E. D., W. S. Hosseine, and A. Yasbek. 1952. *Hospital, Rio* 41:905. [98] Bock, A. V., D. B. Dill, and H. T. Edwards. 1930. *J. Clin. Invest.* 8:533. [99] Bierman, H. R. 1951. *Am. J. M. Sc.* 222:162. [100] Wexler, J., and J. L. Whittenberger. 1946. *J. Clin. Invest.* 25:447. [101] Bornstein, A. 1912. *Verh. deut. Kongr. inn. Med.* 29:457. [102] Gubner, R., S. Schnur, and J. H. Crawford. 1939. *J. Clin. Invest.* 18:395. [103] Doyle, J. T., J. S. Wilson, C. Lepine, and J. V. Warren. 1953. *J. Laborat. Clin. M.* 41:29. [104] Sutton, G. C., J. Karnell, and G. Nylin. 1950. *Am. Heart J.* 39:741. [105] Westermarck, N. 1942. *Acta radiol. Stockh.* 23:473. [106] Newman, H. F., and I. B. Cohen. 1949. *J. Laborat. Clin. M.* 34:674. [107] Gilges, B., and P. E. Teschan. 1952. *Ibid.* 40:537. [108] Grace, J. B., I. J. Fox, W. P. Crowley, Jr., and E. H. Wood. 1957. *J. Appl. Physiol.* 11:405. [109] Gray, J. A. B., and W. D. Paton. 1949. *J. Physiol., Lond.* 110:173. [110] Mukherjee, S. R., and S. Rowlands. 1951. *Lancet, Lond.* 1:1041. [111] Wolff, H. G., and H. L. Blumgart. 1929. *Arch. Neur. Psychiat., Chic.* 21:795. [112] Groom, A. C., W. B. Morris, and S. Rowlands. 1957. *J. Physiol. Lond.* 136:218. [113] Turner, C. W., and H. A. Herman. 1931. *Missouri Agr. Exp. Sta. Res. Bull.* 159. [114] Loevenhart, A. S., B. H. Schlomovitz, and E. G. Seybold. 1922. *J. Pharm., Lond.* 19:221. [115] Winder, C. V., H. O. Winder, and R. Gesell. 1933. *Am. J. Physiol.* 105:311. [116] Olson, W. H., H. Gutmann, S. O. Levinson, and H. Necheles. 1941. *War Med., Chic.* 1:830. [117] Wang, S. C., E. E. Painter, and R. R. Overman. 1946. *J. Exp. M.* 84:549. [118] Franck, O., and W. Alwens. 1910. *Munch. med. Wschr.* 57:950. [119] Hamilton, W. F., J. W. Moore, J. M. Kinsman, and R. G. Spurling. 1928. *Am. J. Physiol.* 84:338. [120] Henriques, V. 1913. *Biochem. Zschr.* 56:230. [121] Wilburne, M., J. G. Schlichter, M. Grossman, and F. Cisneros. 1947. *Am. J. Physiol.* 150:504. [122] Romm, S. O. 1924. *Pflügers Arch.* 202:14. [123] Stewart, G. N. 1894. *J. Physiol., Lond.* 15:1. [124] Fishback, D. B. 1941. *J. Laborat. Clin. M.* 26:1966. [125] Barclay, A. E., K. J. Franklin, and M. M. L. Prichard. 1944. *The foetal circulation*, Blackwell, Oxford. [126] Barclay, A. E., J. Barcroft, D. H. Barron, K. J. Franklin, and M. M. L. Prichard. 1942. *J. Physiol., Lond.* 101:375. [127] Rodbard, S., and A. Fink. 1948. *Am. J. Physiol.* 152:183.

59. CIRCULATION TIME IN PATHOLOGIC CONDITIONS. MAN

Circulation time = time required for an indicator substance introduced into the blood stream to traverse a particular vascular circuit. Values in parentheses are ranges, estimate "c" unless otherwise indicated (cf. Introduction).

Condition (A)	Circuit (B)	Indicator (C)	Registration Time sec (D)	Reference (E)
1 Anemia	Arm to carotid sinus	Lobeline	(3.5-8.0)	1
2 Anemia	Arm to lips	Fluorescein	12	2
3 Anemia	Arm to respiratory center	Theophylline-ethylene-diamine	9.2(6.0-15.9)	3
4 Anemia, hyperthyroidism	Arm to carotid sinus	Lobeline	12.6	4
5 Arteriosclerosis	Arm to conjunctiva	Fluorescein	23	5
6 Arteriosclerosis	Arm to foot	Fluorescein	(71-97)	5
7 Arteriosclerosis	Arm to hand	Fluorescein	(51-54)	5
8 Arteriosclerosis	Arm to lips	Fluorescein	29	5
9 Arteriosclerosis	Arm to tongue	Macasol	18.5	6
10 Arteriosclerosis	Arm to perineum	Macasol	30.0	6
11 Arteriosclerosis	Arm to hand	Macasol	29.5	6
12 Arteriosclerosis	Arm to foot	Macasol	42	6

- [14] Klein, O., and J. Heinemann. 1929. *Zbl. inn. Med.* 50:490. [15] Chapman, C. B., and R. S. Fraser. 1954. *Circulation*, N. Y. 9:57. [16] Ball, J. D., H. Kopelman, and A. C. Witham. 1952. *Brit. Heart J.* 14:363. [17] Chapman, C. B., and R. S. Fraser. 1954. *Circulation*, N. Y. 9:347. [18] Mill, H., and A. G. Kattus, Jr. 1956. *J. Laborat. Clin. M.* 48:413. [19] Wollheim, E. 1950. *Deut. med. Wschr.* 75:482. [20] MacGregor, A. G., and E. J. Wayne. 1951. *Brit. Heart J.* 13:80. [21] Koch, E. 1922. *Arch. klin. Med.* 140:39. [22] Blumgart, H. L., and S. Weiss. 1927. *J. Clin. Invest.* 4:15, 399. [23] Gerlach, J., P. M. Wolf, and H. J. Born. 1942. *Arch. exp. Path., Lpz.* 199:83. [24] McCarthy, F. M., and A. O. Hubbell. 1947. *Anesthesie, Par.* 15:29. [25] Stanojevic, L. 1938. *Zachr. Kreislauforsch.* 30:521. [26] Berliner, K. 1940. *Arch. Int. M.* 65:896. [27] Piccione, F. V., and L. J. Boyd. 1941. *J. Laborat. Clin. M.* 26:766. [28] Hagen, H. 1954. *Munch. med. Wschr.* 96:940. [29] Trimarco C. 1953. *Policlinico, sez. med.* 60:285. [30] Masbernard, A., M. Delestras, and A. Camelin. 1953. *Ann. med., Par.* 54:228. [31] Masbernard, A., and A. Camelin. 1953. *Klin. Wschr.* 31:455. [32] Smith, L. A., E. V. Allen, and W. McK. Craig. 1940. *Arch. Surg.* 41:1366. [33] Gargill, S. L. 1933. *N. England J. M.* 209:1089. [34] Robb, G. P., and S. Weiss. 1933. *Am. Heart J.* 8:650. [35] Bellis, C. J., A. K. Doss, and C. B. Craft. 1943. *Surgery* 13:35. [36] Smith, L. A., and E. V. Allen. 1941. *Proc. Mayo Clin.* 16:53. [37] Stead, E. A., Jr., and P. Kunkel. 1939. *Am. J. M. Sc.* 198:49. [38] Cohen, M. E., and K. J. Thomson. 1936. *J. Clin. Invest.* 15:607. [39] Nahl, M. M., and K. Lange. 1958. *Circulation*, N. Y. 17:922. [40] Fishback, D. B., S. A. Guttman, and E. B. Abramson. 1942. *Am. J. M. Sc.* 203 535. [41] Lange, K., and L. J. Boyd. 1943. *Ibid.* 206:438. [42] Sicher, W. D. 1943. *Proc. Mayo Clin.* 18:515. [43] Miller, B. J. 1948. *J. Laborat. Clin. M.* 33:910. [44] Dees, T. M., J. A. Rumsfeld, W. F. Miller, and C. B. Chapman. 1957. *J. Appl. Physiol.* 10:451. [45] Jablons, B., and J. Cohen. 1943. *Proc. Soc. Exp. Biol., N. Y.* 52:294. [46] Magistretti, M., E. Sartorelli, and E. Peirone. 1957. *Med. lavoro* 48:266. [47] Weiss, S., G. P. Robb, and H. L. Blumgart. 1929. *Am. Heart J.* 4:664. [48] Monge, C. C., A. Cazorla, G. M. Wittembury, Y. B. Sakata, and C. Rizo-Patón. 1955. *Acta physiol. lat. amer.* 5:198. [49] Shipley, R. A., and R. E. Clark. 1956. *Circul. Res., N. Y.* 4:456. [50] Brauner, F., and G. Niebauer. 1955. *Wien. med. Wschr.* 105:338. [51] Krieger, H., J. P. Storaasli, W. J. MacIntyre, W. D. Holden, and H. L. Friedell. 1952. *Ann. Surg.* 136:357. [52] Spier, L. C., I. S. Wright, and L. Saylor. 1936. *Am. Heart J.* 12:511. [53] Meneely, G. R., and A. Segaloff. 1947. *Am. J. M. Sc.* 214:176. [54] Kvale, W. F., and E. V. Allen. 1939. *Am. Heart J.* 18:519, 546. [55] Quimby, E. H., and B. C. Smith. 1944. *Science* 100:175. [56] Robb, G. P., and I. Steinberg. 1939. *Am. J. Roentg.* 41:1. [57] Moros, G., R. Villagordoa, R. Néri, N. Dorbecker, and G. Legault. 1956. *Arch. mal. coeur* 49:250. [58] Knutsen, J. R. B., B. E. Taylor, E. J. Ellis, and E. H. Wood. 1950. *Proc. Mayo Clin.* 25:405. [59] Hitzig, W. M. 1934-35. *Am. Heart J.* 10:1080. [60] Hitzig, W. M. 1934. *Proc. Soc. Exp. Biol., N. Y.* 31:935. [61] Miller, H. R. 1934. *Ibid.* 31:942. [62] Webb, G., W. Sheinfeld, and H. Collin. 1936. *Ann. Surg.* 104:460. [63] Scharf, R. 1957. *Bruxelles méd.* 37:1025. [64] Baer, S., and B. G. Slipakoff. 1936. *Am. Heart J.* 16:29. [65] Hodas, J. H., and C. E. Cucci. 1956. *Ibid.* 51:767. [66] Candel, S. 1938. *Ann. Int. M.* 12:236. [67] Koster, H., and S. J. Sarnoff. 1943. *J. Laborat. Clin. M.* 28:812. [68] Friedfeld, L., H. R. Marcus, and J. J. Vorsimer. 1948. *N. York State J. M.* 48:2047. [69] Elek, S. R., and S. D. Solarz. 1942. *Am. Heart J.* 24:821. [70] Kahler, H. 1929. *Wien. Arch. inn. Med.* 19:1. [71] Glover, N., and P. S. Marcus. 1958. *South. M. J.* 51:478. [72] Herbeuval, R., G. Luny, A. Larcan, and M. Tenette. 1957. *Presse méd.* 65:1498. [73] Greenfield, I. 1950. *Ann. Int. M.* 32:524. [74] Mellerowicz, H., and A. Peterman. 1952. *Medizinische* 31:1010. [75] Rose, D. N. 1951. *Brit. Heart J.* 13:56. [76] Mauro, G., and G. Nattero. 1955. *Fol. card., Milano* 8:241. [77] Neurath, O. 1937. *Zschr. klin. Med.* 132:134. [78] Nathanson, M. H., and S. A. Elek. 1947. *Am. Heart J.* 33:464. [79] Gross, D. 1945. *Ibid.* 30:19. [80] Lubic, L. G., and N. J. Sissman. 1952. *Ibid.* 44:443. [81] Wall, H. C. 1939. *Ibid.* 18:228. [82] Wilburne, M. 1942. *Ibid.* 24:816. [83] Manchester, B., and S. D. Loube. 1946. *Ibid.* 32 215. [84] Forlini, E. 1957. *Friuli med.* 12:91. [85] Tanzariello, R. 1955. *Boll. mal. orecchio* 73:367. [86] Pearce, M. L., A. E. Lewis, and M. R. Kaplan. 1952. *Circulation*, N. Y. 5:583. [87] Malt, R. A.,

58. CIRCULATION TIME. VERTEBRATES (Concluded)

- J. S. Cassells, and D. D. Smith. 1958. *J. Appl. Physiol.* 13:303. [88] Cottrell, J. D., and D. C. Cuddie. 1942. *Brit. M. J.* 1:70. [89] Winternitz, M., J. Deutsch, and Z. Brüll. 1931. *Med. Klin., Berl.* 27:986. [90] Tarr, L., B. S. Oppenheimer, and R. V. Sager. 1933. *Am. Heart J.* 6:766. [91] Bernstein, M., and S. Simkins. 1939. *Ibid.* 17:218. [92] Fishberg, A. M., W. M. Hitzig, and F. H. King. 1933. *Proc. Soc. Exp. Biol., N. Y.* 30:651. [93] Esser, K. H., and K. Berliner. 1945. *Ann. Int. M.* 19:64. [94] Wright, H. P., S. B. Osborn, and D. G. Edmonds. 1948. *Lancet, Lond.* 2:767. [95] Levi, J. E., and E. F. Lewison. 1950. *Bull. Johns Hopkins Hosp.* 56:370. [96] Wright, H. P., S. B. Osborn, and D. G. Edmonds. 1950. *Surg. Gyn. Obst.* 90:481. [97] Oliveira, J. E. D., W. S. Hassne, and A. Yasbek. 1952. *Hospital, Rio* 41:905. [98] Bock, A. V., D. B. Dill, and H. T. Edwards. 1930. *J. Clin. Invest.* 8:533. [99] Bierman, H. R. 1953. *Am. J. M. Sc.* 222:162. [100] Wexler, J., and J. L. Whittenberger. 1946. *J. Clin. Invest.* 25:447. [101] Bornstein, A. 1912. *Verh. deut. Kongr. inn. Med.* 29:457. [102] Gubner, R., S. Schmur, and J. H. Crawford. 1939. *J. Clin. Invest.* 18:395. [103] Doyle, J. T., J. S. Wilson, C. Lepine, and J. V. Warren. 1953. *J. Laborat. Clin. M.* 41:29. [104] Sutton, G. C., J. Karnell, and G. Nylan. 1950. *Am. Heart J.* 39:741. [105] Westermarck, N. 1942. *Acta radiol., Stockh.* 23:473. [106] Newman, H. F., and I. B. Cohen. 1949. *J. Laborat. Clin. M.* 34:674. [107] Giges, B., and P. E. Teschan. 1952. *Ibid.* 40:537. [108] Grace, J. B., I. J. Fox, W. P. Crowley, Jr., and E. H. Wood. 1957. *J. Appl. Physiol.* 11:405. [109] Gray, J. A. B., and W. D. M. Paton. 1949. *J. Physiol., Lond.* 110:173. [110] Mukherjee, S. R., and S. Rowlands. 1951. *Lancet, Lond.* 1:1041. [111] Wolff, H. G., and H. L. Blumgart. 1929. *Arch. Neur. Psychiat., Chic.* 21:795. [112] Groom, A. C., W. B. Morris, and S. Rowlands. 1957. *J. Physiol., Lond.* 136:218. [113] Turner, C. W., and H. A. Herman. 1931. *Missouri Agr. Exp. Sta. Res. Bull.* 159. [114] Loevenhart, A. S., B. H. Schlemovitz, and E. G. Seybold. 1922. *J. Pharm., Lond.* 19:221. [115] Winder, C. V., H. O. Winder, and R. Gesell. 1933. *Am. J. Physiol.* 105:311. [116] Olson, W. H., H. Gutmann, S. O. Levinson, and H. Necheles. 1941. *War Med., Chic.* 1:830. [117] Wang, S. C., E. E. Palster, and R. R. Overman. 1946. *J. Exp. M.* 84:549. [118] Franck, O., and W. Alwens. 1910. *Münch. med. Wochr.* 57:950. [119] Hamilton, W. F., J. W. Moore, J. M. Kinsman, and R. G. Spurling. 1928. *Am. J. Physiol.* 84:338. [120] Henriques, V. 1913. *Biochem. Zentr.* 56:230. [121] Wilburne, M., J. G. Schlichter, M. Grossman, and F. Cisneros. 1947. *Am. J. Physiol.* 150:504. [122] Romm, S. O. 1924. *Physiogr. Arch.* 202:14. [123] Stewart, G. N. 1894. *J. Physiol., Lond.* 15:1. [124] Fishback, D. B. 1941. *J. Laborat. Clin. M.* 26:1966. [125] Barclay, A. E., K. J. Franklin, and M. M. L. Prichard. 1944. *The foetal circulation.* Blackwell, Oxford. [126] Barclay, A. E., J. Barcroft, D. H. Barron, K. J. Franklin, and M. M. L. Prichard. 1942. *J. Physiol., Lond.* 101:375. [127] Rodbard, S., and A. Fink. 1948. *Am. J. Physiol.* 152:383.

59. CIRCULATION TIME IN PATHOLOGIC CONDITIONS. MAN

Circulation time = time required for an indicator substance introduced into the blood stream to traverse a particular vascular circuit. Values in parentheses are ranges, estimate "c" unless otherwise indicated (cf. Introduction).

Condition (A)	Circuit (B)	Indicator (C)	Registration Time sec (D)	Reference (E)
1 Anemia	Arm to carotid sinus	Lobeline	(5.5-8.0)	1
2 Anemia	Arm to lip	Fluorescein	12	2
3 Anemia	Arm to respiratory center	Theophylline-ethylene-diamine	9.2(6.0-15.9)	3
4 Anemia, hyperthyroidism	Arm to carotid sinus	Lobeline		
5 Arteriosclerosis	Arm to conjunctiva	Fluorescein	12.8	4
6 Arteriosclerosis	Arm to foot	Fluorescein	25	5
7 Arteriosclerosis	Arm to hand	Fluorescein	(71-93)	3
8 Arteriosclerosis	Arm to lip	Fluorescein	(51-54)	5
9 Arteriosclerosis	Arm to tongue	Macasol	29	5
10 Arteriosclerosis	Arm to perineum	Macasol	16.5	6
11 Arteriosclerosis	Arm to hand	Macasol	30.0	6
12 Arteriosclerosis	Arm to foot	Macasol	29.5	6
			42	6

59. CIRCULATION TIME IN PATHOLOGIC CONDITIONS- MAN (Continued)

	Condition	Circuit	Indicator	Registration Time sec	Reference
	(A)	(B)	(C)	(D)	(E)
13	Asthma	Arm to tongue	Decholin	110-15	7
14	Bronchial asthma	Arm to lung	Ether	6.9(2-10)	8
15	Bronchial asthma	Arm to tongue	Decholin	13.3(7-17)	8
16	Cardiac asthma	Arm to lips	Fluorescein	120-32	2
17	Cardiac asthma	Arm to lung	Ether	16(5-40)	8
18	Cardiac asthma	Arm to tongue	Decholin	29.2(12-73)	8
	Cardiac disease				
19	Unspecified	Arm to arm	Evans' blue (T-1824)	33.4(22.7-46.1)	9
20	Unspecified	Arm to arm	Fluorescein	33.5(19.0-45.2)	10
21	Unspecified	Arm to arm	Riboflavin	37.5(29.0-55.2)	10
22	Uncompensated	Arm to carotid sinus	Lobeline	30.2	4
23	Uncompensated	Arm to respiratory center	Theophylline ethylene-diamine	28.5(18-39)	11
24	Pseudocompensated	Arm to arm	Evans' blue (T-1824)	32	12
25	Compensated	Arm to arm	Evans' blue (T-1824)	21	12
26	Compensated	Arm to carotid sinus	Cyanide	18.3(12-22)	13
27	Compensated	Arm to carotid sinus	Lobeline	10(7-18)	14
28	Compensated	Arm to carotid sinus	Lobeline	13.4	4
29	Compensated	Arm to carotid sinus	Lobeline	10.3(6.4-19.4)	15
30	Compensated	Arm to respiratory center	Theophylline ethylene-diamine	14.5(13-16)	11
31	Compensated	Arm to tongue	Thiamine HCl	15-20	16
32	Compensated	Arm to tongue	Magnesium sulfate	13.0(6.6-17.0)	17
33	Compensated	Lung to face	Amyl nitrite	17-27	18
34	Decompensated	Arm to carotid sinus	Cyanide	38.3(16-55)	13
35	Decompensated	Arm to carotid sinus	Lobeline	23.7(16-31)	15
36	Decompensated	Arm to conjunctiva	Fluorescein	16-25	19
37	Decompensated	Arm to ear	Methylene blue	20-45	20
38	Decompensated	Rectum to lung	Ether	132-65	21
39	Without decompensation	Arm to ear	Methylene blue	9-15	20
40	Untreated	Arm to respiratory center	Sodium succinate	25.1(21-32)	22
41	Treated	Arm to respiratory center	Sodium succinate	19.2(16.7-22.0)	22
42	Cardiac failure	Arm to ear	Evans' blue (T-1824)	22.1(13-32)	23
43	Cardiac failure	Arm to tongue	Magnesium sulfate	30.3(17.2-42.0)	17
44	Cardiac failure	Arm to tongue	Calcium gluconate	120-40	24
45	Cardiac failure	Arm to tongue	Saccharin	119-34	25
46	Cardiac failure (rest)	Arm to carotid sinus	Lobeline	9.4	26
47	Cardiac failure (rest)	Arm to lung	Ether	6.5	26
48	Cardiac failure (rest)	Arm to tongue	Decholin	11.5	26
49	Cardiac failure (rest)	Arm to tongue	Decholin	27.9(11-69)	27
50	Cardiac failure (exercise)	Arm to tongue	Decholin	20.6(8-39)	27
51	Cardiac failure, mild	Arm to tongue	Thiamine HCl	8-25	16
52	Cardiac failure, moderate	Arm to tongue	Thiamine HCl	10-50	16
53	Cardiac failure, moderate to severe	Arm to femoral artery	Sodium ¹³¹ I	35.1(19-67)	28
54	Cardiac failure, severe	Arm to tongue	Thiamine HCl	113-55	16
55	Congestive cardiac failure	Arm to carotid sinus	Lobeline	19(7-46)	14
56	Congestive cardiac failure	Arm to medulla	Aminophyllin	24-38	29
57	Congestive cardiac failure	Arm to tongue	Calcium gluconate	117.0-60.8	30
58	Congestive cardiac failure	Arm to tongue	Decholin	21-70	30
59	Congestive cardiac failure	Arm to tongue	Decholin	25.4(18-40)	31
60	Congestive cardiac failure	Arm to tongue	Decholin	31(17-55)	32
61	Congestive cardiac failure	Arm to tongue	Fluorescein	32.8(21.2-78.2)	30
62	Congestive cardiac failure	Arm to tongue	Magnesium sulfate	116.8-71.0	30
63	Congestive cardiac failure (before hexamethonium)	Arm to tongue	Decholin	30(16-47)	33
64	Congestive cardiac failure (after hexamethonium)	Arm to tongue	Decholin	23(12-45)	33
65	Congestive cardiac failure, chronic	Arm to tongue	Decholin	35(22-48)	34
66	Congestive cardiac failure, chronic	Arm to lung	Ether	16(11-24)	34

59. CIRCULATION TIME IN PATHOLOGIC CONDITIONS. MAN (Continued)

Condition		Circuit	Indicator	Registration Time sec (D)	Reference (E)
(A)	(B)	(C)			
67	Cardiac disease (concluded)				
	Congestive cardiac failure, untreated	Arm to tongue	Saccharin	36.5(18-62)	35
68	Congestive cardiac failure, treated	Arm to tongue	Saccharin	19.4(13-33)	35
69	Congestive cardiac failure	Pulmonary artery to brachial artery	Evans' blue (T-1824)	24.9(14.5-35.3)	36
70	Left cardiac failure	Arm to lung	Ether	(4.5-7.0)	37
71	Left cardiac failure	Arm to tongue	Saccharin	(13.0-60.5)	37
72	Left cardiac failure	Arm to respiratory center	Theophylline ethylene- diamine	21(15.8-35.0)	3
73	Right cardiac failure	Arm to respiratory center	Theophylline ethylene- diamine	21(13-58)	3
74	Cardiac infarction, healed (rest)	Arm to arm	Evans' blue (T-1824)	18.3(12.0-25.7)	38
75	Cardiac infarction, healed (exercise)	Arm to arm	Evans' blue (T-1824)	14.4(9.5-27.2)	38
76	Rheumatic carditis	Arm to respiratory center	Theophylline ethylene- diamine	10.4(7.6-17.0)	3
77	Left ventricle failure	Antecubital vein to brachial artery	Evans' blue (T-1824)	54.5(35-88)	39
78	Left ventricle recovery	Antecubital vein to brachial artery	Evans' blue (T-1824)	35.5(21-63)	39
79	Mitral stenosis, failing	Antecubital vein to brachial artery	Evans' blue (T-1824)	41.5(35-49)	39
80	Mitral stenosis (recovery)	Antecubital vein to brachial artery	Evans' blue (T-1824)	28.5(24-34)	39
81	Mitral stenosis (rest)	Arm to arm	Evans' blue (T-1824)	27.1(17-40)	40
82	Mitral stenosis (exercise)	Arm to arm	Evans' blue (T-1824)	28.2(16-40)	40
83	Mitral stenosis	Pulmonary artery to brachial artery	Evans' blue (T-1824)	24.6	36
84	Tricuspid insufficiency	Pulmonary artery to brachial artery	Evans' blue (T-1824)	16.3	36
85	Compensated valvular disease	Arm to respiratory center	Theophylline ethylene- diamine	12.4(6.2-16.4)	3
86	Mitral lesions	Arm to throat	Macasol	26.9	41
87	Aortic lesions	Arm to throat	Macasol	20.2	41
88	Mitral and aortic lesions	" " " "	" " " "	24.8	41
89	Septal defects, uncomplicated ¹	" " " "	" " " "	110-133	42
90	Tetralogy of Fallot ¹	" " " "	" " " "	(3.8-4.8)	42
91	Cardiovascular disease	" " " "	" " " "	12(8-46)	43
92	Cardiovascular disease	" " " "	" " " "	" " " "	" " " "
93	Cardiovascular disease	Arm to tongue	Decholin	23.3(15-70)	44
		Arm to tongue	Intra-sul (sulfur and salts)	20(12-59)	44
94	Cardiovascular disease	Arm to tongue	K ⁴²	23(14-66)	44
95	Chronic pulmonary disease, untreated	Arm to tongue	Saccharin	21.6(12-60)	35
96	Chronic pulmonary disease, treated	Arm to tongue	Saccharin	11.4(10-14)	35
97	Cirrhosis of liver	Rectum to lung	Ether	(14-180)	19
98	Cirrhosis of liver (before surgery)	Rectum to lung	Ether	(26-42)	45
99	Cirrhosis of liver (after surgery)	Rectum to lung	Ether	(12-24)	45
100	Emphysema and silicosis	Arm to carotid sinus	Lobeline	14.5(7.8-29.5)	46
101	Fever	Arm to lips	Fluorescein	(9-13)	2
102	Hypertension, essential	" " " "	" " " "	11.5(6.0-26.8)	46
103	Hypertension	" " " "	" " " "	12.8(10-17)	3
104	Hypertension	" " " "	" " " "	" " " "	" " " "
105	Hypertension	Arm to tongue	Calcium gluconate	(8-16)	24
106	Hypertension	Arm to perineum	Macasol	14.4	6
107	Hypertension	Arm to hand	Macasol	22	6
108	Hypertension	Arm to foot	Macasol	23	6
109	Hypertension and arteriosclerosis	Arm to throat	Macasol	36.5	6
110	Hyperthyroidism	Arm to ear	Methylene blue	19.4	6
111	In children,			(8-9)	41
					20

Condition		Circuit	Indicator	Registration Time sec	Reference
(A)	(B)	(C)	(D)	(E)	
111	Hyperthyroidism	Arm to lips	Fluorescein	10.2	2
112	Hyperthyroidism	Arm to medulla	Aminophyllin	(9-11)	29
113	Hyperthyroidism	Arm to tongue	Calcium gluconate	(7-11)	24
114	Hyperthyroidism	Arm to tongue	Macasol	10.4	6
115	Hyperthyroidism	Arm to perineum	Macasol	15	6
116	Hyperthyroidism	Arm to hand	Macasol	16.3	6
117	Hyperthyroidism	Arm to foot	Macasol	21.3	6
118	Hyperthyroidism	Lung to face	Amyl nitrite	(8-13)	18
119	Hypothyroidism	Arm to lips	Fluorescein	26	2
120	Lumbar sympathectomy (before)	Arm to foot	Fluorescein	23(9-48)	47
121	Lumbar sympathectomy (after)	Arm to foot	Fluorescein	9.8(5-16)	47
122	Lung diseases	Lung to face	Amyl nitrite	(15-27)	18
123	Obesity	Arm to carotid sinus	Fluorescein	7.5(5.2-9.4)	46
124	Occlusive arterial disease	Arm to arm	Fluorescein	17.1(11-30)	47
125	Occlusive arterial disease	Foot to foot	Fluorescein	38.8(19-87)	47
126	Osler's disease	Arm to carotid sinus	Lobeline	19.6(7-27)	48
127	Pneumoconiosis	Arm to carotid sinus	Lobeline	(6-9)	1
128	Polycythemia	Arm to respiratory center	Theophylline ethylene-diamine	18.3(17.6-19.0)	3
129	Thromboangitis obliterans	Arm to foot	Fluorescein	(69-82)	5
130	Thromboangitis obliterans	Arm to hand	Fluorescein	55	5
131	Thromboangitis obliterans	Arm to lips	Fluorescein	24	5
132	Thromboangitis obliterans	Arm to tongue	Macasol	14.5(10-19)	6
133	Thromboangitis obliterans	Arm to perineum	Macasol	22.35(10-39)	6
134	Thromboangitis obliterans	Arm to hand	Macasol	24.5(15-39)	6
135	Thromboangitis obliterans	Arm to foot	Macasol	38(20-74)	6
136	Thrombophlebitis migrans	Arm to arm	Fluorescein	15.5(13-18)	47
137	Thrombophlebitis migrans	Arm to foot	Fluorescein	32.6(24-41)	47
138	Thyrototoxicosis, untreated	Arm to respiratory center	Theophylline ethylene-diamine	7.1(5.6-13.6)	3
139	Thyrototoxicosis, treated	Arm to respiratory center	Theophylline ethylene-diamine	14(10.0-17.4)	3
140	Varicose veins ²	Foot to tongue	Calcium gluconate	540	49

/2/ Measured while standing.

Contributors: (a) Rowlands, S., and S. R. Mukherjee, (b) Wright, H. Payling, (c) Edholm, O. G., (d) Ruskin, Arthur

References: [1] Piccione, F. V., and L. J. Boyd. 1941. J. Laborat. Clin. M. 26:766. [2] Lange, K., and L. J. Boyd. 1943. Am. J. M. Sc. 206:438. [3] Ross, D. N. 1951. Brit. Heart J. 13:56. [4] Stanojevic, L. 1938. Zschr. Kreislaufforsch. 30:521. [5] Sicher, W. D. 1943. Proc. Mayo Clin 18:515. [6] Slobody, L. B., G. D. Rook, M. Levbarg, and M. Morcy. 1950. Pediatrics, Springf. 6:254. [7] Cottrell, J. D., and D. C. Cuddle. 1942. Brit. M. J. 1:70. [8] Schuman, C., and H. G. Simmons. 1952. Ann. Int. M. 36:864. [9] Mill, H., and A. G. Katius, Jr. 1956. J. Laborat. Clin. M. 48:413. [10] Wright, H. P., S. B. Osborn, and D. G. Edmonds. 1950. Surg. Gyn. Obst. 90:481. [11] Mauro, G., and G. Nattero. 1955. Fel. card., Milano 8:241. [12] Wollheim, E. 1950. Deut. med. Wschr. 75:482. [13] Heller, B. I., and W. E. Jacobson. 1950. Am. Heart J. 39:180. [14] Berliner, K. 1940. Arch. Int. M. 65:896. [15] Masbernard, A., and A. Camelin. 1953. Klin. Wschr. 31:455. [16] Ruskin, A., and G. M. Decherd. 1947. Am. J. M. Sc. 213:137. [17] Bernstein, M., and S. Simpkins. 1939. Am. Heart J. 17:218. [18] Gross, D. 1945. Ibid. 30:19. [19] Fishback, D. B., S. A. Guttman, and E. B. Abramson. 1942. Am. J. M. Sc. 203:535. [20] Jablons, B., and J. Cohen. 1943. Proc. Soc. Exp. Biol., N. Y. 52:294. [21] Newman, H. F., and I. B. Cohen. 1949. J. Laborat. Clin. M. 34:674. [22] Greenfield, I. 1950. Ann. Int. M. 32:524. [23] Kaufmann, G. 1957. Cardiologia, Basel 30:102. [24] Goldberg, S. J. 1936. Am. J. M. Sc. 192:36. [25] Cohen, M. E., and K. J. Thomson. 1936. J. Clin. Invest. 15:607. [26] Hagen, H., and H. Jensen. 1954. Zschr. Kreislaufforsch

- 43 484. [27] Scharf, R. 1957. *Bruxelles mfd.* 37:1025. [28] Shipley, R. A., and R. E. Clark. 1956. *Circul. Res.* N. Y. 4:456. [29] Friedfeld, L., H. R. Marcus, and J. J. Vorgimer. 1948. *N. York State J. M.* 48:2047. [30] Lubic, L. G., and N. J. Siesman. 1952. *Am. Heart J.* 44 443. [31] Pearce, M. L., A. E. Lewis, and M. R. Kaplan. 1952. *Circulation*, N. Y. 5:583. [32] Moyer, J. H., et al. 1952. *J. Clin. Invest.* 31:267. [33] Kelly, R. T., E. D. Freis, and T. E. Higgins. 1953. *Circulation*, N. Y. 7:169. [34] Groseman, J., R. E. Weston, J. P. Halperin, and J. Lester. 1950. *J. Clin. Invest.* 29:1320. [35] Kopelman, H. 1951. *Brit. Heart J.* 13:501. [36] Doyle, J. T., J. S. Wilson, C. Lepine, and J. V. Warren. 1953. *J. Laborat. Clin. M.* 41:29. [37] Illisig, W. M., F. H. King, and A. M. Fishberg. 1935. *Arch. Int. M.* 55:112. [38] Chapman, C. B., and R. S. Fraser. 1954. *Circulation*, N. Y. 9:347. [39] Kopelman, H., and G. deJ. Lee. 1951. *Clin. Sci., Lond.* 10:383. [40] Ball, J. D., H. Kopelman, and A. C. Witham. 1952. *Brit. Heart J.* 14 363. [41] Nathanson, M. H., and S. R. Elek. 1947. *Am. Heart J.* 33 464. [42] McGuire, J., and F. Goldman. 1937. *Idid.* 14 230. [43] Hodas, J. H., and C. E. Cucchi. 1956. *Idid.* 51:767. [44] Conn, H. L., D. F. Helman, and C. B. Joyner. 1957. *Circulation*, N. Y. 15:245. [45] Gigen, B., and P. E. Teschan. 1952. *J. Laborat. Clin. M.* 40:537. [46] Masbernard, A., M. Delatras, and A. Camelin. 1953. *Ann. mfd., Par.* 54:228. [47] MacGregor, A. G., and E. J. Wayne. 1951. *Brit. Heart J.* 13:80. [48] Djordjevic, B. S., and M. Djordjevic-Joksic. 1953. *Arch. mal. coeur* 46:46. [49] Heller, R. E. 1942. *Surg. Gyn. Obst.* 74:1118.

60. CEREBRAL BLOOD FLOW: MAN, MONKEY

Subjects recumbent, lines I-B Method N_2O = nitrous oxide, intermittent sampling (Kety); Kr^{85} = radioactive krypton, intermittent sampling (Lassen and Munch), I-B = intercalated bubble flow meter. Values determined in vivo Vascular resistance = $\frac{\text{mean arterial pressure}}{\text{blood flow}}$, Cerebral arterio-venous O_2 difference (the quantity of O_2

removed by the brain from each 100 ml of blood flowing through the brain) = $100 \times \frac{\text{column F}}{\text{column E}}$. Values in parentheses are ranges, estimate "e" (cf Introduction).

Condition	Sex	Age yr	Method	Blood Flow ml/100 g/min	O_2 Consumption ml/100 g/min	Vascular Resistance mm Hg/ml/100 g/min	Reference
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)
Man							
1 Normal, alert ¹	♂	25	N_2O	54(40-67)	3.3(2.6-4.2)	1.6(1.2-2.5)	1-3
2 Normal, alert	♂, ♀	29	N_2O	52(40-64)	3.1(2.2-3.9)	1.8	4
3		56	N_2O	46(30-64)	2.9(2.0-3.9)	2.2	
4	10♂, 10♀	38(17-55)	Kr^{85}	52(33-67)	3.4(2.4-4.5)	1.8(1.3-2.3)	5
5 Passive hyperventilation with air	♂	26(23-31)	N_2O	41(36-47)	4.7(4.0-5.1)	2.4(1.7-3.5)	1
6 Inhaling 5-7% CO_2	♂	25	N_2O	93(65-141)	5.3(2.4-3.9)	1.1(0.7-1.4)	3
7 Inhaling 85-100% O_2	♂	25	N_2O	45(34-55)	3.2(2.6-4.4)	2.2(1.8-2.7)	
8 Inhaling 10% O_2	♂	25	N_2O	73(54-93)	3.2(2.6-3.5)	1.1(0.8-1.6)	
9 Cerebral arterio-sclerosis			N_2O	41(31-56)	2.8(1.7-3.6)	3.0(1.9-3.5)	6
10 Anesthetized (thiopental)			N_2O	60(33-117)	2.1(1.5-3.0)	1.3(0.6-2.1)	7
Monkey							
11 Anesthetized ² (light barbiturate)			I-B	47(33-74)	3.7(2.5-4.5)		8
12 Anesthetized (thiopental)			I-B	24(14-46)	2.2(0.9-3.8)		

[1] Approximately same values found in subjects with essential hypertension [9] and with schizophrenia [10].

[2] Active reflexes, spontaneous movements

Contributors: (a) Kety, Seymour S., (b) Schmidt, Carl F., (c) Patterson, John L., Jr.

References: [1] Kety, S. S., and C. F. Schmidt. 1946. J. Clin. Invest. 25:107. [2] Kety, S. S., and C. F. Schmidt. 1948. Ibid. 27:476. [3] Kety, S. S., and C. F. Schmidt. 1948. Ibid. 27:484. [4] Heyman, A., J. L. Patterson, Jr., T. W. Duke, and L. L. Battey. 1953. N. England J. M. 249:223. [5] Lassen, N. A., and O. Munck. 1955. Acta physiol. scand. 33:30. [6] Freyhan, F. A., et al. 1951. J. Nerv. Ment. Dis. 113:449. [7] Wechsler, R. L., et al. 1951. Anesthesiology 12:308. [8] Schmidt, C. F., S. S. Kety, and H. H. Pennes. 1945. Am. J. Physiol. 143:33. [9] Kety, S. S., et al. 1948. J. Clin. Invest. 27:511. [10] Kety, S. S., et al. 1948. Am. J. Psychiat. 104:765.

61. RENAL BLOOD FLOW. MAMMALS

Abbreviations PAH = p-aminohippurate plasma clearance, UV/P = diodrast plasma clearance.

Part I MAN

Effective plasma flow determined from p-aminohippurate, or diodrast, plasma clearance. Effective blood flow calculated from effective plasma flow divided by 1 minus hematocrit ($EBF = \frac{EPF}{1 - \text{hematocrit}}$). Values in parentheses are ranges, estimate "c" unless otherwise indicated (cf. Introduction).

Subjects		No. of Observations	Method	Plasma Flow ml/min/1.73 sq m body surface area	Blood Flow ml/min/1.73 sq m body surface area	Reference
Age	No. and Sex					
(A)	(B)	(C)	(D)	(E)	(F)	(G)
1 2-40	91♂	258	PAH, UV/P	655(492-818) ^b	1166(910-1422) ^b	1
2 2-40	31♀	91	PAH, UV/P	600(498-702) ^b	940(756-1124) ^b	
3 29-29	10♂	27	UV/P	786(529-993)	1339(860-1780)	2
4 30-39	10♂	21	UV/P	676(323-956)	1145(521-1622)	
5 40-49	9♂	14	UV/P	689(509-846)	1150(868-1342)	
6 50-59	7♂	21	UV/P	586(343-804)	987(746-1341)	
7 60-69	2♂	5	UV/P	584(498-680)	997(844-1192)	
8 22-51	11♀	16	UV/P	596(441-782)	973(718-1264)	
9 23-57	2♂, 21♀	13	PAH	497(361-648)	881(631-1179)	3
10 20-29	9♂	9	UV/P	614(481-724)	1077(859-1353)	4
11 30-39	9♂	9	UV/P	649(519-804)	1181(931-1493)	
12 40-49	10♂	10	UV/P	574(396-736)	1008(737-1252)	
13 50-59	11♂	11	UV/P	500(341-617)	849(631-1007)	
14 60-69	10♂	10	UV/P	442(253-534)	775(453-941)	
15 70-79	9♂	9	UV/P	354(234-519)	589(411-845)	
16 80-89	12♂	12	UV/P	289(147-462)	475(237-732)	

Contributors. (a) Riatt, Edwin P., (b) Davies, Dean F., (c) Seikurt, Ewald E., (d) Lauson, Henry D., (e) Blake, William D.

References: [1] Smith, H. W. 1951. The kidney structure and function in health and disease Oxford University Press, New York [2] Goldring, W., H. Chasis, H. A. Ranges, and H. W. Smith 1940. J. Clin. Invest. 19:739 [3] Pfeiffer, J. B., Jr., and H. G. Wolff. 1950. Ibid. 29:1227 [4] Davies, D. F., and N. W. Shock 1950. Ibid. 29:496.

61. RENAL BLOOD FLOW: MAMMALS (Concluded)

Part II. DOG, RABBIT, RAT

Animal	No. of Subjects	Condition	Method	Plasma or Blood	Flow				Reference
					ml/min (both kidneys)	ml/min/kg kidney weight	ml/min/kg body weight	ml/min/sq m body surface area	
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)
1 Dog	6	Unanesthetized	Calculated	Plasma	212 ¹	2.20 ²	15.6		1, 2
2			Thermotromuhr	Blood	384	4.00	28.6		
3	51	Unanesthetized	Calculated	Plasma	158 ¹	1.73 ²	11.5		3-6
4			Direct venous out-flow	Blood	286	3.15	21.0		
5	75	Unanesthetized	PAH, UV/P	Plasma	166	1.96 ²	13.5	266	7
6			Calculated	Blood	302 ¹	3.46	24.6	484	
7	11	Unanesthetized	Calculated	Plasma	156 ¹	2.20 ³	16.0	248	8
8			Urea extraction	Blood	284	4.00	18.0	450	
9	77	Unanesthetized	Calculated	Plasma			9.9 ¹	250	9
10			Phenol red extraction	Blood			18.0	460	
11	13	Pentobarbital anesthesia	Calculated	Plasma	212 ¹	2.42	16.4		8
12			Direct venous out-flow	Blood	376	4.30	29.2		
13	13	Pentobarbital anesthesia	Calculated	Plasma		1.76 ¹			10
14			Rotameter: arterial inflow	Blood		3.20			
15	22	Pentobarbital anesthesia	Calculated	Plasma	130 ¹	1.54	9.6		11
16			Rotameter: venous outflow	Blood	236	2.80	17.6		
17	10	Chloralose anesthesia	Calculated	Plasma	197 ¹	1.82 ²	12.8		12
18			Bubble meter: arterial inflow	Blood	358	3.30	23.4		
19 Rabbit	37	Unanesthetized	PAH, UV/P	Plasma	45	2.40 ⁴	15.4 ⁵	240	13-15
20			Thermotromuhr	Blood	82	4.30	27.6	438	
21 Rat	100	Unanesthetized	PAH, UV/P	Plasma	4.6	3.60 ⁴	27.7	167	16-20
22			Calculated	Blood	8.4 ¹	7.34	51.0	296	

1/1 Calculated for an average hematocrit ratio of 0.45. 1/2 Kidney weight in grams = 7.1 x body weight in kilograms. 1/3 Kidney weight in grams = 96.2 x sq m body surface area. 1/4 Kidney weight in grams = 8.0 x body weight in kilograms. 1/5 Calculated for an average body weight of 2.9 kilograms.

Contributors (a) Selkurt, Ewald E., (b) Abramson, David J.

References [1] Medes, G., and J. F. Herrick 1933. *Proc. Soc. Exp. Biol.*, N. Y. 31:116. [2] Herrick, J. F., F. C. Mann, and H. L. Sheehan. 1939. *J. Pharm. Exp. Ther.* 66:73. [3] Mason, M. F., A. Blalock, and T. R. Harrison 1937. *Am. J. Physiol.* 118:667. [4] Levy, S. E., and A. Blalock. 1937. *Ibid.* 118:368. [5] Levy, S. E., R. A. Light, and A. Blalock 1938. *Ibid.* 122:38. [6] Levy, S. E., and A. Blalock. 1938. *Ibid.* 122:609. [7] Houck, C. R. 1948. *Ibid.* 153:169. [8] Van Slyke, D. D., C. P. Rhoads, A. Miller, and A. S. Alving. 1934. *Ibid.* 109:316. [9] Corcoran, A. C., and L. H. Page. 1939. *Ibid.* 126:354. [10] Shipley, R. E., and R. S. Study 1951. *Ibid.* 167:676. [11] Study, R. S., and R. E. Shipley 1950. *Ibid.* 161:442. [12] Conn, H. L., Jr., and K. Markley 1950. *Ibid.* 160:547. [13] Walker, A. M., C. F. Schmidt, K. A. Elsom, and C. G. Johnston 1937. *Ibid.* 118:95. [14] Forster, R. P. 1947. *Ibid.* 150:523. [15] Brod, J., and J. H. Sirota 1949. *Ibid.* 157:31. [16] Friedman, S. M., J. R. Polley, and C. L. Friedman. 1947. *Ibid.* 150:340. [17] Friedman, M. 1947. *Ibid.* 148:387. [18] Lippman, R. W. 1948. *Ibid.* 152:27. [19] Dicker, S. E., and H. Heller. 1945. *J. Physiol., Lond.* 103:449. [20] Dicker, S. E., and H. Heller. 1946. *Ibid.* 104:353.

62. HEPATIC BLOOD FLOW: MAMMALS

Values in parentheses are ranges, estimate "c" (cf. Introduction).

Part I: MAN

	Condition	Method	Blood Flow		Reference
			ml/min/sq m body surface area	ml/min/subject	
	(A)	(B)	(C)	(D)	(E)
1	Normal	Bromsulphalein	833(600-1200)		1,2
2			836(500-1290)		3
3			865(640-1070)		4
4			883(744-1085)		5
5			895(756-1098)	(1192-1737)	6
6			1530(970-2720) ¹		3
7			798		7
8			(1390-2500)		8
9		Bromsulphalein minus const. inf. ²		676(503-1055)	9
10		Bromsulphalein, urea excretion	851(641-1210)		10
11		Urea excretion	800(600-1260)		1,2
12		Colloidal gold, Au ¹⁹⁸		1425(1120-2010)	11
13		Colloidal gold		731(533-1668)	12
14			597(357-951)	1090(542-1779)	13
15		Galactose clearance	806	1460	14
16		P ³² labelled CrPO ₄		(1500-1800)	15
17	Pregnant	Bromsulphalein	898(621-1425)	(1092-2065)	6
18	Praeanesthetization	Bromsulphalein		1258(746-2140)	16
19	Anesthetized	Bromsulphalein		884(467-1850)	16
20	Anesthetized, low spinal	Bromsulphalein	742(510-1161)		5
21	Anesthetized, high spinal	Bromsulphalein	590(295-919)		5
22	Cirrhosis	Colloidal gold, Au ¹⁹⁸		525(315-775)	13
23		Colloidal gold		437(96-1054)	12
24			212		13
25		Galactose clearance		(490-1660)	14
26	Cirrhosis, Laennec's	Bromsulphalein		695(470-710)	2
27	Hepatitis	Colloidal gold	266		13
28		Galactose clearance		(1920-2557)	14

/1/ Peripheral venous bromsulphalein less than 1 mg/100 ml. /2/ Bromsulphalein minus constant infusion rate, 2.5 mg/min/sq m.

Contributors: (a) Lichtman, S S, (b) Zierler, Kenneth L.

References: [1] Myers, J. D. 1947. J. Clin. Invest. 24:890. [2] Myers, J. D. 1950. Ibid. 29:1421. [3] Sherlock, S., A. G. Bearn, B. H. Billing, and J. C. S. Paterson. 1950. J. Laborat. Clin. M. 35:923. [4] Bradley, S. E., F. J. Ingelfinger, G. P. Bradley, and J. J. Curry. 1945. J. Clin. Invest. 24:890. [5] Mueller, R. P., R. B. Lynn, and S. M. Sancetta. 1952. Circulation. N. Y. 6:894. [6] Munnell, E. W., and H. C. Taylor, Jr. 1947. J. Clin. Invest. 26:952. [7] Wilkins, R. W., J. W. Culbertson, and A. A. Rymut. 1952. Ibid. 31:529. [8] Habib, D. V., E. M. Papper, H. F. Fitzpatrick, P. Lawrence, C. McC. Smythe, and S. E. Bradley. 1951. Surgery 30:241. [9] Culbertson, J. W., T. L. Welton, and J. G. Easton. 1951. J. Laborat. Clin. M. 38:801. [10] Bondy, P. K., D. F. James, and B. W. Farrar. 1949. J. Clin. Invest. 28:238. [11] Vetter, H., R. Falkner, and A. Neumayr. 1954. Ibid. 33:1594. [12] Krook, H. 1956. Acta med. scand., Suppl. 318.6. [13] Burke, J. S., and M. L. Gliedman. 1959. Gastroenterology 36:112. [14] Tygstrup, N., and K. Winkler. 1958. Clin. Sc., Lond. 17:1. [15] Dobson, E. L., G. F. Warner, C. R. Finney, and M. E. Johnston. 1953. Circulation, N. Y. 7:690. [16] Shackman, R., I. G. Graber, and D. G. Melrose. 1953. Clin. Sc., Lond. 12:307.

62. HEPATIC BLOOD FLOW: MAMMALS (Concluded)

Part II: CAT, DOG, RAT

Animal	Condition	Method	Blood Flow			Reference
			ml/min/animal	ml/min/kg body wt	ml/min/100 g liver	
(A)	(B)	(C)	(D)	(E)	(F)	(G)
1	C				133-481	1
2	D					
3						
4		measured				
5		Bromsulphalein; plasma dye clearance	146(72-272)	48.6(37.7-65.4)		4
6	Anesthetized	Blood collected and measured	390(250-534)	44(26-85)	95(64-144)	5
7		Bromsulphalein		35(20-50)		6
8		Bromsulphalein	690(252-1878)	42(16-111)	140(57-358)	7
9		Bromsulphalein	(403-1304)			8
10		Stromuhr (hepatic artery)	143(101.3-187.0)	7.8(5.0-11.8)		9, 10
11		Stromuhr (portal vein)	268(123.5-565.0)	31.7(8.2-53.2)	64	9, 10
12						
13						
14						
15						
16		(hepatic artery)				
17		Thermostromuhr (hepatic artery)	144(43-430)	7.5(2-16)	25(11-44)	14
18		Thermostromuhr (portal vein)	150(80-200)			13
19		Thermostromuhr (portal vein)	147(52-400)	15.5(5-37)		12
20	Anesthetized, pre- Eck fistula	Bromsulphalein	(403-922)			8
21	Anesthetized, Eck fistula	Bromsulphalein	(202-439)			8
22	Basal, 3-da post- operative	Bromsulphalein		42.5		15
23		Thermostromuhr	330(157-520)	26(10-33)	58.5(43-70)	16, 17
24	Unanesthetized	Urea excretion	383(241-705)	32(18-48)		18
25	Anesthetized	Thermoelectric			79(75-92)	19
					42	19

Contributors (a) Lichtman, S. S., (b) Abramson, David L.

References: [1] Barcroft, J. and L. E. Shore 1912-13. *J. Physiol., Lond.* 45 296. [2] Herrick, J. F., F. C. Mann, H. E. Essex, and E. J. Baltes. 1934. *Am. J. Physiol.* 109:52. [3] Bialock, A., and M. F. Mason. 1936. *Ibid.* 117:328. [4] Pratt, E. B., F. D. Burdick, and J. H. Holmes. 1952. *Ibid.* 171:471. [5] MacLeod, J. J., and R. G. Pearce. 1914. *Ibid.* 35 87. [6] Giger, B., W. C. Morse, W. S. Sharon, and J. Wyn. 1953. *Am. J. Med.* 14:506. [7] Werner, A. Y., and S. M. Horvath. 1952. *J. Clin. Invest.* 31:433. [8] Hallett, E. B., G. W. Holton, J. C. S. Paterson, and J. A. Schilling. 1952. *Surg. Gyn. Obst.* 95:401. [9] Burton-Optiz, R. 1910. *Q. J. Exp. Physiol., Lond.* 3:297. [10] Burton-Optiz, R. 1911. *Ibid.* 4:113. [11] Soskin, S., H. E. Essex, J. F. Herrick, and F. C. Mann 1938. *Am. J. Physiol.* 124 558. [12] Grindlay, J. H., J. F. Herrick, and F. C. Mann. 1941. *Ibid.* 132 489. [13] Schweigk, H. 1932. *Arch. exp. Path., Lpz.* 188:693. [14] Grodins, F. S., S. L. Osborne, A. C. Ivy, and L. Goldman. 1941. *Am. J. Physiol.* 132 375. [15] Bollman, J. L., M. Khattab, R. Thors, and J. H. Grindlay 1953. *Arch. Surg.* 66 562. [16] Grab, W., S. Janssen, and H. Rein. 1929. *Klin. Wochr.* 8 1539. [17] Grab, W., S. Janssen, and H. Rein. 1929. *Klin. Wochr.* 8 1539. [18] Lipscomb, A., and L. A. Crandall, Jr. 1947. *Am. J. Physiol.* 148 302. [19] Birnie, J. H., and J. Grayson. 1952. *J. Physiol., Lond.* 116 189.

63. DYNAMICS OF PULMONARY CIRCULATION: MAN, DOG

Pulmonary pressure zero point = 10 cm anterior to back in man; zero point = back in dog

Condition	No. of Observations	Pressure, mm Hg				Blood Flow L/min	Vascular Resistance dynes sec/cm ⁵	Reference
		Systolic	Diastolic	Mean	Capillary Mean			
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)
Man								
1 Resting	3 ¹	25	14	18		6.0	176 ²	1
2	8 ¹			14		7.1	158 ²	2
3	10 ¹	25	11	16		5.4	238 ²	3
4	8	23	9	15	9	7.4	67	4
5	9			16	7	5.5	131	5
6 Exercise, supine	8 ¹			15		10.4	115 ²	6
7	7			21	10	9.6	90	7
8 Exercise, standing	3 ¹	26	9	16		12.0	76 ²	8
9 Hypoxia	9			20	7	5.5	189	9
10	8 ¹	26	12	18	8	8.0	119	10
11 Saline infusion	4 ¹			23	16	6.7	84	11
Drug effect (IV injection)								
12 Epinephrine	5 ¹			19	7	7.9	121	12
13 Norepinephrine	9	28	11	21	15	4.5	118	13
14 Tetraethylammonium chloride	3			15	10	4.7	85	14
Pathologic								
15 Emphysema, mild	6 ¹	29	12	18		5.5	262 ²	15
16 Emphysema, severe	6 ¹	43	16	28		5.6	400 ²	16
17 Cor pulmonale, with hypoxia	11 ¹	43	19	28		5.4	414 ²	17
18	17			35	9	5.4	440	18
19 Cor pulmonale, with congestive failure	4 ¹	74	39	52		7.4	562 ²	19
20 Primary pulmonary hypertension	3 ¹	101	47	66		2.9	1776 ²	20
21 Left ventricular failure	10			45	30	3.5	344	21
22 Mitral stenosis	21			51	29	4.3	491	22
23	22	68	33	47	28	4.0	448	23
24 Mitral insufficiency	1			15	12	5.9	41	24
25 Mitral stenosis and insufficiency	12			36	23	3.8	304	25
26 Aortic stenosis and insufficiency	6			29	16	5.0	208	26
27 Chronic constrictive pericarditis	4	36	22	26	20	4.3	112	27
Atrial septal defect								
28 Left to right shunts	3 ¹	38	12	23	13 ¹	16.5	49	28
29	7 ¹	31	14	21	8	18.0	58	29
30 Right to left shunts	1 ¹	96	53	68	9 ³	2.6	1770	30
31	2 ¹	88	59	73		3.5	3710 ²	31
32 Patent ductus arteriosus	6	43	19	29		12.2	190 ²	32
33	2	34	17	28	14	19.0	53	33
34	3 ¹	46	23	31		7.7	300 ²	34
35	13 ¹			36		12.4	206 ²	35
36 Tetralogy of Fallot	1		8	4	6	2.5	192 ²	36
37 Pulmonary stenosis	8	24	10	17	11	5.4	89	37
38 Eisenmenger syndrome	3	120	65	87	6	3.3	1915	38
39	4			86		3.5	2225 ²	39
40 Ventricular septal defect	15 ¹			37		10.3	248 ²	40
41	2	35	20	27		5.7	386 ²	41
Dog								
42 Resting (anesthetized)	16			18	10	2.3	285	42
43 Resting	16			17	4	3.2	325	43
44 Hypoxia, moderate (anesthetized)	13			22	8 ¹	2.4	467	44
45 Hypoxia, moderate	16			26	4 ¹	3.8	463	45
46 Hypoxia, severe	3			26	9 ³	8.1	167	46

/1/ Pressure zero point corrected to approximately 10 cm. /2/ Total pulmonary resistance, utilizing pulmonary arterial mean minus pulmonary "capillary" mean as pressure gradient /3/ Pulmonary venous or left atrial mean pressure

Contributor: Gorlin, Richard

- References: [1] Riley, R. L., A. Hummelstein, H. L. Motley, H. M. Weiner, and A. Cournand. 1948. *Am. J. Physiol.* 152:372. [2] Hickam, J. B., and W. H. Cargill. 1948. *J. Clin. Invest.* 27:16. [3] Cournand, A. 1950. *Circulation*, N. Y. 2:641. [4] Dexter, L., J. W. Dow, F. W. Haynes, J. L. Whittenberger, B. G. Ferris, W. T. Goodale, and H. K. Hellems. 1950. *J. Clin. Invest.* 29:602. [5] Westcott, R. N., N. O. Fowler, R. C. Scott, V. D. Hauenstein, and J. McGuire. 1951. *Ibid.* 30:957. [6] Dexter, L., J. L. Whittenberger, F. W. Haynes, W. T. Goodale, R. Gorlin, and C. G. Sawyer. 1951. *J. Appl. Physiol.* 3:439. [7] Doyle, J. T., J. S. Wilson, and J. V. Warren. 1952. *Circulation*, N. Y. 5:263. [8] Doyle, J. T., J. S. Wilson, E. H. Estes, and J. V. Warren. 1951. *J. Clin. Invest.* 30:345. [9] Witham, A. C., and J. W. Fleming. 1951. *Ibid.* 30:707. [10] Fowler, N. O., R. N. Westcott, R. C. Scott, and J. McGuire. 1951. *Ibid.* 30:517. [11] Fowler, N. O., R. N. Westcott, V. D. Hauenstein, R. C. Scott, and J. McGuire. 1950. *Ibid.* 29:1387. [12] Harvey, R. M., M. I. Ferrer, D. W. Richards, Jr., and A. Cournand. 1951. *Am. J. Med.* 10:719. [13] Dexter, L., J. L. Whittenberger, R. Gorlin, D. M. Lewis, F. W. Haynes, and R. J. Spiege. 1951. *Tr. Ass. Am. Physicians* 64:22b. [14] Dresdale, D. T., M. Schultz, and R. J. Michtom. 1951. *Am. J. Med.* 11:686. [15] Gorlin, R., F. W. Haynes, W. T. Goodale, C. G. Sawyer, J. W. Dow, and L. Dexter. 1950. *Am. Heart J.* 41:30. [16] Lukas, D. S., and C. T. Dotter. 1952. *Am. J. Med.* 12:639. [17] Gorlin, R., B. M. Lewis, F. W. Haynes, and L. Dexter. 1952. *Am. Heart J.* 43:357. [18] Gorlin, R., M. B. Matthew, I. K. MacMillen, R. Daley, and W. E. Medd. 1953. *Annual Meet. Brit. Cardiac Soc.*, May 21. [19] Sawyer, C. G., C. S. Burwell, L. Dexter, E. C. Eppinger, W. T. Goodale, R. Gorlin, D. E. Harken, and F. W. Haynes. 1952. *Am. Heart J.* 44:207. [20] Hickam, J. B. 1949. *Ibid.* 38:801. [21] Handelsman, J. C., R. J. Bing, J. A. Campbell, and H. E. Greswold. 1948. *Bull. Johns Hopkins Hosp.* 82:615. [22] Taylor, H. E., A. A. Pollack, H. B. Burchell, O. T. Clagett, and E. H. Wood. 1950. *J. Clin. Invest.* 29:745. [23] Cournand, A., J. S. Baldwin, and A. Hummelstein. 1949. *Cardiac catheterization in congenital heart disease*. Commonwealth Fund, New York. [24] Wood, P. 1950. *Brit. M. J.* 2:639. [25] Bing, R. J., L. D. Vandam, and F. D. Gray, Jr. 1947. *Bull. Johns Hopkins Hosp.* 80:323. [26] Lewis, B. M., and R. Gorlin. 1952. *Am. J. Physiol.* 170:574. [27] Stroud, B. C., and H. Rahn. 1953. *Ibid.* 172:211.

64. LIMB BLOOD FLOW: MAN

Part I: EFFECT OF TEMPERATURE

Abbreviations: A = air plethysmography; W = water plethysmography; S = Stewart dye-dilution (Evans' blue dye injected at constant rate into brachial artery; dye concentration determined in continuously sampled venous blood). Values in parentheses are ranges, (cf. Introduction).

Extremity	Condition of Blood Vessels	Temperature °C		Blood Flow		Reference
		Ambient	Bath	Method	ml/min/100 ml segment	
(A)	(B)	(C)	(D)	(E)	(F)	(G)
1 Forearm	Normal		32	W	1.8(1.5-2.5) ^c	1
2		25	32	W	1.8(0.4-3.2) ^b	2
3		15-20	32	W	2.3(1.2-3.0) ^c	3, 4
4		22-25	32	W	2.6(1.6-4.8) ^c	5
5		15-20	33	W	2.7(1.9-3.8) ^c	3, 4
6		15-20	35	W	4.2(1.5-7.0) ^a	3, 4
7		18.5		A	3.1(2.6-3.6) ^b	4
8		25-27		S	4.5(2.8-6.3) ^c	6
9		27-29		A	4.9(1.7-7.3) ^c	7
10		22-25	28-30	W	1.6(1.0-2.5) ^c	8
11	Maximal constriction		10-14	W	0.7(0.5-1.0) ^c	9
12	Maximal dilatation	22-25	17	W	0.7(0.5-1.0) ^c	5
13			45	W	17.6(11.0-25.4) ^c	3
14		22-25	45	W	7.6(3.7-12.7) ^c	5
15 Hand	Normal		32	W	3.5(1.2-5.3) ^c	10
16		21	32	W	9.3(5.1-13.5) ^b	2
17		24	35	W	5.9(2.7-9.6) ^c	11
18		21-24		A	2.1(0.5-6.4) ^c	12
19		25-29		A	9.3(5.8-16.5) ^c	12
20	Maximal constriction		10-14	W	2.5(0.3-4.7) ^c	9
21	Maximal dilatation		45	W	22.0	9
22 Finger	Normal	"Comfortable"		A	(15-40) ^d	13
23	Maximal constriction				0.2	14
24	Maximal dilatation		45	W	90.0	13
25 Leg	Normal		32	W	1.4(0.8-2.6) ^c	9
26	Maximal constriction		10-14	W	1.2(0.4-1.5) ^c	9
27	Maximal dilatation		45	W	3.6	15
28 Calf	Normal	25	32	W	1.4(0.4-2.4) ^b	2
29 Foot	Normal	21	32	W	2.0(0.5-4.4) ^c	10
30		24-26.5	32	W	3.9(1.4-7.8) ^c	16
31			32	W	2.1(1.2-2.8) ^c	15, 17
32	Maximal dilatation		45	W	18.1(11.0-34.0) ^c	16

Contributors: (a) Zierler, Kenneth L., and Reubin Andres, (b) Abramson, David I.

References: [1] Abramson, D. I., H. Zazeela, and J. Marrus. 1939. *Am. Heart J.* 17:194. [2] Abramson, D. I., and S. M. Fierst. 1942. *Ibid.* 23:84. [3] Barcroft, H., and O. G. Edholm. 1943. *J. Physiol., Lond.* 102:5. [4] Barcroft, H., and O. G. Edholm. 1946. *Ibid.* 104:366. [5] Abramson, D. I., A. Kahn, S. Tuck, Jr., G. A. Turman, H. Rejal, and C. J. Fleischer. 1958. *J. Clin. Invest.* 37:1031. [6] Andres, R., H. M. Anderson, W. Stainsby, G. Cader, A. S. Ghayyib, K. L. Zierler, and J. L. Lillenthal, Jr. 1954. *Ibid.* 33:482. [7] Slaughter, O. L., H. S. Brown, and K. G. Wakim. 1948. *J. Laborat. Clin. M.* 33:743. [8] Abramson, D. I., S. Tuck, Jr., Y. Bell, C. Burnett, and H. Rejal. 1959. *J. Clin. Invest.* In press. [9] Abramson, D. I., K. H. Katzenstein, and E. B. Ferris, Jr. 1941. *Am. Heart J.* 22:329. [10] Hoobler, S. W., S. D. Malton, H. T. Ballantine, Jr., S. Cohen, R. B. Neligh, M. M. Peet, and R. H. Lyons. 1949. *J. Clin. Invest.* 28:638. [11] Speelman, C. R. 1945. *Am. J. Physiol.* 145:218. [12] Forster, R. E., II, B. G. Ferris, Jr., and R. Day. 1946. *Ibid.* 146:600. [13] Burton, A. C. 1939. *Ibid.* 127:437. [14] Wilkins, R. W., J. Doupe, and H. W. Newman. 1938. *Clin. Sc., Lond.* 3:403. [15] Abramson, D. I. Unpublished. [16] Scheinberg, P., E. W. Dennis, R. L. Robertson, and E. A. Stead, Jr. 1948. *Am. Heart J.* 35:409. [17] Abramson, D. I. 1942. *J. Mount Sinai Hosp. N. York* 8:328.

64. LIMB BLOOD FLOW: MAN (Concluded)

Part II: EFFECT OF VARIOUS CONDITIONS

Ambient temperature = 21-23°C. (+) = increase in blood flow, (-) = decrease in blood flow.

Condition (A)	Blood Flow ml/min/100 ml segment					Reference (G)
	Forearm (B)	Hand (C)	Digit (D)	Foot (E)	Calf (F)	
1 Anemia	2.5	6.1	12	2.0	2.8	B, C, 1; D-F, 2
2 Aortic insufficiency	2.0	6.7			1.7	3
3 Arteriosclerosis			5	2.1	1.6	2
4 Hypertension	2.9	5.4			2.4	4
5 Hyperthyroidism	4.4	11.6	30	4.5	3.6 ¹	B, C, 5, D-F, 2
6 Mitral stenosis	1.7	6.5				3
7 Raynaud's phenomenon		2.5	1	2.0	1.8	2
8 Schizophrenia	2.3	7.2			1.6	6
9 Sympathectomy, 1-2 da		35.0	100	22.0	7.5	7
10 1-2 wk		11.0	40	8.0	4.4	
11 >3 mo		8.0	28	4.5	3.3	
12 Varicostitis					2.4	8
13 Anesthesia, general		20.0	60	15.0	7.8	9
14 Low spinal		2.0	62, 70 ¹	12.0	6.0	10, 11
15 High spinal		+	60 ² , 100 ³	+	+	10, 11
16 Epinephrine		1.8	-	-	4.0	12
17 Tolazoline HCl		28.0	80	17.0	1.6	13
18 Reflex heating		30.0	90	18.0	4.5	2
19 Exercise, before	1.5				2.3	B, 14, F, 15
20 Maximum during	4				30	
21 Immediately after	22				77	

1/1 & 1 also reported [16]. 12/ Finger. 13/ Toe.

Contributors: (a) Lynn, R. Beverly, (b) Abramson, David L., (c) Zierler, Kenneth L.

References. [1] Abramson, D. L., S. M. Fierst, and K. Flachs. 1943. *Am. Heart J.* 25:609. [2] Abramson, D. L. 1944. Vascular responses in the extremities of man in health and disease. University of Chicago Press, Chicago. [3] Abramson, D. L., S. M. Fierst, and K. Flachs. 1942. *J. Clin. Invest.* 21:747. [4] Abramson, D. L., and S. M. Fierst. 1942. *Am. Heart J.* 23:84. [5] Abramson, D. L., and S. M. Fierst. 1942. *Arch. Int. M.* 69:409. [6] Abramson, D. L., N. Schkloven, and K. H. Katzenstein. 1941. *Arch. Neur. Psychiat.* Chic. 45:973. [7] Smith, H. W. 1951. *The kidney: structure and function in health and disease.* Oxford University Press, New York. [8] Abramson, D. L., and S. M. Fierst. 1942. *Arch. Surg.* 45:964. [9] Lynn, R. B., and R. Shackman. 1951. *Brit. M. J.* 2:333. [10] Lynn, R. B., et al. 1952. *Surgery* 32:195. [11] Sances, S. M., et al. 1952. *Circulation*, N. Y. 6:559. [12] Barcroft, H., and H. J. C. Swan. 1953. Sympathetic control of human blood vessels. *Edward Arnold, London.* [13] Lynn, R. B. 1950. *Lancet*, Lond. 2:676. [14] Grant, R. T. 1938. *Clin. Sc.*, Lond. 3:157. [15] Barcroft, H., and A. C. Dornhurst. 1949. *J. Physiol.*, Lond. 109:402. [16] West, R. O. Unpublished.

Condition: N_2 = norepinephrine infused at rate of 0.018(0.001-0.035) mg/min at room temperature of 26-29°C; V = after vasodilatation by indirect heating for 1 h; V + A = V + A administered via an intravenous drip (5% glucose in water); H = after indirect heating for 1 hour or more, supplemented by autonomic ganglion blockade; S = after spinal anesthesia; T = 3-5 years after thoracolumbar sympathectomy; B = after intravenous benzodioxane. Values in parentheses are ranges, estimate "b" (cf. Introduction).

Condition	Procedure	Blood Flow ¹ ml/min/ sq cm skin	Arterial Blood Pressure Systolic ² mm Hg	Diastolic ³ mm Hg	Reference
(A)	(B)	(C)	(D)	(E)	(F)
Finger					
1 Normal	RT	0.20(0.08-0.32)	104(70-138)	65(39-91)	1,2
2	V	0.28(0.22-0.34)	101(81-121)	58(35-81)	3-6
3	V + A	0.29(0.21-0.37)	87(67-107)	52(32-72)	1
4	V + A ⁴	0.26(0.16-0.36)	91(62-120)	58(40-76)	7,8
5 Normal (N_2)	V + A ¹	0.24(0.12-0.36)	134(100-168)	83(57-109)	7,8
6 Pregnancy, 3rd trimester	RT	0.29(0.15-0.43)	106(83-129)	68(48-88)	9
7	V + A	0.33(0.19-0.47)	91(64-118)	58(32-84)	
8 Primary hypertension	RT	0.18(0.02-0.34)	168(113-223)	103(55-151)	1
9	V + A	0.31(0.21-0.41)	120(74-166)	75(38-112)	
10	V + A ¹	0.21(0.01-0.41)	122(74-170)	77(41-113)	7
11 Primary hypertension (N_2)	V + A ¹	0.16(0.00-0.38)	184(133-235)	105(65-145)	7
12 Hypertension of pregnancy, 3rd trimester	RT	0.29(0.17-0.41)	121(90-152)	73(58-88)	9
13	V + A	0.35(0.23-0.47)	96(56-136)	62(33-91)	
14 Cushing's syndrome ⁴	V + A ¹	0.31(0.17-0.45)	100(67-133)	62(28-92)	8
15 Cushing's syndrome ⁴ (N_2)	V + A ¹	0.27(0.10-0.44)	151(96-206)	94(60-128)	8
16 Clubbing	V	0.35(0.24-0.46)	109(72-135)	72(59-85)	3,5
17 Coarctation of aorta	V + A	0.33	111	64	3
18 Scleroderma	V + A	0.14(0.00-0.36)			
19 Raynaud's disease	V + A	0.23(0.05-0.41)			
20 Acrocyanosis	V + A	0.33			
21 Lymphedema	V + A	0.35(0.26-0.44)			
22 Coronary occlusion	V	0.22(0.09-0.35)			
Great Toe					
23 Normal	H	0.21(0.15-0.29)	87(57-117)		3
24	S	0.13(0.05-0.21)			
25 Primary hypertension	H	0.21(0.08-0.34)			
26	T	0.11(0.00-0.23)			
27	H, T	0.16(0.05-0.26)			
28	H, T, B	0.16(0.04-0.27)			
29 Diabetes mellitus ⁵	H	0.17(0.05-0.28)			
30 Thromboangiitis obliterans	H	0.10(0.01-0.23)			
31 Coarctation of aorta	H	0.24	75		
32 Cold injury	H	0.14(0.05-0.24)	84		
33 Scleroderma	H	0.12			
34 Raynaud's disease	H	0.12(0.01-0.23)			
35 Arteriosclerosis	H	0.08			
36 Thrombophlebitis	H	0.17	91		

/1/ Determined calorimetrically on the terminal phalangeal skin of 3rd or 4th finger, or on the terminal phalangeal skin of the great toe. /2/ Determined with a Gaertner capsule by the "flushing" technique on the great toe or middle phalanx of the 3rd or 4th finger. /3/ Determined from the point of cessation of throbbing on the middle phalanx of the 3rd or 4th finger. /4/ Caused by endogenous secretion of excessive adrenal corticosteroids, or by their exogenous administration. /5/ Of less than ten years' duration and without overt arterial disease.

Contributors (a) Mendlowitz, Milton, (b) Lamport, Harold, (c) Levy, Matthew N.

References [1] Mendlowitz, M., S. M. Torosdag, and L. Sharney 1957. J. Appl. Physiol. 10:436. [2] Weaver, J. C., and D. F. Bohr. 1950. Am. Heart J. 39:413. [3] Mendlowitz, M. 1954. The digital circulation. Grune and Stratton, New York. [4] Arnott, W. M., and J. M. MacFie. 1948. J. Physiol., Lond. 107:233. [5] Wilson, G. M. 1952. Q. J. Med., Oxf. 21:201. [6] Capriglione, L. A., and F. L. V. Duque. 1948. Arg. clin., Rio 7:3. [7] Mendlowitz, M., and N. Naftchi. 1958. J. Appl. Physiol. 13:247. [8] Mendlowitz, M., S. Gilow, and N. Naftchi. 1958. Ibid 13:252. [9] Mendlowitz, M., A. Altchek, and N. Naftchi. 1958. Am. J. Obst. 76:473.

66. RESISTANCE OF ANIMAL TISSUE TO ARRESTED CIRCULATION: MAMMALS

Circulation arrested by obstructing or bypassing total afferent blood supply to organ. Adult tissue, in situ, at normal body temperature, unless otherwise indicated.

Tissue	Animal	Initial Disappearance of Function ¹		Development of Irreversible Changes ²		Reference
		min	Remarks	min	Remarks	
(A)	(B)	(C)	(D)	(E)	(F)	(G)
1 Brain, cerebral cortex	Cat	1/6-1/4		<3		C.1,2,E.3
2				>5-10		4
3	Dog			1-8	Newborn animal	5,6
4				12-20		
5 Brain stem	Dog	3-4	Isolated, perfused head			7
6 Pupillary centers	Dog	4-5	Isolated, perfused head	15-30	Isolated, perfused head	7
7 Cardiorespiratory, vaso-motor, and adreno-secretory centers	Dog			5-10		8
8 Respiratory center	Dog, rat	4-5	Isolated, perfused head	15-30		7
9		1/3-1/2				6,9
10	Rat	20-40	Newborn animal, decapitated			9
11 Spinal cord	Cat	2/3-1		35-45		C.10,E.11
12				90-120		12
13	Rabbit	2/3-2				13
14 Heart	Dog, rabbit, rat	5-20				14
15 Lung	Dog			30-45	Criterion: death of animal	15
16 Kidney	Dog			30-60		16,17
17				>120		18,19
18	Rabbit			60-90		20
19	Rat			<120		19,21
20 Liver	Dog			20-75		22-25
21				>60		24
22 Skeletal muscle	Dog, rabbit, rat	120-360		480		26,27
23 Small intestine	Dog			120-240+	Criteria: electrical reaction and death of animal	28
24 Testis	Rat			10-30	Histological study of spermatogenesis	29

/1/ Interval from the start of circulatory arrest to the disappearance of a specific functional activity; i.e., the time of survival of a function, without implication of reversibility. /2/ Interval from the start of circulatory arrest to the occurrence of irreversible changes in the organ so that the function does not reappear, i.e., the time of survival of an organ before irreversible changes occur /3/ Value for liver at 24-27°C.

Contributors (a) Sonnenschein, Ralph R., Robert Lewis, and Loulae Darling, (b) Van Harreveld, A., (c) Wesolowski, Sigmund A

- References [1] Sugar, O., and R. W. Gerard. 1938. J Neurophysiol. 1:558 [2] Van Harreveld, A. 1947. Ibid. 5:361. [3] Gänshirt, H., and W. Zylka. 1952. Arch Psychiat., Berl. 189:23. [4] Ten Cate, J., and G. P. Horsten. 1954. Arch. internat. physiol., Liège 62:6. [5] Kabat, H., and C. Dennis. 1939. Proc. Soc. Exp. Biol., N. Y. 42:534 [6] Kabat, H. 1940. Am. J. Physiol. 130:588. [7] Heymans, C., et al. 1937. Arch. Neur. Psychiat., Chic. 38:304 [8] Malmeljac, J., and P. Plane. 1954. Afrique fr. chir. 12:435. [9] Selle, W. A., and T. A. Witten. 1941. Proc. Soc. Exp. Biol., N. Y. 47:495 [10] Van Harreveld, A. 1944. Am. J. Physiol. 141:97. [11] Van Harreveld, A., and G. Marmont. 1939. J Neurophysiol. 2:101 [12] Van Harreveld, A., and D. B. Tyler. 1944. Am. J. Physiol. 142:32. [13] Blasius, W. 1951. Zachr. Biol. 104:121. [14] Wesolowski, S. A., et al. 1952. In O. Wangersten, ed. Forum on fundamental surgical problems. W. B. Saunders, Philadelphia. p. 270. [15] Blader, B. 1954. Arch. Surg. 69:525. [16] Guthrie, C. C. 1910. Arch. Int. M. 5:232. [17] McEnery, E. J., et al. 1926. J. Laborat. Clin. M. 12:349 [18] Friedman, S. M., et al. 1954. Circul. Res., N. Y. 2:231. [19] Scheibe, J. R., et al. 1949. Surgery 52:724 [20] Badenoch, A. W., and E. M. Darmedy. 1947. J. Path. Bact., Lond. 59:79. [21] Latorre, G. 1954. J. Urol., Balt. 72:639. [22] Raffucci, F. L. 1953. Surgery 33:342 [23] Drapanas, T., et al. 1955. Ann. Surg. 142:831 [24] Bernhard, W. F., et al. 1955. N. England J. M. 253:159. [25] Hines, J. R., and M. Roncoroni. 1956. Surg. Gyn. Obst. 102:689. [26] Harmon, J. W. 1947. Am. J. Path. 23:551. [27] Harmon, J. W., and R. P. Gwinn. 1949. Ibid. 25:741 [28] Engquist, I. F., and A. J. Kremen. 1952. In O. Wangersten, ed. Forum on fundamental surgical problems. W. B. Saunders, Philadelphia. p. 87. [29] Oettle, A. G., and R. G. Harrison. 1952. J. Path. Bact., Lond. 64:273

67. LYMPH FLOW AND LYMPHOCYTE CONTENT: MAMMALS

For a comprehensive review of the subject, see reference 22. Lymph collections made during period of anesthesia, unless otherwise specified. Condition: F = fasting, NF = non-fasting. Anesthetic: B = barbitalurate, C = chloroform, E = ether, I = inhalation anesthetic, L = chloralose, M = morphine, U = urethane, X = unspecified. Values in parentheses are ranges, estimate "c" (cf. Introduction).

Animal	Subjects		Weight kg	Anesthetic	Lymph Flow ml/hr	Lymphocytes		Refer- ence
	No.	Con- dition				(cells/cu mm) $\times 10^3$	(cells/hr/kg) $\times 10^6$	
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)
Thoracic Duct								
1 Man	5	NF	60.6(48.5-78.5)	X ¹	48(37-83)	11.4(2.1-25.5)	8.7	1
2	1	Fistula	50	X ¹	(53-78)	1.8	3.1	2
3 Calf	4	NF	34.4(29-38)	B, I	115(51-172)	26.1(21.5-30.6)	85(43-117)	3
4 Cat	21	NF ²	1.9	B, U	5.0(1.8-7.3)	14.2(4.3-29.6)	35.5(9.6-92.6)	4
5 Dog	78	NF	10.4(6.3-15.7)	C, E, M	23.6(7.0-68.1)			5
6	21	F ³	10	L			21.2(3.0-87.5)	6
7	10	F ⁴	10.9(5.4-20.8)	M	8.6(4.0-11.6)	7.2(3-20)	5.7	7
8 Goat	4	NF	32(25-35)	B, I	102(30-154)	6.9(2.7-9.0)	23.0(6.3-55.7)	3
9 Guinea pig	40	NF	0.303	B	0.95	14.8	45.7	8
10 Monkey	26	NF	3.5	B	3.0(0.95-8.1)	29.9(2.9-100.3)	26.7(2.8-71.2)	9
11 Mouse	7	NF	0.023	B	0.08		69	10
12 Rabbit	9	NF	2.5(2.1-3.2)	B, E	4.6(1.5-9.0)	36.1	69(19-126)	11
13	7	NF	2.5	B	4.0	32.3	54.5	12
14 Rat	28	NF	0.25(0.20-0.29)	B	0.88(0.5-1.2)	31.5	107(60-181)	13
Intestinal Lacteal Lymphatics								
15 Mouse	5	NF	0.03	B	0.10	47.1	133.7	10
16 Rat	8	NF ⁵	0.18-0.24	X ¹		34.6(18.3-60.5)		14
Abdominal Thoracic Duct								
17 Mouse	20	NF	0.02	B	0.1	11.7	57.0	15
18 Rat	7	NF	0.16-0.18	B ¹			246	16
19	5	NF	0.36-0.40	B ¹			158	16
20	9	NF	0.155(0.135-0.170)	E ¹	0.66(0.39-1.04)	32.5(13.2-61.1)	133(45-212)	17
21	9	NF	0.274(0.21-0.31)	E ¹	1.06(0.81-1.28)	32.1(15.4-55.9)	121(74-198)	17
22	27	NF ⁶	0.172(0.145-0.200)	X ¹	1.6(0.7-2.7)	20.0	185	18
23	8	NF ⁶	0.199(0.16-0.29)	B, E ¹	5.8(2.1-8.2)	3.3(2.2-6.2)	95.6	19
24	6	NF ⁷	0.235(0.20-0.26)	B, E ¹	0.61(0.53-0.66)	11.8(7.2-13.1)	34.5	19
Cervical Lymphatics								
25 Guinea pig	5	NF	0.235	B	0.046	6.5	1.36	8
26 Rabbit	1	NF	2.05	B	0.16	37.2	3.2	9
27 Rat	10	NF	0.25	B	0.038	7.1(1.5-17.5)	1.24	20
Right Duct								
28 Monkey	1	NF	3.8	B	0.17	10.2	0.45	9
29 Monkey ⁸	5	NF	2.49	B	2.07(1.1-3.6)	40.4(32.0-56.4)	42.3(10.8-111.0)	9
30 Rabbit	3	NF		B	0.35(0.24-0.46)	40.9(39.5-43.6)		12
31 Rat	10	NF	0.303(0.280-0.334)	B	0.10(0.07-0.14)	17.3(13.0-23.7)	5.7	21
Subclavian Duct								
32 Monkey	4	NF	2.6	B	0.06(0.04-0.06)	4.2(2.5-7.0)	0.08(0.06-0.10)	9
33 Rabbit	7	NF ⁹	2.3	B ¹	0.69	38.9(13.8-50.8)	11.2	12
34	5	NF ¹⁰	2.3	B ¹	1.33	22.5(19.3-26.4)	14.2	12

/1/ Lymph collected after recovery from anesthetized state. /2/ Fed cream 4 hours before. /3/ 36-48 hours.
/4/ 24-36 hours. /5/ Oral 0.2% saline. /6/ Oral saline. /7/ Oral water. /8/ Right duct functional thoracic duct.
/9/ Prone. /10/ Sitting.


Contributor: Reinhardt, William O.

References [1] Bierman, H. R., R. L. Byron, K. H. Kelly, R. S. Gillfillan, L. P. White, N. E. Freeman, and N. L. Petrakis. 1953. J. Clin. Invest. 32:637. [2] Courtice, F. C., W. J. Simmonds, and A. W. Steinbeck. 1951. Austral. J. Exp. Biol. 29:201. [3] Wingvist, G. 1954. Acta anat., Suppl., Basel 22:21. [4] Adams, W. S., R. H. Saunders, and J. S. Lawrence. 1945. Am. J. Physiol. 144:297. [5] Heidenhain, R. 1891. Arch. ges. Physiol. 49:209. [6] Yoffey, J. M. 1935-36. J. Anat., Lond. 70:507. [7] Chistoni, A. 1909. Arch. fistol., Fir 6:74. [8] Reinhardt, W. O., and J. M. Yoffey. 1956. Am. J. Physiol. 187:493. [9] Reinhardt, W. O. Unpublished. [10] Shrewsbury, M. M. Unpublished. [11] Sanders, A. G., H. W. Florey, and J. M. Barnes. 1940. Brit. J. Exp. Path. 21:254. [12] Hughes, R., A. J. May, and J. G. Widdicombe. 1956. J. Physiol., Lond. 132:384. [13] Hungerford, G. F., W. O. Reinhardt, and C. H. Li. 1952. Blood, N. Y. 7:193. [14] Mann, J. D., and G. M. Higgins. 1950. Ibid. 5:177. [15] Shrewsbury, M. M. 1958. Proc. Soc. Exp. Biol., N. Y. 99:53. [16] Whaler, B. C., and J. G. Widdicombe. 1946. J. Physiol., Lond. 132:41. [17] Hungerford, G. F., and W. O. Reinhardt. 1950. Am. J. Physiol. 160:9. [18] Gowans, J. L. 1957. Brit. J. Exp. Path. 38:67. [19] Shrewsbury, M. M., and W. O. Reinhardt. 1952. Am. J. Physiol. 168:366. [20] Reinhardt, W. O., and C. H. Li. 1945. Proc. Soc. Exp. Biol., N. Y. 58:321. [21] Schooley, J. 1958. Ibid. 99:511. [22] Yoffey, J. M., and F. C. Courtice. 1956. Lymphatics, lymph and lymphoid tissue. Edward Arnold, Lond.


68 ELECTROCARDIOGRAPHIC LEADS IN STANDARD USE: MAN

Part I: STANDARD LIMB LEADS



— and 5 mm squares. Standardisation of the electro-
 graph shows a vertical deflection of
 cental axis represents an
 in potential between
 des on the right arm and
 iographic galvanometer.




Standardization



Lead I



Lead II




Lead III


Reference Rushmer, R. F. 1955. *Cardiac diagnosis*. W. B. Saunders, Philadelphia. p 217.

68. ELECTROCARDIOGRAPHIC LEADS IN STANDARD USE. MAN (Concluded)
Part II UNIPOLAR LIMB LEADS


When the heart is oriented vertically, the QRS complexes are similar in leads aVR and aVL . In the horizontal heart, aVR and aVL are essentially mirror images, a phenomenon which suggests that the wave of excitation actually progresses from right to left. This is confirmed by the diphasic QRS in lead aVF .



Vertical heart




G = galvanometer



Horizontal heart


Part III UNIPOLAR PRECORDIAL LEADS

Unipolar precordial electrodes are routinely placed on six positions. V_1 is in the fourth I.C.S. (intercostal space) at the right sternal edge. V_2 is in the fourth I.C.S. just to left of the sternum. V_4 is in the fifth I.C.S. in the mid-clavicular line, with V_3 midway between V_2 and V_4 . On a horizontal line drawn from V_4 , V_5 is located at the anterior axillary line, and V_6 at the mid-axillary line. Electrocardiographic patterns recorded from the six precordial electrode positions are schematically presented.



Definitions: P wave = first upward deflection in the electrocardiogram; period of atrial excitation. PR interval = period measured from the beginning of the P wave to the beginning of the QRS complex; the time required for excitation to go from the sino-atrial node through the atrial musculature (through the junctional tissues and Purkinje system) to the ventricular muscle. QRS interval = period measured from the beginning to the end of the QRS complex; the time required for excitation waves to spread through ventricular walls. QT interval = duration of ventricular systole; period of uniform ventricular excitation and return of ventricles to resting or excitable state. QT_c = QT interval divided by the square root of the cycle length.

Part I: GRAPHIC



Reference: Rushmer, R. F. 1955 Cardiac diagnosis. W. B. Saunders, Philadelphia. p. 239.

69. DURATION OF ELECTROCARDIOGRAPHIC WAVES AND INTERVALS. VERTEBRATES (Concluded)

Part II. TABULAR

Values are given in milliseconds. Those in parentheses are ranges, estimate "c" (lines 1-13, 19, 24, 33, 38) and estimate "b" (lines 14, 16-18, 20-23, 25-27, 31, 39) (cf. Introduction).

Animal (A)	P Wave (B)	PR Interval (C)	QRS Interval (D)	QTc (E)	Reference (F)
Man				421(366-533)	1
1 Birth-1 da	51(40-64)	99(80-120)	65(40-100)	402(322-476)	1
2 1 da-1 wk	48(36-64)	95(80-120)	56(40-80)	385(345-434)	1
3 1 wk-1 mo	48(40-60)	95(80-120)	55(40-70)	397(304-500)	1
4 1-3 mo	54(48-64)	96(80-160)	62(50-80)	392(348-458)	1
5 3-6 mo	54(48-64)	101(80-120)	68(50-80)	408(350-455)	1
6 6 mo-1 yr	59(48-70)	106(80-140)	65(40-80)	394(338-435)	1
7 1-3 yr	65(48-88)	113(80-160)	71(40-90)	402(349-465)	1
8 3-5 yr	72(40-96)	125(100-160)	72(40-90)	414(350-514)	1
9 5-8 yr	72(50-96)	134(100-180)	75(50-96)	419(350-536)	1
10 8-12 yr	77(50-96)	139(100-200)	77(40-100)	408(302-455)	1
11 12-16 yr	82.5(60-104)	149(100-200)	77(40-100)	397(337-433)	2-5
12 Adult ♂	90(70-120)	160(110-210)	83(50-100)	415(380-456)	3-6
13 Adult ♀	90(70-120)	160(110-210)	72(50-100)	430(400-460)	7
14 Ass	80	200	60	350(400-460)	7
15 Cat	40	80	30	180	7
16 Cattle, ♂	80	160(160-200)	50	310(300-320)	7
17 Cattle, ♀	170(100-120)	225(220-230)	80	350(260-440)	7
18 Dog	70(60-80)	120(110-130)	75(70-80)	220(200-240)	7
19 Elephant		350(280-410)	150(120-180)	600(390-790)	8,9
20 Goat	80	130(120-140)	50	290(260-320)	7
21 Guinea pig	40	65(60-70)	30	120	7
22 Hamster		43(40-46)	23(19-23)		10
23 Horse	195(170-220)	370(320-420)	100(80-120)	520(440-600)	7
24 Monkey		71(54-88)	31(20-44)	212(160-284)	11
25 Mouse, white		43(39-47)	22(20-24)		10
26 Mouse, wild		32(27-37)	22(20-24)		10
27 Pig		130(120-140)	60	250(220-280)	7
28 Rabbit	40	60	30	120	7
29 Rat, white		42	10	70	12
30 Sheep	80	80	40	240	7
31 Spermophile		48(38-58)	28(26-30)		10
32 Whale, beluga		320	190(120)		13
33 Chicken	41(35-47)	79(72-80)	23.3(20-27)		14
34 Duck		90	28		15
35 Pigeon		69	21		15
36 Sea gull		48	26		15
37 Frog		440	200	1550	16
38 Elasmobranch		250(200-440)	50(40-160)	750(400-1000)	17
39 Teleost		190(120-260)	50(40-60)		17

Contributors (a) Rosenbaum, Francis F., (b) Modell, Walter, (c) Lepeschkin, Eugene, (d) Sturkie, Paul D., (e) Kossman, Charles E., (f) Kibler, Hudson H., (g) Ross, Lucille J.

- References [1] Ziegler, R. F. 1951. Electrocardiographic studies in normal infants and children. C. C. Thomas, Springfield, IL [2] Graybiel, A., R. A. McFarland, D. C. Gates, and F. A. Webster. 1944. Am. Heart J. 27 524. [3] Lepeschkin, E. 1951. Modern electrocardiography. Williams and Wilkins, Baltimore. v. 1. [4] McGinn, S., and P. D. White. 1934. Am. Heart J. 9 642. [5] Shipley, R. A., and W. R. Holloran. 1936. Ibid. 11:325. [6] Rosenbaum, F. F. Unpublished. [7] Luisada, A., A. Weiss, and H. W. Hantman. 1944. Cardiologia 8 65. [8] White, P. D., J. L. Jenks, and F. G. Benedict. 1938. Am. Heart J. 16 744. [9] Forbes, A., S. Cobb, and McK. Cattell. 1921. Am. J. Physiol. 55 385. [10] Lombard, E. A. 1952. Ibid. 171:189. [11] De Waart, A., and C. J. Storm. 1934. Acta brevis neerl. 4 130. [12] Hundley, J. M., L. L. Ashburn, and W. H. Sebrell. 1945. Am. J. Physiol. 144 404. [13] King, R. L., J. L. Jenks, Jr., and P. D. White. 1953. Circulation, N. Y. 8:387. [14] Sturkie, P. D. 1949. Am. J. Vet. Res. 10:168. [15] Klisch, B. 1951. Exp. M. Surg. 9:103. [16] De Boer, S. 1933. Arch. internat. pharm. dyn., Par. 46 477. [17] Klisch, B. 1948. Exp. M. Surg. 6:31.

70. AMPLITUDES OF ELECTROCARDIOGRAPHIC WAVES VS AGE: MAN

Values are in millimeters. Values in parentheses are ranges, estimate "c" (cf. Introduction).

Age	Wave				
	P	Q	R	S	T
(A)	(B)	(C)	(D)	(E)	(F)
Lead I					
1 Birth-1 da	0.82(0-1.80)	0.3(0-0.5)	1.5(0-5.5)	6.4(0-12.5)	2.0(0-3.0)
2 1 da-1 wk	0.88(0-1.80)	0.8(0-1.0)	2.0(0-6.0)	5.0(0-10.0)	1.0(0.6-3.0)
3 1 wk-1 mo	0.90(0-1.80)	1.0(0-3.0)	3.5(1.0-8.5)	4.6(0-9.3)	2.1(1.0-4.0)
4 1-3 mo	0.37(0-2.00)	1.1(0-2.5)	6.0(2.0-14.0)	3.6(0-9.0)	2.2(0.2-4.3)
5 3-6 mo	1.00(0-2.00)	1.2(0-2.0)	7.8(2.0-13.0)	3.3(0-7.0)	2.6(0.2-5.2)
6 6 mo-1 yr	1.05(0.50-2.50)	1.6(0-4.5)	8.0(3.0-17.0)	3.7(0-10.0)	2.4(1.0-6.0)
7 1-3 yr	1.10(0.50-2.50)	1.5(0-3.0)	8.0(3.0-17.0)	3.1(0-8.0)	2.9(1.0-6.6)
8 3-5 yr	1.07(0.50-2.00)	1.2(0-2.0)	6.5(3.0-16.0)	2.1(0-6.0)	2.5(1.0-6.0)
9 5-8 yr	1.05(0.50-2.00)	1.1(0-2.5)	6.2(1.0-15.5)	2.0(0-7.0)	2.6(1.0-5.5)
10 8-12 yr	1.01(0-2.00)	1.3(0-3.0)	6.5(1.0-16.0)	1.5(0-8.0)	2.9(1.0-6.0)
11 12-16 yr	0.94(0-2.00)	1.0(0-3.0)	5.9(0-12.0)	1.3(0-10.0)	2.5(1.0-6.0)
12 16-20 yr	0.66(0.20-1.00)	0.3(0-1.3)	4.3(1.8-9.5)	1.0(0-3.2)	2.1(0.2-3.7)
13 20-30 yr	0.74(0.20-1.00)	0.2(0-1.4)	4.9(0.9-11.2)	1.0(0-4.9)	2.5(0.4-5.6)
14 30-40 yr	0.74(0.15-1.15)	0.3(0-1.6)	5.3(0.7-11.3)	1.0(0-3.6)	2.2(0.6-4.2)
15 40-50 yr	0.70(0.20-1.10)	0.3(0-1.7)	5.6(1.6-9.8)	0.8(0-4.0)	2.2(0.5-4.2)
16 >50 yr	0.72(0.45-0.94)	0.3(0-1.1)	5.5(2.2-11.7)	1.0(0-6.4)	1.8(0.9-3.3)
Lead II					
17 Birth-1 da	1.45(0-2.50)	1.5(0-5.0)	5.5(0-23.0)	2.0(0-7.0)	1.0(-1.0 to 3.0)
18 1 da-1 wk	1.80(0-2.60)	1.8(0-4.0)	5.0(2.0-12.0)	2.3(0-8.0)	1.6(0-4.0)
19 1 wk-1 mo	1.50(0.50-2.00)	1.6(0-4.0)	7.5(2.0-14.0)	1.4(0-8.5)	2.7(0-5.6)
20 1-3 mo	1.50(0.50-2.00)	1.8(0-5.0)	11.5(5.0-19.0)	1.9(0-6.0)	3.1(0.5-5.6)
21 3-6 mo	1.60(0.50-2.50)	2.3(0-5.5)	13.5(3.0-23.0)	1.7(0-6.0)	3.4(0.2-6.7)
22 6 mo-1 yr	1.70(0-2.50)	2.4(0-5.0)	13.0(3.5-25.0)	2.2(0-6.0)	3.2(0.3-5.8)
23 1-3 yr	1.60(0.50-3.00)	2.1(0-4.5)	12.5(1.5-23.0)	1.6(0-6.0)	3.0(0.2-5.7)
24 3-5 yr	1.66(0.50-3.00)	1.6(0-4.5)	12.5(3.0-21.0)	1.3(0-5.0)	3.1(0.1-5.9)
25 5-8 yr	1.72(0.50-3.00)	1.4(0-5.0)	12.5(3.5-21.5)	1.7(0-6.0)	3.3(0.6-5.9)
26 8-12 yr	1.54(1.50-3.00)	1.4(0-3.0)	13.5(4.0-23.5)	1.5(0-6.0)	3.6(-0.5 to 8.2)
27 12-16 yr	1.62(1.00-3.00)	1.2(0-2.5)	13.5(3.5-24.5)	1.4(0-7.0)	3.3(-0.2 to 6.7)
28 16-20 yr	0.96(0-2.10)	0.4(0-2.8)	9.5(2.9-15.8)	1.6(0-6.3)	2.7(0.2-5.7)
29 20-30 yr	1.01(0.20-2.19)	0.3(0-1.9)	8.8(0.5-17.2)	1.3(0-8.2)	2.9(0.5-6.1)
30 30-40 yr	0.98(0-1.90)	0.3(0-1.8)	7.1(1.8-16.8)	1.2(0-4.9)	2.6(0.6-5.5)
31 40-50 yr	0.99(0-1.90)	0.3(0-2.2)	6.9(2.1-15.3)	1.1(0-4.7)	1.9(0.3-4.2)
32 >50 yr	1.00(0.66-1.30)	0.2(0-1.2)	6.5(2.4-14.0)	0.8(0-3.1)	2.4(1.1-5.2)
Lead III					
33 Birth-1 da	0.62(-1.00 to 2.00)	2.5(0-9.0)	9.0(2.0-23.0)	0.2(0-3.0)	1.0(-1.2 to 3.0)
34 1 da-1 wk	0.82(0-2.00)	2.5(0-5.0)	7.5(3.0-15.0)	0.3(0-4.0)	0.5(-2.3 to 3.1)
35 1 wk-1 mo	0.80(-0.50 to 2.00)	2.4(0-5.0)	7.9(2.0-17.0)	0.2(0-2.0)	0.6(-2.2 to 3.5)
36 1-3 mo	0.65(-1.00 to 2.00)	3.3(0-7.0)	8.5(4.0-15.0)	0.6(0-5.0)	0.9(-1.7 to 3.7)
37 3-6 mo	0.81(0-2.00)	4.5(0-9.0)	10.2(3.0-22.0)	0.6(0-6.0)	1.2(-2.1 to 4.6)
38 6 mo-1 yr	0.85(0-2.00)	4.1(0-8.0)	9.8(2.0-22.0)	0.8(0-6.0)	0.7(-2.7 to 4.0)
39 1-3 yr	0.74(-1.50 to 2.00)	3.5(0-8.0)	7.5(1.0-19.0)	0.8(0-8.0)	0.2(-3.2 to 3.6)
40 3-5 yr	0.58(-1.00 to 2.00)	2.5(0-6.0)	8.5(1.0-18.0)	0.8(0-7.0)	0.3(-2.2 to 3.4)
41 5-8 yr	0.65(-1.00 to 2.00)	2.2(0-5.5)	8.5(1.0-20.0)	0.8(0-8.0)	1.6(-2.2 to 4.4)
42 8-12 yr	0.55(-1.50 to 2.00)	1.9(0-5.0)	9.2(1.0-24.0)	1.0(0-5.0)	0.7(-1.8 to 3.5)
43 12-16 yr	(-0.50 to 2.00)	1.6(0-5.0)	9.0(1.0-26.0)	1.0(0-9.0)	0.8(-1.6 to 3.5)
44 16-20 yr	0.43(-1.00 to 1.80)	0.6(0-4.6)	6.2(1.2-15.0)	1.2(0-4.9)	0.8(-1.9 to 3.9)
45 20-30 yr	0.48(-0.75 to 2.00)	0.5(0-3.1)	5.3(0-16.4)	1.1(0-7.3)	0.7(-2.0 to 3.8)
46 30-40 yr	0.42(-0.73 to 1.30)	0.4(0-2.8)	3.8(0.3-13.1)	1.2(0-5.5)	0.5(-0.6 to 3.1)
47 40-50 yr	0.48(-0.65 to 1.42)	0.3(0-1.9)	3.3(0.6-12.7)	1.6(0-8.8)	0.2(-1.9 to 3.2)
48 >50 yr	0.58(0-1.42)	0.4(0-1.4)	3.3(0.8-11.0)	1.5(0-7.4)	0.7(-0.8 to 2.2)
Lead aVR					
49 Birth-1 da	-0.90(-2.00 to 1.50)	2.4(0-4.0)	3.9(0-9.0)	0.3(0-9.5)	-0.5(-3.2 to 2.4)
50 1 da-1 wk	-1.23(-2.00 to 0)	2.7(0-4.0)	2.7(0-7.0)	1.9(0-7.0)	-1.6(-3.2 to 0)
51 1 wk-1 mo	-1.03(-2.00 to 1.00)	4.0(0-9.0)	2.2(0-6.0)	2.0(0-8.0)	-2.3(-3.9 to -0.6)
52 1-3 mo	-1.12(-2.00 to 0)	7.1(0-12.0)	2.1(0-4.0)	4.3(0-14.5)	-2.3(-4.7 to -0.2)
53 3-6 mo	-1.06(-2.00 to 0)	8.0(0-12.0)	1.8(0-5.0)	6.0(0-15.0)	-2.9(-4.5 to -0.3)
54 6 mo-1 yr	-1.22(-2.50 to 0)	8.1(0-13.0)	2.3(0-5.0)	5.0(0-18.0)	-2.8(-5.1 to -0.4)
55 1-3 yr	-1.31(-2.00 to 0)	7.3(0-14.0)	2.2(0-6.0)	5.0(0-15.0)	-2.9(-5.2 to -0.6)
56 3-5 yr	-1.20(-2.00 to -1.00)	7.4(0-13.5)	1.7(0-5.0)	5.0(0-14.0)	-2.6(-4.9 to -0.6)
57 5-8 yr	-2.00(-2.08 to 0)	8.2(0-13.0)	1.6(0-5.0)	5.2(0-15.0)	-3.2(-5.3 to -0.7)
58 8-12 yr	-1.24(-2.50 to -0.50)	8.2(0-12.5)	1.3(0-4.0)	1.0(0-16.0)	-2.9(-5.2 to -0.5)
59 12-16 yr	-1.14(-2.00 to -0.50)	7.9(0-14.0)	1.2(0-3.0)	0.8(0-14.0)	

70. AMPLITUDES OF ELECTROCARDIOGRAPHIC WAVES VS AGE- MAN (Continued)

70. AMPLITUDES OF ELECTROCARDIOGRAPHIC WAVE							
Age	P		Q		R	S	T
	(A)	(B)	(C)	(D)	(E)	(F)	
Lead aVR (concluded)							
60	16-20 yr	-0.84(-1.75 to -0.25)	2.1(0-8.9)	1.9(0-4.7)	4.3(0-11.1)	-1.9(-4.8 to -0.1)	
61	20-30 yr	-0.78(-1.54 to -0.10)	2.5(0-11.2)	1.1(0-3.6)	3.4(0-11.1)	-2.7(-5.3 to -0.6)	
62	30-40 yr	-0.81(-1.79 to -0.10)	2.1(0-9.0)	1.0(0-3.3)	3.5(0-14.7)	-2.4(-5.4 to -0.3)	
63	40-50 yr	-0.79(-1.38 to -0.28)	2.0(0-8.5)	0.9(0-4.1)	3.5(0-13.2)	-2.3(-4.7 to -0.2)	
64	>50 yr	-0.89(-1.15 to 0.54)	2.9(0-10.2)	1.0(0-2.8)	2.9(0-9.5)	-2.2(-4.6 to -1.5)	
Lead aVL							
65	Birth-1 da	0.10(-1.50 to 1.50)	1.3(0-2.0)	2.0(1.0-6.0)	6.5(0-16.0)	0(-2.4 to 2.6)	
66	1 da-1 wk	0.11(-1.50 to 1.00)	0.7(0-1.0)	1.6(0-3.0)	5.5(0-13.0)	0.9(-0.8 to 2.7)	
67	1 wk-1 mo	0.26(-1.00 to 1.50)	1.3(0-1.5)	2.3(0-7.0)	6.0(0-11.0)	1.0(-1.2 to 3.1)	
68	1-3 mo	0.18(-1.00 to 1.00)	1.7(0-2.0)	3.6(1.0-9.0)	4.8(0-9.5)	0.7(-1.5 to 3.1)	
69	3-6 mo	0.35(-0.50 to 1.00)	1.1(0-1.5)	3.9(0-8.0)	4.0(0-7.0)	0.8(-0.6 to 2.4)	
70	6 mo-1 yr	0.29(-1.50 to 1.00)	1.8(0-5.5)	4.6(1.0-10.0)	4.2(0-11.0)	1.2(-1.0 to 3.6)	
71	1-3 yr	0.36(-1.00 to 1.00)	1.3(0-4.0)	4.3(1.0-10.0)	3.8(0-9.0)	1.4(-0.5 to 3.4)	
72	3-5 yr	0.44(-1.00 to 1.00)	1.3(0-3.0)	3.3(0-7.0)	3.2(0-10.0)	1.2(-1.0 to 3.6)	
73	5-8 yr	0.30(-1.00 to 1.50)	1.2(0-4.0)	2.6(0-7.0)	3.0(0-14.0)	1.0(-1.2 to 3.3)	
74	8-12 yr	0.47(-1.00 to 1.00)	1.2(0-6.0)	2.7(0-8.0)	3.0(0-20.0)	1.3(-0.7 to 3.6)	
75	12-16 yr	0.28(-1.00 to 1.00)	1.3(0-6.0)	2.4(0-12.0)	2.0(0-7.7)	1.1(-1.1 to 3.6)	
76	16-20 yr	0.25(-0.85 to 1.42)	0.3(0-2.3)	1.9(0-2.5)	2.1(0-10.1)	0.6(-1.8 to 3.6)	
77	20-30 yr	0.22(-0.95 to 1.22)	0.3(0-3.5)	2.2(0-7.6)	1.3(0-9.0)	0.9(-1.5 to 3.7)	
78	30-40 yr	0.71(-0.85 to 1.40)	0.3(0-2.2)	2.6(0-7.5)	1.4(0-11.3)	0.9(-1.6 to 2.7)	
79	40-50 yr	0.77(-0.95 to 1.35)	0.3(0-1.3)	3.2(0.3-10.1)	1.4(0-6.3)	0.9(-1.9 to 3.8)	
80	>50 yr	0.07(-0.95 to 0.70)	0.4(0-1.5)	2.7(0.4-5.7)		0.6(-1.1 to 2.7)	
Lead aVF							
81	Birth-1 da	1.06(-1.00 to 2.00)	1.9(0-6.0)	6.5(2.0-20.0)	0.8(0-7.0)	0.9(-1.2 to 3.0)	
82	1 da-1 wk	1.18(0-2.00)	2.0(0-4.0)	5.8(2.0-13.0)	0.7(0-3.0)	0.2(-1.6 to 3.8)	
83	1 wk-1 mo	1.23(-1.00 to 3.00)	1.9(0-4.0)	6.8(2.0-15.0)	0.5(0-5.0)	0.5(-0.5 to 3.7)	
84	1-3 mo	1.16(0-2.00)	2.5(0-5.0)	8.8(0.5-17.8)	0.8(0-4.0)	2.0(-0.1 to 3.9)	
85	3-6 mo	1.13(0.50-2.00)	2.7(0-6.0)	10.1(0.5-20.2)	0.9(0-6.0)	2.2(0-4.2)	
86	6 mo-1 yr	1.25(0-2.00)	2.9(0-6.0)	9.4(0-17.2)	1.1(0-4.0)	1.9(-0.1 to 3.7)	
87	1-3 yr	1.24(0-2.00)	2.2(0-5.0)	8.2(1.0-20.0)	0.8(0-5.0)	1.7(-0.6 to 4.1)	
88	3-5 yr	1.15(0-2.00)	1.7(0-5.0)	9.8(0.5-18.8)	0.7(0-3.5)	2.1(-0.7 to 4.9)	
89	5-8 yr	1.24(0-2.00)	1.5(0-5.0)	9.7(0.5-18.7)	0.9(0-7.0)	2.0(-0.6 to 4.5)	
90	8-12 yr	1.12(0-2.00)	1.5(0-3.0)	10.2(0.7-20.0)	1.0(0-5.0)	2.3(-0.5 to 4.7)	
91	12-16 yr	1.18(0-2.00)	1.3(0-3.0)	10.5(0.1-21.8)	1.0(0-4.0)	2.4(-0.7 to 5.4)	
92	16-20 yr	0.76(-0.80 to 1.80)	0.4(0-3.8)	7.7(1.8-14.0)	1.3(0-4.9)	1.9(-0.5 to 5.4)	
93	20-30 yr	0.71(-1.77 to 1.61)	0.4(0-2.8)	6.8(0.6-16.1)	1.0(0-5.9)	2.4(-0.6 to 5.2)	
94	30-40 yr	0.77(-0.60 to 1.60)	0.3(0-1.9)	5.3(0.2-14.8)	1.0(0-7.1)	1.7(-0.4 to 4.6)	
95	40-50 yr	0.73(0-1.52)	0.3(0-1.8)	4.5(0.5-12.1)	1.1(0-5.2)	1.4(-0.5 to 4.0)	
96	>50 yr	0.74(-0.30 to 1.25)	0.3(0-1.0)	4.0(0.8-13.2)	0.8(0-3.2)	1.6(0.3-4.9)	
Lead V ₁							
97	Birth-1 da	0.96	0	16.0(6.0-27.0)	10.0(0-28.0)	0.6(-4.0 to 5.0)	
98	1 da-1 wk	1.10	0	17.0(4.0-30.0)	11.0(4.0-25.0)	-3.6(-6.0 to -2.0)	
99	1 wk-1 mo	0.63(-1.00 to 2.00)	0	13.5(2.6-24.0)	7.0(0-17.0)	-2.8(-8.0 to -0.8)	
100	1-3 mo	0.42(-1.00 to 1.50)	0	10.0(1.5-19.0)	6.5(0-20.0)	-2.7(-7.0 to 2.0)	
101	3-6 mo	0.39	0	12.0(5.0-19.0)	7.0(3.0-12.0)	-4.2(-7.5 to -0.4)	
102	6 mo-1 yr	0.73(-1.00 to 1.00)	0	12.8(1.5-20.0)	8.0(3.0-18.0)	-3.9(-7.4 to 0)	
103	1-3 yr	0.81(-1.00 to 2.00)	0	9.0(3.9-18.0)	7.0(0-28.0)	-3.7(-6.0 to -2.0)	
104	3-5 yr	1.03	0	7.0(2.0-16.0)	13.0(6.0-33.0)	-3.8(-6.2 to -1.0)	
105	5-8 yr	0.94	0	7.0(1.2-20.0)	14.0(4.0-26.0)	-3.5(-6.0 to -0.8)	
106	8-12 yr	0.68	0	6.0(1.8-16.0)	13.0(5.0-26.0)	-2.9(-5.6 to -0.5)	
107	12-16 yr	0.69	0	5.5(0-16.0)	14.5(5.0-26.0)	-1.5(-4.0 to 7.0)	
108	16-20 yr	0.65(-0.22 to 2.20)	0	4.6(0.4-16.7)	11.7(1.8-25.1)	0.7(-3.3 to 6.0)	
109	20-30 yr	0.62(-1.11 to 1.80)	0	3.7(0.07-11.80)	11.5(2.5-26.2)	1.3(-3.6 to 7.6)	
110	30-40 yr	0.61(-0.80 to 1.80)	0	3.3(0-9.4)	8.4(0.8-21.3)	0.9(-2.0 to 12.2)	
111	40-50 yr	0.46(-0.70 to 1.20)	0	3.2(2.2-10.6)	9.3(0.9-20.0)	1.1(-2.3 to 5.6)	
112	>50 yr	0.38(-0.38 to 1.40)	0	2.3(0.8-15.5)	7.3(2.0-17.3)	0.6(-3.7 to 3.7)	
113	Birth-1 da	1.05					
114	1 da-1 wk	1.22					
115	1 wk-1 mo	1.30(0.50-2.00)					
116	1-3 mo	0.81					
117	3-6 mo	1.36(0.50-2.00)					
118	6 mo-1 yr	1.00					
119	1-3 yr	1.39					
120	3-5 yr	1.05	0	13.5(1.0-25.0)	22.0(16.0-34.0)	-4.1(-7.0 to -1.0)	

70. AMPLITUDES OF ELECTROCARDIOGRAPHIC WAVES VS AGE: MAN (Concluded)

Age	Wave				
	P	Q	R	S	T
	(B)	(C)	(D)	(E)	(F)
Lead V ₆ (concluded)					
181 3-6 mo	0.79(0-1.50)	1.8(0-4.0)	13.0(0.5-17.0)	1.7(0-5.0)	2.7(0-5 to 6.0)
182 6 mo-1 yr	0.84(0-1.50)	2.4(0-5.0)	13.0(4.5-21.0)	2.0(0-3.0)	2.9(0.7-5.0)
183 1-3 yr	0.84(0.50-1.50)	1.7(0-4.5)	12.0(2.5-21.5)	1.2(0-4.0)	3.3(0.6-5.6)
184 3-5 yr	0.75(0.50-1.00)	1.8(0-4.0)	13.0(3.5-22.0)	0.7(0-3.0)	3.6(0.8-6.4)
185 5-8 yr	0.74(0-1.50)	1.7(0-5.0)	14.0(6.5-21.5)	0.9(0-5.0)	4.0(0.3-7.7)
186 8-12 yr	0.70(0-1.50)	1.7(0-4.0)	13.5(10.0-22.0)	1.1(0-3.0)	4.4(0.5-8.4)
187 12-16 yr	0.66(0-1.50)	1.3(0-2.5)	13.5(7.0-21.0)	1.2(0.2-5.0)	4.0(0.8-7.2)
188 16-20 yr	0.50(0-1.00)	0.6(0-4.2)	9.8(3.5-24.4)	1.5(0-11.2)	2.9(0-7.1)
189 20-30 yr	0.53(0-1.00)	0.5(0-2.6)	10.7(2.2-22.6)	1.6(0-14.3)	3.3(0.5-6.9)
190 30-40 yr	0.56(0-1.00)	0.4(0-2.7)	9.4(2.5-26.0)	1.3(0-8.4)	2.8(0-6.7)
191 40-50 yr	0.46(0-1.11)	0.4(0-2.6)	8.8(3.6-19.0)	0.9(0-7.0)	2.8(0-6.8)
192 >50 yr	0.52(0.16-0.95)	0.4(0-1.1)	8.6(2.0-15.5)	0.8(0-4.7)	2.5(0.7-5.6)

Contributor Lepeschkin, Eugene

References: From birth-16 yr: [1] Ziegler, R. F. 1951. Electrocardiographic studies in normal infants and children. C. C. Thomas, Springfield, Ill. [2] Nadrai, A. 1938. Zschr. Kinderh. 60 285. [3] Romero-Neto, M. M. 1949. Derivacoes unipolares do precordio em criancas normais. São Paulo [4] Yu, P. N. G., H. A. Joos, and C. P. Katsampas. 1951. Am. Heart J. 41:91. [5] Allimurung, M. M., L. G. Joseph, A. S. Nadas, and R. F. Massell. 1951. Circulation, N. Y. 4:420. From 16->50 yr: [6] Vaquero, M., R. Limon, and A. Limon. 1948. Mem. II Congr. Interamer. Card. 2 887.

71. R WAVE EXPRESSED AS PERCENTAGE OF RS DEFLECTION MAN

Age	Lead					
	V ₁	V ₂	V ₃	V ₄	V ₅	V ₆
	(B)	(C)	(D)	(E)	(F)	(G)
1 Birth-1 da	62(26-96)	48(22-75)	41(25-56)	44(15-72)	43(10-77)	50(0-100)
2 1 da-1 wk	58(35-81)	48(22-75)	53(20-85)	56(12-100)	60(20-100)	65(25-100)
3 1 wk-1 mo	66(34-94)	52(29-75)	55(34-76)	63(45-78)	66(33-100)	76(30-100)
4 1-3 mo	63(27-100)	54(26-79)	56(33-81)	62(32-92)	71(45-100)	78(60-100)
5 3-6 mo	60(33-86)	57(36-76)	59(32-86)	71(50-100)	78(50-100)	86(60-100)
6 6 mo-1 yr	56(28-84)	50(21-78)	56(33-83)	70(45-100)	80(55-100)	88(70-100)
7 1-3 yr	45(32-67)	42(23-65)	54(36-83)	75(42-100)	90(56-100)	94(75-100)
8 3-5 yr	42(21-67)	40(18-63)	54(22-83)	78(52-100)	85(65-100)	95(68-100)
9 5-8 yr	35(10-55)	33(5-62)	48(12-84)	76(58-96)	86(67-100)	94(69-100)
10 8-12 yr	34(16-50)	32(10-54)	46(11-81)	75(38-96)	88(58-100)	95(80-100)
11 12-16 yr	27(10-55)	27(10-56)	49(13-85)	78(50-100)	87(70-100)	90(70-100)
12 Adults	25(0-50)	17(9-93)	42(9-90)	80(17-95)	88(50-96)	90(70-96)

Contributor Lepeschkin, Eugene

References: From birth-16 yr: [1] Ziegler, R. F. 1951. Electrocardiographic studies in normal infants and children. C. C. Thomas, Springfield, Ill. Adults [2] Lepeschkin, E. 1951. Modern electrocardiography. Williams and Wilkins, Baltimore, v. 1.

72. AMPLITUDES OF ELECTROCARDIOGRAPHIC WAVES: FARM ANIMALS

Animals were tested in standing position. Values are in millimeters. Ranges are estimate "c" (cf. Introduction).

Animal ¹	P		Wave			T	
	Monophasic	Diphasic	Q	R	S	Monophasic	Diphasic
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)
Lead I							
Cattle, dairy							
1 Calf	0.3-0.5	0.3-0.5	0.6-1.2	0.2-5.5		-1.5 to 1.5	0.5-1.0
2 Cow	0.6-1.0	0.4-1.5	0.3-2.3	0.4-2.5	0.2, 5.0	-2.0 to -1.0	0.5-3.0
3 Goat	0-0.1	0.5	0.5-1.2	0.5-2.2		0.5-1.5	
4 Horse, draft	0.1-0.5	2.0	1.0	0.2-5.0	0.5	-3.5 to -1.2	2.0
5 Mule	0.1-0.5			1.0-4.0		-3.0 to 4.5	
6 Pig	0.5-1.0		1.0-1.4	1.5-4.5		-2.5 to 1.0	
7 Sheep	0.2-0.5	0.2	0.7-3.0	1.2-4.5	6.0	-3.0 to 2.0	
Lead II							
Cattle, dairy							
8 Calf	0.4-1.2	0.8-1.2	1.0-5.2	0.8-4.2	0.2-1.5	-3.0 to 3.2	
9 Cow	0.4-1.2	0.3, 1.2	0.4-1.9	0.7-8.0	0.2, 1.5	-3.0 to 4.0	2.0-3.5
10 Goat	0-0.2		0.4-1.2	1.3-3.0		0.7-3.0	
11 Horse, draft	0.6-3.0	3.0-4.0	0.2-1.5	1.0-9.0	0.2-1.5	-5.5 to 5.0	1.5-4.0
12 Mule	1.2-2.5			8.0-12.2	0.5-3.0	-3.0 to 2.5	
13 Pig	0.6-1.2		0.3-1.5	3.2-6.0	1.0	-4.0 to -2.0	1.2-4.0
14 Sheep	0.2-0.6		0.5-4.0	0.7-5.0	4.0	-4.0 to 0.5	
Lead III							
Cattle, dairy							
15 Calf	0.2-1.0	8.0	1.0	0.8-4.0	1.4-1.5	1.2-2.5	1.5
16 Cow	0.5-1.0	0.3	1.2-2.7	0.5-9.0	5.0	0-3.0	1.2-2.0
17 Goat	0-0.3		1.0	1.0-5.3	0.9	0.5-2.0	
18 Horse, draft	2.2-2.7		0.5-1.0	2.0-8.0	1.5-3.0	1.5-4.5	2.5-5.0
19 Mule	1.0-2.0			6.0-9.0	2.5-4.0	1.5-4.0	2.8
20 Pig	0.3-0.8		0.5-0.8	0.5-4.0	2.5	2.0-3.7	0.7
21 Sheep	0.1-0.5		0.3-3.0	0.5-4.0	2.0	-2.0 to 2.0	

/1/ Subjects: 13 calves, 10 cows, 5 goats, 13 horses, 4 mules, 6 pigs, 12 sheep.


Contributor: Brody, Samuel

Reference: Plainer, W. S., H. H. Kibler, and S. Brody. 1948. Missouri Agr. Exp. Sta. Res. Bull. 419, p. 9.

73. EVENTS DURING THE CARDIAC CYCLE- MAN, DOG

Relative time intervals of events within the cardiac cycle may vary with change in heart rate.

Part I. SCHEMATIC



/1/ Schematic presentation of phases of the cardiac cycle, not an exact depiction of the duration of events.

Contributor: Wiggers, Carl J.

Reference: Wiggers, C J. 1952 Circulatory dynamics. Grune and Stratton, New York.

Part II. TABULAR

Phase	Interval	Events at Onset of Phase	Main Events during Phase	Events at Close of Phase	Phase Duration ¹ sec	
					Man (F)	Dog (G)
1 Isometric contraction	I	Onset of ventricular contraction	Closure of atrio-ventricular valves, rapid rise of intraventricular pressure	Opening of semilunar valves	0.05	0.05
2 Maximum ejection	II	Opening of semi-lunar valves	Rapid outflow of blood from ventricles	Peak of intraventricular pressure	0.09	0.12
3 Reduced ejection	III	Peak of intraventricular pressure	Declining outflow of blood from ventricles	Onset of ventricular relaxation	0.13	0.10
4 Protodiastole	IV	Onset of ventricular relaxation	Rapid drop in intraventricular pressure	Closure of semilunar valves	0.04	0.02
5 Isometric relaxation	V	Closure of semi-lunar valves	Continued ventricular relaxation with no volume change	Opening of atrio-ventricular valves	0.08	0.05
6 Rapid inflow	VI	Opening of atrio-ventricular valves	Rapid flow of blood from atria to ventricles	Slowing of inflow rate	0.11	0.06
7 Diastasis	VII	Slowing of inflow from atria to ventricles	Continued slower flow from atria to ventricles	Onset of atrial contraction	0.19	0.29
8 Atrial systole	VIII	Onset of atrial contraction	Increased flow from atria to ventricles	Termination of atrial, and onset of ventricular, contraction	0.11	0.11

^{1/1} Assumed heart rate: 75 beats/min.

Contributors: (a) Gleckler, William J., (b) Soloff, Louis A.

References [1] Wiggers, C. J. 1923. Circulation in health and disease. Lea and Febiger, Philadelphia. [2] Wiggers, C. J. 1949. Physiology in health and disease. Ed. 5. Lea and Febiger, Philadelphia. [3] Rushmer, R. F. 1956. Physiol Rev. 36:400. [4] Rushmer, R. F. 1955. Cardiac diagnosis. W. B. Saunders, Philadelphia.

74. DURATION OF HEART SOUNDS AT VARIOUS AGES. MAN

Method. A = recorded at apex with stethoscopic microphone, B = recorded at apex with logarithmic microphone; C = recorded at aortic area with stethoscopic microphone. Values in parentheses are ranges, estimate "c" (cf. Introduction).

Age yr	Method	Sound, sec			Interval Between 2nd and 3rd Sound sec	Reference
		1st (C)	2nd (D)	3rd (E)		
1 <4	A	0.085	0.055			1
2	C	0.070	0.060			
3 4-10	A	0.120	0.065	0.050	0.12 ¹	1
4	C	0.145	0.110	0.040	0.14 ¹	1
					0.14 ¹	1
					(0.18-0.24) ²	2
					(0.16-0.19) ²	
					0.16 ¹	1
					0.18 ¹	
					0.18 ¹	1
					0.19 ¹	1

^{1/1} Interval between the peak of the largest oscillations of the 2nd sound and the beginning of the 3rd sound.
^{2/2} Interval between the beginning of the 2nd heart sound and the mid-point of the 3rd heart sound.

Contributor: Kuo, Peter T.

References: [1] Luisada, A. A., F. Mendoza, and M. M. Alimurung 1949. Brit. Heart J. 11:41. [2] Rappaport, M. B., and H. B. Sprague. 1942. Am. Heart J. 23:591.

75. CHARACTERISTICS OF FUNCTIONAL HEART MURMURS: MAN

Variable	Area				Reference
	Pulmonary	Precordial ¹	Parasternal ²	Mitral	
(A)	(B)	(C)	(D)	(E)	(F)
1 Age	Children and young adults.	Children and young adults.	Children and adults.	All ages.	B,1-13;C,5-14;D,E,14
2 Intensity	Faint or loud.	Faint or loud.	Up to loud.	Loud.	B,1,2,5-13;C,5-14,D,14,15,E,14
3 Effect of posture	Louder with recumbency.	Usually louder with recumbency, but may become louder in erect position.	Little effect.	Little or variable effect.	B,5,6,C,5,6,14,D,14,E,2,14
4 Effect of respiration	Disappears or becomes faint with deep inspiration.	Either unaffected or faint with deep inspiration.	Persists with deep inspiration.	Persists with deep inspiration.	B,5,6,C,5,6,14,D,E,14
5 Position of murmur in cardiac cycle	Early or mid-systolic.	Mid-systolic.	Mid-systolic.	Late systolic (may be due to organic mitral valve incompetence).	B,1,2,5,6,C,5,6,14;D,14,E,14,b
6 Character of murmur	Usually of low intensity, blowing (may be high pitched)	Musical, vibratory, blowing, or twanging	Blowing, or rough and musical.	Blowing, rough, and partially musical.	B,1,2,5-13,C,2,5-14,D,15,E,2,14
7 Transmission of murmur	Often also audible at apex.	If loud, sometimes transmitted toward either apex or base, or both.	If loud, may be audible at base and near apex.	Audible over precordium.	B,5,6;C,5,6,14;D,E,14

/1/ Includes space immediately medial to the mitral area and extending to the left sternal border in the 4th and 5th interspaces /2/ Soft mid-systolic parasternal or pulmonic murmurs occur in atrial septal defect.

Contributors: (a) Kuo, Peter T., (b) Perloff, Joseph K.

References. [1] Wood, F. C Unpublished. [2] McKusick, V. A. 1958. Cardiovascular sound in health and disease. Williams and Wilkins, Baltimore. [3] Paulin, S., and E. Mannheim. 1957. Acta paediat., Upps. 46 439 [4] Levine, S. A., and W. P. Harvey. 1949. Clinical auscultation of the heart. W. B. Saunders, Philadelphia. [5] Friedman, S., W. A. Robie, and T. N. Harris. 1949. Pediatrics, Springf. 4 782. [6] Wollerth, C C Unpublished. [7] Ash, R 1948. Am. Heart J. 36:89. [8] Harris, T. N., and S. Friedman. 1952. Ibid. 43 707. [9] Harris, T. N. 1955. Ibid. 50 805. [10] Harris, T. N., S. Friedman, and C. F. Haub. 1949. Pediatrics, Springf. 3 845. [11] Measehoff, C. R. 1949. Am. J. M. Sc. 217.17. [12] Lynxwiler, C. P., and J. L. Donahoe. 1955. South. M. J. 48 164. [13] Stuckey, D 1957. Med. J. Australia 44.38. [14] Evans, W. 1947. Brit Heart J. 9 1. [15] Wells, B 1957 Ibid. 19.129.

76. PHYSIOLOGIC HEART SOUNDS AND MURMURS; MAN

Heart Sound or Murmur (A)	Probable Mechanism of Production (B)	Frequency cps (C)	Duration sec (D)	Phonocardiographic Registration			Audible Manifestation		Reference
				Linea ²	Stethoscopic ³	Logarithmic ⁴	In Normal Subj. (H)	Elicitation (I)	
1 Atrial sound	Tensing of atrium or vibration of distended ventricles.	28-33	0.04-0.06	Yes	88%	21%	No	Pathologic conditions; occasionally in normal children.	B,1,2,C,D,3; E-G,4,H,1,5 (J)
2 1st heart sound	Muscular, valvular, and vascular factors.	33-111	0.06-0.16	Yes	Yes	Yes	Yes	Intensity varies with many physiologic and pathologic conditions.	B,1,6,C,D,7; E-G,4,H,1,5
3 1st component	Pre-isometric ventricular and valvular factors.	35	0.03-0.06	Yes	Attenuated	No	No	None.	B-D,6;E-G,4; H,1,a
4 2nd component	Ventricular muscle tension and contraction, closure of atrioventricular valves.	34-39	0.03-0.05	Attenuated	Yes	Yes	Yes	Normally heard as a single sound, but sometimes is split due to asynchronous closure of atrioventricular valves under physiologic and pathologic conditions. Physiologically: mitral closure before tricuspid closure.	B-D,1,E-G,4; H,a,1,5
5 3rd component	Opening of semilunar valves, onset of ventricular ejection.	37-40	0.05-0.06	Attenuated	Yes	Yes	Yes		
6 4th component	Acceleration of blood in arterial vessels.	31-33	0.06-0.07	Yes	Attenuated	No	No	Pathologic conditions.	B-D,1,E-G,4; H,1,8
7 2nd heart sound	Valvular and vascular factors.	31-111	0.04-0.11	Yes	Yes	Yes	Yes	Intensity varies with many physiologic and pathologic conditions.	B,1,7,C,D,7;E-G,4,H,1,5
8 1st component	Relaxation of ventricles.		Yes	Yes	Attenuated	Attenuated	No	None.	B,E-G,4;H,1,9
9 2nd component	Closure of semilunar valves.		Yes	Yes	Yes	Yes	Yes	Sometimes is split under physiologic and pathologic conditions. Physiologically: aortic closure before pulmonic closure.	B,E-G,4;H,9,1,5,10
10 3rd component	Vibrations of arterial wall and blood column.		Yes	Yes	Attenuated	No	No	None.	B,E-G,4;H,1,9
11 4th component	Opening of atrioventricular valves.		Yes	Yes	Attenuated	No	No	Pathologic conditions.	B,E-G,4;H,1,5,9
12 3rd heart sound (diastolic)	Distention of ventricles during period of rapid filling.	32-50	0.02-0.09	Yes	82%	30%	6-20%	In children and young adults: at apex, during inspiration in recumbent posture; pathologic conditions.	B,1,C,3-D-G,4,H,1,1,2,1,5
13 Systolic click	Extracardiac factors.		Short				0.5%	At apex and lower left sternal edge, varying with posture and respiration.	B,D,1,5,10;H,12
14 Apical systolic murmur	Accelerated blood flow, acoustical characteristics of chest wall, respiratory factors.	120			58%	85%	2.3-9%	Acceleration of heart rate; deep expiration, recumbent posture, asthenia.	B,5,10,a,C,11; F,G,4;H,1,1,13,1,5,10,14

15	Precordial "vibratory" murmur	Vibration of cardiac tissue	128	0.114				Yes	In children and young adults; recumbent posture.	B, C, D, 15, H, C, 1, 12, 13
16	Pulmonary systolic murmur	Accelerated blood flow, position of pulmonary artery, acoustical characteristics of chest wall	Medium	Moderate				Yes	Acceleration of heart rate; deep expiration, recumbent posture.	B, 10, a, C, D, a, H, C, 1, 10, 14

11/ Cycles per second. 12/ Records the heart sounds on the surface of the chest, as well as pulsations of the chest wall; actually an "apex cardiogram." 13/ Records the sound vibrations perceived by an observer of normal hearing, using an average acoustic bell stethoscope (light touch). 14/ Records the sound vibrations (audible heart sounds) perceived by an observer of normal hearing, using an average acoustic diaphragm stethoscope.

Contributors. (a) Porter, Reno R., (b) Kuos, Peter T., (c) McKusick, Victor A.

References: [1] Orlos, O., and E. Braun-Menendez. 1939. The heart sounds in normal and pathological conditions. Oxford University Press, New York. [2] McKusick, V. A. 1958. Cardiovascular sound in health and disease. Williams and Wilkins, Baltimore. [3] Boyer, N. R., R. W. Eckstein, and C. J. Wiggers. 1940. *Am. Heart J.* 19:257. [4] Happort, M. B., and H. B. Sprague. 1942. *Ibid.* 23:591. [5] Levine, S. A., and W. P. Harvey. 1949. Clinical auscultation of the heart. W. B. Saunders, Philadelphia. [6] Counihan, T., A. L. Messer, M. B. Reppasport, and H. B. Sprague. 1951. *Circulation*, N. Y. 3:710. [7] Wiggers, C. J. 1944. *Physiology in health and disease*. Ed. 4. Lea and Febiger, Philadelphia. [8] Leatham, A., and L. Vogelbeil. 1954. *Brit. Heart J.* 16:21. [9] Luisado, A. A. 1947. *A physiologic and clinical study of cardiovascular disease*. Williams and Wilkins, Baltimore. [10] White, P. D. 1951. *Heart disease*. Ed. 4. Macmillan, New York. [11] McKee, M. M. 1938. *Am. Heart J.* 16:305. [12] Friedman, S., W. A. Robie, and T. N. Harris. 1949. *Pediatrics*, Springf. 4:782. [13] Sachs, H. A., and S. C. Roberts. 1939. *Am. Heart J.* 16:305. [14] Examination of the heart. 1931. American Heart Association, New York. [15] Harris, T. N., S. Friedman, and C. F. Haub. 1949. *Pediatrics*, Springf. 3:445.

77. CONGENITAL MALFORMATIONS OF THE HEART: MAN

Murmur: Parentheses indicate infrequent occurrence. QRS complex: RVH = right ventricular hypertrophy; RBBB = right bundle branch block; LVH = left ventricular hypertrophy; CVH = combined ventricular hypertrophy. Hemodynamics: RA = right atrium; RV = right ventricle; PA = pulmonary artery; AO = aorta or systemic artery; PF = pulmonary blood flow, CO = cardiac output. (0) = normal, (+) = increase, (-) = decrease.

Cardiac Malformation	Murmur	Roentgenographic Findings			Electrocardiographic Findings			Hemodynamics										Prospect of Surgical Improvements
		Size and Shape of Heart (C)	Pulmonary Conus (D)	Pulmonary Vascularity (E)	QRS Axis (F)	QRS Complex (G)	Pressure RA ¹ RV ¹ PA ¹ RA ¹ RV ¹ PA ¹ AO ¹	O ₂ Saturation (H) (I) (J) (K) (L) (M) (N)	Ratio ² (O)	(P)								
1 Tetralogy of Fallot	Systolic ⁴ (Continuous) ⁵	Boot shaped, usually not enlarged	Concave	-	Right	RVH	0	+	-	0	0	0	-	<1	Good			
2 Pulmonary stenosis with atrial septal defect	Systolic	Right ventricle enlarged	Usually convex	-	Right	RVH	+	+	-	0	0	0	-	<1	Good			
3 Transposition of great vessels with pulmonary stenosis	Systolic	Both ventricles enlarged; supracardiac vessels	Concave	-	Right	RVH	0	+	-	0	0	0	-	<1	None			
4 Complete transposition of great vessels	Systolic (Diastolic)	Enlarged, supracardiac vessels	Convex	+	Usually right	RVH or CVH	0	+	+	0	+	+	-	>1	None			
5 Ebstein's anomaly, may be associated with a patent foramen ovale, an intact atrial septum, or an atrial septal defect	Systolic Diastolic None	Enlarged right atrium which is recessed into right ventricular cavity.	Convex	-	Right	RBBB (usually)	+	0	0	0	0	0	-	<1	Poor			
6 Tricuspid atresia	Systolic	Right ventricle diminutive	Concave	-	Left	LVH	+	0	0	0	0	0	-	<1	Fair to poor			
7 Eisenmenger's complex	Systolic (Diastolic)	Both ventricles enlarged	Convex	+	Right	RVH or CVH	0	+	+	0	+	+	-	<1	Poor			
8 Persistent truncus arteriosus	Systolic Diastolic None	Both ventricles enlarged	Concave	+	(may or may not)	RVH or CVH	0	+	+	0	+	+	-	>1 (usually)	None			
9 Patent ductus arteriosus with reversed shunt	Systolic Diastolic ⁷ None	Both ventricles enlarged	Convex	+	Right	CVH	0	+	+	0	0	+	-	>1	Poor			
10 Complete anomalous pulmonary venous drainage	Systolic Diastolic (Continuous)	Characteristic "figure eight" silhouette	Convex	+	Right	RVH	0	+	+	0	0	+	-	>1	Poor			
11 Isolated pulmonary stenosis	Systolic	Right ventricle enlarged	Concave or convex	-	Right	RVH	+	+	-	0	0	0	0	<1	Good			
12 Atrial septal defect, secundum type	Systolic (Diastolic)	Right ventricle enlarged	Convex ⁸	+	Right	RBBB or RVH	0	0	0	+	+	+	0	>1	Good			


13	Atrial septal defect, ostium primum type Ventricular septal defect	Systolic Diastolic	Both ventricles enlarged	Convex	+	Right or left	CVII	0	+	+	+	+	-	>1	Fair to poor
14	Small	Systolic	Normal	Convex	0	0	0	0	0	0	0	0	0	>1	Good
15	Large	Systolic None (Diastolic)	Both ventricles enlarged	Convex	+	Right	RVII, RDBB or CVII	0	+	0	+	0	+	>1	Fair
16	Patent ductus arteriosus	Continuous (Diastolic)	Normal or left ventricle enlarged	Convex	+	Left	0 or LVII	0	0	0	0	0	0	>1	Good

11/ Indicated increase in pressure refers to height of A wave of atrial pulses; mean pressure is usually normal. 12/ Does not take into account collateral circulation to lungs when pulmonary stenosis is present. 13/ Only the more common pathologic combinations are listed. The hemodynamics in each complicated anomaly [determined by relative severity of component abnormalities] vary widely, and are discussed in detail in the references. 14/ Across the pulmonary valve. 15/ Continuous murmur originates in bronchial collaterals and is generally confined to individuals with pulmonary atresia. 16/ Ventricle is functionally divided into 2 chambers with different pressures. 17/ Of pulmonary insufficiency only. 18/ Normal when shunt is small.

Contributors: (a) Witham, A. Cathoun, (b) Glenn, William W. L., (c) Perloff, Joseph K.

References: [1] Kjellberg, S. R., E. Mannheimer, O. Rudhe, and B. Jonason. 1955. Diagnosis of congenital heart disease. Year Book Publishers, Chicago. [2] Brown, J. W. 1950. Congenital heart disease. Ed. 2. Staples Press, London. [3] Nadas, A. 1957. Pediatric cardiology. W. B. Saunders, Philadelphia. [4] Witham, A. C., and R. Ellison. 1957. Am. J. Med. 22:593. [5] Gould, S. E. 1953. Pathology of the heart. C. C. Thomas, Springfield, Ill. [6] McKusick, V. A. 1950. Cardiovascular sound in health and disease. Williams and Wilkins, Baltimore. [7] Bing, R. J., T. A. Lombardo, L. M. Burgeron, M. Taeschler, and S. Tulug. 1952. Ann. Int. M. 37 664. [8] Cournaud, A., J. S. Baldwin, and A. Himmelfeld. 1949. Cardiac catheterization in congenital heart disease. Commonwealth Fund, New York.


Only those murmurs with typical configuration are presented; schematic drawings simulate shape but not frequency. Abbreviations: A = aortic valve closure, P = pulmonic valve closure, OS = opening snap of mitral stenosis, 1st = 1st heart sound, 2nd = 2nd heart sound, 3rd = 3rd heart sound [rapid filling].



(1) Labels for semilunar valve closure indicate pulmonary circuit obstruction and should be reversed for aortic lesions.
Contributor: Witham, A. Calhoun

References: [1] Levine, S. A., and W. P. Harvey. 1949. Clinical auscultation of the heart. W. B. Saunders, Philadelphia. [2] Wells, B. G., M. B. Rappaport, and H. B. Sprague. 1949. Am. Heart J. 37:586. [3] Wells, B. G. 1952. Brit. Heart J. 14:261. [4] Ongley, P. A., H. B. Sprague, and M. B. Rappaport. 1955. N. England J. M. 253:1049. [5] Bridgen, W., and A. Leatham. 1953. Brit. Heart J. 15:55. [6] Kjellberg, S. R., E. Mannheim, O. Rude, and B. Jonsson. 1955. Diagnosis of congenital heart disease. Year Book Publishers, Chicago. [7] McKusick, V. A. 1958. Cardiovascular sound in health and disease. Williams and Wilkins, Baltimore.

79. RELATION OF NORMAL AND PATHOLOGIC HEART SOUNDS TO THE CARDIAC CYCLE. MAN




Sound	Appearance	Cause	Remarks
(A)	(B)	(C)	(D)
1	Near end of P wave	Atrial contraction	Usually observed with atrial hypertrophy.
2	End of QRS	Mainly closure of atrio-ventricular valves	Normal heart sound. May be split. May be delayed and accentuated in mitral stenosis.
3	ES	Sudden distension of aorta or pulmonary artery ("ejection sound")	Observed with dilatation of either the aorta or pulmonary artery. May be delayed 4th component of 1st sound. Pulmonic ejection sound often selectively decreases with inspiration and increases with expiration.
4	C	Uncertain, extra-cardiac.	Usually changes with posture and respiration. Often observed in individuals with history of pleuroperticardial disease.
5	A	Closure of aortic valve	Normal part of 2nd sound. Intensity varies with pressure in aorta. Delayed sound in left bundle branch block, valvular stenosis.
6	P	Closure of pulmonic valve	Normal part of 2nd sound. Intensity varies with pressure in pulmonary artery. Delayed sound in right bundle branch block, valvular stenosis, and high right ventricular stroke volume.
7	OS	Opening snap of mitral or tricuspid valve	Observed with stenosis, either valve. Time of appearance after 2nd sound varies inversely with respective atrial pressures.
8	3	Rapid distension of either ventricle	Normal heart sound. Accentuated in heart failure, rapid ventricular inflow. With rapid heart rate, sounds 4 and 3 may occur simultaneously ("summation gallop rhythm").

Contributors: (a) Witham, A. Calhoun, (b) Perloff, Joseph K.


References: [1] Leatham, A., and L. Vogelpoel. 1954. *Brit Heart J* 16:21. [2] Rappaport, E., and H. B. Sprague. 1942. *Am Heart J* 23:591. [3] Mounsey, P. 1953. *Brit Heart J* 15:135. [4] Levine, S. A., and W. P. Harvey. 1949. *Clinical auscultation of the heart*. W. B. Saunders, Philadelphia. [5] Well, B. G. 1954. *Brit Heart J* 16:261. [6] McKusick, V. A. 1958. *Cardiovascular sound in health and disease*. Williams and Wilkins, Baltimore.

80. PULSE CONFIGURATIONS: MAN


Part I: INTRACARDIAC PULSE




Normal ECG




Normal Right Atrial Configuration




Giant A wave, Right Atrial Configuration
in Right Ventricular Hypertension or Tricuspid Stenosis




Right Atrial Tracing in Fibrillation
(Inconspicuous X descent, no A wave)




Left Atrial Configuration in Mitral Stenosis



Right Atrial Tracing in Tricuspid Insufficiency
(High systolic plateau followed by deep Y descent)




Early Diastolic Collapse, Right Atrial Configuration
in Constrictive Peri- or Endocarditis, Amyloidosis




In Valvular Pulmonary Stenosis

(Continuous recording, withdrawal of catheter, pulmonary artery (PA) to right ventricle (RV))



In Infundibular Pulmonary Stenosis



Right Ventricular Pulse in Constrictive Pericarditis
(Accentuated early diastolic dip, high diastolic plateau)


Contributor: Witham, A. Calhoun

References: [1] Broadbent, W. H. 1890. *The pulse*. Cassell, London. [2] MacKenzie, J. 1902. *The study of the pulse*. Young J. Pentland, Edinburgh and London. [3] Marey, E. J. 1878. *La methode graphique*. O. Masson, Paris. [4] Wood, P. 1956. *Diseases of the heart and circulation*. Ed 2, rev. J. B. Lippincott, Philadelphia.

80. PULSE CONFIGURATIONS: MAN (Concluded)
Part II ARTERIAL RADIAL PULSE



Normal Radial Pulse



Corrigan Pulse
In Aortic Insufficiency
(Rapid upstroke and diastolic collapse)



Plateau Pulse
In Aortic Stenosis
(Slow rise and fall, low pulse pressure)




Pulsus Bisferiens
In Combined Aortic Stenosis and Insufficiency
(Double systolic peak)



Pulsus Alternans
In Myocardial Insufficiency
(Alternating weak and strong beats)



Anacrotic Pulse



Dicrotic Pulse
In Low Peripheral Resistance
(Increased dicrotic wave)

Contributor, Witham, A. Calhoun

References [1] Broadbent, W. H. 1890. The pulse. Cassell, London. [2] MacKenzie, J. 1902. The study of the pulse. Young & Pentland, Edinburgh and London. [3] Marey, E. J. 1878. La methode graphique. G. Masson, Paris. [4] Vierordt, K. 1855. Die Lehre von Arterienpuls in Gesunden und Kranken Zuständen. F. Vieweg, Brunswick. [5] Wood, P. 1956. Diseases of the heart and circulation. Ed. 2, rev. J. B. Lippincott, Philadelphia.

81. EFFECT OF PREGNANCY ON CIRCULATORY FUNCTION: MAN
Values in parentheses are ranges, estimate "b" (cf. Introduction).

		Lunar Month of Pregnancy												First Month Post-partum
		Sub-jects	Non-pregnant	Second	Third	Fourth	Fifth	Sixth	Seventh	Eighth	Ninth	Tenth		
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)	(K)	(L)	(M)	(N)	
1 Heart rate, beats/min	Pulse	68	68	78	83	84	85	87	89	85	83	77		
2 Cardiac output, L/min	Catheterization	46	5.5											
3 Cardiac index, L/min/sq m	Dye dilution	31	6.3											
4 Cardiac output, L/min/sq m	Acetylene	6												
5 Cardiac output, L/min/sq m	Direct Fick	46	3.1											
6 Cardiac output, L/min/sq m	Direct Fick	46	3.1											
7 Arterial pressure, mm Hg	Sphygmo-manometer	68	121/75 (96/58-138/98)	115/70 (104/64-121/74)										
8 Venous pressure, cm H ₂ O	Water manometer	6												
9 Femoral pressure, mm Hg	Water manometer	158	11.4 (4-18)	9.1 (2-16)	14.0 (0-28)	14.6 (6-21)	17.8 (12-24)	21.0 (16-26)	22.3 (17-27)	21.3 (15-29)	24.4 (9-5)			
10 Antecubital pressure, mm Hg	Catheterization	46	<5	7.9 (4-12)	8.1 (1-15)	7.6 (1-14)	8.9 (4-14)	10.8 (5-17)	7.2 (2-12)	7.0 (1-13)	7.6 (3-12)			
11 Pulmonary artery mean pressure, mm Hg	Catheterization	42												
12 Right ventricular pressure, mm Hg	Catheterization	46	<10/5											
13 Total peripheral resistance, dynes sec/cm ⁵	Direct Fick, catheterization, calculation	33	1200											
14 Left ventricular work, kg-m/min/sq m	Direct Fick, catheterization, calculation	33												
15 Blood volume, L	p32-labelled RBC	157	3.7 (2.8-4.6)	3.6 (2.2-4.6)	3.9 (3.1-4.7)	4.2 (3.0-5.4)	4.5 (3.6-5.4)	4.4 (2.8-6.0)	4.4 (3.4-5.4)	4.2 (1.5-6.9)	4.2 (2.6-4.8)	6		
16 Plasma volume, L	Evans' blue + Fe ⁵⁵ labelled RBC	12	4.2	3.9	4.3	4.9	5.0	5.4	5.8	5.9	6.1	5.7		
17 Plasma volume, L	p32-labelled RBC	157	2.1 (1.5-2.7)	2.3 (1.6-3.0)	2.4 (1.7-2.9)	2.7 (1.7-3.1)	2.8 (2.3-3.3)	2.9 (2.8-4.0)	2.9 (2.5-3.3)	3.1 (2.6-4.1)	3.3 (2.6-4.1)	3.3 (2.6-4.1)		
18 Evens' blue		14	2.9	2.8	2.6	2.7	3.0	3.3	3.7	4.0	3.8	2.9		
19														
20														

21	RBC volume, L	157	1.6	1.4	1.3	1.4	1.4	1.4	1.5	1.5	1.5	1.5	1.6	1.6	1.4	7
	labelled RBC	(1.2-2.0)	(0.8-2.0)	(1.0-1.7)	(1.1-1.7)	(1.0-1.7)	(1.0-1.7)	(1.0-1.7)	(1.0-1.7)	(1.0-1.7)	(1.0-1.7)	(1.0-1.7)	(1.0-1.7)	(1.0-1.7)	(1.0-1.7)	
22	Evans' blue + hematocrit	12	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	8
23		14														9
24	Blood flow Uterine, ml/100 g/min	11													15	10
25		14													15	11
26	Effective renal p-Amino-hippurate clearance	11	950												15	12
27	1.73 ag m Hepatic, Bromsulphalein clearance	15	1554												15	13
28	1.73 ag m Cerebral Nitrous oxide	105	54												15	14
29	Renal plasma flow	8	2.06												15	15
30	Renal plasma flow	58	4.56												15	16
31	Hand, ml/100 Venous occlusion	58	1.0												15	17
32	Finger, Calorimetric	16	0.20												15	18
33	Leg, ml/100 Plethymo-grammetry	58	1.87												15	19
34	Apparent viscosity of whole blood relative to water	362	4.61												15	20

1/1 4 subjects 12/1 subject. 3/2 subjects.

Contributors (a) Asaali, N. S., R. Jensen, and S. Larkin, (b) Chesley, Leon C.

References: [1] Hamilton, H. F. H. 1949. *J. Obst. Gyn. Brit. Empire* 56:546. [2] Burwell, C. S., et al. 1938. *Arch. Int. M.* 62:979. [3] Bader, R., et al. 1955. *J. Clin. Invest.* 34:1324. [4] Adams, J. Q. 1954. *Am. J. Obst.* 45:568. [5] McLennan, C. E. 1945. *Am. J. Obst.* 45:568. [6] Rose, B. J., M. E. Bader, R. A. Bader, and E. Brauwald. 1956. *Ibid.* 72:233. [7] Berlin, N. L., et al. 1953. *Surg. Gyn. Obst.* 97:173. [8] Catton, W. J., et al. 1951. *Am. J. Obst.* 61:1207. [9] Tysoe, F. W., and L. Lowenstein. 1950. *Ibid.* 60:1187. [10] Asaali, N. S., et al. 1955. *Ibid.* 66:248. [11] Metcalfe, J., et al. 1955. *J. Clin. Invest.* 34:1332. [12] Sims, E. A. H., and K. E. Krantz. 1958. *Ibid.* 37:1764. [13] Munnell, E. W., and H. C. Taylor, Jr. 1947. *Ibid.* 26:952. [14] McCall, M. L. 1953. *Am. J. Obst.* 66:1015. [15] Durt, C. C. 1950. *Edinburgh M. J.* 57:118. [16] Herbert, C. M., E. A. Darnier, and K. G. Wakim. 1956. *Am. J. Obst.* 76:742. [17] Mendlowitz, M., A. Altschek, and N. Naftchi. 1959. *Ibid.* 76:473. [18] Hamilton, H. F. H. 1950. *J. Obst. Gyn. Brit. Empire* 57:530.

82. HEMODYNAMIC EFFECTS OF HYPOXIA (LOW O₂)

Part I. HEART SIZE: MAMMALS

Animal (A)	Condition Producing Hypoxia (B)	Measurement (C)	Effect (D)	Reference (E)
1 Man	12,000 ft altitude, 3 hr daily for 4 wk	X-ray examination	No change in size.	1
2	13,000 and 23,000 ft in altitude chamber	X-ray examination	No definite relationship between altitude changes and changes in X-ray size of heart.	2
3	O ₂ mixture equivalent to that at 18,000-20,000 ft altitude	Electrocardiography	No change in size.	3
4	Gradually increasing daily exposures up to 27,000 ft over a total of 35 da	X-ray examination	Decrease in size.	4
5	Up to 30,800 ft altitude	X-ray examination	Changes in X-ray contour not consistent (increases and decreases).	5
6	18,000 ft altitude	X-ray kymography	Increase in size, systolic volume from 149 sq cm at sea level to 160 sq cm, and diastolic volume from 157 sq cm at sea level to 171 sq cm.	6
7	21,300 ft altitude	X-ray kymography	Decrease from heart size at 18,000 ft (line 6).	7
8 Flying personnel	Intermittent exposure to altitude for 1 yr or longer	X-ray examination	Increase in size; muscle hypertrophy.	8
9	Intermittent exposure to altitude for 1-10 yr	X-ray examination	Increase in size.	9
10	Intermittent exposure to altitude with supplementary O ₂	X-ray examination	No change in size.	9
11	16,400-19,700 ft altitude without supplementary O ₂	X-ray examination	Increased right heart.	9
12 Residents	10,000-15,000 ft altitude		Increase in size, transverse diameter 15% above normal, frontal area 16.3% above normal; no further enlargement with age.	10
13	15,000 ft altitude		Increase in size; transverse diameter increased in 50-65% of healthy adults; abnormal diameter of aortic pedicle and prominence of pulmonary conus.	11
14	High altitude		Increase in size	12
15 Cat	Reduced inspiratory O ₂ content		Increase in size; most drastic in slight hypoxia.	13
16 Dog	100% nitrogen for 2-3.5 min		Increase in size, dilated approximately 75 sec after start of inhalation.	14
17	10% O ₂ -gas mixture		Increase in size; diastolic volume increased independently of change in venous pressure.	15
18 Rabbit	Acute exposure following rapid ascent to 31,000-36,000 ft altitude	X-ray kymography	Increase in size; beginning at 14,710 ft, all 4 sectors of heart dilated, especially laterally.	16
19	Several hr at 32,500 ft after rapid ascent	X-ray kymography	Initially dilated heart (line 18) became smaller.	17
20	Several hr at 32,500 ft after slow ascent	X-ray kymography	Usually no change, decrease in some cases.	17

Contributor: Marbarger, John P.

References: [1] Clinton, M., Jr., G. W. Thorn, and V. D. Davenport. 1946. Bull. Johns Hopkins Hosp. 79:70. [2] Meister, W., and H. Dertlich. 1938. Luftfahrtmedizin 3:314. [3] Keys, A. J., P. Stapp, and A. Violante. 1943. Am. J. Physiol. 136:763. [4] Graybiel, A., J. L. Patterson, Jr., and C. S. Houston. 1948. Proj. NM 001-013, Rep. 6. U. S. Naval School of Aviation Medicine, Pensacola, Fla. [5] Syber, C. 1931. Arbeitsphysiologie 4:390. [6] Heddus, C. 1940. Luftfahrtmed. Abh. 3:20. [7] Schaare, U. 1938. Luftfahrtmedizin 3:104. [8] Hurtado, A., et al. 1947. J. Aviat. Med. 18:406. [9] Percador, L. 1941. Rev. clin. espan. 2:449. [10] Kerwin, A. J. 1944. Am. Heart J. 28:69. [11] Miranda, A., and A. Rotta. 1944. An. Fac. med. Lima 28:69. [12] Monge, C. 1943. Physiol. Rev. 23:166. [13] Takeuchi, K. 1925. J. Physiol., Lond. 60:208. [14] Feldman, M., Jr., et al. 1948. Am. J. Physiol. 154:391. [15] Strughold, H. 1930. Ibid. 94:641. [16] Weitz, G. A., et al. 1938. Luftfahrtmedizin 2:27. [17] Weitz, G. A., and H. Kottenhoff. 1938. Ibid. 2:33.

Part II ARTERIAL, VENOUS, AND PULSE PRESSURES MAN

Blood pressure values obtained by arterial and venous catheterization, except for lines 1 and 3.

Pressure	Condition Producing Hypoxia	Blood Pressure after Exposure for:										Reference
		0 min		5 min		10 min		15 min		20 min		
		mm Hg (C)	% Change (E)	mm Hg (D)	% Change (F)	mm Hg (G)	% Change (H)	mm Hg (I)	% Change (J)	mm Hg (K)	% Change (L)	
1	Systolic arterial	129	+4.7	135	+6.2	137	+6.2	136	+5.4	135	+4.7	1
2	Diastolic arterial	123.7	-0.6	123.0	-0.6	121.9	-1.5	114.1	-7.8	112.3	-5.2	1
3	Pulse pressure	81	+2.5	83	+2.5	82	+1.3	81	0	78	-3.7	1
4	Arterial, mean	67.7	-2.3	66.3	-2.3	65.3	-1.5	59.0	-14.9	61.4	-11.7	2
5	Pulse pressure	48	+6.3	52	+6.3	55	+6.3	53	+1.6	57	+18.7	2
6	Arterial, mean	54.6	-2.0	53.3	-2.0	54.4	-0.3	55.1	+1.0	55.9	+2.4	2
7	Arterial, mean	91.2	-2.0	89.4	-2.0	87.5	-4.0	78.2	-14.3	80.6	-11.6	2
8	Venous, mean	48	+6.2	51	+6.2	53	+10.4	57	+16.7	52	+8.3	2

Contributor: Marbarger, John P.

References: [1] Mathers, J. A. L., and R. L. Levy. 1950. Circulation, N. Y. 1:426. [2] Marbarger, J. P., et al. 1952. Proj. 21-23-019 (1). USAF School of Aviation Medicine, Randolph Field, Tex.

Part III SYSTEMIC, PULMONARY, AND RIGHT VENTRICULAR PRESSURES: MAN

Pressure	Condition Producing Hypoxia			Exposure Time min	Blood Pressure Changes, mm Hg			Reference
	O ₂ Content %	Altitude Equivalent ft	Systolic (E)		Diastolic (F)	Pulse (G)	Venous (Mean) (H)	
Simulated Altitude								
1 Systemic	14.4	10,000	-0.1	-0.4	+0.3	-1.4	-4.0	1
2 Systemic	13.4-10.4	15,000-18,000	0	-10.0	+10.0	-1.0	-1.0	2
3 Systemic	10.4	18,000	-1.8	-3.0	-0.2	+3.7	+1.0	1
4 Systemic	10.4	18,000	-6.4	-7.9	+1.3	-10.6	-10.6	1
Oxygen Mixture								
5 Systemic	10.0	19,000	+6.0	+1.0	+7.0	-1.0	-1.0	3
6 Systemic	10.0	19,000	+6.0	-3.0	+9.0	-1.0	-1.0	3
7 Systemic	10.0	19,000	-4.0	-7.0	+3.0	-5.0	-5.0	4
8 Pulmonary	10.0	19,000	+1.0	+1.0	+6.0	+1.0	+1.0	5
9 Pulmonary	10.0	19,000	+2.0	+3.0	+6.0	+6.0	+6.0	5
10 Pulmonary	10.0	19,000	+13.2	+3.0	+9.9	+9.9	+9.9	5
11 Right ventricular	10.0	19,000	+13.0	+2.8	+6.1	+6.1	+6.1	5

Contributor: Marbarger, John P.

References: [1] Marbarger, J. P., et al. 1952. Proj. 21-23-019 (1). USAF School of Aviation Medicine, Randolph Field, Tex. [2] Starr, L., and M. McMichael. 1948. J. Appl. Physiol. 1:430. [3] Mathers, J. A. L., and R. L. Levy. 1950. Circulation, N. Y. 1:426. [4] Doyle, J. T., et al. 1951. Proj. NN 001 050.01.04. U. S. Naval School of Aviation Medicine, Pensacola, Fla. [5] Motley, H. L., et al. 1947. Am. J. Physiol. 150 315.

82. HEMODYNAMIC EFFECTS OF HYPOXIA (LOW O₂)

Part I. HEART SIZE- MAMMALS

Animal (A)	Condition Producing Hypoxia (B)	Measurement (C)	Effect (D)	Reference (E)
1 Man	12,000 ft altitude, 3 hr daily for 4 wk		No change in size.	1
2	13,000 and 23,000 ft in altitude chamber	X-ray examination	No definite relationship between altitude changes and changes in X-ray size of heart.	2
3	O ₂ mixture equivalent to that at 18,000-20,000 ft altitude	Electrocardiography	No change in size.	3
4	Gradually increasing daily exposures up to 27,000 ft over a total of 35 da	X-ray examination	Decrease in size.	4
5	Up to 30,600 ft altitude	X-ray examination	Changes in X-ray contour not consistent (increases and decreases).	5
6	18,000 ft altitude	X-ray kymography	Increase in size, systolic volume from 149 sq cm at sea level to 160 sq cm, and diastolic volume from 157 sq cm at sea level to 171 sq cm.	6
7	21,300 ft altitude	X-ray kymography	Increase in size; muscle hypertrophy.	7
8 Flying Personnel	Intermittent exposure to altitude for 1 yr or longer	X-ray examination	Decrease from heart size at 18,000 ft (line 6).	8
9	Intermittent exposure to altitude for 1-10 yr	X-ray examination	Increase in size; muscle hypertrophy.	9
10	16,400-19,700 ft altitude with supplementary O ₂	X-ray examination	Increase in size.	10
11	16,400-19,700 ft altitude without supplementary O ₂	X-ray examination	No change in size.	11
12 Residents	10,000-15,000 ft altitude	X-ray examination	Increased right heart.	12
13	15,000 ft altitude		Increase in size, transverse diameter 15% above normal, frontal area 16.3% above normal; no further enlargement with age.	13
14	High altitude		Increase in size; transverse diameter increased in 50-65% of healthy adults, abnormal diameter of aortic pedicle and prominence of pulmonary conus.	14
15 Cat	Reduced inspiratory O ₂ content		Increase in size.	15
16 Dog	100% nitrogen for 2-3.5 min		Increase in size; most drastic in slight hypoxia.	16
17	10% O ₂ -gas mixture		Increase in size; dilated approximately 75 sec after start of inhalation.	17
18 Rabbit	Acute exposure following rapid ascent to 31,000-36,000 ft altitude	X-ray kymography	Increase in size, diastolic volume increased independently of change in venous pressure.	18
19	Several hr at 32,500 ft altitude	X-ray kymography	Increase in size, beginning at 14,710 ft, all 4 sectors of heart dilated, especially laterally.	19
20	Several hr at 32,500 ft after rapid ascent	X-ray kymography	Initially dilated heart (line 18) became smaller.	20
Contributor: Marbarger, John P.				21

References. [1] Clinton, M. J., G. W. Thorn, and V. D. Davenport. 1946. Bull. Johns Hopkins Hosp. 79:70. [2] Meister, W., and H. Derlich. 1938. Luftfahrtmedizin 3:314. [3] Krys, A., J. P. Stapp, and A. Violante. 1943. Am. J. Physiol. 138:763. [4] Graybiel, A., J. L. Patterson, Jr., and C. S. Houston. 1948. Proc. NM 001-013, Rep. 6. U. S. Naval School of Aviation Medicine, Pensacola, Fla. [5] Spycher, C. 1931. Arbeitsphysiologie 4:390. [6] Heedus, C. 1940. Luftfahrtmedizin 3:20. [7] Schaare, U. 1938. Luftfahrtmedizin 3:104. [8] Hurtado, A., et al. 1947. J. Aviat. Med. 18:406. [9] Pescador, L. 1941. Rev. clin. españ. 2:445. [10] Kerwin, A. J. 1944. Am. Heart J. 28:69. [11] Miranda, A., and A. Rotta. 1944. Am. Fac. med. J. Physiol. 154:391. [12] Monger, C. 1943. Physiol. Rev. 23:166. [13] Takenochi, K. 1925. J. Physiol., Lond. 60:208. [14] Feldman, M. J., et al. 1948. Am. J. Physiol. 154:391. [15] Straghold, H. 1930. Ibid. 94:641. [16] Weltz, G. A., et al. 1938. Luftfahrtmedizin 2:27. [17] Weltz, G. A., and H. Kottenhoff. 1938. Ibid. 2:33.

Part VII CIRCULATION OF PERIPHERAL BLOOD- MAN, DOG

Body Part or Region (A)	Condition Producing Hypoxia (B)	Man	Effect (C)	Reference (D)
1 Finger	6.4% O ₂ -gas mixture		Increase at distal, decrease later.	1
2	7% O ₂ -gas mixture		Effect varied.	2
3	10% O ₂ -gas mixture		30% decrease.	3
4 Hand	10% O ₂ -gas mixture for 10-21 min		Decrease.	4
5 Forearm and leg	10% O ₂ -gas mixture for 10-21 min		Increase.	4
6 Ear	6.4% O ₂ -gas mixture		Increase in peripheral blood, independent of blood pressure.	1
7 Scalp region	7% O ₂ -gas mixture		Increase in peripheral blood with increase in temperature of ear.	2
8 Capillaries	Altitude of 2200 ft		Decrease in peripheral blood. Mean resistance increased 25 mm Hg; return to sea level values during continued stay at 5200 ft.	3
9	High altitudes (acclimatized)		Dilated.	6
10 Splanchnic region	Esclatinal		Decreased in intestinal vessels.	7
11	Unspecified		Increased in hepatic artery.	3
12	Unspecified		Circulation changed locally in proportion to arterial blood pressure, especially in splanchnic region.	3
13 Skin	Breathing "normal" O ₂ from demand system at 35,000 ft	Dog	Decrease in peripheral blood indicated by significantly reduced temperature.	9
14 Skin and muscle	O ₂ -gas mixture		Initially strong increase.	10
15 Internal carotid artery	O ₂ -N ₂ mixture		Transient increase.	11
16 Kidney	Unspecified		Decrease in peripheral blood, independent of blood pressure.	12

Contributor: Marbarger, John P.

References: [1] Matthes, K., and K. Schlaudraff. 1944. *Luftfahrtmedizin* 8:161. [2] Matthes, K., and R. Falk. 1941. *Verh. deut. Ges. Kreislauforsch.* 14:109. [3] Steele, J. M. 1942. *Nat. Res. Coun. Proc. OSMemo* 130 (Rep. 7). [4] Abramson, D. L., H. Landt, and J. E. Benjamins. 1943. *Arch. Int. M.* 71:593. [5] Schmid, A. 1949. *Helvet. physiol. pharm. acta* 7:167. [6] Monge, C. 1943. *Physiol. Rev.* 23:166. [7] Rein, H. 1943. *Verh. deut. Ges. Kreislauforsch.* 24:680. [8] Jensen, K., W. Kratz, and W. Schoedel. 1940. *Luftfahrtmedizin* 5:40. [9] Lippin, J. L., and W. V. Whitcomb. 1931. *J. Aviat. M.* 22:278. [10] Noell, W., and M. Schneider. 1941. *Luftfahrtmedizin* 5:234. [11] Dendinger, T., E. Opitz, and W. Schoedel. 1938. *Ibid.* 3:46. [12] Kressberg, W., L. Prokop, and T. Schiffer. 1949. *Fliegern Arch.* 25:167.

82. HEMODYNAMIC EFFECTS OF HYPOXIA (LOW O₂) (Continued)

Part IV ARTERIAL BLOOD PRESSURE DURING PROGRESSIVE ADAPTATION TO ALTITUDE: MAN

Values in parentheses are ranges, estimate "b" (cf. Introduction).

Altitude ft (A)	Systolic Blood Pressure			Diastolic Blood Pressure			Mean Blood Pressure		
	Value mm Hg (B)	Change mm Hg (C)	% (D)	Value mm Hg (E)	Change mm Hg (F)	% (G)	Value mm Hg (H)	Change mm Hg (I)	% (J)
1 Sea level (control)	112.50(92.76-132.24)			69.16(58.34-79.98)			90.83(76.43-105.23)		
2 12,140	118.33(98.53-138.13)	35.83	+5.2	84.16(73.28-95.04)	+15.0	+21.7	101.25(88.75-113.75)	+10.42	+11
3 13,120	125.00(98.60-151.40)	+12.5	+11.1	88.33(77.19-99.47)	+19.17	+27.2	107.08(87.58-126.58)	+16.25	+17
4 19,360	130.00(105.40-154.60)	+17.5	+15.6	93.33(74.43-112.23)	+24.17	+35.0	111.66(90.46-132.86)	+20.83	+22

Contributor: Marbarger, John P.

Reference: Hartman, H., G. Hepp, and U. C. Luft. 1941. Luftfahrtmedizin 6:1. (Nanga Parbat expedition).

PART V. PULSE RATE CHANGE, IN PERCENT OF GROUND LEVEL VALUE, DURING PROGRESSIVE EXPOSURE: MAN

Pulse rate index at ground level = 100.

Simulated Altitude ft (A)	O ₂ Content		Pulse Rate Change, in % of Ground Level Value, after Exposure for:							
	(B)	%	10 min (C)	20 min (D)	30 min (E)	40 min (F)	50 min (G)	60 min (H)	70 min (I)	80 min (J)
1 12,000	13.3	11.4	113	115	113	106	104	99		
2 16,000	11.4	10.5	111	110	103	103	101	105		
3 18,000	10.5	9.6	107	109	108	111	108	104		
4 20,000	9.6	8.8	124	112	117	107				
5 22,000	8.8		131	126	124					

Contributor: Marbarger, John P.

Reference: Bahn, H., and A. B. Ode. 1947. Am. J. Physiol. 150:202.

Part VI PULSE RATE CHANGE DURING PROGRESSIVE EXPOSURE: MAN

Condition Producing Hypoxia (A)	O ₂ Content		% Change in Pulse Rate after Exposure for:							
	(B)	%	5 min (C)	10 min (D)	15 min (E)	20 min (F)	30 min (G)	40 min (H)	50 min (I)	60 min (J)
1 18,000 ft simulated altitude	10.4	10.4	25	30	30	28				
2 Gas mixture	10.0	10.5	17	20	21	20				
3 Gas mixture	10.5	10.5	14	31	31	32	33			

Contributor: Marbarger, John P.

References: [1] Marbarger, J. P., et al. 1952. Proj. 21-23-019 (1). USAF School of Aviation Medicine, Randolph Field, Tex. [2] Mathers, J. A. L., and R. L. Levy. 1950. Circulation, N. Y. 1 426. [3] Schaefer, K. E., and H. J. Alois. 1951. Bur. Med. Proj. NM 002 015.03.02. U. S. Naval Marine Base, New London, Conn.

Part X AMPLITUDES OF THE ELECTROCARDIOGRAM, MAN

For additional information on the ECG, see pages 137-143.

Wave (A)	Condition Producing Hypoxia (B)	Effect (C)	Reference (D)
1 P	O ₂ gas mixture (arterial O ₂ saturation = 95-70%) ¹	• Inversion for P ₁ and P ₂	1
2	Short exposures at simulated altitude of 25,000 and 35,000 ft	• Slight increase from 0.05-0.15 mv.	2
3	Simulated altitude of 19,685 ft	• Slight increase from 0.05-0.15 mm (sitting).	3
4	Simulated altitude of 23,000-26,250 ft	• Slight increase.	4
5	1% O ₂ gas mixture	• Slight decrease without consistency.	5
6	Continuous exposure to low O ₂	• Slight decrease.	6
7 QRS complex	Short exposures at simulated altitude of 25,000 and 35,000 ft	• R-potential in some instances (not consistently).	7
8	Simulated altitude of 19,685 ft	• Slight increase from 0.05-0.15 mm (sitting).	8
9	Simulated altitude of 23,000-26,250 ft	• Slight increase.	9
10	1% O ₂ gas mixture	• Slight decrease without consistency.	10
11	Up to 20,000 ft altitude	• Slight decrease.	11
12 T	O ₂ gas mixture (arterial O ₂ saturation = 95-70%) ¹	• T wave flattened	12
13	Short exposures at simulated altitude of 25,000 and 35,000 ft	• T wave related to degree of decrease in O ₂ saturation. T wave may become biphasic, or with progressive anoxemia	13
14	Simulated altitude of 35,000 ft ²	• Voltage decrease. Onset of significant changes: a 30-40 sec. 15,000 ft = 10-30 sec.	14
15	Simulated altitude of 19,685 ft	• Height before hypoxia = 0.675 mv; mean	15
16	Approximately 24,750 ft altitude (alveolar O ₂ pressure = 50-60 mm Hg)	• Height during hypoxia = 0.151 mv (1-23%). Wave is biphasic.	16
17	Simulated altitude of 23,000-26,250 ft	• Difference from sea level = -0.83 (sitting).	17
18	1% O ₂ gas mixture	• Onset of wave began at alveolar O ₂ pressure of 50-60	18
19	Up to 20,000 ft altitude	• Flattening proportional to increase in heart	19
20	Flights at 15,000 and 26,000 ft ⁴	• Absolute loss of height greater in initially higher	20
21	Continuous exposure to low O ₂	• Is.	21
22 ST segment	O ₂ gas mixture (arterial O ₂ saturation = 95-70%) ¹	• Quickly reduced to approximately half the previous potential. Flattening increased with increased heart rate and increased hypoxia.	22
23	Simulated altitude of 19,685 ft	• Lowered amplitude.	23
24	Simulated altitude of 23,000-26,250 ft	• Occasionally very slight depression.	24
25	1% O ₂ gas mixture	• Diminished in all 4 leads	25
26	Simulated altitude of 16,250 and 20,000 ft	• Inverted wave.	26
27	Any instance of sinus arrhythmias disappeared when heart rate increased by more than 23 beats per minute. 1/2 Occasionally extrasystoles. 1/4 In many cases, U wave appeared.	• Total ST deviation in Leads I-IV increased slightly (up to 25 mm) with increased anoxemia. Most commonly, the incidence of Lead II depression (0.5 mm) increased with increasing hypoxemia.	27
28	Shifting pacemaker, and premature atrial and ventricular beats.	• ST above isoelectric line.	28
29	Contributor: Marbarger, John P.	• ST depressed with progressive hypoxia.	29
30	References: [1] Penney, R., and C. D. Thomas. 1950. Circulation, N. Y. 1415. [2] King, B. G., and M. Henson. 1947. J. Aviat. Med. 18 3. [3] Doetsch, W. 1938. Luftfahrtmedizin 2 354. [4] Tittel, S. 1940. Ibid. 4:322. [5] Müller, H. 1940. Dissertation, Erlangen. [6] Monge, C. 1943. Physiol. Rev. 23:166. [7] Benson, O. O. 1940. J. Aviat. Med. 11:167. [8] Hemingway, A. 1944. Ibid. 15:286. [9] Cohn, E., and F. Palmer. 1944. Physiolog. Arch. 246:387. [10] White, M. S. 1940. J. Aviat. Med. 11:166. [11] Rühl, A. 1948. In P. H. Rehn, ed. Physiologie. Part I, p. 201.	• Lowered amplitude.	30

82. HEMODYNAMIC EFFECTS OF HYPOXIA (LOW O₂) (Continued)
Part VIII. PULMONARY CIRCULATION: MAN, DOG

Specification	Pulmonary Arterial Pressure				Pulmonary Capillary Pressure (E)	Pulmonary Pressure Gradient (F)	Pulmonary Resistance (G)	Pulmonary Blood Volume (H)	Reference
	Systolic (B)	Diastolic (C)	Mean (D)						
1 Control (20.96% O ₂)	16 mm Hg	6 mm Hg	10 mm Hg	Man	5 mm Hg	5 mm Hg	66 dynes sec/cm ⁵		(1)
2 Low O ₂ (10%) ¹	23 mm Hg	9 mm Hg	15 mm Hg		5 mm Hg	10 mm Hg	119 dynes sec/cm ⁵	660 ml/sq m	
3 Change	+44%	+50%	+50%		0	+100%	+80%	670 ml/sq m	1
4 Control (20.96% O ₂)	21.9 mm Hg	6 mm Hg	13.1 mm Hg				1216 dynes sec cm ⁻⁵		
5 Low O ₂ (10%) ¹	35.1 mm Hg	13 mm Hg	23.0 mm Hg				1407 dynes sec cm ⁻⁵		2
6 Change	+60.3%	+116.7%	+75.6%				+15.7%		
7 Control (20.96% O ₂)	29 mm Hg	12 mm Hg		Dog					
8 Low O ₂ (10%) ¹	26 mm Hg	12 mm Hg							
9 Change	-10.3%	0							3

1/1 O₂-N₂ mixture.

Contributor: Marbarger, John P

References: [1] Doyle, J. T., J. S. Wilson, and R. B. Warren. 1951. Proj. NM 001 050.01.04. U. S. Naval School of Aviation Medicine, Pensacola, Fla.
[2] Moilety, M. L., et al. 1947. Am. J. Physiol. 150.315. [3] Beard, E. F., et al. 1951. Proj. 21-26-003 (1). USAF School of Aviation Medicine, Randolph Field, Tex

Part IX: INTERVALS OF THE ELECTROCARDIOGRAM: MAN
For additional information on the ECG, see pages 137-141.

Interval	Condition Producing Hypoxia	Effect	Reference
1 PR	Sudden elevation to 23,000 and 26,250 ft	Interval unaltered.	(D)
2 PQ	Continuous exposure to altitude	Interval shortened.	
3 PT	7% O ₂ gas mixture	Uniform shortening of interval, from 0.01-0.05 sec.	1
4 QRS complex	Unspecified	At first, interval increased with increasing hypoxia.	2
5 QT	19,685 ft simulated altitude	later increased with increasing heart rate.	3
6	Unspecified	Mean difference from sea level = +0.03 sec.	4
7	Unspecified	Increase to more than 0.1 sec.	
8	Unspecified altitude	At first, interval increased with increasing heart rate,	5
9	Flights at 15,000 and 20,000 ft	later increased with increasing hypoxia.	6
Contributor: Marbarger, John P.		Interval relatively unaltered during adaptation to altitude.	4
		Interval prolonged.	

References [1] Tittel, S. 1940. Luftfahrtmedizin 4.328. [2] Monger, C. 1943. Physiol. Rev. 23:166. [3] Müller, H. 1940. Dissertation. Erlangen. [4] Van Tavel, F. 1943. Helvet. physiol. pharm. acta, Suppl. 1:65. [5] Doetsch, W. 1938. Luftfahrtmedizin 2:354. [6] Schütz, E. 1938. Ibid 2:192 [7] White, M. S. 1940. J. Aviat M. 11:166.

Part XII CARDIAC WORK AND BLOOD PRESSURE AT REDUCED ARTERIAL O₂ SATURATION. MAN

Normal subjects 12 males and 4 females, 19-33 years of age. Anoxemia was induced with oxygen-nitrogen mixtures administered through an anesthesia machine to give the desired levels of arterial O₂ saturation, as measured with the Mullikan oximeter. Blood pressure was determined by the standard auscultatory method. The ballistocardiograph used to derive stroke volume, cardiac index, left ventricular work, and maximum cardiac force, is of the Starr high-frequency, horizontal type. Values in parentheses are ranges, estimate "c" (cf Introduction).

Variable	Arterial O ₂ Saturation																			
	96%				95%				80%				75%				70%			
	Room Air Control	No of Subjects	Mean Value	% Change	Room Air Control	No of Subjects	Mean Value	% Change	Room Air Control	No of Subjects	Mean Value	% Change	Room Air Control	No of Subjects	Mean Value	% Change	Room Air Control	No of Subjects	Mean Value	% Change
1 Pulse rate, beats/min	68.9	3	62.6	+19.9	10	87.2	+26.7	15	91.7	+33.0	13	94.3	+36.8							
2 Stroke volume, ml/beat	92.3	3	90.8	-1.6	10	93.1	+0.9	15	92.1	+0.2	13	94.3	+2.2							
3 Cardiac index, l/min/m ²	3.48	3	4.14	+18.8	10	4.45	+27.8	15	4.71	+35.2	13	4.97	+42.7							
4 Left ventricular work, kg-m/min	8.12	2	10.43	+28.5	9	10.97	+5.1	14	11.32	+37.4	12	11.26	+31.0							
5 Maximum cardiac force/beat	33.1	3	36.5	+10.1	10	36.7	+1.0	15	36.7	+16.8	13	43.2	+30.7							
6 Maximum cardiac force/min	2275	3	3010	+32.5	10	3230	+42.2	15	3710	+63.3	13	4165	+83.3							
7 Pulse pressure, mm Hg	44.6	2	46.2	+3.5	8	49.6	+11.2	14	57.3	+28.5	12	61.2	+37.2							
8 Mean blood pressure, mm Hg	99.4	2	108.9	+8.2	8	101.8	+2.2	14	100.0	+0.7	12	103.9	+4.5							

1/1 MCF/beat = $I + J + J_2$, where I_2 and J_2 are amplitudes of the I and J waves in a large complex, and where I and J are amplitudes of the I and J waves in a small complex (see Appendix II, page 360). 2/2 MCF/min = MCF/beat x pulse rate.

Contributor: Penney, Raymond

Reference: Scarborough, W. R., R. Penney, C. B. Thomas, B. M. Baker, Jr., and R. E. Mason. 1951. Circulation, N. Y. 4 190.

82. HEMODYNAMIC EFFECTS OF HYPOXIA (LOW O₂) (Concluded)

Part XL THE ELECTROCARDIOGRAM, HEART RATE, AND BLOOD PRESSURE AT REDUCED ARTERIAL O₂ SATURATION: MAN

Normal subjects: 73 males and 3 females, 22-28 years of age. Anoxemia was induced with oxygen-nitrogen mixtures administered through an anesthesia machine to give the desired levels of arterial O₂ saturation, as measured with the Millikan oximeter. Blood pressure was determined by the standard auscultatory method. Before inhaling gas mixture, subject rested on a bed until blood pressure and pulse rate became stable. "Absolute change" is from the value for the control on air. Values in parentheses are ranges, estimate "c" (cf. Introduction).

Variable	80%			Arterial O ₂ Saturation			70%
	(A)	No of Observations (B)	Absolute Change (C)	No. of Observations (D)	Absolute Change (E)	No. of Observations (F)	
T wave height, mm							
1 Lead I		64	-0.68(-2 to +0.5)	67	-0.77(-2.0 to +0.9)	28	-1.19(-2.7 to -0.1)
2 Lead II		64	-1.04(-3.7 to +0.2)	67	-1.31(-4.4 to +1.0)	28	-1.99(-4.7 to -0.5)
3 Lead III		64	-0.37(-1.5 to +0.8)	67	-0.66(-2.6 to +0.9)	28	-1.17(-3.2 to 0)
4 Lead IV		64	-0.75(-2.9 to +3.3)	67	-0.96(-4.3 to +1.9)	28	-1.15(-5.0 to +1.3)
5 RS-T deviation, mm ¹		64	0.60(0-1.5)	67	0.73(0-2)	28	0.86(0-2.5)
6 Heart rate, beats/min		69	+14.0(+1 to +33)	73	+18.3(0 to +39)	30	+23.4(+6 to +39)
Blood pressure, mm Hg							
7 Systolic		67	+4.1(-11 to +23)	70	+4.1(-13 to +22)	28	+6.3(-4 to +16)
8 Diastolic		67	+0.9(-12 to +13)	70	+0.5(-12 to +12)	28	-0.1(-12 to +10)
1/1 Indicates total for all 4 leads regardless of direction of change from control electrocardiogram.							

Contributor: Penneys, Raymond

Reference: Penneys, R., and C. B. Thomas. 1950. Circulation, N. Y. 1:415.

30	Cat, dog 1 dog	12 cats.	Rubber bag	100%	1-4 min	Mean pressure = -2.2	During anesthesia (chloral)	12
31	Rabbit	4			1-1 min	Mean pressure = -1	During anesthesia (urethane)	13

/1/ Mask check. /2/ Errors of method increased by high O₂ pressure; decrease of pulse rate noted. /3/ Arterial O₂ saturation = 95%. /4/ Approximate figure from graphs /5/ Due mainly to factors other than flow.

Contributor: Shepherd, Roy J.

References [1] Dripps, R. D., and J. H. Comroe 1947. *Am. J. Physiol.* 149:277. [2] Dauterhand, L., and J. S. Italdane. 1921. *J. Physiol.* Lond 55:296. [3] Whitehorn, W. V., A. Edelmann, and F. A. Hillecock. 1946. *Am. J. Physiol.* 146:61. [4] Miller, A. T., H. L. Perdue, E. L. Teague, and J. S. Ferebee. 1952. *J. Appl. Physiol.* 5:165. [5] Keys, A., J. P. Stapp, and A. Violante. 1943. *Am. J. Physiol.* 138:763. [6] Benedict, F. G., and H. L. Higgins. 1921. *Ibid.* 26:1. [7] Chiodi, H., J. C. Pasciolo, J. R. Sudres, and A. C. Taquini. 1948. *J. Appl. Physiol.* 11:148. [8] Alveryd, A., and S. Brody. 1948. *Acta physiol. scand.* 25:140. [9] Comroe, J. H., R. D. Dripps, P. R. Dunke, and M. Deming. 1945. *J. Am. Med. Ass.* 128:10. [10] Becker-Freyung, H., and H. G. Clamann. 1939. *Klin. Wochr.* 18:1382. [11] Richards, D. W., Jr., and A. L. Barach. 1934. *Q. J. Med.* 27:147. [12] Von Euler, U. S., and G. Liljestrang. 1942. *Acta physiol. scand.* 4:34. [13] Lieberman, E. 1943. *Ibid.* 6:335. [14] Kety, S. S., and C. F. Schmidt. 1948. *J. Clin. Invest.* 27:484. [15] Ott, A. B., H. Rahn, M. Dronman, L. J. Mullins, and W. O. Fenn. 1946. *Ibid.* 25:415. [16] Grollman, A. 1930. [17] Rosenthal, C. N. 1939. *Arch. Neur. Psychiat.* 42:197. [18] Lennox, W. G., and E. L. Gibbs. 1932. *J. Clin. Invest.* 11:155. [19] Lamberstein, C. J., R. H. Kough, D. V. Cooper, G. L. Emmet, H. H. Lorschke, and C. F. Schmidt. 1952. *J. Appl. Physiol.* 5:471. 487, 803. [20] Wolff, H. G., and W. G. Lennox. 1930. *Arch. Neur. Psychiat.* 42:197. [21] Dunke, P. R., and C. F. Schmidt. 1943. *Am. J. Physiol.* 138:421. [22] Von Euler, U. S., and G. Liljestrang. 1946. *Acta physiol. scand.* 12:301.

83. HEMODYNAMIC EFFECTS OF HYPEROXIA (HIGH O₂): MAMMALS

All controls breathing air. Values in parentheses are ranges, estimate "c" (cf. Introduction).

Animal (A)	No. of Subjects (B)	Method of Administration (C)	O ₂ Concentration (D)	Exposure Time (E)	Pulse Rate	Observed Change (% of Resting Value) (F)	Remarks (G)	Reference (H)
1 Man	33	Face mask and demand valve	100%	1-2 min	-3.7 (-12.5 to +4.6)			1
2	2	Douglas bag	100%	5 min	-7.0			2
3	16	Mask and demand valve	100%	5 min	-12.2			3
4	33	Face mask and demand valve	100%	6-8 min	-5.5 (-14.4 to +4.8)			1
5	26		100%	10-20 min	No change		During exercise (4 min at 7 mph)	4
6	7	Anesthesia mask	100%	15-20 min	-10 (-17.5 to -5)			5
7	6	Spirometer	90%	15-30 min	-6.7 (-10.0 to -3.3)			6
8	10		100%	15-30 min	-15.2 (-21 to -10)		Anemic patients	7
9	15	Spirometer	100%	20-30 min	-9.1 (-17.4 to -4.3)			8
10	16	Mask and demand valve	100%	60 min	-13.9			3
11	28	Mask and demand valve	100%	24 hr	No change			9
12	2	Decompression chamber	90%	65 hr	+15			10
13	2	Oxygen chamber	45%	168 hr	(-13 to -12)		CO ₂ rose to 0.5% During anesthesia (chloral)	11
14	12 cats, 1 dog	Rubber bag	100%	1-4 min	-3.3			12
15	Rabbit		100%	1-1 min	-0.6		During anesthesia (urethane)	13
16 Man	16	Oxygen mask and demand valve	100%	5 min	Cardiac Output -13.0		Ballistocardiogram	3
17	33	Mask and demand valve	100%	6-8 min	-8		Ballistocardiogram	1
18	7	Anesthesia mask	100%	15-20 min	-10 (-20 to +19)		Roentgenkymograph	5
19	6		85-100%	15-30 min	-4.9		Ballistocardiogram	14
20	5	Spirometer	100%	30 min	-8.5		Ballistocardiogram	15
21	1	Douglas bag	100%	45 min	Probably no decrease		Acetylene method	16
22	16	Oxygen mask and demand valve	100%	60 min	-19.4		Ballistocardiogram	3
23	2	Oxygen chamber	40-50%	1 wk	No change		CO ₂ rose to 0.5%	11
24 Man	16	Mask and demand valve	100%	5 min	Systemic Blood Pressure			3
25	7	Anesthesia mask	100%	15-20 min	Systolic pressure = -2.3 Diastolic pressure = +2.1 Increase in systolic and diastolic pressure			5
26	6	Oxygen chamber	85-100%	15-30 min	Systolic pressure = +9.3 Diastolic pressure = +16.7			14
27	15	Spirometer	100%	20-30 min	Systolic pressure = +1.6 (-2.5 to +5.7) Diastolic pressure = +5.3 (-1.5 to +10.9)			8
28	16	Mask and demand valve	100%	60 min	Systolic pressure = +2.7 Diastolic pressure = +6.9			3
29	28	Mask and demand valve	100%	24 hr	No significant change			9

20		4	BLB mask	5% in air	3	Systolic pressure = $+4.2(-2.5 \text{ to } +10.0)$ Diastolic pressure = $+5.0(+1.2 \text{ to } +7.5)$ Mean pressure = $+4.5(+1.2 \text{ to } +4.8)$	Direct recordings from aorta in subjects with patent ductus arteriosus.	2
21		8	Mouthpiece and demand valve	2-16% in air	8-10	Mean pressure = -1		
22				4-31%	8-10			
23				5-48%	8-10			
24		2		7% in air	15-30	Systolic pressure = $+10.7$ Diastolic pressure = $+14.6$		4
25		6		5% in air	15-30	Systolic pressure = $+10.7$ Diastolic pressure = $+14.6$		3
26	Cat		Gas bag	7-14.9% in O ₂	0-1	Systolic pressure = $+7.8$ Diastolic pressure = $+8.8$ -50	6 min after injection of ergotamine	3
27	Dog	7	Rubber bag and respirator	8-15% in O ₂	<20	Normal = $+12 \text{ to } +14$ With curare = $-50(-65 \text{ to } -33)$		9
28	Man	16	Rebreathing	0-8% in 30% O ₂	17-32	Blood flow, 1% CO ₂ = -9.8 2% CO ₂ = -10.0 3% CO ₂ = -16.7 4% CO ₂ = -19.3 5% CO ₂ = -31.5 6% CO ₂ = -39.8	Hand blood flow.	11
29						Primary contractions = -9.3 Secondary dilatations = $+16.3$ After ceasing to breathe CO ₂ = 429.4	Mesenteric vessels (superior or mesenteric, common iliac), during urethane anesthesia. Concentration of 0.5-1.0% CO ₂ adequate to produce changes.	12
30		30		0.2-10% in air	2-5			
31								
32								
33								
34	Cat							
35	Dog			4-10%	2-4	Decreased flow in both innervated and denervated leg, but effect reversed by prolonged anesthesia (6 hr). Increased flow; effect unaltered by denervation. Quite marked increase.	Femoral artery, brachial artery, during urethane anesthesia. Superior mesenteric artery. Aortic arch.	13
36								
37								

1/1 Boothby, Lovelace, Bulbunan. 1/2/ Approximate figure from graphs. 1/3/ Cardiac output may be initially increased; postulated later failure of venous return. 1/4/ Functional venesection.

Contributors. (a) Shephard, Roy J., (b) Spencer, Joseph N.

References: [1] Dripps, R. D., and J. H. Comroe. 1947. *Am. J. Physiol.* 149 43. [2] Shephard, R. J. 1954. *Brit. Heart J.* 16:491. [3] Kelly, S. S., and C. F. Schmidt. 1948. *J. Clin. Invest.* 27:464. [4] Lambertsen, C. J., R. H. Kough, D. V. Cooper, G. L. Emmet, H. H. Loeschcke, and C. F. Schmidt. 1952. *J. Appl. Physiol.* 5 471, 487, 803. [5] Bidtek, G., R. Johansson, and H. Schmidt. 1956. *Acta physiol. scand.* 37:71. [6] Grollman, A. 1930. *Am. J. Physiol.* 94:287. [7] Armussen, E. 1943. *Acta physiol. scand.* 6:176. [8] Keys, A., J. P. Sapp, and A. Violante. 1943. *Am. J. Physiol.* 138:763. [9] Lippert, A. 1946. *Acta physiol. scand.* 15:198. [10] Page, I. H., and F. O. O'Rourke. 1931. *Circulation*, N. Y. 3:801. [11] Schmidt, C. F., and D. Truesdell. 1922. *Am. J. Physiol.* 63 155. [12] Fletcher, A., I. Sibul, and V. Ponomarev. 1932. *Flugers Arch.* 230:814. [13] Bernthal, T. G. 1930. *Am. J. Physiol.* 94:446.

84. HEMODYNAMIC EFFECTS OF HYPERCAPNIA (HIGH CO₂)
 Part I CHANGE IN PULSE RATE, CARDIAC OUTPUT, BLOOD PRESSURE, AND BLOOD FLOW: MAMMALS
 Values in parentheses are ranges, estimate "c" (cf. Introduction).

Animal	No. of Subjects	Method of Administration	CO ₂ Concentration	Exposure Time min	Pulse Rate (% of Resting Value)	Observed Change (% of Resting Value)	Remarks	Reference
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	
1 Man	29	Regulator, reservoir, and mask	10.4% in O ₂	2-6	Pulse Rate +19.5(-5 to +45)			
2	42	Regulator, reservoir, and mask	7.6% in O ₂	2-8	+20.9(-12.5 to +85)			1
3	18	BLB mask	5% in air	3-4	+4.8			1
4	2		7% in air	5-10	+3		16 cases of congenital heart disease, 2 normal subjects. Effect observed only in recumbent position.	2
5	6		5% in air	5-10	+9			
6	8	Douglas bag	2.16% in air	8-10	+6.4			
7			4.31% in air	8-10	+0.6			3
8			5.48% in air	8-10	+2.1			3
9 Hedgehog	8	Gas chamber	3% in O ₂	70-240	No change			4
10			5% in O ₂	70-240	Hibernating = +202			
11			9.5% in O ₂	70-240	Non-hibernating = +232			5
12 Man	16	BLB mask	5% in air	Cardiac Output	Hibernating = +25			
13					Non-hibernating = +14			
14	4	Douglas bag	0-9% in air	4-40	+14(-8.9 to +72.8)		Subjects with congenital heart disease, in recumbent position. Modified direct Fick method.	2
15	2	Cylinder mixtures	2.5-7.5% in air	10-15	No effect on circulation rate at 0-6% CO ₂ ; moderate increase with higher concentrations.		Acetylene method.	6
16	6	Gasometer and anesthesia mask	2 6% in 10% O ₂	15-20	Recumbent, 6.5% CO ₂ = +66, +60		Acetylene method.	7
17	2				Upright = -17.5, -20			
18 Man	6		7% in air	15-30	Legs "empty" = +44			
19	34	Regulator, reservoir, and mask	7.6% in O ₂	15-30	Legs "full" = +20			
			10.4% in O ₂	+1.6	+14(+3 to +45)			
				+8.5	Diastolic volume = -2.5(-6 to 0)		Röntgenokymograph.	8
				Systemic Blood Pressure				
				2-8	Systolic pressure = +25.6(+10 to +53)			3
					Diastolic pressure = +28(+5 to +59)			
				2-8	Systolic pressure = +28(+6 to +43)			1
					Diastolic pressure = +31(+4 to +77)			

Part III CHANGE IN PULSE RATE, BLOOD PRESSURE, AND THE ELECTROCARDIOGRAM: DOO

Cardiovascular effects of breathing high CO₂ concentrations in the environmental air vary with the kind of anesthetic, depth of anesthesia, the state of the nervous system, experimental procedure, duration of exposure, oxygenation, and, to a lesser degree, with the animal species. Symbols: (+) = increase or elevation, (-) = decrease, (0) = no change

CO ₂ Concentration (A)	Exposure Time min (B)	Preparation or Method (C)	Anesthetic (D)	Arterial pHi (E)	Pulse Rate (F)	Blood Pressure (G)	QRS Interval (H)	PR Interval (I)	ST Segment (J)	Wave (K)	Remarks (L)	Reference (M)
7.5 0%		Heart-lung			Marked decrease ¹						Ventricular dilatation ² and decrease in coronary blood flow ³	1
2	15		Nembutal		-4%	0					Decrease in pulmonary resistance	2
3	9	Open chest	Pentobarbital		-1%	-14%	0	0	Starting	+	25% increase in systolic excursion; 15% decrease in contraction force.	3
4	20		Pentobarbital		0	+21%						4
5	20	Artificial respiration	Pentobarbital		+	+6%						4
6	10	Heart-lung	Nembutal and succinylcholine	7.2	-						Cardiac dilatation; decrease in cardiac output and systolic excursion	5
7	13	Open chest	Pentobarbital		+15%	-18%	0	0	Starting	+	21% decrease in systolic excursion; 24% decrease in contraction force; increased skeletal muscle tonus; increased heart rate and irregularities on return to room air.	3
8	45		Pemopal	7.24	0		0	0	0	+	100% increase in cerebrospinal fluid pressure.	6, 7
9	37.5%	Decerebrate			-		0	+	0	+	Marked decrease in atriocentricular conduction.	8
10	15 0%	Heart-lung	Nembutal		-						Cardiac dilatation; marked reduction in cardiac output, rise in central venous pressure; sympathoadrenal system; unaccounted effect of CO ₂ on heart.	5
11	10	Artificial respiration	Nembutal and succinylcholine	7.1	+	0					Vagotomy resulted in 5-10% increase in blood pressure response to CO ₂	5

1/1 True also of cat.

84. HEMODYNAMIC EFFECTS OF HYPERCAPNIA (HIGH CO₂) (Continued)

Part II. CHANGE IN PULSE RATE AND BLOOD PRESSURE. MAN

Cardiovascular effects of breathing high CO₂ concentrations in the environmental air vary with the kind of anesthetic, depth of anesthesia, the state of the nervous system, experimental procedure, duration of exposure, oxygenation, and, to a lesser degree, with the animal species. Symbols: (+) = increase or elevation, (0) = no change.

CO ₂ Concentration	Exposure Time min	Anesthetic	Arterial pH	Pulse Rate	Blood Pressure	Remarks	Reference
1 2.5%	(B)	(C)	(D)	(E)	(F)	(G)	(H)
2 3.5%	15-20	None	7.39	0	0	No effect on cerebral circulation.	1
3 5.0%	15-20	None	7.35	0	0	Cerebral vasodilatation.	1
	3-5	None	7.35	0	0	25% increase in cardiac output; peripheral vasodilatation.	2
4 7.6%	2-7	None		0	Systolic = +6% Diastolic = +4%	increased metabolic rate.	
5 10.0%	4-20	None		+20 to +30%	Systolic = +22 to +24% Diastolic = +13 to +20%		
6 20.0%	15	Pentothal	7.15	+	Systolic = +25 to +55% Diastolic = +19 to +38%	Return to room air produced slight fall in systolic pressure and marked fall in diastolic pressure.	3, 4
7 25.0%	15	Pentothal		+		Transient extrasystoles; decrease in plasma volume; fall in blood pressure with return to air.	5
8 30.0%	15	Pentothal	6.98	+		Transient extrasystoles; decrease in plasma volume.	5
9	10-26	Pentothal	6.8	+		Extrasystoles; 10.5% increase in plasma volume; fall in blood pressure with return to room air.	5
10		None				Marked fall in blood pressure with return to room air.	6
						Muscular tremors; decreased coagulation time.	7

Contributors (a) Spencer, Joseph N., (b) Shephard, Roy J.

- References [1] Patterson, J. L., Jr., A. Heyman, L. L. Battey, and R. W. Ferguson. 1955. *J. Clin. Invest.* 34:1857. [2] Shephard, R. J. 1955. *J. Physiol.* Lond. 129:393. [3] Dripps, R. D., and J. H. Comroe, Jr. 1947. *Am. J. Physiol.* 149:43. [4] Seevers, M. H. 1944. *N. York State J. M.* 44:597. [5] Buesse, E. W., T. M. Parry, E. L. Goldensohn, R. W. Whitehead, and W. B. Draper. 1952. *Dis. Nerv. Syst.* 13:35. [6] Clowers, G. H., Jr., A. L. Hopkins, and F. A. Simcone. 1955. *Ann. Surg.* 142 446. [7] Leake, C. D., and R. M. Waters. 1929. *Current Res. Anesth.* 8:17.

22	40 0%	45	Pentothal	6.79	47%	0	0	0	+	synpharyngeal drugs, increased plasma potassium and coronary flow. 150% increase in cerebrospinal fluid pressure; decreased ventilation time. Ventricular fibrillation, circulatory collapse, and death with return to room air. 68% decrease in systolic excursion; 50% decrease in contraction force, marked decrease in cardiac output, epicardial hemorrhage; marked cardiac dilatation.	6,7
23		120	Pentothal	6.67							16
24	50 0%	4	Pentobarbital		-28%	-37%	0	0	Slurring	+	3
25	60 0%	190	Pentothal	6.46	-43%	-6%	+	+			12
26	70 0%	90	Artificial respiration	6.44	-40%	-28%	0	+			12
27	80 0%	66	Artificial respiration	6.40	-40%	-40%	0	+			12

Contributors. (a) Spencer, Joseph N. (b) Shephard, Roy J.

References [1] Vaughan Williams, E. M. 1955. *J. Physiol.*, Lond. 129:90. [2] Stroud, R. C., and H. Rahm. 1953. *Am. J. Physiol.* 172:211.

[3] Bonface, K. J., and J. M. Brown. 1953. *Ibid.* 172:752. [4] Page, I. H., and F. Olmsted. 1951. *Circulation*, N. Y. 3:801. [5] Nahas, G. G., and H. M. Cavert. 1957. *Fed. Proc.*, Balt. 16:92. [6] Spencer, J. N., T. M. Parry, R. W. Whitehead, and W. B. Draper. 1950. *J. Pharm. Exp. Ther.* 98:366. [7] Goldschohn, E. S., R. W. Whitehead, T. M. Parry, J. N. Spencer, R. F. Grover, and W. B. Draper. 1951. *Am. J. Physiol.* 165:334. [8] Whitehorn, W. V., and J. W. Bean. 1952. *Ibid.* 168:528. [9] Li, T. H., J. W. Dow, and M. Yamamoto. 1957. *J. Pharm. Exp. Ther.* 119:160. [10] Richards, J. B., and S. N. Stein. 1957. *Am. J. Physiol.* 188:1. [11] Leake, C. D., and R. M. Waters. 1929. *Current Res. Anesth.* 8:17. [12] Brown, E. B., Jr., and F. Miller. 1952. *Am. J. Physiol.* 170:550. [13] Billings, H. H., and E. R. Brown, Jr. 1953. *Ibid.* 180:363. [14] Houle, D. B., M. H. Well, E. B. Brown, Jr., and C. S. Campbell. 1957. *Proc. Soc. Exp. Biol. N. Y.* 94:561. [15] Brown, E. B., Jr. 1955. *Ibid.* 90:319. [16] Brown, E. B., Jr., and F. Miller. 1952. *Am. J. Physiol.* 169:56.

84. HEMODYNAMIC EFFECTS OF HYPERCAPNIA (HIGH CO₂) (Continued)

Part III: CHANGE IN PULSE RATE, BLOOD PRESSURE, AND THE ELECTROCARDIOGRAM: DOG (Concluded)

CO ₂ Concen- tration (A)	Exposure Time min (B)	Preparation or Method (C)	Anesthetic (D)	Arterial pH (E)	Pulse Rate (F)	Blood Pressure (G)	QRS Interval (H)	PR Interval (I)	ST Segment (J)	T Wave (K)	Remarks (L)	Refer- ence (M)
12 15.0%	10		Pentothal and chloralose	7.1							18% decrease in cardiac output and 20% increase in peripheral resistance due to change in pH.	9
13	20		Pentobarbital		+	+20%					Vagotomy, atropine, and carotid sinus denervation had no effect on response; para- vertebral ganglion- ectomy, total sympac- tomy, and adrenalectomy doubled response to CO ₂ .	4
14	20	Artificial respiration	Pentobarbital and curare		+	-30%					Marked rise in cerebro- spinal fluid pressure; maximal adrenocortico- cal stimulation.	6,7,10
15 20.0%	45		Pentothal	7.05	0		0	0	0	+	28% decrease in systolic excursion; 42% de- crease in contraction force; ventricular dilatation; depression of sino-atrial node and bundle of His; cardiac irregularities with return to room air.	3
16			None		Marked increase -20%	Marked increase -22%					Clonic convulsions; anesthesia; decreased coagulation time; increased resistance of RBC to hypotonic solution.	11
17 25.0%	12	Open chest	Pentobarbital				0	0	Slurring	+	Refractory to sym- -thombiotic drugs.	4
18 30.0%	15		None		0	0					Increased cerebrospinal fluid pressure; de- creased coagulation time; contraction of spleen; refractory to	12 6,7,13- 15
19	20	Artificial respiration	Pentobarbital and curare	6.93	+	-36%						
20	30		Pentothal		-18%	0	0	0				
21	45		Pentothal	6.83	+17%		0	0				

84. HEMODYNAMIC EFFECTS OF

Part IV: MEAN PRESSURE, pH, CO₂ PRESSURE,

Abbreviations: D-M = measured with a damped mercury manometer; S-T = measured directly by Stratham trans-appropriate corrections for body temperature; P-V = calculated by nomograms of Peters and Van Slyke; S-H = by bubble method of Riley; V-S = measured with Van Slyke manometric apparatus; N-G = obtained from nomogram

No.	Subjects Condition	CO ₂ Concentration ¹	Exposure Time min	Mean Blood Pressure		
				Method	Control (Air)	Hypercapnia
(A)	(B)	(C)	(D)	(E)	(F)	(G)
					mm Hg	
16 ²	Supine and fasting	5.0%	(15-30)	D-M	81(78-84)	90(83-94)

/1/ In air. /2/ Young males. /3/ Young, normotensive, non-arteriosclerotic subjects; mean age, 30 yr. /4/ arteriosclerosis; mean age, 51 yr. /6/ Hypertensive, arteriosclerotic subjects; mean age, 55 yr. /7/ Mean value

Contributors: Patterson, John L., Jr., and E. L. Hardie

References: [1] Kety, S. S., and C. F. Schmidt. 1948. J. Clin. Invest. 27:484. [2] Novack, P., et al. 1953, and A. Wasserman. Unpublished.

Part V. CARDIAC OUTPUT, BLOOD

Subjects unanesthetized, unless otherwise specified. Blood pressure determined by auscultatory method. estimate "c" (cf. Introduction).

No.	Subjects Condition	CO ₂ Concentration	Exposure Time min	Cardiac Output		
				Method	Control (Air)	Hypercapnia
(A)	(B)	(C)	(D)	(E)	(F)	(G)
					L/min	
16 ¹		3.0% ^{2,3}	23.5(17-32)			
2		6.0% ^{2,3}	23.5(17-32)			
3 ⁴	Supine and fasting	1.52% in air	15	A-C	4.4	4.3
4		1.82% in air	37	A-C	3.6	3.5
5		3.52% in air	40	A-C	4.0	3.8
6		5.93% in air	15	A-C	4.1	4.8
7		6.05% in air	8	A-C	4.4	4.6
8		6.90% in air	15	A-C	5.0	6.2
9		7.49% in air	15	A-C	4.0	5.4
10		7.61% in air	25	A-C	4.0	5.7
11		9.25% in air	5	A-C	4.0	5.7
12	Recumbent	10.0% in O ₂	5			
13	Recumbent	10.0% in O ₂	5			
14	Recumbent	10.0% in O ₂	5			
15	Recumbent	6.5% in air	15	A-C	5.15(5.07-5.22)	8.38(8.34-8.42)
16	Tilted +60°	6.5% in air	15	A-C	5.25	4.18
17	Supine and fasting	5.0% in air	(15-30)	B-C	4.7(3.5-6.0)	5.9(3.5-6.3)
18	Supine and fasting	7.0% in air	(15-30)	B-C	5.0(4.3-5.6)	5.1(4.5-5.7)
19	Local anesthesia	7.0% in air	7	D-D	6.40(3.48-18.28)	8.27(5.99-10.78)

/1/ 46 observations. /2/ O₂ concentration of rebreathed air maintained at about 30% /3/ Rebreathing in Larson-experiments. /5/ Patients with essential hypertension, some of whom showed narrowing of arterioles in ocular 1-hour rest period in horizontal position prior to testing. /8/ Young males. /9/ 17 observations.

Contributors: Patterson, John L., Jr., and E. L. Hardie

References: [1] Schneider, E. C., and D. Truesdell. 1922. Am. J. Physiol. 63:155. [2] Grollman, A. 1930. 1943. Acta physiol. scand. 6:176. [5] Kety, S. S., and C. F. Schmidt. 1948. J. Clin. Invest. 27:484. [6] Richardson,

HYPERCAPNIA (HIGH CO₂) (Continued)

AND HEART RATE: CAT, DOG, FROG

Ringer's solution equilibrated with a CO₂-O₂ mixture, M-M = artery cannulated and direct blood pressure measured
 Hastings and Sendroy, G-E = measured by glass electrode pH meter. Values in parentheses are ranges, estimate

Blood Pressure				Heart Rate		Arterial pH			Reference	
Systolic Change	Mean Arterial		Change	Control (Air)	Hypercapnia	Method	Control (Air)	Hypercapnia		
	(K)	(L)		(M)	(N)		(O)	(P)	(Q)	(R)
+70 mm Hg										1
+54 mm Hg										2
+91 mm Hg										3
+120 mm Hg										4
						Slows slightly				5
						Slows markedly				6
					91	78				7
					88	78				8
					69	68	H-S	7.34	7.35	9
					86	80	H-S	7.35	7.30	10
					75	64				11
					67	61	H-S	7.34	7.35	12
					72	56				13
			(+10 to +28)							14
			(-10 to -30) ⁵							15
			-20(-40 to +12)							16
			-40(-106 to +33)							17
Varies ⁸				162	70(52-88)	G-E	7.39	6.41		5
				154	120	G-E	7.34	6.60		19
				184	107	G-E	7.29	6.46		20
				198	126					21
					75	G-E	7.34	6.40		22
-14%					-7% ⁹					6
-18%					-16% ⁹					23
-22%					-20% ⁹					24
-37%					-28% ⁹					25
	135(107-152)	141(119-152)	+6(0 to +12)	206	184(101-216)					7
				206	179(106-211)					26
					61(147-205)					27
					83(136-213)					28
					53(129-176)					29
					52(130-172)					30
					51(134-167)					31
				20	19					7
				26	26					2
				28	25					34
										35

maximum pressure change attained /5/ Followed by a 10-20 minute increase. /6/ Average concentration at time of varying from 1% every 3 minutes to 2% every minute /8/ At 30% CO₂ in O₂, blood pressure fell sharply, but above control level and remained high until 50-70% CO₂ was reached, but fell precipitously as CO₂ increased above

Ibid 40 279. [3] Marshall, E. K., Jr 1926 J. Pharm Exp Ther. 29:167. [4] Page, I. H., and A. B. Olmsted. K. J., and J. M. Brown. 1953. Ibid 172 752. [7] Brickner, E. W., et al. 1956. Ibid 184 275.

Abbreviations: H-P = measured with glass heart plethysmograph, F-P = Fick principle, R-S = perfusion with mercury manometer, G-M = measured with a Gregg manometer; H-S = measured by calorimetric method of "c" (cf. Introduction).

Subjects			Anesthetic	CO ₂ Concentration	Exposure Time min	Cardiac Output			Method
Animal	No.	Condition				Method	Control (Air)	Hypercapnia	
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)
1	Cat	Decerebrate	Curare and chloroform	4.7% ¹	1.83				M-M
2				7.0% ²	1.33				M-M
3				10.8% ³	1.58				M-M
4				10.0% ⁴	0.67				M-M
5	6	Isolated heart-lung	A.C.E. mixture or chloroform	2-8%		H-P		+	
6				12-20%		H-P		-	
7	Dog	1 Trained	None	4.0%	5	F-P	2.37	2.42	
8				3.8%	5	F-P		2.32	
9				5.4%	5	F-P	2.40	2.27	
10				7.9%	5	F-P	2.65	1.98	
11				5.0%	5	F-P	1.19	1.16	
12				5.5%	5				
13				7.6%	5				
14		Ventilated normally	Pentobarbital	5-15% in O ₂	Maximum ⁴				M-M
15		Ventilated artificially	Pentobarbital and curare	5-15% in O ₂					M-M
16									M-M
17									M-M
18	7	Tracheotomized	Sodium pentothal						M-M
19	1	Tracheotomized	Sodium pentothal	50% in O ₂	270				
20				60% in O ₂	190				
21				70% in O ₂	90				
22				80% in O ₂	66				
23	19	Bilateral cervical vagotomy; heart in situ	Pentobarbital and barbital	5% in O ₂	9				M-M
24				10% in O ₂	13				M-M
25				25% in O ₂	12				M-M
26				50% in O ₂	4				M-M
27	6	Splenectomized	Sodium pentobarbital	2.3%	23				G-M
28				6.3%	33				G-M
29				10.4%	32				G-M
30				14.1%	26				G-M
31	3	Splenectomized	Sodium pentobarbital	6.5%	33				G-M
32				10.3%	30				G-M
33				14.3%	36				G-M
34	Frog	Isolated heart		16% in O ₂		R-S	0.234	0.048	
35				12% in O ₂		R-S	0.208	0.039	
36				7% in O ₂		R-S	0.280	0.113	

/1/ 20% O₂ in gas mixture. /2/ 25% O₂ in gas mixture. /3/ 19% O₂ in gas mixture. /4/ Inhalation continued until death of animal. /7/ CO₂ concentration was started at 30% and increased by adding 100% CO₂ to mixture at rates immediately recovered to control value or higher. As CO₂ concentration increased, blood pressure usually rose 70%. /9/ From control value.

Contributors: Patterson, John L., Jr., and E. L. Hardie

References: [1] Mathison, G. C. 1911. J. Physiol. Lond. 42:283. [2] Jerusalem, E., and E. H. Starling. 1910. 1951. Circulation, N. Y. 3 801. [5] Brown, E. B., Jr., and F. Miller 1952. Am. J. Physiol. 170:550. [6] Boniface.

HYPERCAPNIA (HIGH CO₂) (Continued)

VASCULAR RESISTANCE: CAT, DOG

branch of left coronary artery, I-H = isolated heart, coronaries perfused with blood through which 100% CO₂ bubbled, cannula in anterior descending coronary artery, orifice meter.

Vascular Resistance			Arterial CO ₂ Content		Arterial pH		Reference
Control (Air)	Hypercapnia	Change %	Control (Air)	Hypercapnia	Control (Air)	Hypercapnia	
mm Hg/ml/100 g/min	mm Hg/ml/100 g/min		vol %		(O)	(P)	
(J)	(K)	(L)	(M)	(N)	(O)	(P)	(Q)
					7.2	6.6	1
45	43	-4					2
							2
15	10	-33					3
20	19	-9	15	25	7.7	7.6	4
21	13	-38	15	38	7.7	7.4	5
	No change						5
							6

J. E., et al. 1947. Am. J. Physiol. 148:582. [3] Anrep, G. V., and R. S. Stacey. 1927. J. Physiol., Lond. 64:187, 135-271.

VASCULAR RESISTANCE. MAN, DOG

eter. Values in parentheses are ranges, estimate "c" (cf. Introduction).

Vascular Resistance			CO ₂ Pressure		Reference
Control (Air)	Hypercapnia	Change %	Control (Air)	Hypercapnia	
mm Hg/ml/100 ml/min	mm Hg/ml/100 ml/min		mm Hg		
(J)	(K)	(L)	(M)	(N)	(O)
					1
13.2	21.6	+64	412	552	2
13.2	27.5	+108	412	642	3
30	33	+3			4
45	42	-9	482	642	5
6.4	4.6	-28	405	565	6
17.4	28.3	+60			7
12.6	8.6	-28			8
					9

aining 8 5% CO₂ /4/ Sympathetic innervation blocked by intra-arterial phenoxybenzamine. /5/ Arterial CO₂ Ganglionic blockade with intravenous tetraethylammonium chloride.

D. Truesdell 1922. Am. J. Physiol. 63:155. [3] Clarke, R. S. J. 1957. J. Physiol., Lond. 118:537. [4] Richardson,

Part VII: CORONARY BLOOD FLOW AND

Method: C-C = closed chest, spontaneous respiration, bubble flowmeter between carotid artery and circumflex
 M-C = Morowitz cannula in coronary sinus, heart-lung preparation; O-C = open chest, artificial respiration,

Subjects		Anesthetic	Method	CO ₂ Concentration	Exposure Time min	Blood Flow		
Animal	No.					Control (Air) ml/100	Hypercapnia g/min	Change %
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)
1 Cat	1		I-H		5	50	75	+50
2 Dog	6	Nembutal, or chloralose and morphine	C-C	5-7%		23	23	0
3	1		M-C	7%		68	98 ¹	+45
4	1	Chloralose and morphine	M-C	3%	20	45	50	+12
5				7%	15	45	70	+55
6	5 ²	Barbital and morphine	O-C	5-8%			Slight decrease	

/1/ Heart volume increased 7.5 ml during hypercapnia. /2/ 14 observations.

Contributors: Richardson, David W., and John L. Patterson, Jr.

References: [1] Trethewie, E. R., and M. M. Hodgkinson. 1955. Q. J. Exp. Physiol., Lond. 40.1. [2] Eckenhoff, [4] Hilton, R., and F. Eichholtz. 1925. Ibid. 59:413. [5] Wegria, R., and H. D. Green. 1942. Am. J. Physiol.

Part VIII: LIMB BLOOD FLOW AND

Method: V-P = venous occlusion plethysmography; S-C = Stewart calorim-

Subjects		Extremity	Method	CO ₂ Concentration	Exposure Time min	Blood Flow		
Animal	No.					Control (Air) ml/100	Hypercapnia ml/min	Change %
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)
1 Man	4	Hand	V-P	5.4%	5-6	2(1.5-2.5)	1(0.5-1.5)	-50
2		Hand ¹	V-P	5.4-6.4%	5-7	1.5(1-2)	2(1.0-2.5)	+33
3	6	Hand	S-C	5%	15	6.3(0.9-12.7)	4.3(0.4-8.7)	-32
4				7%	20	6.3(0.9-12.7)	3.6(0.8-6.8)	-43
5	8	Forearm	V-P	6.5% ³	3	3(1.5-4.0)	3(2-5)	0
6	10	Forearm	V-P	8%	3	2(1.5-3.0)	3(2-4)	+50
7	8	Forearm ⁴	V-P	5 or 7%	10 or 5	14(9-21)	24(12-35)	+71
8 Dog	4	Leg ⁵	V-P	5%	Maximum ⁷	9.5(5-16)	6.8(3-12)	+30
9	5	Leg ^{6,8}	V-P	5%	Maximum ⁷	9.1(4-12)	12.5(5-23)	+37

/1/ Sympathectomized. /2/ Alveolar CO₂ pressure. /3/ Vigorous voluntary hyperventilation of gas mixture compressure. /6/ Anesthetized with chloralose. /7/ CO₂ inspired until maximum circulatory effects noted. /8/

Contributors: Richardson, David W., and John L. Patterson, Jr.

References: [1] Gellhorn, E. 1943. Autonomic regulations. Interscience, New York. [2] Schneider, E. C., and D. W., et al. 1958. Clin. Res. 6:127. [5] Patterson, J. L., Jr., et al. Unpublished.

HYPERCAPNIA (HIGH CO₂) (Continued)

AND VASCULAR RESISTANCE MAN

flow. Values in parentheses are ranges, estimate "c" (cf. Introduction).

Mean Arterial Blood		Arterial CO ₂ Pressure		Arterial pH		Arterio-venous		
(L)	(M)	(N)	(O)	(P)	(Q)	(R)	(S)	(T)
82(78-88)	93(83-110)	43	52	7.38	7.33	6.1	3.8	1
91(79-111)	95(82-111)	43	50	7.36	7.30	6.6	4.7	2
87(61-107)	90(74-105)	38.4	42.5	7.44	7.39	5.9	6.1	3
98(72-144)	95(78-126)	39.4	44.7	7.40	7.35	6.0	5.4	3
								4
								5
								6
								7
								8
132(110-169)	143(92-193)							5
112(86-168)	130(95-193)							6
								10
95(83-108)	115(90-193)							5
								11
136(119-160)	150(110-200)	40	48	7.41	7.35	6.0	4.3	2
								12
94(72-115)	98(76-115)	46	53	7.34	7.30	6.3	4.9	2
								13
132(117-156)	159(110-172)	38	48	7.41	7.33	7.7	5.1	2
								14

32 696. [3] Patterson, J. L., Jr., et al 1955. *Ibid.* 34:1857. [4] Schieve, J. F., and W. P. Wilson. 1953. *Am. Sc* 223:245

VASCULAR RESISTANCE MAN, DOG

estimate "c" (cf. Introduction)

Change %	Vascular Resistance ¹		Change %	Mean Blood Pressure			Reference
	Control (Air)	Hypercapnia		Control (Air)	Hypercapnia	Change %	
	mm Hg/ml/min			mm Hg			
(J)	(J)	(K)	(L)	(M)	(N)	(O)	(P)
-13(0 to -39)			+46(7-110)			+21	1
-23(-9 to -66)			+75(23-228)			+16	2
	0.56	0.56		150	152	+1.3	3
	0.56	0.51	-8.9	150	142	-5.3	4
	0.56	0.50	-10.7	150	139	-7.3	5
	0.56	0.60	+7.1	150	135	-10	6

progressive hypercapnia, stage IV = 16.8% average CO₂ concentration.

Exp Biol. N. Y. 84:15.

Nitrous oxide sampling method employed in measuring cerebral blood

Subjects			CO ₂ Concentration ²	Exposure Time min	Blood Flow			Vascular Resistance			
No.	Age ¹ yr	Condition			Control (Air)	Hypercapnia	Change %	Control (Air)	Hyper- capnia	Change %	
					ml/100 g brain/min			mm Hg/ml/100 g brain/min			
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)		
4	19	35	Convalescent	3.5%	15-30	52(30-68)	57(33-74)	+9.6	2.0(1.2-3.6)	1.8(1.1-3.4)	-10.0
5	29	44	Normal	5%	4	59(55-62)	89(77-105)	+50.8			
6				7%	4	59(55-62)	129(119-139)	+101.7			
7	10	59	Stroke	5%		35	49	+40.1			
8				7%		35	61	+74.2			
9	18	69	Hypertension	5%	10-15	39(25.5-67.6)	47(28.4-77.0)	+20.5	3.6(2.1-4.7)	3.2(1.6-5.1)	-11.1
10	27	71	Cerebral vascular disease	5%	10-15	35(29.6-55.0)	51(35.4-89.9)	+45.7	3.5(2.0-4.3)	2.8(1.0-4.0)	-18.9
11	13	74	Normotension	5%	10-15	35(20.7-63.6)	55.5 (32.7-94.5)	+58.7	2.9(1.4-4.4)	2.2(1.0-3.2)	-24.1
12	6		Essential hypertension	5%	10	52(31-67)	86(72-103)	+65.4	2.6(1.9-3.5)	1.7(1.0-2.3)	-34.6
13	10		Arterioscle- rosis	5%	10	47(29-98)	55(33-99)	+17.1	2.1(0.8-3.2)	1.9(0.8-3.0)	-9.8
14			Hypertension, arterioscle- rosis	5%	10	36(23-51)	49(24-94)	+36.2	3.7(2.5-6.3)	3.0(1.4-6.0)	-18.9

/1/ Mean. /2/ Lines 1-9, 11-14, CO₂ in 21% O₂, line 10, CO₂ concentration assumed also to be in 21% O₂.

Contributors: Patterson, John L., Jr., and C. L. Carter

References: [1] Kety, S. S., and C. F. Schmidt. 1948. J. Clin. Invest. 27:484. [2] Novack, P., et al. 1953. Ibid. J. Med. 15:171. [3] Fazekas, J. F., et al. 1953. J. Geront. 8:137. [4] Fazekas, J. F., et al. 1952. Am. J. M.

Part X. RENAL BLOOD FLOW AND

Values in parentheses are ranges.

Animal	No. of Subjects	Anesthetic	Method	CO ₂ Concentration in O ₂	Exposure Time min	Blood Flow	
						Control (Air)	Hypercapnia
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)
1 Man, normotension	6			10%	10		
2 Man, hypertension	5			10%	10		
3 Dog	8	Pentobarbital, intravenous	p-Amino- hippu- rate and mannitol	Stage I ²	30	275	275
4				Stage II ²	40	275	275
5				Stage III ²	40	275	275
6				Stage IV ²	40	275	230

/1/ Renal vascular resistance = mean blood pressure/renal blood flow. /2/ Roman numerals indicate stages of

Contributor: Patterson, John L., Jr

References: [1] Little, W. J., et al. 1949. Fed. Proc., Balt. 8:98. [2] Dowds, E. G., et al. 1953. Proc. Soc.

HYPERCAPNIA (HIGH CO₂) (Concluded)

PULMONARY ARTERIAL PRESSURE: MAMMALS

spontaneous breathing, I-R = isolated, perfused lung, open-circuit respiration; I-T = isolated, perfused lung, 20 ml P-A = isolated, perfused lung, 8% CO₂ equilibrated with perfusate. Values in parentheses are ranges, estimate "c"

Blood Flow			Mean Arterial Pressure			Reference
Control (Air)	Hypercapnia	Change	Control (Air)	Hypercapnia	Change	
L/min			mm Hg			
(G)	(H)		(J)	(K)		
			19(12-25)	20(16-27)	+4%	1
5.3(4-10)	8.0(6-11)	+45%				2
9.1(4-17)	10.8(7-23)	+18%	31(11-77)	34(12-84)	+10%	3
4.6(2-9)	5.1(2-10)	+11%	14(8-30)	15(7-30)	+11%	4
			13	15	+10% ¹	5
			20	18	-8% ¹	6
			20	18	-8%	7
			19	21	+12%	8
	Increased		14	15	+11%	9
	Decreased		21	16	-25%	10
	Decreased		18	16	-9%	11
3.1(2.6-3.7) ^b	3.3(2.7-3.9) ^b	+6%	27(19-34) ^b	27(21-34) ^b	+2%	7
			30	31(29-34)	+3%	8
			30	30(29-33)	+1%	14
					+15-16) mm Hg	9
					+9-12) mm Hg	15
					+12-13) mm Hg	16
			12(9-22)	15(10-25)	+25%	17
						18

after anesthesia. /5/ Observations, not subjects. /6/ 21 observations.

Res 7:136. [3] Hebb, C. O., and R. H. Nimmo-Smith. 1949. Q. J. Exp. Physiol., Lond. 34:159. [4] Shephard, R. J. 1951. Ibid. 23:85, 361. [7] Stroud, R. C., and H. Rahn. 1953. Am. J. Physiol. 172:211. [8] Duke, H. H. 1950. Q. J.

RESISTANCE, AND HEPATIC VASCULAR RESISTANCE. DOG

spirometer never decreased below 21 volumes per cent. CO₂ concentration is mean per cent obtained from 3-7. Values in parentheses are ranges, estimate "c" (cf. Introduction).

Mesenteric Vascular Resistance ²			Inferior Vena Caval Pressure		Hepatic Vascular Resistance ³		
Control (Air)	Hypercapnia	Change %	Control (Air)	Hypercapnia	Control (Air)	Hypercapnia	Change %
mm Hg/ml/min			mm Hg		mm Hg		
(L)	(M)	(N)					

/3/ Calculated from: $\frac{\text{portal venous pressure} - \text{inferior vena caval pressure}}{\text{blood flow}}$ /4/ 6% CO₂ in spirometer at start of

Part XI: PULMONARY BLOOD FLOW AND MEAN

	Animal	No. of Subjects	Anesthetic	Method	CO ₂ Concentration	Exposure Time min
	(A)	(B)	(C)	(D)	(E)	(F)
1	Man	5	Seconal	C-C	5%	5-10
2		10		D-D	7%	7
3	Man, congenital heart disease	81		C-C	5%	3
4		12 ²		C-C	5%	3
5	Cat	1	Chloralose	S-B	3%	
6					6.5%	
7					18.7%	
8					20.5%	
9		2		I-R	8%	4
10		1		P-X	10%	6
11		1		P-A	8%	6
12	Dog	5	Nembutal	C-C	5%	10-15
13		10 ⁴	Nembutal	I-R	5%	21-73
14		60 ^{4,5}	Nembutal	I-R	10%	21-73
15				I-R	10%	31-45
16					20%	31-45
17					30%	31-45
18	Monkey, rhesus	24, 6	Nembutal	I-T	100%	

Contributors: Richardson, David W., and John L. Patterson, Jr.

Part XII: MESENTERIC BLOOD FLOW AND VASCULAR

[illegible]

Contributors: Patterson, John L., Jr., and E. L. Hardie

Reference: Brickner, E. W., E. G. Dowds, B. Willets, and E. E. Selkurt. 1956. *Am. J. Physiol.* 184: 275.

88. EFFECT OF BREATHING AIR AT 4 ATMOSPHERES ON PULMONARY ARTERIAL PRESSURE. RAT

Following prolonged exposure (4-6 weeks) in compressed air (O_2 pressure = 0.7-1.0 atmosphere), pulmonary arteriolar lesions (thickening and hyalinization leading to thrombosis) develop and later degenerative lesions of the large arteries of the lungs. The obliterative vascular changes are followed by a rise in pulmonary pressure, enlargement of the right ventricle, and a fall in systolic pressure. O_2 pressure of 4 atmospheres = 615 mm Hg. Values in parentheses are ranges, estimate "c" (cf. Introduction).

No. of Rats	Duration of Exposure da	Pulmonary Arterial Pressure ¹ cm H_2O
(A)	(B)	(C)
1/5	3	207(133-231)
2/5	10	244(176-304)
3/5	17	255(163-425)
4/5	24	318(279-394)
5/5	31	409(322-597)
6/3	38	423(322-529)

1/1 Control average for 34 rats = 256 cm H_2O

Contributor: Behnke, A. R.

Reference: Bennett, G. A., and F. J. C. Smith 1934. J. Exp. M. 59 181.

89. EFFECT OF RECOMPRESSION, WITH O_2 AND WITH AIR, ON ARTERIAL BLOOD PRESSURE. DOG

All values are for blood pressure, in mm Hg. Compression period = 105 minutes at 65 lb/sq in. gauge pressure

Control Period ¹	Asphyxial Period ²		Recompression Period ³	Postrecompression Period ⁴
(A)	High (B)	Low (C)	(D)	(E)
Breathing O_2				
1 127	166	64	104	100
2 102	110	62	74	74
3 117	136	60	115	
4 116	140	30	90	
5 150	172	68	90	100
6 158	166	84	60	80
Breathing Air				
7 147	154	80		65
8 97	104	62	105	
9 142	130	78	80	102
10 115	128	54	122	85
11 128	162	86	100	80
12 132	137	80	124	108
13 146	120	64	86	125
			90	114
				90

1/1 In air at 1 atmosphere 1/2 Following rapid decompression in 5-6 seconds. 1/3 Breathing at 30 lb/sq in. for 1 1/2 hours. 1/4 1 hour following recompression period.

Contributor: Behnke, A. R.

Reference: Behnke, A. R. 1945. Medicine, Balt. 24 381

85. PRESSURE EQUIVALENTS

Atmospheres		mm Hg	Absolute lb/sq in.	Gauge lb/sq in.	Diving Depth ft
(A)	(B)	(C)	(D)	(E)	
1	1	760	14.7	0	0
2	2	1520	29.4	14.7	33
3	3	2280	44.1	29.4	66
4	4	3040	58.8	44.1	99
5	5	3800	73.5	58.8	132
6	6	4560	88.2	73.5	165
7	7	5320	102.9	88.2	198
8	8	6080	117.6	102.9	231
9	9	6840	132.3	117.6	264
10	10	7600	147.0	132.3	297

Contributor: Behnke, A. R.

86. EFFECT OF COMPRESSION ON PULSE RATE,
SYSTOLIC BLOOD PRESSURE, AND PULSE PRESSURE: MAN

Tests conducted in pressure chamber. Values expressed as per cent of sea level values.

Atmospheres		Pressure mm Hg	Pulse Rate			Blood Pressure		Pulse Pressure	
(A)	(B)	(C)	Reclining	Standing	Exercising	Reclining	Standing	Reclining	Standing
1	1	760	98.8	100.0	100.0	113.8	119.3	48.2	38.3
2	2	1520	90.7	91.6	92.8	106.2	113.2	37.2	26.9
3	4	3040	86.6	86.6	88.4	108.9	113.9	41.6	32.5
4	6	4560	89.4	89.9	89.3	107.2	113.7	34.9	26.8
5	7	5320	83.8	87.4	85.2	108.5	112.0	36.7	24.0
6	10	7600	90.7	84.0	85.2	109.0	126.0	30.0	30.0

Contributor: Behnke, A. R.

Reference: Shilling, C. W., J. A. Hawkins, and R. A. Hansen. 1936. U. S. Nav. M. Bull. 34 39.

87. EFFECT OF BREATHING O₂ AT 3-4 ATMOSPHERES ON RESPIRATORY RATE,
PULSE RATE, AND BLOOD PRESSURE: MAN

Period: C = control period of air-breathing at one atmosphere, E = experimental period of O₂-breathing at increased ambient pressure. Other effects: O = none discernible, P = pallor, T = twitching movements of a myoclonic nature, M = mental confusion, G = generalized type of convulsions, S = sweating. Values in parentheses are ranges, estimate "c" (cf. Introduction).

No. of Subjects	Period	O ₂ Pressure atmospheres	Respiratory Rate breaths/min	Pulse Rate beats/min	Blood Pressure		Other Effects	
					Arterial mm Hg	Internal Jugular mm Hg		
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	
1	1	C	0.2	10	73	96	11.3	
2		E	3.0	11	61	102	9.0	O
3	7	C	0.2	15(12-19)	67(53-87)	78(59-97)	10.7(8.2-17.2)	
4		E	3.5	16(11-20)	59(49-71)	86(75-108)	6.9(5.5-13.2)	O
5	1	C	0.2	11	49	77	9.6	
6		E	3.5	10	48	79	7.9	P
7	3	C	0.2	14(10-17)	68(56-76)	79(72-85)	8.3(8.0-8.7)	
8		E	3.5	21(10-32)	63(61-66)	80(76-82)	6.1(4.1-7.4)	T
9	1	C	0.2	20	75	87	10.0	
10		E	3.5	12	51	91	9.6	P, T, G, S
11	1	C	0.2	19	79	85	6.5	
12		E	4.0	19	78	89	5.4	P, T, G, S
13	1	C	0.2	7	63	79	6.0	
14		E	4.0	12	59	86	4.8	P, M
15	1	C	0.2	13	79	76	5.5	
16		E	4.0		66	85		T, M, S

Contributor: Behnke, A. R.

Reference: Lambertsen, C. J., R. H. Kough, D. Y. Cooper, G. L. Emmel, H. H. Loeschke, and C. F. Schmidt. 1953. J. Appl. Physiol. 5 471

88. EFFECT OF BREATHING AIR AT 4 ATMOSPHERES ON PULMONARY ARTERIAL PRESSURE. RAT

Following prolonged exposure (4-6 weeks) in compressed air (O_2 pressure = 0.7-1.0 atmosphere), pulmonary arteriolar lesions (thickening and hyalinization leading to thrombosis) develop and later degenerative lesions of the large arteries of the lungs. The obliterative vascular changes are followed by a rise in pulmonary pressure, enlargement of the right ventricle, and a fall in systolic pressure. O_2 pressure of 4 atmospheres = 635 mm Hg. Values in parentheses are ranges, estimate "c" (cf. Introduction).

No. of Rats	Duration of Exposure da	Pulmonary Arterial Pressure ¹ cm H ₂ O
(A)	(B)	(C)
1 5	3	207(133-234)
2 5	10	244(176-304)
3 5	17	255(163-425)
4 5	24	318(279-394)
5 5	31	409(322-597)
6 3	38	423(322-529)

^{1/1} Control average for 34 rats = 256 cm H₂O

Contributor: Behnke, A R

Reference: Bennett, G A., and F J. C. Smith 1934 J. Exp. M. 59:181.

89. EFFECT OF RECOMPRESSION, WITH O_2 AND WITH AIR, ON ARTERIAL BLOOD PRESSURE. DOG

All values are for blood pressure, in mm Hg. Compression period = 105 minutes at 65 lb/sq in. gauge pressure.

Control Period ¹	Asphyxial Period ²		Recompression Period ³	Postrecompression Period ⁴
	High	Low		
(A)	(B)	(C)	(D)	(E)
			Breathing O_2	
1 127	166	64	104	100
2 102	110	62	74	74
3 117	136	60	115	
4 116	140	30	90	
5 150	172	68	90	100
6 158	166	84	60	80
			Breathing Air	65
7 147	154	80	106	
8 97	104	62	80	102
9 142	130	78	122	85
10 115	128	54	100	80
11 128	162	86	124	108
12 132	137	80	86	125
13 146	120	64	90	114
				90

^{1/1} In air at 1 atmosphere ^{2/2} Following rapid decompression in 5-6 seconds. ^{3/3} Breathing at 30 lb/sq in. for ^{4/4} 1 hour following recompression period.

Contributor: Behnke, A R

Reference: Behnke, A R. 1945 Medicine, Balt. 24 381.

90. EFFECT OF EXPLOSIVE DECOMPRESSION ON HEART RATE AND ARTERIAL PRESSURE: MAMMALS

Values in parentheses are ranges, estimate "c" (cf. Introduction).

Animal	No. of Subjects	Rate of Decompression mm Hg/sec	Range of Decompression mm Hg	Heart Rate	Arterial Pressure	Remarks	Reference
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)
1 Man	150 ¹	(38-1310)	(564-179) to (253-111)	Occasional sinus tachycardia associated with apprehension.	No reported or observed circulatory signs or symptoms.	No significant changes in configuration of ECG.	1
2 Dog	45 ²	(1,200-33,650)	(522-87)	Transient bradycardia at rates greater than 8520 mm Hg/sec, followed by recovery.	Transient fall, proportional to rate and range of decompression, followed by recovery.	Anesthetized (nembutal).	2
3	8 ³	3280	(522-30)	Bradycardia.	Fall to 60 mm Hg, then constant during period of decompression.	Anesthetized (nembutal).	3
4	15	33,650	(560-30)	Bradycardia following moment of decompression.	Fall to 70 mm Hg 30 sec after decompression, then essentially unchanged during 1-min exposure to low pressure.	Anesthetized (nembutal). Disappearance of arterial pulse within 30 sec after decompression; recovery usual after recompression.	4
5	8 ⁴		(520-30) or (180-30)	Bradycardia, premature beats, paroxysmal tachycardia, heart block.	Gradual fall to 50-70 mm Hg within 15 sec of decompression, then constant for remainder of 1-min exposure. ⁵	Unanesthetized. Gradual decrease and disappearance of arterial pulse pressure; occasional atypical elevation of arterial pressure.	5
6 Monkey	105 ⁶	Approximately (1000-4000)	(349-27), (522-76), or (750-93)	Transient bradycardia followed by recovery and secondary "anoxic fall," if animal retained at low pressure.		Anesthetized and unanesthetized animals.	6
7 Rat	250	Approximately 3000	(349-186) to (349-27)	Transient bradycardia followed by recovery and secondary "anoxic fall."		Anesthetized and unanesthetized animals.	7

¹/1 554 observations. ²/2 52 observations. ³/3 24 observations. ⁴/4 8 observations. ⁵/5 Venous pressure increases suddenly to 25-35 mm Hg during period of decompression. ⁶/6 202 observations.

Contributor. Whitehorn, William V.

References [1] Hitchcock, F. A., W. V. Whitehorn, and A. Edelmann. 1948. *J. Appl. Physiol.* 1:418. [2] Whitehorn, W. V., A. Lein, and A. Edelmann. 1946. *Am. J. Physiol.* 147:289. [3] Vail, E. G. 1952. *J. Aviat. M.* 23:577. [4] Whitehorn, W. V. 1948. *Fed. Proc.* Balt 7:133. [5] Kempf, J. P., B. H. Burch, F. M. Beman, and F. A. Hitchcock. 1954. *J. Aviat. M.* 25:107. [6] Gellan, S. 1950. *J. Appl. Physiol.* 3:254. [7] Gellan, S., L. F. Nims, and R. B. Livingston. 1950. *Am. J. Physiol.* 162:37.

91. EFFECT OF EXPLOSIVE DECOMPRESSION ON ARTERIAL PRESSURE: DOG

Values in parentheses are ranges, estimate "c" (cf. Introduction).

Part I. FINAL PRESSURE GREATER THAN THEORETICAL VAPOR PRESSURE OF BODY FLUIDS

Section 1: Effect of Decompression from 522-87 mm Hg (10,000-50,000 ft) at Varying Rates

No. of Dogs	No. of Observations	Rate of Decompression mm Hg/sec	Blood Pressure mm Hg	Cardiac Slowing
(A)	(B)	(C)	(D)	(E)
1 7	8	1200	24(0-40)	None
2 13	19	4800	28(0-50)	None
3 10	10	8520	28(5-65)	Occasional
4 10	10	13,460	50(32-80)	Present
5 5	5	33,650	81(75-100)	Present

Section 2. Effect of Decompression at Rate of 4800 mm Hg/sec and at Varying Intervals in Range of Decompression

No. of Dogs	No. of Observations	Interval in Range of Decompression mm Hg	Blood Pressure mm Hg
(A)	(B)	(C)	(D)
1 1	1	675	70
2 13	19	435	28(0-50)
3 5	5	375	14(0-25)
4 5	5	300	9(0-12)
5 5	5	275	3(0-12)

Section 3 Effect of Increased Expiratory Resistance at Decompression Rate of 4800 mm Hg/sec and Range of 522-87 mm Hg

Diameter of Expiratory Opening mm	Blood Pressure mm Hg	Cardiac Slowing
(A)	(B)	(C)
1 15 (normal)	0	None
2 15 (normal)	10	None
3 15 (normal)	18	None
4 4 5	45	None
5 2 5	55	Present
6 1 0	60	Present
		Marked

Contributor Whitehorn, William V.

Reference Whitehorn, W. V., A. Lein, and A. Edelmann. 1946 Am. J. Physiol 147:289.

Part II. FINAL PRESSURE LESS THAN THEORETICAL VAPOR PRESSURE OF BODY FLUIDS

Decompression from 560-30 mm Hg Arterial pressures for 14 controls systolic = 164(95-190) mm Hg, diastolic = 124(85-165) mm Hg

No. of Observations	Time after Decompression sec	Mean Arterial Blood Pressure mm Hg	Pulse Pressure Observation	
(A)	(B)	(C)	Present (D)	Absent (E)
1 14	10	73.6(50-97)	6	8
2 13	30	70.7(50-97)	3	10
3 12	45	76.1(52-97)	2	10
4 5	60	71.8(60-75)	1	6

Contributor Whitehorn, William V.

Reference Whitehorn, W. V. 1948. Fed. Proc., Balt 7 133.

92. EFFECT OF ACCELERATION ON CIRCULATORY AND RESPIRATORY FUNCTION: MAMMALS

agents used, depth of anesthesia, differences in strength and duration of acceleration patterns, and variations in the rates of acceleration. G = the acceleration of a freely falling body due to the attraction of gravity, expressed as the rate of increase of velocity per unit of time.

Variable (A)	Animal (B)	Effect (C)	Reference (D)
Positive (headward) Acceleration			
1 Blood pressure Arterial	Man	At 2.5 G, minimum systolic pressure for clear vision (eye level) = >50 mm Hg. At 3.5 G, minimum systolic pressure for dimming of vision (eye level) = 45 mm Hg. At 4.0 G, minimum systolic pressure for "grayout" (eye level) = 25 mm Hg. At 4.5 G, minimum systolic pressure for "black-out" (eye level) = <20 mm Hg. At 5.0 G, systolic pressure for unconsciousness (eye level) = 0.	1-3
2		Reflex blood pressure recovery due to carotid-sinus-induced vasoconstriction abolished by administration of tetraethylammonium ion.	4
3	Cat, monkey, anesthetized	Carotid pressure falls to zero at 3.4 G or more.	5,6
4	Dog; anesthetized	Carotid pressure falls 1-2 sec after onset of G; at heart level, after 3-8 sec lag Reflex blood pressure recovery in 9-20 sec. Vascular collapse in susceptible subjects.	7
5	Dog, monkey, unanesthetized	Inverse relation between carotid pressure and pulse rate (Marey's Law) well developed in primate, less perfect in dog Man and monkey have a functional advantage of 0.5-1.0 G over dog, because of greater circulatory shifts in dog.	8,9
6	Monkey, anesthetized	Carotid pressure falls with no lag Femoral pressure at first increased and then decreased below normal.	10
7 Venous	Man	Jugular bulb. subatmospheric pressures as great as -60 mm Hg maintain cerebral arterio-venous pressure gradient. Forearm and saphenous venoconstriction demonstrated by isolated venous segment technique.	11,12
8	Cat, monkey, anesthetized	Abdominal venous pressure obeys hydrostatic laws. In peripheral leg veins, pressure builds up slowly due to effect of valves. Parallel changes in sagittal sinus and subarachnoid pressure Jugular collapse.	6,13
9	Dog, anesthetized	Saphenous venoconstriction demonstrated with miniature balloon technique	14
10	Dog, monkey, unanesthetized	Jugular vein pressure falls 25-50%	8
11 Atrial	Cat, rabbit, anesthetized	Atrial pressure falls as much as 43 mm Hg below atmospheric; partially offset by fall in intrapleural pressure.	5,15
12 Renal	Goat, anesthetized	Arterial pressure decreases, venous pressure increases with reduced arterio-venous pressure gradient.	16
13 Heart rate and rhythm Rate	Man	Immediate increase to maximum values of 160-180 beats/min. depending on stress; marked vagal slowing following test, with secondary increase. Increase nearly independent of initial rate	4,7,17,18
14		(3 min or more).	19
15	Monkey, anesthetized	No change during 5-sec exposure	10

92 EFFECT OF ACCELERATION ON CIRCULATORY AND RESPIRATORY FUNCTION MAMMALS (Continued)

Variable (A)	Animal (B)	Effect (C)	Reference (D)
Positive (headward) Acceleration (continued)			
Heart rate and rhythm (concluded)	Rabbit; anesthetized	Decreases with long exposure	15
Rate (concluded)	Monkey, rabbit, unanesthetized	PR, QT, and cycle length consonant with heart rate change; R and S wave changes associated with changes in position of heart in chest; rarely T-wave changes (flattening, biphasic). Displacement of pacemaker during deceleration. In severe cases, atrioventricular block or ventricular rhythm. Electrical axis shifts 10-15° with G-exposure in monkeys.	7,17,19
ECG	Man	Marked sinus arrhythmia, ventricular extrasystoles, bundle branch block.	7,17
Arrhythmias	Man	Marked sinus arrhythmia, ventricular extrasystoles, bundle branch block.	7,17
Blood flow	Man	Estimated by X-ray marked reduction in cardiac shadow.	17
Cardiac output	Cat, monkey, anesthetized	Estimated by X-ray cinematography with thorotrast: decrease in systolic and diastolic heart shadow, lack of movement of heart contours (principal change is in ventricular length); decreased systolic and diastolic residual. Ultimately heart remains in contracted state	7
Cerebral circulation	Man	Cerebral venous O ₂ saturation preserved in spite of markedly diminished carotid arterial pressure (30 mm Hg), suggests compensatory effects maintain cerebral blood flow.	11
	Cat, monkey, anesthetized	Motion pictures of cerebral vessels through a Forbes window: blanching of cerebral surface, retention of blood (stasis) in large vessels, reactive hyperemia afterward	6
	Dog, monkey; unanesthetized	Venturimeter in carotid: in standard 10-sec run, flow reaches minimum in 4-6 sec, generally corresponding to carotid pressure. Flow = 0 at 3-4 G, may be negative at higher G, depression of flow proportional to $\Delta G/G$, rebound after run to 25% above normal	19
	Rabbit, anesthetized	Thermistor: decrease in carotid flow.	20
Retinal circulation	Man	Ophthalmoscopic examination. arteriolar pulsation followed by arteriolar exsanguination and collapse at blackout. Transient venous distension during recovery.	21
	Monkey, anesthetized	Direct cannulation of retinal artery: decrease in retinal blood pressure.	6
Pulmonary circulation	Man	Indirect evidence for pulmonary blood pooling. Changes in vital capacity demonstrate pulmonary blood reservoir function.	22,23
	Cat, monkey, anesthetized	X-ray cinematography: apical clearing, increased density at bases.	7
Ear lobe	Man	Ear opacity diminished (amount related to G level). Ear pulse: increased amplitude at low G; amplitude bears close relation to systolic pressure at head level when pressure is below 90 mm Hg.	2,3,17,18
Peripheral circulation	Man	Plethysmography: increase in volume of lower leg, 500 ml stored at normal temperature in thighs and buttocks during orthostasis (1 G), double this amount at elevated temperatures. Following administration of tetraethylammonium ion, increase in rate of pooling but no increase in amount pooled	17

92. EFFECT OF ACCELERATION ON CIRCULATORY AND RESPIRATORY FUNCTION. MAMMALS (Continued)

Variable (A)	Animal (B)	Effect (C)	Reference (D)
Positive (headward) Acceleration (concluded)			
32 Blood flow (concluded) Peripheral circula- tion (concluded)	Cat; anesthetized	X ray with contrast media: vascular engorgement of liver and lung bases and distensions of iliac and femoral veins.	24
33	Goat; anesthetized	Short-duration, high-magnitude G produces edema, congestion and hemorrhagic changes.	22
34 Blood constituents	Man	3.5-5.0 G for 3-5 min (blackout level) produces a fluid loss of 3.6-4.5 ml/100 ml blood. Humans less sensitive than animals to G-induced fluid loss.	25
35	Goat, anesthetized	Following onset of G, venous hemoconcentration occurs (red cell and plasma protein concentration increased 10-15% above control); arterial hemoconcentration (3-6%) lags venous changes 1.5-2.5 min; during G, protein leakage and red cell packing reported, increase in tissue pressure; rapid post-run dilution.	22
36	Cat, monkey, rat, anesthetized	Hyperglycemia without glycosuria; liver, heart and skeletal muscle glycogen values reduced, serum K and NaCl unaffected; no change in organ water content, blood cell volume increased 20%; blood volume decreased 2%.	24
37 Respiratory effects			
38 Rate of breathing	Man	Increases.	7,22
	Cat, rabbit; anesthetized	Decreases.	5,15
39	Dog, monkey; unanesthetized	Brief increase followed by decrease of 50% post-run increase of 5-10% over control	19
40 Depth of breathing	Man	Early respiratory amplitude large; later minute volume maintained by rate increase.	7,22
41	Cat; anesthetized	Shallow.	5
42	Dog, monkey; unanesthetized	Brief early increase in depth, then decrease.	19
43 Intrapleural pressure	Cat; anesthetized	Decreases.	5
44 Lung volumes	Man	Shift in midposition of chest due to visceral movement, increased end-expiratory volume and decreased vital capacity, changes in vital capacity related in part to fluctuation in pulmonary blood volume during orthostasis.	7,26,27
45 O ₂ intake	Man	Markedly decreased values	22
Negative (footward) Acceleration			
46 Blood pressure			
Arterial	Man	In absence of cardiac arrhythmias, radial arterial pressure (wrist at head level) increases 70-90 mm Hg at 3 negative G for 15 sec, increases rapidly at first, then gradually falls. (See line 64.)	28
47	Cat, anesthetized	Carotid pressure increases more than venous pressure.	13
48	Dog, anesthetized	Prolonged asystole results in gross fall in carotid arterial pressure, when asystole is abolished by vagotomy, arterial pressure increases	29
49	Goat, anesthetized	Carotid pressure increases At 12-15 negative G for 6-30 sec, arterial pressure may increase to 6 times normal.	29
50	Monkey, anesthetized	More resistant to arterial pressure effects of negative acceleration than other animals, response most nearly resembles that in man.	9
51 Venous	Man	Frontal vein pressure increases linearly with G due to hydrostatic column based at heart level Initial rapid increase followed by slow steady rise due to blood drainage from legs.	28-30

92. EFFECT OF ACCELERATION ON CIRCULATORY AND RESPIRATORY FUNCTION. MAMMALS (Continued)

Variable (A)	Animal (B)	Effect (C)	Reference (D)																				
Negative (footward) Acceleration (continued)																							
52 Blood pressure (concluded) Venous (concluded)	Cat; anesthetized	Jugular pressure directly and linearly related to cerebrospinal fluid pressure. Level of zero venous pressure change in jugular and cerebral spinal fluid located at diaphragm in cat and goat.	13,29																				
53	Dog, monkey, anesthetized	Jugular pressure more rapid, and greater but more irregular pressure changes occur under negative than under positive G.	8																				
54	Goat; anesthetized	Jugular: 12-15 negative G for 6-30 sec increases venous pressure to 400-500 mm Hg	29																				
55	Monkey, anesthetized	Jugular: Initial transient venous pressure fall before hydrostatic rise.	31																				
56 Heart rate and rhythm Rate	Man	Bradycardia in proportion to amount of negative G and its duration.	20,28,29																				
57	Dog, monkey; anesthetized	Heart rate varies inversely and specifically in relation to carotid sinus pressure changes, particularly in the primate.	8																				
58	Dog, goat, monkey, anesthetized	Bradycardia At 2-6 negative G for 5-15 sec, 15 monkeys showed no change, 16 monkeys an average decrease of 8 beats/min, and 41 monkeys an average increase of 3 beats/min. No significant change observed in dogs.	19,31																				
59 ECG	Man	Bradycardia, ectopic beats. Atrioventricular nodal rhythm with ectopic coupled beats. Prolonged PR interval proceeding to atrioventricular block, with increasing negative G or increased duration. Extrasystoles ventricular in origin, longest asystole = 9 sec.	28,29,32																				
60	Dog, monkey, anesthetized	Bradycardia, extrasystoles, sinus arrhythmia, large amplitude T waves. Signs of vagal heart block abolished by vagotomy.	28,29,33																				
61 Blood flow Cardiac output	Man	Estimated by X ray: no measurable cephalad displacement of heart at 3 negative G.	28																				
62	Dog, goat, anesthetized	Prolonged asystole in animals with intact cardiovascular reflexes results in fall in cardiac output. Peripheral reflex changes absent in goat.	29,31																				
63 Cerebral circulation	Man	Prolonged asystole results in cessation of cerebral blood flow similar to carotid sinus syncope.	29																				
64	Man, dog, goat, monkey, anesthetized	Changes in arterio-venous pressure gradient under negative G: <table border="1"> <thead> <tr> <th>Animal</th> <th>Initial Change</th> <th>Later Change</th> <th>Post-run</th> </tr> </thead> <tbody> <tr> <td>Man</td> <td>+</td> <td>-30%</td> <td>-65%</td> </tr> <tr> <td>Dog</td> <td>-</td> <td>-16%</td> <td>-60%</td> </tr> <tr> <td>Goat</td> <td>+</td> <td>+65%</td> <td>0%</td> </tr> <tr> <td>Monkey</td> <td>+</td> <td>-12%</td> <td>-40%</td> </tr> </tbody> </table> Transient post-run cerebral ischemia except in goat. Later changes in arterio-venous gradient attributed to carotid sinus reflex activity or decrease in effective blood volume and failure of cardiac filling.	Animal	Initial Change	Later Change	Post-run	Man	+	-30%	-65%	Dog	-	-16%	-60%	Goat	+	+65%	0%	Monkey	+	-12%	-40%	29,31
Animal	Initial Change	Later Change	Post-run																				
Man	+	-30%	-65%																				
Dog	-	-16%	-60%																				
Goat	+	+65%	0%																				
Monkey	+	-12%	-40%																				
65	Goat, anesthetized	Arterio-venous (carotid-jugular) pressure difference reported by other investigators as decreased. When increased, arterio-venous gradient not greater than that sustained by normal vessels, and no cerebral hemorrhages found in absence of asphyxia or extraneous trauma.	13,29,34,35																				
66	Monkey; anesthetized	Cerebral circulation time increased 30% (photo-fluorographic technique); evidence for continued circulation at 12 negative G for 40 sec	33																				
67 Pulmonary circulation	Man, monkey	X ray with contrast media: increased opacity at lung apexes associated with fall in arterial O ₂ saturation after several sec under negative G.	29																				

92. EFFECT OF ACCELERATION ON CIRCULATION AND RESPIRATORY FUNCTION. MAMMALS (Continued)

Variable (A)		Animal (B)	Effect (C)	Reference (D)
Negative (footward) Acceleration (concluded)				
68	Blood flow (concluded)			
	Renal circulation	Goat; anesthetized	Both arterial and venous renal pressures decrease.	16
69	Peripheral circulation	Man	Occlusive cuffs at thighs decrease venous pressure rise in frontal vein during negative G.	29
70		Dog; anesthetized	Petechial threshold. 11 negative G for 0.2 sec, 7 negative G for 1 sec, 3 negative G for 3-4 sec.	36
71		Goat; anesthetized	Boggy, pitting edema of head and neck (5 negative G for 5-10 sec), swelling of tongue, edema of pharynx and upper trachea which may lead to asphyxia; conjunctival, sinus, anterior eye chamber and middle ear hemorrhages.	29
72	Blood constituents	Goat, anesthetized	5 negative G for 15-30 sec post-run hematocrit peak = 118% of control, plasma protein peak = 107% of control, fluid loss = 15 ml/100 ml blood; protein loss = 1 g/100 ml blood; short rapid concentration phase followed by slower dilution phase.	35
73			Progressive stagnation of cerebral blood flow with repeated exposures as evidenced by anaerobic metabolic changes, such as initial elevation followed by depression of blood glucose, increased arterio-venous O ₂ content difference (decreased venous O ₂ content, constant arterial O ₂ content); reduction in arterial and venous CO ₂ content, increase in lactic acid and pyruvic acid content; increasing lactate-pyruvate ratio.	34
74		Monkey, anesthetized	Six animals, 12 negative G for 40 sec. arterial O ₂ fell from 13 vol % to 6.1 vol %, venous O ₂ (confluens sinuum) fell from 7.7 vol % to 2.4 vol %; arterio-venous O ₂ difference not significantly changed; arterio-venous CO ₂ difference not significantly changed.	33
75	Respiratory effects			
76	Rate of breathing	Man	Slight increase.	26
		Monkey, anesthetized	12 negative G for 40 sec: apnea with sometimes incomplete inspiratory gasps.	33
77	Tidal volume	Man	Decreases.	26
78	Lung volume	Man	Decreases (measured at end-expiration).	26
79	Vital capacity	Man	Decreases.	26

Contributors. (a) Lawton, Richard W., and Herbert G. Shepler, (b) Bondurant, Stuart, and George D. Zuidema

References. [1] Lambert, E. H., and E. H. Wood. 1946. Med. Clin. N. America 30:833. [2] Wood, E. H., E. H. Lambert, E. J. Baldes, and C. F. Code. 1946. Fed. Proc., Balt. 5:327. [3] Wood, E. H., E. H. Lambert, and C. F. Code. 1947. J. Aviat. Med. 18:471. [4] Brown, G. E., Jr., E. H. Wood, and E. H. Lambert. 1949. J. Appl. Physiol. 2:117. [5] Greenfield, A. D. M. 1945. J. Physiol., Lond. 104:5P. [6] Jasper, H. H., and A. J. Cipriani. 1945. Ibid. 104:6P. [7] Gauer, O. 1950. German Aviation Medicine, World War II. Dep. of the Air Force. v. 1, chap. VI-B. [8] Britton, S. W. 1949. Am. J. Physiol. 156:1. [9] Pertzoff, V. A., and S. W. Britton. 1948. Ibid. 152:492. [10] Lawton, R. W., L. C. Green, G. H. Kydd, L. H. Peterson, and R. J. Crosbie. 1956. Aviation Medical Acceleration Laboratory. U. S. Naval Air Development Center, Johnsville, Pa. MA-5611. [11] Henry, J. P., O. H. Gauer, S. S. Kety, and K. Kamer. 1951. J. Clin. Invest. 30:292. [12] Hiett, E. P., S. D. Leverett, and S. Bondurant. 1958. Fed. Proc., Balt. 17:70. [13] Rushmer, R. F., E. L. Beckman, and D. Lee. 1947. Am. J. Physiol. 151:355. [14] Salzman, E. W., and S. D. Leverett. 1956. Circul. Res., N. Y. 4:540. [15] Stauffer, F. R., E. L. Beckman, and J. I. Thorn. 1949. U. S. Naval School of Aviation Medicine. Pensacola, Fla. Proj. No.

- NM-001-048. Rep. No. 1. [16] Ames, W. W., S. Rosenfeld, and C. F. Lombard. 1951. *J. Appl. Physiol.* 3:399.
- [17] Franks, W. R., W. K. Kerr, and B. Rose. 1945. *J. Physiol., Lond.* 104:9P. [18] Lambert, E. H. 1949. *J. Aviat. M.* 20:308. [19] Britton, S. W., V. A. Pertzoff, C. R. French, and R. F. Kline. 1947. *Am. J. Physiol.* 150:7.
- [20] Jongbloed, J., and A. K. Noyons. 1933. *Pflügers Arch.* 233:67. [21] Duane, T. D. 1954. *Arch. Ophth., Chic.* 51:343. [22] Raulston, B. O., and C. F. Lombard. 1951. ONR. U. S. Navy, Dep. of Aviation Medicine. University of Southern California. Final Rep. Contract N6ori77. Proj. NR-161-014, Task 1. [23] Henry, J. P. 1955. Orthostasis and the kidney. WADC Tech. Rep. No. 55-478, WPAFB [24] Britton, S. W., E. L. Corey, and G. A. Stewart. 1946. *Am. J. Physiol.* 146:33. [25] Clark, W. G., I. D. R. Gardiner, A. K. McIntyre, and H. Jorgenson. 1945. *Nat. Res. Council, Canada. Rep. No. 468.* [26] Lombard, C. F., H. P. Roth, and D. R. Drury. 1948. *J. Aviat. M.* 19:355. [27] Morrison, R. B., and J. L. Patterson, Jr. 1946. U. S. Naval School of Aviation Medicine, Pensacola, Fla. Rep. No. X-723 [AV-379-R]. [28] Gamble, J. L., Jr., R. S. Shaw, J. P. Henry, and O. H. Gauer. 1949. *J. Appl. Physiol.* 2:133 [29] Henry, J. P. 1950. Air Force Tech. Rep. No. 5953, WPAFB.
- [30] Shaw, R. S., J. P. Henry, J. L. Gamble, and O. Gauer. 1948. *J. Appl. Physiol.* 1:441. [31] Rosenfeld, S., and C. F. Lombard. 1950. *J. Aviat. M.* 21:293. [32] Ryan, E. A., W. K. Kerr, and W. R. Franks. 1950. *Ibid.* 21:173. [33] Duane, T. D., R. L. Wechsler, J. E. Zeigler, and E. L. Beckman. 1952. *Ibid.* 23:479. [34] Pogrud, R. S., S. W. Ames, and C. F. Lombard. 1951. *Ibid.* 22:50. [35] Stauffer, F. R., and C. Hyman. 1948. *Am. J. Physiol.* 153:64. [36] Gamble, J. L., and R. S. Shaw. 1947. Mem. Rep. TSEAA-695-74B, WPAFB.

Abbreviations and definitions: kv = kilovolt, a unit of electrical potential equal to 1000 volts; kvp = kilovolt peak, the crest value of the potential wave in air produces, in air, ions carrying one electrostatic unit of electrical charge of either sign, rem = roentgen equivalent for man (or mammal), the absorbed dose equal to the dose in rads multiplied by the RBE; rad = rad unit, equivalent to 100 ergs/g energy absorption, RBE = relative biological effectiveness, i.e., the appropriate value of the biological effectiveness of the radiation in question relative to that of X radiation with an average specific ionization of 100 ion pairs per micron of water, for the particular biological system and biological effect under consideration and for the condition under which the radiation is received; rep = roentgen equivalent physical, equivalent to 93 ergs/g energy absorption; n = neutron, a nuclear particle of zero charge and mass number 1; mc = millicurie, the quantity of radionuclide disintegrating at the rate of 3.7×10^{10} atoms per second, μ c = microcurie, 3.7×10^4 disintegrations per second.

Animal	Exposure or Administration	Type of Radiation	Dose	Accumulated Dosage or Exposure Time	Effect	Initial Symptom Manifestation	Reference
1 Man	Entire body, single exposure (accident)	Mixed X ray 80 kv equivalent r plus gamma	480 r		Decreased blood pressure. Tachycardia.	24 hr (G)	1
	Hands, single exposure (accident)	X ray 80 kv	110 r		Cardiac hypertrophy. Increased heart rate.	15th da	
	Entire body, single exposure (accident)	equivalent r	25,000-55,000 rem		Endocardial fibrosis and pericarditis.	16th da	
2	Entire body, single exposure (accident)	Mixed X ray 80 kv	1930 r		Hypotension. Circulatory collapse and myocardial hemorrhage.	22nd da	
	Hands, single exposure (accident)	equivalent r plus gamma	114 r			24th da (last da of life)	
3	Entire body, single exposure (accident)	Mixed X ray 80 kv	3,000-30,000 rem			14 hr	
4	Entire body, single exposure (accident)	equivalent r	390 r		Temporary fall in blood pressure.	9th da (last da of life)	
5	Entire body, single exposure (military)	Atomic bomb (Japan, 1945)	264 r			14 hr	
6	Entire body, single exposure (military)	Atomic bomb (Japan, 1945)	Lethal				
7	Entire body, repeated exposure (therapy)	X ray 200 kv	10-25 r		Epicardial hemorrhage, endocardial extravasation.	Terminal 3-6 wk	2
8	Thorax, repeated exposure (therapy)	X ray 250 kv	40-100 r		Epicardial petechiae, perivascular edema of coronary vessels.	1st wk	3
	Thorax, repeated exposure (therapy)	X ray	4500 r		Expansion of hemorrhage. Reduced intracellular fluid.	2 mo	4
			52 da		Interstitial fluid and plasma volume.		
			7 da		Tachycardia.		
			175-250 r/da		Right heart failure, hypertrophy of right side of heart, pericarditis.	1-6 mo	5
							6

9	Thorax, repeated exposure (therapy)	X ray	Intensive			T waves ² lowered and inverted, myocardial degeneration.	7
10	Thorax, repeated exposure (therapy)	X ray	Intensive			Premature atrial and ventricular beats, myocardial degeneration.	8
11	Thorax, repeated exposure (therapy)	X ray	60 r		540 r	Scarring of myocardium.	9
12	Thorax, repeated exposure (therapy)	X ray	200 kv		5 mμ	Increased blood pressure, tachycardia, accentuated aortic second sound, telangiectasia of skin in irradiated of abdomen	10
13	Left upper quadrant of abdomen (therapy)	X ray	200 kv		3000 r	Increased blood pressure, tachycardia, telangiectasia of skin in irradiated of abdomen	11
14	Left renal region, repeated exposure (therapy)	X ray	200 kv		3000 r	Bounding pulse, increased blood pressure, malignant hypertension.	12
15	Kidney region, repeated exposure (therapy)	X ray	200 kv		4700 r	Subendothelial proliferation of fibrous connective tissue in walls of medium and small renal arteries.	13
16	Head, single and repeated exposure (therapy)	X ray	250 kv		2300 r	Hypertension.	14
17	Tongue (therapy)	X ray	8250 r		3 wk	Collagenous thickening, fibrinoid necrosis and thrombosis of smaller blood vessels of brain	15
18	Skin (forearm), single exposure	X ray	300 r		2300 r	Sudden hypertension.	16
19	Skin, single exposure	X ray	1000 r or more		2300-10,500 r	Capillary dilatation.	17
20	Skin, single exposure	X ray	400 r soft rays, 525 r medium rays, 600 r hard rays			Erythema.	18
21	Skin, single exposure	X ray	175-1 skin unit dose			Edema.	19
22	Finger nail fold area, single exposure	X ray	200 r			Inflammation.	20
23	Inhalation, repeated exposure (mining)	Radium	60 μg			Telangiectasia.	21
24	Myocardium (accident)	Radium	2 mg			Erythema.	22
25	Pelvis, repeated exposure (therapy)	Radium	600-1000 rad			Fibrosis of media of pulmonary artery.	23
26	Skin, single exposure	Beta	170-250 rep			Myocardial degeneration.	24
27	Forearm, single exposure	Beta	140 rep			Myocardial degeneration and necrosis.	25

1/1 Terminal refers here to period shortly before death when animal is moribund. 1/2 Slow final wave of the ventricular complex resulting in restitution of the resting state in ventricles.

93. HEMODYNAMIC EFFECTS OF IONIZING RADIATION: VERTEBRATES (Continued)

Animal	Exposure or Administration	Type of Radiation	Dose	Accumulated Dosage or Exposure Time	Effect	Initial Symptom Manifestation	Reference
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)
28 Bat	Entire body, single exposure	X ray 250 kv	10,000-60,000 r		WBC adherence to endothelium, intra-vascular clumping of RBC, slowing of blood flow.	(G)	26
29	Wing, local, single exposure	X ray 50 kv	50,000-5,000,000 r		WBC adherence to endothelium, intra-vascular clumping of RBC, slowing of blood flow.		
30 Burro	Entire body, single exposure (external)	Gamma Zirconium ⁹⁵ Niobium ⁹⁵	500-900 r		Increased capillary fragility.	3-4 wk	27
31	Entire body, single exposure (external)	Tantalum ¹⁸²	200 r 400 r		Hemoconcentration.		28
32	Entire body, single exposure (external)	Gamma Tantalum ¹⁸² Cobalt ⁶⁰	780 r (LD-50/30) 710 r		Venous pulsations, accelerated pulse rate (arterial).	Terminal	29
33	Entire body, repeated exposure (external)	Gamma Cobalt ⁶⁰	25 r/da (LD-50/20)	50-75 da	T wave ² inversion, alteration of QRS complex ³ , increased heart rate.		30
34 Cat	Abdomen, single exposure (under anesthesia)	X ray 260 kvp	5000 r		Increased carotid blood pressure.	10 min-2 hr	31
35	Thorax, Abdomen, entire body, single exposure (under anesthesia)	X ray	Unfiltered X ray of medium hardness		Decreased carotid blood pressure, cardiac failure.	Within min	32
36 Dog	Entire body, single exposure	X ray 2 mev	600 r (LD-100/30)		Subepicardial petechiae, ecchymosis, hemorrhage, fibrous anginous pericarditis.	Terminal	33
37	Entire body, single exposure	X ray 2 mev	1800 r		Dehydration.	3-4 da	34
38	Entire body, single exposure	X ray 2 mev	2000-3000 r		Vascular collapse.	Terminal	35
39	Entire body, single exposure	X ray 250 kvp	250 r, 500 r		Increased heart rate, decreased blood pressure, hemoconcentration.	Terminal (3 da)	36
40	Entire body, single exposure	X ray 250 kv	350 r		Abnormal reactivity of isolated segments of carotid arteries.		37
41	Entire body, single exposure	X ray 250 kv	350 r		Increased capillary permeability.		38
42	Entire body, single exposure	X ray 250 kv	450-550 r		Increased vascular fragility.		39
43	Entire body, single exposure	X ray 250 kv	Massive exposure		Hemorrhage, plasma volume increased, total body water, blood volume decreased.	6-16 da	40
					Inversion of RBC into lymph.		

93. HEMODYNAMIC EFFECTS OF IONIZING RADIATION: VERTEBRATES (Continued)

Animal	Exposure or Administration	Type of Radiation	Dose	Accumulated Dose or Exposure Time	Effect	Initial Symptom Manifestation ¹	Reference
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)
28 Rat	Entire body, single exposure	X ray 250 kv	10,000-60,000 r		WBC adherence to endothelium, intra-vascular clumping of RBC, slowing of blood flow.	(G)	26
29	Wing, local, single exposure	X ray 50 kv	50,000-5,000,000 r		WBC adherence to endothelium, intra-vascular clumping of RBC, slowing of blood flow.		
30 Burro	Entire body, single exposure (external)	Gamma Zirconium ⁹⁵ Niobium ⁹⁵	500-900 r		Increased capillary fragility.	3-4 wk	27
31	Entire body, single exposure (external)	Gamma Tantalum ¹⁸²	200 r		Hemodilution.		
32	Entire body, single exposure (external)	Gamma Tantalum ¹⁸² Cobalt ⁶⁰	400 r		Hemoconcentration.		28
33	Entire body, single exposure (external)	Gamma Tantalum ¹⁸² Cobalt ⁶⁰	760 r (LD-50/30) 710 r		Venous pulsations, accelerated pulse rate (arterial).	Terminal	29
34 Cat	Entire body, repeated exposure (external)	Gamma Cobalt ⁶⁰	25 r/da	50-75 da	T wave ² inversion, alteration of QRS complex ³ , increased heart rate.		30
35	Abdomen, single exposure (under anesthesia)	X ray 260 kvp	5000 r		Increased carotid blood pressure.	10 min-2 hr	31
36 Dog	Thorax, abdomen, entire body, single exposure (under anesthesia)	X ray	Unfiltered X ray of medium hardness		Decreased carotid blood pressure, cardiac failure.	Within min	32
37	Entire body, single exposure	X ray 2 mev	600 r (LD-100/30)		Subepicardial petechiae, ecchymosis, hemorrhage, fibrous anguineous pericarditis.	Terminal	33
38	Entire body, single exposure	X ray 2 mev	1800 r		Dehydration.	3-4 da	34
39	Entire body, single exposure	X ray 2 mev	2000-3000 r		Vascular collapse.	Terminal	35
40	Entire body, single exposure	X ray 250 kvp	250 r, 500 r		Increased heart rate, decreased blood pressure, hemoconcentration.	Terminal (3 da)	36
41	Entire body, single exposure	X ray 250 kv	350 r		Abnormal reactivity of isolated segments of carotid arteries.		37
42	Entire body, single exposure	X ray 250 kv	350 r		Increased capillary permeability.		38
43	Entire body, single exposure	X ray 250 kv	450-550 r		Increased vascular fragility.		39
	Entire body, single exposure	X ray 250 kv	Massive exposure		Hemorrhage, plasma volume increased, total body water, blood volume decreased.	5-16 da	40
					Dilation of RBC into lymph.		

74	Entire body, repeated exposure	X ray 200 kv	(60 r at 15-da intervals)	900 r total	Cardiac hemorrhage.	67
75	Entire body, single exposure (external)	Barium 140 Lanthanum 140	2500 R (1000 R/min)		Blood pressure depressed, then increased	68
76	Entire body, single exposure	X ray 250 kv	410 r		Increased amount of ground substance in tunica media of arteries.	69
77	Entire body, single exposure	X ray 250 kv	575 r (LD-50/30)		Decreased blood volume (plasma and cell), increased capillary permeability.	70, 71
78	Entire body, single exposure	X ray 250 kv	575 r		Increased vascular fragility and capillary permeability.	72
79	Entire body, single exposure	X ray 250 kv	575 r		Decreased RBC mass.	73
80	Entire body, single exposure	X ray 250 kv	575 r		Decreased plasma volume.	74
81	Entire body, single exposure	X ray 250 kv	575 r		Blood perfusion through liver allowed.	75
82	Entire body, single exposure	X ray 250 kv	575 r		Walls of blood vessels thinned.	76
83	Entire body, single exposure	X ray 250 kv	575 r		Disappearance of tunica adventitia.	77, 78
84	Entire body, single exposure	X ray 250 kv	575 r		Inflammation of eyelids.	79
85	Entire body, single exposure	X ray 250 kv	575 r		Erythema on ears and feet.	80
86	Entire body, single exposure	X ray 250 kv	575 r		Hypertrophy of myocardium.	81
87	Entire body, single exposure	X ray 250 kv	575 r		Swelling of myocardium.	82
88	Entire body, single exposure	X ray 250 kv	575 r		Hypotension, circulatory collapse.	83
89	Entire body, single exposure	X ray 250 kv	575 r		Plasma volume decreased.	84
90	Entire body, single exposure	X ray 250 kv	575 r		Plasma volume increased.	85
91	Entire body, single exposure	X ray 250 kv	575 r		Slight increase in plasma volume.	86
92	Entire body, single exposure	X ray 250 kv	575 r		Increased vascular fragility and capillary permeability.	87
93	Entire body, single exposure	X ray 250 kv	575 r		Decreased blood volume, increased capillary permeability.	88
94	Entire body, single exposure	X ray 250 kv	575 r		Decreased plasma volume followed by increase.	89
95	Entire body, single exposure	X ray 250 kv	575 r		RBC mass decreased.	90
96	Entire body, single exposure	X ray 250 kv	575 r		Decreased blood pressure followed by compensatory rise.	91
97	Entire body, single exposure	X ray 250 kv	575 r		Transient decrease in blood pressure.	92
98	Entire body, single exposure	X ray 250 kv	575 r		Increased capillary permeability and number of patent blood vessels.	93
99	Entire body, single exposure	X ray 250 kv	575 r		Peripheral vascular collapse.	94
100	Entire body, single exposure	X ray 250 kv	575 r		Decreased peripheral circulation, constriction of peripheral blood vessels.	95

1/1 Terminal refers here to period shortly before death when animal is moribund. 1/2 Slow final wave of the ventricular complex resulting in restitution of the resting state in ventricles.

Animal	Exposure or Administration	Type of Radiation	Dose	Accumulated Dosage or Exposure Time	Effect	Initial Symptom Manifestation	Reference
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)
60 Dog (concluded)	Intravenous, 3 injections	Colloidal gold	65,000 equivalent β r to liver	6 wk	Disappearance of endothelium, and marked increase in connective tissue surrounding hepatic blood vessels.	Terminal 62-119 da	55
61	Sino-atrial node implantation	Gamma Radon	0.6-5 mc	8 hr-8 mo	Engorgement and then rupture of wall of small vessels of region, occlusion of vasa vasorum, myocardial degeneration.		56
62	Intramuscular, single injection	Beta Strontium 89	Lethal doses (LD-50/30 = 1.0 μ c/g of body weight, or 0.5 μ c/g of body weight retained dose for 10 da)		Increased plasma volume. Increased heart rate.	Terminal	57
63	Intravenous, single injection	Alpha Plutonium 239	Lethal dose (LD-50/30 = 0.3 μ c/g of body weight)		Increased heart rate and plasma volume.	3-4 da before death	58
64	Entire body, repeated exposure	Neutron cyclotron		400 n over 4 da period	Decreased venous pressure; weak, rapid heart action; myocardial damage.	Terminal	59
65	Entire body, repeated exposure	Neutron	1.7 n/da	312 da	Myocardial hemorrhage.		60
66 Guinea pig	Entire body, single exposure	X ray	2000 r		Bradycardia.		61
67	Entire body, head only or body with head shielded, single exposure	X ray 200 kv	50 r/min	25,000 r	Decreased heart rate, ventricular extrasystole, T wave 2 inversion.		62
68	Thorax, abdomen, entire body, single exposure	X ray	Unfiltered X ray of medium hardness		Decreased blood pressure, cardiac failure.	Within min	32
69 Hamster	Entire body, single exposure	X ray 130 kv	1000-1500 r		Bradycardia, vasoconstriction, T waves 2 depressed and inverted terminally.		63
70	Abdomen, single exposure	X ray 200 kv	2000 r		Decreased blood volume.	3rd da	64
71	Cheek pouch implantation, repeated exposure	Beta Strontium 90	140 μ c		Increased vascularity, thrombo-embolism, petechiae, edema (of cheek pouch).	24 hr	65
72	Cheek pouch, single exposure	X ray	15,000-20,000 r		Increased vascularity, thrombo-embolism, petechiae.		
73 Monkey	Entire body, single exposure	X ray	400-500 r		Expanded plasma volume.		66

107	Rat	Entire body, single exposure	X ray 250 kvp	400-600 r 750 r	Plasma volume increased. Plasma volume decreased 20%, blood volume decreased 12%. Plasma and blood volumes increased 0-20%. Plasma volume increased 43% 1-3 hr and 4-6 da Decreased blood pressure.	4-10 da 72 hr 96 hr 10-20 da 1-3 hr and 4-6 da 2nd da	96
108		Entire body, single exposure	X ray 250 kvp	600 r	Vasodpressor material (VDM), decreased blood flow.	2nd da	98
109		Entire body or abdomen, single exposure	X ray 250 kvp	600 r	Peripheral vascular paralysis. Vasodictor material (VEM), increased blood flow.	1st wk 2nd wk	
110		Entire body, single exposure	X ray 250 kvp	650-750 r	Increased capillary permeability.		37
111		Entire body, single exposure	X ray 250 kvp	Massive	Diversion of RUC into lymph.	Immediate	40
112		Entire body, single exposure	X ray 250 kvp	600 r	Plasma volume increased.	10-8 da	99
113		Entire body, single exposure	X ray 220 kv	600 r (LD-50/30)	Plasma volume decreased.	10th da	100
114		Entire body, single exposure	X ray 200 kvp	700 r	VDM - vasoconstriction. VEM - vasoconstriction. Increased plasma volume and extracellular fluid volume.	3, 6, 18 da 10th da 2nd and 20th da	101
115		Intestine shielded, entire body, single exposure	X ray 200 kv	1000 r	Decreased blood and plasma volumes.	4-6 da	102
116		Entire body, single exposure	X ray 140 kv	700 r (LD-50/30)	Increased venous pressure. Decreased venous pressure. Increased plasma volume. Decreased plasma volume. Decreased pulse rate. Cardiac arrhythmia.	3 hr and 5-12 da 2-4 da 3-12 hr 12 hr-3 da 4-8 da 8-10 da	103
117		Entire body, single exposure	X ray 140 kv	700 r (LD-50/30)	Cardiac failure. Increased venous pressure, length and amplitude of 1st heart sound; irregular heart beat; increase in height of T wave.	Terminal 12 hr and 5-12 da	104
118		Entire body, anoxia, single exposure	X ray 250 kvp	600-1400 r	Increased plasma volume.	6-10 da 3 mo-1 yr	105
119		Entire body, anoxia, single exposure	X ray 250 kvp	800 r	Hypertension.	1-1 1/2 yr	106
120		Thorax, single exposure	X ray 200 kv	1500 r 3000 r 7500 r 10,000 r	Pulmonary capillary engorgement. "Fibby" myocardium. Round cell infiltration in myocardium. Thickened coronary arteries. Myocardial edema and congestion with capillary engorgement and hemorrhage, coronary bed dilated.	8th da	107

/1/ Terminal refers here to period shortly before death when animal is moribund. /2/ Slow final wave of the ventricular complex resulting in restitution of the resting state in ventricles

93. HEMODYNAMIC EFFECTS OF IONIZING RADIATION: VERTEBRATES (Continued)

Animal	Exposure or Administration	Type of Radiation	Dose	Accumulated Dosage or Exposure Time	Effect	Initial Symptom Manifestation	Reference
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)
91 Rabbit (concluded)	Entire body, single exposure	X ray 180 kv	1250 r		Blood pressure fell after short period of fluctuation.		83
92	Entire body, single exposure (1 ear shielded)	X ray 250 kv	1500 r		Circulating blood volume decreased.	2 hr	
93	Abdominal skin, single exposure	X ray 20 kv	6000 r		Increased blood viscosity.	2-3 hr	84
94	Localized small areas of skin on side and back, single exposure	X ray 86 kv	680-2720 r		Shielded ear - sludging of RBC. Unshielded ear - WBC adhered to endothelium, interstitial WBC exudation.	Immediate	85
95	Head, single exposure	X ray 260 kv	14,000 r (240 r/min)		Blood pressure rose then fell during irradiation (later returned to normal).	1 hr	86
96	Head, single exposure	X ray 200 kv	2850 r		Capillary dilatation, increased capillary permeability.	30 min	87
97	Ear, single exposure	X ray	10,000 r		Minute hemorrhage in walls of blood vessels of brain.	100 da	88
98	Ear, single exposure	X ray 30 kv	1000 r		Diffuse redness surrounding blood vessels in ear.	1 hr	89
99	Thorax, single and repeated exposure	X ray	2000 r		Erythema.	5 da	90
100	Thorax, repeated exposure	X ray 200 kv 130 kv	600 r, 300 r at monthly intervals	1200 r	Erythema.	30 da	48
101	Isolated ear, single exposure	X ray 260 kv	1000-8000 r		Proliferation of connective tissue around coronary vessels, increased thickness of wall of right atrium, numerous petechiae in wall of right atrium.	8-32 da after last exposure	91
102	Head, single exposure (external)	Gamma Cobalt-60	1000 r		Zenker's necrosis in myocardium.	Immediate but transient	92,93
103	Total body surface, single exposure (external)	Beta Phosphorus-32	1 3-25 x 10 ³ rep		Rate of blood flow decreased, then returned to normal.	Decrease in frequency of cardiac contractions.	54
104	Skin, single exposure	Beta Phosphorus-32	1000 rep		Inflammation of eyelids, erythema on ears and feet.	2nd wk	76
105	Ear, single exposure	Beta Phosphorus-32	3000 rep		Erythema.	6-10 da	94
106	Skin of back, single exposure	Beta Phosphorus-32	5000 rep		Erythema.	18 da	90
		Phosphorus-32			Erythema.	3-5 da	95

137	Chick	Entire body, single exposure	X ray 200 kv	1000 r	<30 min	Hypotension.	Usually within 2 hr, always within 6 hr	120
-----	-------	------------------------------	--------------	--------	---------	--------------	---	-----

1/1 Terminal refers here to period shortly before death when animal is moribund.

Contributor: Michelson, Sol M.

- References** [1] Hempelmann, L. H., et al. 1952. *Ann Int. M.* 36:279. [2] Leibow, A. A., et al. 1949. *Am. J. Path.* 25:653. [3] Bull, U. S. Army M. Dep 1946. 6:350. [4] Greenberg, J., and D. Laaslo 1955. *J. Clin. Invest.* 34:405. [5] Cosgriff, S. W., and M. M. Kligerman. 1951. *Radiology* 57:536. [6] Fried, J. R., and H. Goldberg. 1940. *Am. J. Roentg* 43:877. [7] Hartman, F. W., et al. 1927. *Bull. Johns Hopkins Hosp* 41:36. [8] Emery, E. S., and B. Gordon, Jr. 1925. *Am. J. M. Sc.* 170:889. [9] Levy, R. L., and R. Golden. 1937. *Ibid* 194:597. [10] Dean, A. L., and J. C. Abels. 1944. *J. Urol.* 51:497. [11] Levitt, W. M., and S. Oram 1956. *Brit. M. J.* 2:910. [12] Davey, P. W., et al. 1952. *Canad. M. Ass. J.* 67:648. [13] Paterson, R. 1952. *J. Fac. Radiologists.* Lond. 3:270. [14] Penzabacker, J., and D. S. Russell. 1948. *J. Neur., Lond.* 11:183. [15] Dotti, G. 1949. *Radioterapia, Bologna* 2:253. [16] Pendergrass, E. P., et al. 1944. *Am. J. Roentg.* 52:123. [17] Borak, J. 1942. *Radiology* 58:481. [18] Warren, S. 1942. *Arch. Path.* 43:340. [19] Pohle, E. A. 1926. *Radiology* 6:236. [20] Nickson, M. 1947. *USAECD MDDC-1269.* [21] Rajewsky, B. 1939. *Radiology* 32:57. [22] Rose, J. M. 1932. *J. Path. Bact.* Lond. 35:899. [23] Sheehan, J. F. 1949. *Am. J. Clin. Path.* 19:30. [24] Witten, V. H., et al. 1954. *J. Invest. Derm.* 23:271. [25] Wirth, J. B., and J. R. Raper 1951. *In R. E. Zirkle, et al. Biological effects of external beta radiation.* NNEC 22-E. Chap. 12. [26] Smith, D. E., et al. 1951. *Physiol. Zool.* 24:249. [27] Agr. Res. Prog. Semi-Ann. Prog. Rep. ORO-145. 1955. *University of Tennessee, Knoxville.* [28] Rust, J. H., et al. 1953. *Radiology* 60:579. [29] Trum, D. F. 1955. *Mil. Surgeon* 112:335. [30] Rust, J. H., et al. 1955. *Rad. Res.* 2:475. [31] Brooks, P. M., and E. B. Konecni. 1954. *Fed. Proc.* 13:118. [32] Swann, M. B. R. 1924. *Brit. J. Radiol.* 29:195. [33] Gleiser, C. A. 1954. *Am. J. Vet. Res.* 15:329. [34] Conrad, R. A., et al. 1956. *Proc. Rad. Res. Soc.* p. 12. [35] Brecher, G., and E. P. Cronkite. 1951. *Am. J. Path.* 22:676. [36] Mason, W. B., et al. 1953. *USAECD UR-278.* [37] Bigelow, R. E., et al. 1953. *Proc. Soc. Exp. Biol., N. Y.* 76:734. [38] Field, J. D., and P. E. Rakers 1949. *Am. J. M. Sc.* 218:1. [39] McDonald, R. E., et al. 1953. *Am. Surgeon* 19:587. [40] Rose, M. H., et al. 1952. *Blood.* N. Y. 7:417. [41] Prosser, C. L., et al. 1947. *USAECD MDDC-1272.* [42] Painter, E. E., and A. M. Bruce. 1949. *N. England J. M.* 240:871. [43] Muntz, J. A., et al. 1949. *Arch. Biochem.* N. Y. 23:434. [44] Soberman, R. J., et al. 1951. *Am. J. Physiol.* 164:450. [45] Haley, T., et al. 1952. *USAECD UCLA-180.* [46] Skvortsov-Lukin, L. 1955. Thesis. Faculty of Pure Science, Col. University. [47] Prosser, C. L., and M. C. Moore. 1946. *USAECD MDDC-1271.* [48] Davis, K. S. 1924. *Radiology* 3:301. [49] Hartman, F. W., et al. 1926. *Am. J. M. Sc.* 172:487. [50] Moon, V. H., et al. 1940. *Arch. Path., Chic.* 29:723. [51] Moon, V. H., et al. 1941. *J. Am. M. Ass.* 116:489. [52] Kaplan, M. T., and D. N. Mohlitsky. 1932. *Frankf. Zschr. Path.* 43:434. [53] Lyman, R. S., et al. 1931. *Arch. Neur. Psychiat., Chic.* 29:556. [54] Arkusky, Y. I., and M. M. Mintz. 1938. *Bull. Roentg. Path.* 20:38. [55] Hahn, P. F., et al. 1931. *Science* 114:303. [56] Borman, M. C. 1927. *Am. J. Roentg.* 10:111. [57] Swift, M. N., et al. 1946. *USAECD MDDC-1388.* [58] Painter, E. E., et al. 1946. *USAECD-2042.* [59] Rose, M. H., and J. G. Ely. 1947. *In E. McDonald, ed. Neutron effects on animals.* Williams and Wilkins, Baltimore. Chap. 19. [60] Ely, J. O., et al. 1954. *In H. A. Blair, ed. Biological effects of external radiation.* NNEC V1-2. McGraw-Hill, New York. Chap. 17, p. 419. [61] Brace, K. C., and H. L. Andrews. 1955. *Proc. Internat. Conf. Peaceful Uses Atomic Energy, Geneva 1955.* United Nations, New York. [62] Andrews, H. L., and K. C. Brace. 1953. *Am. J. Physiol.* 175:138. [63] Fulton, G. P., and F. P. Sucka. 1954. *Ibid.* 179:135. [64] Swift, M. N., and S. T. Taketa. 1957. *Fed. Proc., Balt.* 16:126. [65] Fulton, G. P., et al. 1953. *Anat. Rec.* 115:446. [66] Ferrell, C. B., et al. 1954. *Fed. Proc., Balt.* 13:43. [67] Moon, L. E., et al. 1955. *Science* 122:663. [68] Brooks, P. M. 1956. *Rad. Res.* 4:206. [69] Smith, C., and L. A. Lowenthal. 1950. *Proc. Soc. Exp. Biol., N. Y.* 75:859. [70] Furth, J., et al. 1951. *South. M. J.* 44:85. [71] Furth, J., et al. 1951. *USAECD ORNL-878.* [72] Wish, L., et al. 1952. *Am. J. Roentg.* 67:628. [73] Supplee, H., et al. 1951. *USNRDL AD 318(B).* [74] Dobson, E. L. 1950. Thesis. University of California,

93. HEMODYNAMIC EFFECTS OF IONIZING RADIATION: VERTEBRATES (Continued)

Animal	Exposure or Administration	Type of Radiation	Dose	Accumulated Dosage or Exposure Time	Effect	Initial Symptom Manifestation	Reference
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)
Rat (concluded)	Thorax, single exposure (concluded)	X ray	20,000 r		Vacuoles in coronary arterial wall.	8th da	107
121	Thorax, single exposure	X ray	1100 r		Congestion of coronary vessels.	10 da (sacrifice)	108
122	Thorax, single exposure	X ray	700 r		Marked dilatation of right ventricle, congestion of myocardial capillaries, pyknotic of capillary endothelium.	1 wk (sacrifice)	109
123	Thorax, single exposure	X ray	10,000 r		Myocardial damage, degeneration of muscle fibers, intermuscular edema, capillary hemorrhage.	7th da	110, 111
124	Thorax, repeated exposure	X ray	600 r, 300 r, 300 r at monthly intervals	1200 r	Zenker's necrosis of myocardium.	8-32 da after last exposure	91
125	Intravenous, single injection	Radium	270-5900 μ c/kg of body weight		Arterial calcification.		112
126	Peritoneum	Radon obtundent	200 μ c/ml		Increased capillary fragility.		113
127	Head, single exposure	Radium	14.63 mg Ra	3 hr	Blood vessels of brain engorged.		114
128	Intraperitoneal, single injection	Radium	0.5 μ c/g of body weight		Degeneration of tunica adventitia.		75
129	Intravenous, single injection	Polonium	0.9-3.6 μ c/kg of body weight		Myocardial hypertrophy and fibrosis.	375 da	115
130	Total body surface, single exposure	Beta	4000 rep		Inflammation of eyelids, erythema on ears and feet.	4th da	76
131	Abdominal skin, single exposure	Mixed. Strontium 90 (Sr-90) Phosphorus 32 (P-32)	90 rep/sec (Sr-90) 4500 rep/hr (P-32)	28,000 rep	Inflammatory reaction in small arteries and veins of irradiated area.	10th da	116
132	Total body surface, repeated exposure	Beta	625 rep	2 mo	Early collapse and later disintegration of blood vessels.		75
133	Anterior thoracic wall over heart, repeated exposure	Phosphorus 32 X ray	8-12 HED (human erythema dose)	1-5 mo	Hydropericardium, hyperemia of coronary vascular bed, dilatation of pericardial sac, heart distended.		7
134	Entire body, single exposure	X ray	600 r		Degeneration of arterial endothelium.	50 min-184 hr (sacrifice)	117
135	Entire body, single exposure	X ray	275 r (LD-50/30)		Dilatation of capillaries, impairment of circulation, hemostasis, increased capillary permeability, tissue anoxia.		118
136	Entire body, single exposure	Atomic bomb source (gamma)	700 r		Enlarged right ventricle.	7th da (sacrifice)	119
					Cardiac dilatation.	7th da (sacrifice)	

94. EFFECT OF IONIZING RADIATION ON HEMATOPOIETIC TISSUE VERTEBRATES

Abbreviations and definitions: kv = kilovolt, a unit of electrical potential equal to 1000 volts, the crest value of the potential wave in kilovolts, r = roentgen, the quantity of X or gamma radiation such that the associated corpuscular emission 0.001293 g of air produces, in air, ions carrying one electrostatic unit of electrical charge of either sign, n = neutron, a nuclear particle of zero charge and mass number 1; c = curie, the quantity of radionuclide disintegrating at the rate of 3.7×10^{10} atoms per second, mc = millicurie, 3.7×10^7 disintegrations per second; μ = microcurie, 3.7×10^4 disintegrations per second

Animal	Tissue	Exposure or Administration (C)	Type of Radiation (D)	Dose (E)	Effect (F)	Initial Symptom Manifestation (G)	Recovery Following Exposure (H)	Reference (I)
1 Man	Lymphoid tissue	Entire body, single exposure (Japanese victims)	Atom bomb	Lethal or sublethal doses	Atrophy	3 da (first examination).	5 da. (Initial attempt at recovery resulted in production of predominantly atypical mononuclear cells.)	1
2	Bone marrow	Entire body, single exposure (Japanese victims)	Atom bomb	Lethal or sublethal doses	Decreased cellularity	Maximum decrease 6-10 da after exposure, 6 da	Some indication of recovery in all of survivors by 7 da.	1
3	Bone marrow	Therapeutic atmosphere of 4×10^{-9} c/L of air, 1 hr per treatment	Radon	10 or more treatments	Proliferation of reticuloendothelium, increased reticulum cells, plasma cells, macrophages, and lymphocytes. Erythroid hyperplasia, increased lymphocytes, decreased granulocytes, relative increase in basophil WBC in marrow.	6 wk		2
4	Sternal bone marrow	Localized, therapeutic, single exposure	X ray 150-175 kv	100 r	Decreased erythroid cells.	2 da	Complete by 1 mo.	3
5	Sternal bone marrow	Localized, therapeutic, single exposure	X ray 150-175 kv	200 r	Decreased erythroid cells.	2 da	Complete by 1 mo.	3
6	Sternal bone marrow	Localized, therapeutic, single exposure	X ray 150-175 kv	300 r	Decreased erythroid cells, and myeloid cells.	2 da	4 da	3
7	Sternal bone marrow	Localized, therapeutic, single exposure	X ray 150-175 kv	2000 r	Transient aplasia.	1 da	1-2 mo	3
8	Sternal bone marrow	Localized, therapeutic, single exposure	X ray 150-175 kv	3000 r or more	Permanent aplasia.	As early as 2 da after radiation.	24-28 da	4
9 Dog	Bone marrow	Entire body, single exposure	X ray 250 kv	300 r	Decreased cellularity.	2 da	24 da	4
10	Bone marrow	Intraperitoneal injection	Gold-198	244-504 mc	Decreased myeloid cells.	2 da	4 da	5
					Reversed myeloid-erythroid ratio.	1-9 da	40 da	
					Marked decrease in all cellular elements of marrow.	Weeks		

- Berkeley. [75] Rhoades, R. S. 1948. In M. A. Bloom, ed. *Histopathology of irradiation from external and internal sources*. McGraw-Hill, New York. Chap. 16. [76] Raper, J. R., and K. K. Barnes. 1951. In R. E. Zirkle, ed. *Biological effects of external beta radiation*. NRES IV 22-E. Chap. 4. 1952. Blood. N. Y. 7:404. [80] Storey, R. H., et al. 1950. *Proc. Soc. Exp. Biol.*, N. Y. 74:242. [81] Palmer, E. E., et al. 1946. USAEC MDDC-761. Am. J. Path. 28:568. [85] Frank, S. M. 1955. *Proc. Internat. Conf. Peaceful Uses Atomic Energy*, Geneva 11:93. United Nations, New York. [86] Rigdon, R. H., and H. Curi. 1945. Am. J. Roentg. 49:250. [87] Gertner, H. B., et al. 1956. *Rad. Res.* 5:318. [88] Russell, D. S., et al. 1949. *J. Neur.*, Lond. 12:187. [89] Downing, V., et al. 1940. Am. J. Roentg. 43:249. [90] Bishop, F. W. 1954. In R. A. Blair, ed. *Biological effects of external radiation*. NRES VI-2. McGraw-Hill, New York. Chap. 6, p. 194. [91] Warthin, A. S., and E. A. Pohle. 1929. *Arch. Int. M.* 43:125. [92] Gerstner, H. B., et al. 1955. Am. J. Physiol. 162:459. [93] Gertner, H. B., et al. 1955. *Fed. Proc.*, Balt. 14:356. [94] Sinclair, W. K., and H. Blondal. 1952. *Brit. J. Radiol.* 25:360. [95] Raper, J. R., et al. 1951. In R. E. Zirkle, ed. *Biological effects of external beta radiation*. NRES IV 22-E. Chap. 2. [96] Supplee, H., et al. 1952. USAEC CII-3889. [102] Swift, M. N., and T. Taketa. 1956. *Ibid.* 185:446. [100] Bethell, F. H. 1954. USAECU-3030. [101] France, O. 1946. J. Am. Med. Ass. 96:1825. [103] Leach, J. E., and K. Sugiyama. 1941. Am. J. Roentg. 45:414. [106] Warthin, A. S., and E. A. Pohle. 1927. J. Radiol. 10:549. [115] Griffith, J. Q., et al. 1947. *Proc. Soc. Exp. Biol.*, N. Y. 64:331. [114] Colwell, H. A., and R. J. Gladstone. 1937. *Brit. J. Radiol.* 10:549. [117] Tullie, J. L. 1950. U. S. Nav. Med. Res. Rep. NM006-012.04.38. [118] Tullie, J. L. 1949. Am. J. Path. 25:829. [119] Tullie, J. L., et al. 1955. *Ibid.* 31:71. [120] Searner, S. P., et al. 1955. Am. J. Physiol. 182:407.

23	Bone marrow nodes and spleen	Intraperitoneal irradiation single exposure	Sirontium 89 2.0 μ c	Decreased hemopoiesis (all cell series).	3 da	Sustained.	12
24	Lymph nodes	Entire body single exposure	X ray 200 kvp	Occurrence of giant cells. Increased debris in follicles. Necrotic foci in germinal follicles.	4 hr 4 hr 8-12 hr		8
25	All lymphoid tissues	Entire body single exposure	X ray 250 kvp	Increased activity of adenine triphosphatase and 5-nucleotidase.	2-3 hr	4 da	10
26	Lymph nodes	Entire body single exposure	X ray 186 kv	Increased debris. Cytological damage. Disappearance of secondary nodules.	2-3 hr 48 hr	4 da 1-14 da	
27	Lymph nodes	Entire body single exposure	Fast neutrons (plate)	Increased extramedullary myelopoiesis.	3-4 wk	3-9 da	7
28	Thymus	Entire body single exposure	X ray of thermal neutrons plus γ	Complete destruction of nodules.	1 hr		14
29	Thymus	Entire body single exposure	X ray of thermal neutrons plus γ	Induction of thymic lymphoma.	1 yr. (More common in females.)		
30	Thymus and spleen	Entire body single exposure	X ray	Involution.	3 da		13
31	Spleen	Entire body single exposure	X ray	Weight loss (proportional to dose).	120 hr	1-10 da	10
32	Lymphoid elements of spleen	Entire body single exposure	X ray 2000 kvp	Decreased mitosis, lymphocytes, erythropoiesis; cytological damage.	3-5 hr		
33	Erythrocytic elements of spleen	Entire body single exposure	X ray 2000 kvp	Increased debris. Increased megakaryocytes. Increased extramedullary myelopoiesis.	2-3 hr 3-3 hr 10 da	10-14 da	
34	Myeloid elements of spleen	Entire body single exposure	X ray 2000 kvp	Cellular injury (morphological).	1 hr	60 hr	17
				Cellular injury (morphological).	1 hr	66 hr	17
				Cellular injury (morphological).	Few days (sublethal onset).	80 hr	17

94. EFFECT OF IONIZING RADIATION ON HEMATOPOIETIC TISSUE: VERTEBRATES (Continued)

Animal	Tissue	Exposure or Administration	Type of Radiation	Dose	Effect	Initial Symptom Manifestation	Recovery Following Exposure	Reference
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)
11 Dog (concluded)	Regional lymph nodes	Interstitial injection into anterior chest wall	Gold ¹⁹⁸	2 mc	Increased debris, decreased cellularity, abnormal mitosis, many abnormal plasmacytoid cells (predominant cell type).	3 da		6
12	Regional lymph nodes	Interstitial injection into anterior chest wall	Gold ¹⁹⁸	2 mc	Congestion of sinusoids by large phagocytic mononuclear cells.	3 da		6
13	Regional lymph nodes	Interstitial injection into anterior chest wall	Gold ¹⁹⁸	5-10 mc	Decreased cellularity, increased debris, areas of necrosis, many plasmacytoid cells, thrombocytopenia of blood vessels.	3 da		6
14		Interstitial injection into anterior chest wall	Gold ¹⁹⁸	30-85 mc	Decreased cellularity, increased debris, necrosis of 90% or more of node, occasional islands of viable lymphoid tissue, numerous fibroblasts, dense scar tissue.	3-6 wk		6
15 Guinea pig	Lymph nodes	Entire body, single exposure	X ray	175 r	Increased debris.	5-6 wk	24 hr	7
16 Mouse	Bone marrow	Entire body, single exposure	X ray	200 kvp	Slight decrease in cellularity.	3 hr	4 da	
17	Bone marrow	Entire body, single exposure	X ray	50 r	Increased heterophils.	3-8 hr	48 hr	
18	Bone marrow	Entire body, single exposure	X ray	200 r	Increased myeloid cells.	8 hr		8
19	Bone marrow	Entire body, single exposure	X ray	350 r	Increased myeloblasts.	12 hr		9
	Bone marrow	Entire body, single exposure	X ray	400 r	Visible chromosome aberrations.	18 hr		7
	Bone marrow	Entire body, single exposure	X ray	400 r	Increased debris.	3 hr	5 da	
	Bone marrow	Entire body, single exposure	X ray	400 r	Decreased erythropoiesis.	3 hr	5-9 da	
	Bone marrow	Entire body, single exposure	X ray	400 r	Decreased myelopoiesis.	3 hr	5 da	
	Bone marrow	Entire body, single exposure	X ray	400 r	Decreased lymphocytes.	1 da		10
	Bone marrow	Entire body, single exposure	X ray	400 r	Decreased normoblasts, cellularity.	1 da	7 da	
	Bone marrow	Entire body, single exposure	X ray	400 r	Decreased myelopoiesis.	2 da	7-14 da	
20	Bone marrow	Entire body, single exposure	Mixed (detonation of nuclear device)	128 r	Shift to left present in myelocytic series.	10 da		11
	Bone marrow	Entire body, single exposure	Mixed (detonation of nuclear device)	128 r	Threshold dose for induction of myelogenous leukemia.			
21	Bone marrow	Entire body, single exposure	Fast neutrons (pne)	96 n	Increased debris.	3 hr		7
22	Bone marrow	Entire body, single exposure	Fast neutrons (pne)	117 n	Cell death.	3 hr	3 da	
	Bone marrow	Entire body, single exposure	Fast neutrons (pne)	117 n	Decreased cellularity.	3 hr	5-9 da	7
	Bone marrow	Entire body, single exposure	Fast neutrons (pne)	117 n	Increased debris.	3 hr		
	Bone marrow	Entire body, single exposure	Fast neutrons (pne)	117 n	Cytological damage, megakaryocytes.	1 da		
	Bone marrow	Entire body, single exposure	Fast neutrons (pne)	117 n	Increased spindle cells.	9 da		

70	Lymphoid tissue	Entire body, single exposure	X ray	2000 kvp	600 r	Fragmentation of lymphocytes Increased nuclear debris Increased phagocytosis of debris, infiltration of granulocytes. Destruction of almost all lymphocytes of central portion of nodes. Destruction of almost all lymphocytes of entire node (only reticular cells remain). Cytological damage (lympho- cytes).	50 min 60 min 3 hr 36-45 hr (recovery very gradual).	26
71	Chick	Bone marrow	Entire body, single exposure	X ray	200 kvp	25 r	29 hr	27
72		Bone marrow	Entire body, single exposure	X ray	200 kvp	100 r	30 min	27
73		Bone marrow	Entire body, single exposure	X ray	200 kvp	400 r	13 hr	27
74		Bone marrow	Entire body, single exposure	X ray	200 kvp	800 r	7 hr	27
75		Spleen	Entire body, single exposure	X ray	200 kvp	25 r	45 min	27
76		Spleen	Entire body, single exposure	X ray	200 kvp	100 r	1 hr	27
77		Spleen	Entire body, single exposure	X ray	200 kvp	400 r	1-2 hr	27
78		Spleen	Entire body, single exposure	X ray	200 kvp	800 r	1-2 hr	27
79	Salmon	Kidney	Entire body, single exposure	X ray	200 kvp	500 r	2 hr	28

Contributor: Ingram, Marylou

- References: [1] Oughterson, A. W., and S. Warren, eds. 1956. Medical effects of the atomic bomb in Japan, NINES VIII-8, McGraw-Hill, New York. [2] Inama, K. 1952. *Wien. Zsche. Inn. Med.* 33 241. [3] Denstad, T. 1943. Acta radiol., Stockholm, Suppl. 327. [4] Rekers, P. E. 1948. UR-11. Technical Information Service, Oak Ridge, Tenn. [5] Jackson, A. H., and P. Hahn. 1955. Cancer, Phila. 8-480. [6] Christopherson, W. M., and H. F. Berg. 1955. *Biol. 8:1263*. [7] Bloom, W. 1948. The histopathology of irradiation, NINES IV-21. McGraw-Hill, New York. [8] Neille, A. 1949. Radiology 42 64. Acta radiol., Stockholm, Suppl. 116-469. [9] Jacobson, L. O., et al. 1954. Cancer Res. 14:582. [10] Brecher, G., et al. 1948. Blood, N. Y. 3:1259. [11] Furr, J., and A. C. Upton. 1954. *Physiol.* 176 282. [12] Union, A. C., et al. 1954. Cancer Res. 14:582. [13] Part, R. M., et al. 1955. J. Cellul. Physiol. 42:327. [14] Storer, J. B., and W. H. Langham. 1954. AECU-5099. Technical Information Service, Oak Ridge, Tenn. [15] Lamm, B. G., and J. L. Tullis. 1951. *Mil. Surgeon* 109:281. [16] Hagen, C. W., et al. 1944. MDOC-999. Technical Information Service, Oak Ridge, Tenn. [17] Zirkle, R. E., ed. 1956. Biological effects of external X and gamma radiation. Part 2, TID-5228. Technical Information Service, Oak Ridge, Tenn. [18] Metcalf, R. G., et al. 1950. UR-123. Technical Information Service, Oak Ridge, Tenn. [19] Hsu, C. L., and W. C. Ma. 1940. *Am. J. Cancer* 39:319. [20] Fink, R. M., ed. 1950. *Physiological studies with Polonium, radium, and plutonium*. NINES VI-3, McGraw-Hill, New York. [21] Sloof, P. G., et al. 1950. *Anat. Rec.* 108 537. [22] McDonald, E. 1947. Neutron effects on animals. Williams and Wilkins, Baltimore. [23] Trowell, O. A. 1952. J. Path. Bact., Lond. 64:637. [24] Tullis, J. L., et al. 1951. *Mil. Surgeon* 109:271. [25] Murray, R., et al. 1940. CIL-3873. Technical Information Service, Oak Ridge, Tenn. [26] Donham, K., et al. 1948. Growth, Rhaca 12:107.

94. EFFECT OF IONIZING RADIATION ON HEMATOPOIETIC TISSUE. VERTEBRATES (Concluded)

Animal	Tissue	Exposure or Administration	Type of Radiation	Dose	Effect	Initial Symptom Manifestation	Recovery Following Exposure	Reference
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)
60 Rat (concluded)	Lymph nodes	Entire body, single exposure	X ray 200 kvp	600 r	Cytological damage (lymphocytes). Degenerative change (nodules). Disappearance of nodules. Infiltration with heterophils. Hemorrhage.	1 hr 14 hr 24 hr 8 hr	14 hr 21 da 21 da	7
61	Lymph nodes	Single intra-venous injection	Plutonium	300-3000 μ c/kg body weight	Hypoplasia (more marked with greater doses).	At autopsy, 14-260 da.		22
62	Lymph nodes	Single intra-venous injection	Radium	17-8000 μ c/kg of body weight	Hypoplasia (more marked with higher doses), neoplastic changes in 3 of 20 rats examined.	At autopsy, 11-407 da.		22
63	Thymus, lymph nodes, and spleen	Entire body, single exposure	X ray 250 kvp	50-400 r (sublethal)	Increased activity of adenosine triphosphatase and 5-nucleotidase.	3 hr	72 hr	13
64	Spleen and lymph nodes	Single intra-venous injection	Polonium	50-170 μ c/kg of body weight	Extreme decrease in cellularity.	At autopsy, 7-300 da (depending on dose).		22
65	Spleen	Single intra-venous injection	Plutonium	300-3000 μ c/kg of body weight	Definite decrease in cellularity, hyperplasia of red pulp of spleen (extramedullary myelopoiesis).	At autopsy, 14-260 da.		22
66	Spleen	Single intra-venous injection	Radium	17-8000 μ c/kg of body weight	Complete or nearly complete atrophy of lymphoid elements at all dosage levels; extramedullary myelopoiesis more marked with higher doses.	At autopsy, 11-407 da.		22
67	Spleen	Entire body, single exposure	X ray 250 kvp	550 r	Increased debris. Destruction of lymphocytes.	30 min 2 hr	8 hr	20
68	Spleen	Entire body, single exposure	X ray 200 kvp	600 r	Disappearance of nodules. Decreased cellularity. Decreased megakaryocytes. Cytological damage (erythroblasts).	30 hr 2 hr 3 da 3 hr	15 da 9 da 3 wk	7
69	Bone marrow	Entire body, single exposure	X ray 2000 kvp	600 r	Decreased erythropoiesis. Destruction of lymphocytes. Decreased mitosis. Decreased cellularity of nodules. Decreased erythroid elements. Destruction of virtually all nucleated red blood cells. Morphological damage to myeloid cells.	8 hr 3 hr 1 hr 3 hr 9 hr 13 hr 13 hr	9 da 8 hr 14 da No evidence of regeneration during 164-hr period of observation.	26

7	Entire body, 5-10 r/da	X ray 200 kvp 400 kvp 1000 kvp	200-300 r	Decreased lymphocytes	Cancer patients with normal hemato-	5
8	Entire body, 5-20 r/da	X ray 200 kvp 400 kvp 1000 kvp	88-300 r	Decreased lymphocytes, neutrophils, WBC, RBC, hemoglobin.	Cancer patients with normal hemato-	5
9	Occupational, entire body, 0.2 r/wk	High energy gamma	17 wk	Decreased WBC, absolute lymphocytes, absolute neutrophils		6
10	Tumor therapy, 5-6 times/wk, 1-50 megagram-r (integral doses), 3000-6000 r (tumor doses)	X ray 200 kvp 400 kvp	7 megagram-r	Reduced absolute lymphocyte count by 25%.	Increased injurious effect/megagram-r, with decreasing integral dose/da.	7
11	Therapeutic radiation, daily 50, 75, 100 r/da (30 x 30 cm or 45 x 45 cm fields)	X ray 250 kv	3-6 da	Transient increase in platelet count, followed by decrease.	Rate of decrease greater in patients with impaired hematopoiesis and greater with increased size of field.	8
12	Local irradiation of hands during fluoroscopy	X ray (portable machine) 55-80 kv	Approximately 800-1000 r (4 min) in 3 mo	Decreased absolute lymphocytes. Increased % lymphocytes on differential counts. Many abnormal monocytes	Normal when blood re-examined 6 mo later.	9
13	Occupational	Cyclotron workers	Approximately 3 mo; probably below MPE (maximum permissible exposure)	Increased incidence of lymphocytes with bilobed nuclei. Picture returned to normal after extra shielding installed.		10
14	Experimental	20 radium workers	Exposure not known, presumably daily	Persistent decrease in lymphocytes, less marked decrease in neutrophils (as compared with 40 controls)	Reported in 1921.	11
15	Occupational	Particle accelerators (predominantly X and gamma rays)	Tolerance range approximately 200 mr/wk	Increased incidence of lymphocytes with bilobed nuclei.		12
16	Occupational	Radium, several hundred luminifers	Body burden less than 0.2 µg	Almost all workers showed presence of abnormal early mononuclear cells. These cells no longer present after individuals ceased luminizing	Examined each yr from 1941-1947.	13
17	Occupational	193 employees	0.05 r/da, 1943-1946	Increased refractile neutral red bodies in lymphocytes		14
18	Occupational	15 employees of radium institute	1 mo-33 yr	All showed "radiological blood picture," viz. high % of lymphocytes (relative lymphocytosis), normal RBC, and normal or high color index.	Studied in 1921.	15
19	Occupational	Employees of radium institute		All personnel showed counts differing very little from normal.	Studies made in 1923.	15

95. EFFECT OF IONIZING RADIATION ON PERIPHERAL BLOOD: VERTEBRATES

Part I: MAN

Abbreviations and definitions: kv = kilovolt, a unit of electrical potential equal to 1000 volts; kvp = kilovolt peak, the crest value of the potential wave in kilovolts; r = roentgen, the quantity of X or gamma radiation such that the associated corpuscular emission per 0.001293 g of air produces, in air, ions carrying one electrostatic unit of electrical charge of either sign.

Exposure or Administration (A)	Type of Radiation and/or Subject (B)	Accumulated Dosage or Exposure Time (C)	Effect (D)	Remarks (E)	Reference (F)
1 Entire body (single or few closely-spaced exposures)	Atomic bomb detonation in Japan, 1945	Lethal and sublethal ("severe cases")	Single Exposure Rapidly developing leukopenia.	WBC virtually absent from blood of many of the fatally injured.	1
			Steadily decreasing RBC and hemoglobin.	Extremely severe anemia within 3-4 wk.	
			Decreased platelets.	Purpura and other hemorrhagic manifestations common.	
2 Entire body (single or few closely-spaced exposures)	Atomic bomb detonation in Japan, 1945	Sublethal (moderately severe and mild cases)	Bizarre giant neutrophils, atypical lymphocytes, and occasional plasma cells. Leukopenia.	Minimal WBC during 4th wk. Lowest average WBC: 1500-2500/cu mm.	1
			Decreased RBC and hemoglobin.	Lowest RBC and hemoglobin levels 6-9 wk after exposure. Tendency towards macrocytosis.	
			Decreased platelets.	Definite increase in absolute eosinophil count 8-12 wk after exposure.	
3 Surface deposition, inhalation and ingestion	Fail-out from detonation of nuclear device	175 r (calculated)	Increased eosinophils.	Drop to 55% of control level by 3rd da after exposure. Remained stable at this level (greater depression in younger group).	2
			Decreased lymphocytes.	Drop to 70-80% of control level by 2nd wk. Second decrease began 5th wk; counts fell to 50% of control level (greater depression in younger group).	
			Decreased neutrophils.	Minimum level (30% of control value) during 4th wk. Second depression during 7th and 8th wk.	
4 Head and neck	Dental X ray	35-315 r	Decreased platelets.	First measured on 22nd da.	3
			Decreased hematocrit.	Observed during 12-hr post-exposure period.	
5 Therapeutic dose to sacroiliac joints and spine	X ray 250 kv	300 r (mean surface dose)	Decreased absolute lymphocytes.	Lymphocytes decreased 1-2 da after exposure, neutrophils 5th da (after temporary increase), WBC 2nd da, venous hematocrit 2nd da.	4
6 Entire body. 15-20 r/dose	X ray 200 kvp 400 kvp 1000 kvp	60 r	Repeated Exposures Decreased lymphocytes.	Cancer patients with normal hematopoiesis.	5

31	Intravenous thorotrast containing 2.5 g thorium/12 ml solution (890 r/yr/g thorium), injected in 1931	Thorotrast	50 ml. Estimated total dose, 1780 r/yr for 25 yr	Chronic myeloid leukemia developed in 1954. Clinical course mild. Paradoxical two-fold increase in WBC when spleen irradiated with X rays.	25-yr latent period	25
32	Single exposure	Laboratory accident	186 r of 80 kv X rays. Plus 10 7 r y	Decreased absolute lymphocyte count increased refractile neutral red bodies in lymphocytes.	Duration, 1 yr. Present for 18 mo.	26
33	Single exposure	Laboratory accident	390 r of 80 kv X rays. Plus 26 4 r y	Decreased absolute lymphocytes. Increased neutral red bodies in lymphocytes	2 yr after exposure.	26
34	Entire body, single exposure, epilation	Hiroshima atomic bombing	Approximately 400 r	Slight decrease in % lymphocytes in peripheral blood. Slight increase in eosinophils in peripheral blood.	1 yr later.	27
35	Entire body	Full-out after detonation of nuclear device in Marshall Islands (Hongtsepo), spring, 1954	175 r (calculated)	Greater variability in blood picture of the 904 survivors than in blood picture of matched control group.	1 yr later.	28
36	Single exposure	Survivors of atomic bombing at Hiroshima, 1945		Decreased lymphocytes, monocytes, eosinophils, and platelets	1 yr after exposure (compared with carefully selected control groups).	28
37	Single exposure, <1500 meters from impact	4 (of 5075) survivors of Nagasaki atomic bombing		Increased incidence of leukemia among survivors at distance <1500 meters from impact.	Highest incidence among those exposed <1500 meters from impact. Peak incidence 1950-1952. More than 50% of cases acute or subacute (total number of cases, 92).	29
38	Intravenous injection	Thorium X		Refractory (fatal) anemia characterized by leukopenia and thrombocytopenia.	Latent period, 4-7 yr.	30
39	Occupational	Probably X rays and radium	Dosage not known, exposure probably over many yr	Myeloid leukemia developed 1 yr after intravenous Thorium X for tuberculosis of knee joint.		31
40	Occupational	Probably X rays and radium, dermatologists dying between 1935-1944	Dosage not known, exposure probably over many yr	Deaths due to leukemia 1.5 times higher among physicians than among general male population.		32
41	Occupational	X ray, physicians dying between 1938-1942	Dosage not known, exposure of long duration	Increased incidence of leukemia.		33
42	Occupational	X ray, physicians generally, and radiologists	Dosage not known, repeated exposure	Deaths due to leukemia 1.25 times greater among physicians than among general male population. 14 leukemia deaths/299 deaths in radiologists. 344 leukemic deaths/65,992 in physicians other than radiologists.	Lower incidence of death due to cancer in general.	34
						35

95. EFFECT OF IONIZING RADIATION ON PERIPHERAL BLOOD: VERTEBRATES (Continued)

Part I MAN (Continued)

Exposure or Administration (A)	Type of Radiation and/or Subject (B)	Accumulated Dosage or Exposure Time (C)	Effect (D)	Remarks (E)	Reference (F)
20 Occupational	1405 employees in Institute of Radiophysics, Stockholm	Repeated Exposures (concluded)	Hyperregeneration of polymorphonuclear WBC nuclei, pathological lymphocytes, granulocytopenia, left shift in granulocytes	Study reported in 1946.	16
21 Occupational	Employees	0.02-0.05 r/da (estimated doses)	"In all probability, radiation quantities as low as 0.02-0.05 r/da can, after a comparatively short time, give rise to blood changes." Based on data from Nordenson (1946) and Helde (1946).		16-18
22 Experimental	Employees	125 megagram-r/wk	WBC lower than in controls and those receiving 100 mr/wk.		19
23 Experimental	Employees	100 megagram-r/wk	12 of 17 showed definite depression of WBC, particularly neutrophils, with relative lymphocytosis and often a slight increase in eosinophils.	Reported in 1935.	19
24 Occupational	17 "new" radium workers	Few wk	Blood picture same as that of new workers after several weeks.	Reported in 1935.	20
25 Occupational	15 "old" radium workers	6 mo.-1 yr	Shown significant increase in lymphocyte count after 6 and 12 mo of work.	Not statistically significant when compared with very slight lymphocyte increase in control group during same period.	21
26 Occupational	Metallurgical laboratory workers: 205 worked 12 mo, 453 worked 6 mo	Less than 0.1 r/da	22-40% of workers examined showed relative lymphocytosis without neutropenia. Counts returned to normal after individuals ceased luminizing.	Several hundred individuals examined each yr from 1941-1947.	13
27 Occupational	Radium luminizers	Body burden less than 0.2 µg	Latent Effects		
28 Environmental	Adult population	Lifetime exposure	Leukemia 6 times more frequent among adults living on uraniferous terrain than among adults on neighboring, low-radioactive terrain		22
29 Occupational	X ray	Many years	Physician had mycosis fungoides for many years after working with X ray. Had "temporary leukemia" from time to time throughout 14-yr period.	"Temporary leukemia" (WBC approximately 33,000/cu mm, 90% lymphocytes), responded well to X-ray therapy.	23
30 Prenatal exposure during diagnostic X-ray examination of mother during pregnancy	X ray	Dose to fetal gonads frequently greater than 2.5 r	Increased incidence of leukemia in children of irradiated mothers.		24

Abbreviations and definitions kv = kilovolt, a unit of electrical potential equal to 1000 volts, kvp = kilovolt peak, the crest value of the potential wave in kilovolts, r = roentgen, the quantity of X or gamma radiation such that the associated corpuscular emission per 0.01293 g of air produces, in air, ions carrying one electrostatic unit of electrical charge of either sign, rad = rad unit, equivalent to 100 ergs/g energy absorption, n = neutron, a nuclear particle of zero charge and mass number 1, mc = millicurie, the quantity of radionuclide disintegrating at the rate of 3.7×10^{10} atoms per second, μ = microcurie, 5.7×10^4 disintegrations per second

Animal	Exposure or Administration (B)	Type of Radiation (C)	Accumulated Dosage or Exposure Time (D)	Effect (E)	Remarks (F)	Reference (G)
1 Rat	Entire body, single exposure (at room temperature) 400 r/da	X ray 250 kv	500 r	Decreased lymphocytes and granulocytes.	No decrease in RBC.	1
2 Burro	Entire body, 400 r/da	Cobalt-60 gamma	10 da	Pronounced leukopenia (lymphocytes and granulocytes), decreased platelets, hyperferremia	Lymphocytes less than 8% of normal 1 hr after exposure. Minimal granulocyte count on 6th da	2
3 Dog	Entire body, single exposure	X ray 1000 kvp	5 r	Increased incidence of lymphocytes with bilobed nuclei for 4 wk.	Peak incidence 2nd and 3rd wk after exposure.	3
4	Entire body, single exposure	X ray 200 kvp	20 r	Decreased absolute lymphocytes.		4
5	Entire body, single exposure	X ray 200 kvp	50 r	Decreased absolute lymphocytes, platelets, WBC, granulocytes.	Decrease in WBC and platelets more marked with increasing dose to approximately LD ₁₀₀ .	4
6	Entire body, single exposure	X ray 200 kvp	250 r	Decreased absolute lymphocytes, platelets, WBC, granulocytes, reticulocytes.		5
7	Entire body, 0.5 r/da, 6 da/wk	X ray 200 kvp 1000 kvp	2 yr	Decreased lymphocytes.		5
8	Entire body, 1.0 r/da, 6 da/wk	X ray 200 kvp 1000 kvp	1 yr	Decreased lymphocytes, WBC, neutrophils.		5
9	Entire body, 3 r/da, 6 da/wk	X ray 250 kvp 1000 kvp	1-2 mo	Decreased absolute lymphocytes, reticulocytes, platelets.		5
10	Entire body, 6 r/da, 6 da/wk	X ray 250 kvp 1000 kvp	1-2 mo	Decreased WBC, absolute lymphocytes, absolute neutrophils, platelets.		5
11	Entire body, 10 r/da, 6 da/wk	X ray 200 kvp	1 mo	Decreased absolute lymphocytes, absolute neutrophils, WBC, hemoglobin, reticulocytes, RBC.		6
12	Entire body, 12.5 r/da	X ray 200 kvp	1278 r	Aplastic anemia. Drop in absolute lymphocytes to 700/cu mm.		7
13	Entire body, 25 r/da	X ray 200 kvp	900 r	Absolute lymphocytes reached level of 300/cu mm, severe anemia.		7
14	Entire body, 50 r/da	X ray 200 kvp	900 r	Absolute lymphocytes reached level of approximately 100/cu mm.		7

95. EFFECT OF IONIZING RADIATION ON PERIPHERAL BLOOD: VERTEBRATES (Continued)
Part II: MAN (Concluded)

Exposure or Administration (A)	Type of Radiation and/or Subject (B)	Accumulated Dosage or Exposure Time (C)	Effect (D)	Remarks (E)	Reference (F)
43 Occupational	X ray, physicians dying during 1947-1951		Latent Effects (concluded) Increased incidence of deaths due to leukemia.		
44 Therapy for ankylosing spondylitis (1940-1954)	X ray	Repeated exposure	Deaths due to leukemia 5-10 times expected number among patients receiving 1 course of therapy; at least 9 times expected number among patients receiving more than 1 course. Doses less than 200 r associated with increased incidence of leukemia (larger doses with thyroid cancer). After therapy, 5 instances of myeloid leukemia (one had antecedent aplastic anemia), and 2 instances of aplastic anemia.	Relatively short latent period (incidence increased within first 5 yr of exposure). Some, but not all, of increased incidence may reflect increased susceptibility of patients with ankylosing spondylitis to leukemia.	36
45 Therapy for enlarged thymus	X ray, children, 1925-1951	50-1500 r		Total incidence of neoplasia significantly higher among patients studied than among non-irradiated siblings. Latent period, 2½ mo-6 yr.	37
46 X radiation of spine, therapeutic for ankylosing spondylitis	X ray	900-5950 r			38
47 Internal deposition	Chronic radium poisoning		Profound anemia and leukopenia. Fatal.		39

Contributor: Ingram, Marylou

References [1] Oughterson, A. W., and S. Warren, ed. 1956. Medical effects of the atomic bomb in Japan. NNES VIII-8. McGraw-Hill, New York.

[2] Cronkite, E. P., et al. 1955. J. Am. M. Ass. 159:430. [3] Dobson, L. R., and M. Chupp. 1957. UCRL-1574. [4] Brown, W. M., and J. D. Abbott. 1955. Service, Oak Ridge, Tenn. [7] Kohn, H. I. 1955. Radiology 64:382. [8] Brown, W. M. 1949. Acta radiol., Stockh. 32:407. [9] Huitberg, S. 1949. Ibid.

[10] Ingram, Marylou, et al. 1952. Science 116:706. [11] Mottram, J. C., and J. R. Clark. 1920. Proc. R. Soc. M., Lond. 23:25. [12] Dobson, D. R. 1935. Ibid. 8:669. [13] Nordenson, N. G. 1946. Acta radiol., Stockh. 37:416. [14] Dickie, A., and L. H. Hempelmann. Technical Information Service, Oak Ridge, Tenn. [22] Stewart, J., et al. 1956. Lancet, Lond. 2:447. [23] Netousek, M., J. Borek, and K. Dvorak. 1957. Blood, N. Y.

[31] Schaffer, E. L., and H. Grevel. 1952. Münch. med. Wschr. 94:158. [32] Henshaw, P. F., and J. W. Hawkins. 1944. J. Nat. Cancer Inst. 4:339. [33] Simpson, C. L., et al. 1955. Radiology 64:840. [36] Feller, S., and P. Pick. 1951. J. Am. M. Ass. 147:893. [37] Brown, W. M., and J. D. Abbott. 1955. Lancet, Lond. 1:1293.

[38] Simpson, C. L., et al. 1955. Radiology 64:840. [39] Von Sway, H. 1955. Lancet, Lond. 269:225. [40] Matland, H. S. 1929. J. Am. M. Ass.

92:466

32	Entire body, 4.4 r/da	Radium (gamma)	Approximately 1 mo	Decreased lymphocytes, WBC, heterophils.	15
33	Entire body, 4.4 r/da	Radium (gamma)	Approximately 1 yr	Decreased lymphocytes, WBC, heterophils, platelets, RBC, hemoglobin.	15
34	Entire body, 8.8 r/da	Radium (gamma)	10 da	Decreased lymphocytes, WBC, heterophils, platelets.	15
35	Entire body, 8.8 r/da	Radium (gamma)	Approximately 1 mo	Decreased lymphocytes, WBC, heterophils, RBC, hemoglobin (precipitous)	15
36	Entire body, 8.8 r/da	Radium (gamma)	19 wk	Decreased lymphocytes, granulo- cytes, platelets, RBC, and hemoglobin.	16
37	Entire body, single exposure	X ray	600 r	Adhesiveness of WBC and platelets to endothelium not increased. RBC and hemoglobin decrease began about 9th da after exposure. Maximum depression 5-7 da after exposure, returned to normal by 15th da.	17
38	Single exposure	X ray	825 r	Decreased granulocytes.	18
39	Entire body, single exposure	X ray 250 kv	50-100 r	Decreased lymphocytes, granulo- cytes, platelets, and RBC packed volume. WBC seldom <3000/cu mm.	19
40	Entire body, single exposure	X ray 250 kv	260 r	Decreased lymphocytes, granulo- cytes, platelets, RBC, and reticulocytes.	18
41	Entire body, single exposure	X ray 250 kv	400 r	Decreased lymphocytes, granulo- cytes, platelets, RBC packed volume.	19
42	Entire body, single exposure	X ray 250 kv	500-550 r	Decreased lymphocytes, granulo- cytes, platelets, RBC, and reticulocytes.	19
43	Entire body, single exposure	X ray 250 kv	600-650 r	Decreased lymphocytes, granulo- cytes, platelets, RBC, and reticulocytes.	19
44	Entire body, single exposure	X ray 250 kv	800 r	Decreased lymphocytes, granulo- cytes, platelets, and RBC packed volume.	18
45	Single intramuscular injection	Strontium 90 (in equilibrium with Yttrium 90)	0.1-0.2 mc/kg of body weight	WBC at levels below 500/cu mm. 8th da after exposure. Recovery trend began on 15th da (no sur- vivors after 17 da).	20
46	Single intramuscular injection	Strontium 90 (in equilibrium with Yttrium 90)	0.47-0.5 mc/kg of body weight	Lymphocyte fall began immediately after injection; minimal WBC levels in 3rd wk. Reticulocytes normal 11-15 da after exposure, reduced but not absent at 17-20 da. RBC decrease began 10th da after injection.	20

Part II. VERTEBRATES OTHER THAN MAN (Continued)

Animal	Exposure or Administration (B)	Type of Radiation (C)	Accumulated Dosage or Exposure Time (D)	Effect (E)	Remarks (F)	Reference (G)
15 Dog (concluded)	100 r/wk	X ray 175 kv	Weeks	Decreased fragility of RBC in hypotonic saline and allokan, except for sudden increase in fragility in terminal stage of X-ray sickness	Decreased fragility greater in splenectomized than in non-splenectomized dogs.	8
16	Entire body, 1.7 n/da, 6 da/wk	Fast neutrons	1 yr	Decreased WBC, absolute lymphocytes, absolute neutrophils, platelets, RBC.		9
17	Entire body, 1.7 n/da, 6 da/wk	Fast neutrons	3-4 wk	Decreased WBC, absolute lymphocytes, absolute neutrophils.		9
18	Repeated intraperitoneal injections	Gold 198	244-504 mc	Decreased WBC.	Reached approximately 4800 WBC/cu mm; returned to normal 5-8 wk after initial injection.	10
19	Intravenous injection	Gold 198	1 μ c/kg of body weight	Moderate leukopenia, decreased hemoglobin and hematocrit, increased RBC sedimentation rate.	Hemoglobin and hematocrit depressed for several wk.	11
20	Intraperitoneal injection	Gold 198	25.9-42.9 mc/kg of body weight	Definite leukopenia.	WBC in low normal range terminally.	10
21	Repeated intravenous injections	Gold 198		Aplastic anemia.		10
22	Parenteral injection	Sr-90	2 or 7 mc	Decreased granulocytes, WBC, hematocrit.	Hematocrit depression greater with 7 mc. minimum level, 21-25 da.	12
23	Parenteral injection	Sr-90	20 mc	Decreased lymphocytes, granulocytes, WBC, hematocrit.	Hematocrit depression greater than that following 2 or 7 mc dose.	12
24 Goat	Entire body, single exposure	X ray 200 kvp	300 r	Decreased lymphocytes, neutrophils, WBC, RBC, hemoglobin.		13
25 Guinea pig	Entire body, single exposure	X ray 200 kv	220 r	Decreased all WBC, RBC, and hemoglobin.		14
26	Entire body, 0.11 r/da	Radium (gamma)	Approximately 1 mo	Decreased lymphocytes, WBC, heterophils.		15
27	Entire body, 1.1 r/da	Radium (gamma)	Approximately 1 mo	Decreased lymphocytes, WBC, heterophils.		15
28	Entire body, 1.1 r/da	Radium (gamma)	Approximately 1 yr	Decreased lymphocytes, WBC, heterophils; decreased platelets in females only.		15
29	Entire body, 2.2 r/da	Radium (gamma)	Approximately 1 mo	Decreased lymphocytes, WBC, heterophils.		15
30	Entire body, 2.2 r/da	Radium (gamma)	Approximately 1 yr	Decreased lymphocytes, WBC, heterophils.		15
31	Entire body, 2.2 r/da	Radium (gamma)	79 wk	Decreased lymphocytes, WBC, heterophils, platelets.		15
	Entire body, 2.2 r/da	Radium (gamma)		Decreased lymphocytes, WBC, heterophils, RBC and hemoglobin decreased in males only.		15

32	Hamster	Entire body, 4.4 r/da	Radium (gamma)	Approximately 1 mo	Decreased lymphocytes, WBC, heterophils.	15
33		Entire body, 4.4 r/da	Radium (gamma)	Approximately 1 yr	Decreased lymphocytes, WBC, heterophils, platelets, RBC, hemoglobin	15
34		Entire body, 8.8 r/da	Radium (gamma)	10 da	Decreased lymphocytes, WBC.	15
35		Entire body, 8.8 r/da	Radium (gamma)	Approximately 1 mo	Decreased lymphocytes, WBC, heterophils, platelets	25
36		Entire body, 8.8 r/da	Radium (gamma)	19 wk	Decreased lymphocytes, WBC, heterophils, RBC, hemoglobin (precipitous).	15
37		Entire body, single exposure	X ray	600 r	Decreased lymphocytes, granulocytes, platelets, RBC, and hemoglobin.	16
38		Single exposure	X ray	825 r	Decreased granulocytes.	17
39	Monkey	Entire body, single exposure	X ray 250 kv	50-100 r	Decreased lymphocytes, granulocytes, platelets, and RBC packed volume. WBC seldom <5000/cu mm.	18
40		Entire body, single exposure	X ray 250 kv	260 r	Decreased lymphocytes, granulocytes, platelets, RBC, and reticulocytes	19
41		Entire body, single exposure	X ray 250 kv	400 r	Decreased lymphocytes, granulocytes, platelets, RBC packed volume.	18
42		Entire body, single exposure	X ray 250 kv	500-550 r	Decreased lymphocytes, granulocytes, platelets, RBC, and reticulocytes.	19
43		Entire body, single exposure	X ray 250 kv	600-650 r	Decreased lymphocytes, granulocytes, platelets, RBC, and reticulocytes.	19
44		Entire body, single exposure	X ray 250 kv	800 r	Decreased lymphocytes, granulocytes, platelets, and RBC packed volume.	18
45		Single intramuscular injection	Strontium 90 (in equilibrium with Yttrium 90)	0.1-0.2 mc/kg of body weight	Decreased lymphocytes, granulocytes, platelets, and RBC.	20
46		Single intramuscular injection	Strontium 90 (in equilibrium with Yttrium 90)	0.47-0.5 mc/kg of body weight	Decreased lymphocytes, granulocytes, platelets, RBC, and reticulocytes.	20

95. EFFECT OF IONIZING RADIATION ON PERIPHERAL BLOOD: VERTEBRATES (Continued)

Part II: VERTEBRATES OTHER THAN MAN (Continued)

Animal	Exposure or Administration	Type of Radiation	Accumulated Dosage or Exposure Time	Effect	Remarks	Reference
(A)	(B)	(C)	(D)	(E)	(F)	(G)
47 Mouse	Entire body, single exposure	X ray 200 kvp	50 r	Decreased lymphocytes, WBC.		21
48	Entire body, single exposure	X ray	50-250 r	Iron ⁵⁹ uptake by RBC decreased.	Depression more marked with greater doses.	22
49	Single exposure	X ray 250 kvp	100-500 r	Increased incidence of leukemia.	After injection with virulent AK4 leukemia cells, probit of % leukemic deaths plotted against log of dose in r gives regression line of type Y = (a + b log X).	22
50	Entire body, 1 or 2 exposures	X ray 250 kvp	119 rad	Increased incidence of myeloid leukemia.	More common in males; peak incidence, 6-15 mo of age.	23
51	1 or 2 exposures	Cyclotron, fast neutrons	110 rad	Increased incidence of myeloid leukemia.	More common in males; peak incidence, 10-14 mo of age.	23
52	Entire body, single exposure	X ray or thermal neutrons, plus gamma rays (pile)	128 r or more	Increased incidence of myeloid leukemia. Necrosis of bone marrow in more than 50% of mice with myeloid leukemia.	Peak incidence, 12-16 mo of age.	24
53	Entire body, single exposure	Thermal neutrons, plus gamma rays (pile)	80-min exposure, 3.3-4.3 x 10 ¹² n/sq cm, plus 520 r gamma	Decreased WBC, RBC, reticulocytes.	Recovery began after approximately 2 wk.	24
54	Entire body, single exposure	X ray 140 kv	400-600 r	WBC decreased to same minimum level at same time (5-8 da after exposure) in both strains.	Recovery in both strains began 7-13 da after exposure.	25
55	Entire body, 2	Radium (gamma)	29 wk	Decreased WBC, lymphocytes.		15
56	Entire body, 2	Radium (gamma)	29 wk	Decreased WBC, lymphocytes.		15
57	Entire body, 2	Radium (gamma)	79 wk	Decreased WBC, lymphocytes; decreased RBC, hemoglobin, platelets in males only.		15
58	Entire body, 2	Radium (gamma)	29 wk	Decreased WBC, lymphocytes.		15
59	Entire body, 2	Radium (gamma)	29 wk	Decreased WBC, lymphocytes, RBC, hemoglobin, platelets.		15
60	Entire body, 0.07 n/da, 5 da/wk	Fast neutrons (cyclotron)	30 n	Decreased WBC.		26
61	Entire body, 1.4 n/da, 5 da/wk	Fast neutrons (cyclotron)	230 n	Decreased absolute lymphocytes, WBC, RBC, hemoglobin. Increased frequency of WBC below 15,000/cu mm. Shift to left present in heterophils.		26
62	Intraperitoneal injection	Strontium-89	0.068 µc/g of body weight	Decreased heterophilia.	Effect seen in ABC male mice, not in CP-1 mice.	27

63	Intraperitoneal infection	Sironium 89	0.068-14.5 µc/g of body weight	Decreased heterophils and lymphocytes.	Both cell types decreased about equally, decrease greater with greater doses	27
64	Intraperitoneal infection	Sironium 89	2.0 µc/g of body weight	Sustained anemia in splenectomized mice only.	Sufficient myeloid metaplasia in spleen of non-splenectomized mice to prevent anemia.	27
65	Intravenous, intraperitoneal, or intracardial infection	Radium	0.02-0.03 µg	Decreased WBC.		28
66	Intravenous, intraperitoneal, or intracardial infection	Radium	0.1-0.2 µg	Macrocytic anemia.		28
67	Intravenous or intramuscular infection	Plutonium	0.0062 µc/g of body weight	Moderately severe decrease in WBC.	Effect sustained.	28
68	Intravenous or intramuscular infection	Plutonium	More than 0.0062 µc/g of body weight	Decreased WBC, platelets, and reticulocytes; anemia.		28
69	Entire body, single exposure	X ray 200 kvp	25 r	Decreased lymphocytes.	Return to normal by 48 hr after exposure.	30
70	Entire body, single exposure	X ray 200 kvp	100 r	Decreased lymphocytes, heterophils, reticulocytes, and platelets.	WBC depression tends to be greater and recovery slower with increasing doses.	30
71	Entire body, single exposure	X ray 200 kvp	200 r	Decreased lymphocytes, heterophils, platelets, reticulocytes, morphological abnormalities.	Increased neutral red granules in lymphocytes. Occasional phagocytic monocyte containing engulfed nuclear debris 12-24 hr after exposure.	30, 31
72	Entire body, single exposure	X ray 200 kvp	500 r	Decreased WBC, platelets, reticulocytes, RBC, hemoglobin. Morphological abnormalities in leukocytes. Decreased WBC migration <i>in vitro</i> .	Anemia maximal on 14th da after exposure; RBC returned to normal by 23rd da.	30
73	Entire body, single exposure	X ray 200 kvp	800 r or more	Decreased lymphocytes, heterophils, reticulocytes, platelets, RBC, hemoglobin. Increased number of nucleated RBC.		32
74	Entire body, single exposure	Fast neutrons (Ipile)	9 n	Decreased absolute lymphocytes.		32
75	Entire body, single exposure	Fast neutrons (Ipile)	26 n	Decreased absolute lymphocytes, heterophils, platelets.		32
76	Entire body, single exposure	Fast neutrons (Ipile)	66-76 n	Decreased absolute lymphocytes, heterophils, platelets, RBC, hemoglobin, reticulocytes.		32
77	Entire body, single exposure	Fast neutrons (Ipile)	128 n	Decreased absolute lymphocytes.		15
78	Entire body, 1.1 r/da (females only)	Radium (gamma)	Approximately 2 mo	Decreased absolute lymphocytes.		15
79	Entire body, 2.2 r/da (females only)	Radium (gamma)	Approximately 2 mo	Decreased absolute lymphocytes.		15

1/1/ High and low WBC strains. 12/ LAF strain.

95. EFFECT OF IONIZING RADIATION ON PERIPHERAL BLOOD: VERTEBRATES (Continued)

Part II: VERTEBRATES OTHER THAN MAN (Continued)

Animal	Exposure or Administration (B)	Type of Radiation (C)	Accumulated Dosage or Exposure Time (D)	Effect (E)	Remarks (F)	Reference (G)
80 Rabbit (concluded)	Entire body, 2 r/ds Radium (females only)	Radium (gamma)	Approximately 9 mo	Decreased absolute lymphocytes, platelets.		15
81	Entire body, 4 r/ds Radium (females only)	Radium (gamma)	Approximately 2 mo	Decreased absolute lymphocytes.		15
82	Entire body, 4 r/ds Radium (females only)	Radium (gamma)	Approximately 9 mo	Decreased absolute lymphocytes, platelets.		15
83	Entire body, 8 r/ds Radium (females only)	Radium (gamma)	Approximately 2 mo	Decreased absolute lymphocytes.		15
84	Entire body, 8 r/ds Radium (females only)	Radium (gamma)	Approximately 9 mo	Decreased absolute lymphocytes, platelets.		15
85	Entire body, 1 r/ds, 6 da/wk (cyclotron)	Fast neutrons (cyclotron)	15-45 n	Decreased absolute lymphocytes, WBC.		9
86	Entire body, 1 r/ds, 6 da/wk (cyclotron)	Fast neutrons (cyclotron)	50-75 n	Decreased absolute lymphocytes, absolute neutrophils, WBC.		9
87	Entire body, 1.7 n/ds, 6 da/wk (cyclotron)	Fast neutrons (cyclotron)	325 n	Decreased absolute lymphocytes, absolute neutrophils, WBC, RBC, hemoglobin.		9
88	Entire body, 10 r/ds, 6 da/wk	X ray 1000 kvp	60-120 r	Decreased absolute lymphocytes, absolute neutrophils, WBC.		5
89	Entire body, 10 r/ds, 6 da/wk	X ray 250 kvp	720-960 r	Decreased WBC, absolute lymphocytes, absolute neutrophils, RBC, platelets.		5
90	Intraperitoneal, intravenous, or intracardial injection	Radium	0.1 µg	Decreased WBC.		28
91	Intraperitoneal, intravenous, or intracardial injection	Radium	0.1-0.2 µg	Macrocytic anemia.		28
92	Intravenous or intramuscular injection	Plutonium	0.0062 µc/g of body weight	Decreased WBC.	Sustained	28
93	Intraperitoneal injection	Strontium 89	1.0 µc/g of body weight	Decreased heterophils.		28
94	Intraperitoneal injection	Strontium 89	3.0 µc/g of body weight	Decreased heterophils, lymphocytes, RBC, hemoglobin. Morphological changes in WBC and RBC.	RBC and hemoglobin returned quickly to normal.	27
95 Rat	Entire body, single exposure	X ray 250 kvp	5 r	Decreased absolute lymphocytes.		33
96	Entire body, single exposure	X ray 250 kvp	10 r	Decreased absolute lymphocytes, WBC.		33
97	Entire body, single exposure	X ray 250 kvp	50 r	Decreased neutrophils, WBC, absolute lymphocytes, reticulocytes.		33

Entire body, single exposure	X or gamma rays	2-75 r	Increased incidence of lymphocytes with bionized nuclei.	Maximum incidence 1 wk after exposure.	34
Entire body, single exposure	X ray 200 kv	85 r	Lymphocytes decreased by 12th hr after exposure. Neutrophils increased transiently.	Recovery began 2nd da after exposure. Approached normal in 4 wk.	35
Entire body, single exposure	X ray	5-100 r	Decreased iron ⁵⁹ uptake by RBC.	Decrease more marked with greater doses.	36
Entire body, single exposure	X ray	100 r	Decreased absolute lymphocytes, neutrophils, WBC, reticulocytes, platelets		33
Entire body, single exposure	X ray 250 kv	175 r	Lymphocytes decreased by 4th hr. All WBC decreased after 24 hr.	Recovery approximately 12-16 da after exposure.	35
Entire body, single exposure	X ray 200 kv	50-250 r	Decreased iron ⁵⁹ uptake by RBC.	Depression varies as inverse function of radiation dose.	36
Entire body, single exposure	X ray	300 r	Decreased absolute lymphocytes, neutrophils, WBC, reticulocytes, platelets, RBC, hemoglobin.		33
Entire body, single exposure	X ray 250 kv	600 r	Marked decrease in rate and extent of WBC migration in vitro 5 da after exposure.	Effect on WBC, not on plasma.	31
Entire body, single exposure	X ray 200 kv		Surface phagocytosis by WBC in vitro decreased 72 hr after exposure.	Returned to normal by 4th or 5th da after exposure.	5
			Extracts of WBC obtained 3 da after exposure had lost bactericidal activity against test organism (<i>Micrococcus aureus</i>).		5
Entire body, 0.5 r/da	X ray	2 yr	Decreased absolute lymphocytes.		9
Entire body, 1 r/da	X ray	1 yr	Decreased absolute lymphocytes.		9
Entire body, 6 da/wk	Fast neutrons	1 mo	Decreased absolute lymphocytes.		37
Entire body, 1.7 n/da, 6 da/wk	Fast neutrons	1 yr	Decreased absolute lymphocytes, RBC, hemoglobin.	Minimum levels lower with greater doses. Rate of WBC decrease per r greater with smaller doses.	38
Entire body, 1.7 n/da, 6 da/wk	Fast neutrons	0.1-4.5 μ C/kg of body weight	Prompt leukopenia with minimum levels 12-14 da after exposure.	Lymphocyte depression more marked with greater doses. Neutrophils reached minimum level of only 4% of normal 14 da after 17 μ C/kg.	38
Single intraperitoneal injection	Phosphorus ³²	17 μ C/kg of body weight	Decreased WBC, platelets (all).	Spiking reticulocytosis 160 da after injection.	38
Single intravenous injection	Radium chloride	300 μ C/kg of body weight	Decreased WBC, platelets, anemia.	Platelet depression maximum about 7 da after injection, then markedly elevated between 15-50 da.	38
Single intravenous injection	Radium chloride	100 μ C/kg of body weight	Decreased WBC, platelets, RBC, and reticulocytes.		38
Single intravenous injection	Plutonium citrate	16.9 μ C/kg of body weight	Decreased lymphocytes, neutrophils, platelets.		38
Single intravenous injection	Plutonium citrate	47 μ C/kg of body weight	Decreased WBC, platelets, RBC, hemoglobin.		38

95. EFFECT OF IONIZING RADIATION ON PERIPHERAL BLOOD: VERTEBRATES (Concluded)

Part II. VERTEBRATES OTHER THAN MAN (Concluded)

Animal	Exposure or Administration (B)	Type of Radiation (C)	Accumulated Dosage or Exposure Time (D)	Effect (E)	Remarks (F)	Reference (G)
116 Rat (concluded)	Single intravenous injection	Plutonium citrate	95 $\mu\text{C/kg}$ of body weight	Decreased WBC, platelets, RBC, hemoglobin, reticulocytes.	With doses above 126 and 189 $\mu\text{C/kg}$, precipitous fall in RBC and hemoglobin suggests hemolytic process.	38
117	Radium, single intraperitoneal injection	Radium chloride	10-50 μg	Transient increase in platelets on 3rd da after injection, followed by progressive decrease.		39
118	Radium, parenteral injection	Radium chloride	0.02-0.03 μg	Decreased WBC.		28
119	Radium, parenteral injection	Radium chloride	0.1-0.2 μg	Macrocytic anemia.		28
120	Intravenous or intramuscular injection	Plutonium	0.0062 $\mu\text{C/g}$ of body weight	Moderate, sustained WBC decrease.		28
121	Intravenous or intramuscular injection	Plutonium	More than 0.0063 $\mu\text{C/g}$ of body weight	Decreased WBC, RBC, reticulocytes, platelets.	All decreases sustained.	28
122	Single intraperitoneal injection	Strontium ⁸⁹	0.22-0.25 $\mu\text{C/g}$ of body weight	Decreased neutrophilia.	More marked neutrophil decrease; also lymphopenia with higher doses.	27
123	Gavage	Yttrium ⁹¹	10 $\mu\text{C/g}$ of body weight	Transient decrease in absolute lymphocytes.		40
124	Gavage	Yttrium ⁹¹	20 $\mu\text{C/g}$ of body weight	Transient initial increase in RBC and hemoglobin followed by severe anemia; maximal 20 da after administration of Yttrium ⁹¹ .		40
125	Ingestion	Yttrium ⁹¹	0.3-2.0 $\mu\text{C/g}$ of body weight	Decreased lymphocytes; minimum levels at 90 da.	Prompt return to normal levels after Yttrium ⁹¹ feeding discontinued.	40
126 Swine	Entire body, single exposure	X ray 1000 kv	200 r	Decreased WBC.		41
127	Entire body, single exposure	Nuclear explosion (Bikini tests)	Near lethal dose (survived 17 mo)	Macrocytic anemia.	One of two swine under observation also had achlorhydria.	42
128	Entire body, single exposure	Nuclear explosion (Bikini tests)	1300 r in 4-da period, or 1500 r in 5-da period	Decreased WBC and platelets. Many degenerative and abnormal WBC.	WBC less than 50% of normal on 4th da; platelets virtually absent on 9th da.	43
129 Chick	Entire body, single exposure	X ray 85 kv	360 r	Decreased lymphocytes.	Fully recovered from anemia in 4 wk; marked reticulocytosis 4 wk after exposure.	44
130	Entire body, single exposure	X ray 200 kvp	600 r	Decreased lymphocytes, heterophils, thrombocytes, RBC, hemoglobin, reticulocytes.		7

131	Intraperitoneal injection, 1.5 μ C/g of body weight every 3 wk	Phosphorus-32	Total dose, 4.5 μ C/g of body weight	Anemia (most marked 24th post-injection wk).	45
132	Single subcutaneous injection ³	Phosphorus-32	235 μ C	Decreased lymphocytes, WBC, and RBC.	46
133	Single subcutaneous injection ⁴	Phosphorus-32	300 μ C	Fatal anemia.	47
134	Single subcutaneous injection ⁵	Phosphorus-32	760 μ C	Fatal anemia.	47
135	Salmon	Entire body, single exposure	X ray 200 kv	Decreased nucleated cells.	48

1/1 6-da-old chick. 7/4 19-da-old chick. 1/5 28-da-old chick.

Contributor: Ingram, Marylou

- References: [1] Smith, D. E., et al. 1951. ANL-4794. Technical Information Service, Oak Ridge, Tenn. [2] Trum, B. F., et al. 1933. Am. J. Physiol. 174:57. [3] Ingram, Marylou, 1956. Proc. Internat. Conf. Peaceful Uses Atomic Energy, Geneva 13-210. United Nations, New York. [4] Prosser, C. L., et al. 1947. Plutonium Project Record, v. 22B 2D. CH-3738. Technical Information Service, Oak Ridge, Tenn. [5] Ingram, Marylou, and W. B. Mason. Hill, New York. [6] Prosser, C. L., et al. MDDC 631, MDDC 1046, Technical Information Service, Oak Ridge, Tenn. Also 1947. Radiology 49:299. [7] Zirkle, R. E., ed. 1950. Biological effects of external X and gamma radiation. Part I, TID-5220. Technical Information Service, Oak Ridge, Tenn. [8] Constant, M. A., and P. F. Hahn. 1955. Cancer, Phila. 8:450. [9] Ely, J. O., et al. 1950. UR-92. Technical Information Service, Oak Ridge, Tenn. [10] Jackson, A. A., and P. F. Hahn. 1955. Cancer, Phila. 8:450. [11] Wheeler, B., et al. 1951. Am. J. Physiol. 166:363. [12] Nelson, J. E., et al. 1949. Acta Univ. internat. cancer, Brux. 6:819. [13] Smith, M. N., et al. 1946. CH-3868. Technical Information Service, Oak Ridge, Tenn. [14] Lorentz, E. 1951. J. clin. phys., Par. 48:264. [15] Lorentz, E., et al. 1946. CH-3868. Technical Information Service, Oak Ridge, Tenn. [16] Fulton, G. P., et al. 1954. Blood, N. Y. 6:622. [17] Smith, W. W. 1955. Ann. N. York Acad. Sc. 59:815. [18] French, A. B., et al. 1953. Am. J. Physiol. 182:469. [19] Hahn, M. V., and E. Patterson. 1956. Brit. J. Radiol. 29:148. [20] Edington, G. M., et al. 1956. J. Path. Bact., Lond. 71:277. [21] Nettleship, A. 1949. Radiology 42:64. [22] Storer, J. B., and W. H. Langham. 1954. USAECU-3099. Technical Information Service, Oak Ridge, Tenn. [23] Upton, A. C., et al. 1954. Cancer Res. 14:682. [24] Swanson, H. D., et al. USAECU-3130. Technical Information Service, Oak Ridge, Tenn. [25] Swanson, H. D., et al. USAECU-3130. Technical Information Service, Oak Ridge, Tenn. [26] Evers, T. 1948. Radiology 50:811. [27] Simmons, R. L., and L. O. Jacobson. 1946. MDDC-1387 CH-3797. Technical Information Service, Oak Ridge, Tenn. [28] Jacobson, L. O., and E. L. Simmons. [29] Shochmeister, I. C., and P. Fishman. 1950. J. Exp. Med. 101:259. [30] Jacobson, L. O., and E. L. Simmons. [31] Shochmeister, I. C., and P. Fishman. 1950. J. Exp. Med. 101:259. [32] Jacobson, L. O., and E. L. Simmons. [33] Jacobson, L. O., and E. L. Simmons. [34] Delbeck, S., and J. B. Storer. 1954. Am. J. Roentg. 71:846. [35] Hennessy, T. G., and R. L. Huff. 1950. Proc. Soc. Exp. Biol., N. Y. 73:436. [36] Hennessy, T. G., and R. L. Huff. [37] Mitra, S., et al. 1953. Acta radiol., Stockh. 40:593. [38] Fink, R. M., ed. 1950. Biological studies with polonium, radium, and plutonium. NNEs VI-2. McGraw-Hill, New York. [39] Hoofe, P. G., et al. 1950. Anat. Rec. 106:537. [40] Jacobson, L. O., and E. L. Simmons. 1946. USAECU-2037. Technical Information Service, Oak Ridge, Tenn. [41] Cronkite, E. P., et al. 1949. NMR, NM-007-039. [42] Larson, P. D., and E. P. Cronkite. 1949. Yale J. Biol. 22:57. [43] Cronkite, E. P., et al. 1948. Growth, Ithaca 12:107. [44] Warren, S. L., and D. J. Dixon. 1949. Radiology 52:669. [45] Dehard, W. F., et al. 1950. Proc. Centr. Soc. Clin. [46] Bonham, K., et al. 1948. Growth, Ithaca 12:107. [47] Dixon, F. J. 1948. Proc. Soc. Exp. Biol., N. Y. 68:505. [48] Bonham, K., et al. 1948. Growth, Ithaca 12:107.

96. EFFECT OF IONIZING RADIATION ON BLOOD PRESSURE AND PULSE RATE: VERTEBRATES

Blood pressure values for man given as systolic/diastolic. Blood pressure values for other animals are mean arterial pressure, except in line 12. Method: *S* = sphygmomanometer; *H* = saline-mercury manometer; *G* = strain gauge; *P* = piezometer.

Animal	No. of Sub-jects	Radiation and Dosage	Arterial Blood Pressure, mm Hg										Pulse Rate, beats/min					Refer-ence				
			Time After Exposure										Time After Exposure									
			Con-trol	hr					da					Con-trol	hr							
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)	(K)	(L)	(M)	(N)	(O)	(P)	(Q)	(R)	(S)	(T)	(U)	(V)	
1	12	Accidental nuclear reaction. 114 r y, 1930 r X rays (soft)	120/78	110/70				100/65	120/65						S		104	95	85	130		1
2	12	100 r y, 380 r X rays (soft)	120/80	120/70				100/60	120/80					70/50	S		68	75	78	105	250	1
3	1	X rays (soft) 26.4 r y, 390 r X rays (soft)	140/90	136/96				120/80		140/90				135/90	S			80	75	82	85	1
4	1	1 r y, 31 r X rays (soft)	138/90	140/90						120/80					S		60					1
5	1	2.7 r y, 42 r X rays (soft)	120/82					108/72							S							1
6	8	X ray therapy to chest region. 100-700 r (single exposure)	140/	124/	130/										S	84	76					2
7	35	1,000-15,000 r (fractionated) ³	149/											141/	S							2
8	2	X ray, 300 r	117	122	105										H	88	97	84	90	130	150	3
9	1	α particles (Pu ²³⁹), 0.358 μg/g, intra-venous	123											105	H	95	93	84	88	135	4	
10	Rabbit	3-13 X ray, 800 r	99	58	64	74									H	198	298					5
11	4-11 X ray, 50-60 r	98	70	83	77										H							5
12	21 X ray, 630 r	109/66	58/28	64/34											G	282	282					6
13	Rat	7-8 X ray, 1500 r	95												H							7
14	5 X ray, 600 r	107	68	86											H							8
15	Chick, 5 X ray, 600 r	85	76	65	59										H							9
16	3-4 d-215	1000 r	85	67	36	24									H							9
17	Pigeon 6-7 X ray, 2200-2400 r	138	141	137	147										H							10
18	Rooster 6-7 X ray,	150	117	108	120										H							11,12
19	13-17 d 1200 r	150	127	117	108/117										H							11,12

/1/ Dosages given in equivalent terms of total body radiation. Some regions received extremely large amounts of radiation. /2/ Non-survivors at 30 days. /3/ 10-20 daily exposures. Blood pressure during therapy period. /4/ 24-hour survivors. /5/ Non-survivors at 10 hours. /6/ Non-survivors at 24 hours. Contributor: Stearns, S. P., H. Lisco, and J. G. Hoffman. 1952. Ann. Int. M. 36:280. /2/ Bedärfing, G., and G. Grüssner. 1949. Strahlentherapie 78:443. /3/ Prosser, C. L., E. E. Painter, and M. N. Swift. 1956. In R. E. Zirkle, ed. Biological effects of external X and gamma radiation. Part 2, TID-5228. Dep. of Commerce, Office of Technical Services, Washington. pp. 1-99. /4/ Painter, E. E., E. Russell, C. L. Prosser, M. N. Swift, W. Kistelski, and G. Sacher. 1946. Clinical physiology of dogs injected with plutonium. USAEC-2042. Technical Information Service, Oak Ridge, Tenn. /5/ Painter, E. E., C. L. Prosser, and M. C. Moore. 1956. In R. E. Zirkle, ed. Biological effects of external X and gamma radiation. Part 2, TID-5220. Dep. of Commerce, Office of Technical Services, Washington. pp. 147-181. /6/ Brooks, P. M., H. B. Gerstner, and S. A. Smith. 1956. Am. J. Physiol. 186:532. /7/ Montgomery, P. O., and S. Warren. 1951. Proc. Soc. Exp. Biol., N. Y. 77:803. /8/ Weber, R. P., and F. R. Siegel. 1949. Ibid. 70:261. /9/ Stearns, S. P., A. M. Brues, M. H. Sanderson, and E. J. Christian. 1955. Am. J. Physiol. 182:407. /10/ Stearns, S. P., M. H. Sanderson, and E. J. Christian. 1957. The initial post irradiation period in the pigeon. ANL-5732. Q. Rep. Biol. Med. Res. Div., Argonne National Laboratory. /11/ Stearns, S. P., M. H. Sanderson, E. J. Christian, and A. M. Brues. 1956. Am. J. Physiol. 184:134. /12/ Stearns, S. P., et al. 1957. Unpublished.

97. EFFECT OF IONIZING RADIATION ON TOTAL BLOOD VOLUME AND PLASMA VOLUME: VERTEBRATES

Abbreviations TBV = total blood volume; PV = plasma volume; RBCM = red blood cell mass. Values for blood and plasma volume are in % of body weight.

Abbreviations			Time After Exposure			Time After Exposure												Method		Reference																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																														
Animal			Radiation and Dosage			Measure-ment			Control			hr			da									Method		Reference																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																								
(A)			(B)			(C)			(D)			(E)			(F)			(G)			(H)			(I)			(J)			(K)			(L)			(M)			(N)			(O)			(P)			(Q)			(R)																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																															
1	Dog	8	X ray,	200-400 r	TBV	9.1																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																												

1/1 Corrected for T-1824 removed from blood; factor = 1.2. 1/2 Corrected for T-1824 removed from blood; factor = 1.08.

Contributor: Stearns, S. Phyllis

References: [1] Prosser, C. L., E. E. Painter, and M. N. Swift, 1956, In R. E. Zirkle, ed. Biological effects of external X and gamma radiation, Part 2, TID-5226. Dep. of Commerce, Office of Technical Services, Washington, pp. 1-99. [2] Soberman, R. J., R. P. Keating, and R. D. Maxwell, 1951, Am. J. Physiol., 164 450. [3] Painter, E. E., E. Russell, C. L. Prosser, M. N. Swift, W. Mileleski, and G. Sacher, 1946, Clinical physiology of dogs injected with plutonium, USAEC-D-2042, Technical Information Service, Oak Ridge, Tenn. [4] Storey, R. H., L. Wish, and J. Furth, 1950, Proc. Soc. Exp. Biol., N. Y., 74-242. [5] Painter, E. E., C. L. Prosser, and M. C. Moore, 1956, In R. E. Zirkle, ed. Biological effects of external X and gamma radiation, Part 2, TID-5226. Dep. of Commerce, Office of Technical Services, Washington, pp. 147-181. [6] Montgomery, P. O., and E. Warren, 1951, Proc. Soc. Exp. Biol., N. Y., 77:601. [7] Supplee, H., J. D. Hauschildt, and C. Entenman, 1952, Am. J. Physiol., 161:183 [8] France, O., 1956, In R. E. Zirkle, ed. Biological effects of external X and gamma radiation, Part 2, TID-5226. Dep. of Commerce, Office of Technical Services, Washington, pp. 411-427. [9] Stearns, S. P., M. H. Suberson, and E. J. Christian, 1956, Changes in organ blood volume in X-irradiated chicks, ANL-5676, Q. Rep. Biol., Med. Res. Div., Argonne National Laboratory, pp. 31-34. [10] Stearns, S. P., et al., 1957, Unpublished.

98. EFFECT OF IONIZING RADIATION ON ORGAN BLOOD VOLUME: CHICKEN

	Organ	Condition	Organ Weight g (C)	RBC Mass ¹		Plasma Volume (F)	Method (G)	Reference (H)
				Active (D)	Inactive ² (E)			
Chick ³								
1	Liver	Control	1.69	7.4	4.0	15.7	RBC mass: Cr ⁵¹ , p ³² . Plasma volume: I ¹³¹ human serum albumin	1
2		Irradiated ⁴	1.52	3.6	9.2	7.2		
3	Spleen	Control	0.022	2.9	2.7	16.8		
4		Irradiated ⁴	0.020	1.5	4.8	7.1		
5	Kidney	Control	0.245	7.9	1.2	24.7		
6		Irradiated ⁴	0.259	5.2	7.5	9.4		
7	Lung	Control	0.210	17.1	2.1	40.3		
8		Irradiated ⁴	0.217	18.0	2.1	20.9		
9	Proventriculus	Control	0.380	1.0	0	6.25		
10		Irradiated ⁴	0.449	0.74	1.76	4.05		
11	Duodenum	Control	0.257	1.14	0.46	6.64		
12		Irradiated ⁴	0.205	1.87	1.16	3.06		
13	Skeletal muscle	Control		0.91	0.19	2.85		
14		Irradiated ⁴		0.94	0.31	1.70		
Rooster ⁵								
15	Liver	Control	56.84	19.4	7.1	34.8	RBC mass Cr ⁵¹ . Plasma volume: I ¹³¹ human serum albumin	2
16		Irradiated ⁶	44.33	14.3	7.6	16.6		
17	Spleen	Control	4.68	16.5	6.4	22.1		
18		Irradiated ⁶	1.79	6.5	12.7	8.4		
19	Kidney	Control	10.11	20.0	4.7	28.1		
20		Irradiated ⁶	10.02	14.0	4.0	15.4		
21	Lung	Control	10.34	28.8	8.2	27.5		
22		Irradiated ⁶	9.10	31.3	4.3	17.7		
23	Proventriculus	Control	4.44	2.63	0.03	6.80		
24		Irradiated ⁶	4.91	1.64	0.15	3.80		
25	Duodenum	Control	3.69	2.86	0	7.40		
26		Irradiated ⁶	3.53	2.83	0.12	4.10		
27	Skeletal muscle	Control		0.25	0.06	0.63		
28		Irradiated ⁶		0.45	0.17	0.59		

/1/ % of total organ weight. /2/ RBC mass which is pooled or inactively circulating and does not contribute to the actively circulating RBC mass. /3/ 10 chicks, 3 to 4 days after hatching. /4/ 3-5 hours after irradiation with 1000 r X rays. /5/ 7-9 animals. /6/ 6-8 hours after irradiation with 1200 r X rays.

Contributor: Stearner, S. Phyllis

- References: [1] Stearner, S. P., M. H. Sanderson, and E. J. Christian. 1956. Changes in organ blood volume in X-irradiated chicks. ANL-5696. Q. Rep. Biol. Med. Res. Div., Argonne National Laboratory. pp. 31-34.
[2] Stearner, S. P., et al. 1957. Unpublished.


99. BLOOD COAGULATION

Differing theories for Stages II and III on the mechanics of blood coagulation are presented in Parts II and III, respectively.


Part I: STAGES OF BLOOD COAGULATION (According to Monkhouse)

Synonymous terms for clotting factors: PTC (plasma thromboplastin component) = Christmas factor, platelet cofactor-2, autoprothrombin-2; AHF (antithemophilic factor) = antithemophilic globulin, thromboplastinogen; Ac-globulin = labile factor, factor V, proaccelerin; Factor VII = stable factor, autoprothrombin-1, proconvertin, cothromboplastin. Plasminogen = profibrinolysin; Plasmin = fibrinolysin. Symbols. \longrightarrow gives rise to, \dashrightarrow acts on, \longleftarrow inhibits or destroys, $\boxed{}$ present in blood.


Stage I. Formation of Plasma Thromboplastin



Stage II: Formation of Thrombin




Stage III. Change of Fibrinogen to Fibrin



Part II FORMATION OF THROMBIN (According to Seegers)

Prothrombin, which is regarded as being the only source of thrombin, is converted to thrombin by an enzyme also called thrombin. Substances that support the enzyme in this function are known as procoagulants and include derivatives of prothrombin. Substances that interfere with the activation of prothrombin, or of its derivatives, are known as anticoagulants. Fibrin-s is soluble in urea solution, whereas Fibrin-i is insoluble.




/1/ Related to prothrombin production.

Contributor: Seegers, Walter H.

Part III CHANGE OF FIBRINOGEN TO FIBRIN (According to Coon)

Symbols: \longrightarrow gives rise to, \dashrightarrow acts on, \longrightarrow inhibits or destroys, $\boxed{}$ present in blood.



Contributor: Coon, William W.

References: [1] Lorand, L. 1954. *Physiol. Rev.* 34 742. [2] Sherry, S., et al. 1954. *Ibid* 34 736. [3] Astrup, T. 1956. *Blood*, N. Y. 11:781.

100. NATURAL-OCCURRING COAGULANTS AND ANTICOAGULANTS. PREPARATION AND ASSAY

For additional information on preparation and assay, see references 44-47.

Substance	Distribution	Preparation		Assay	
		Method	Reference (D)	Method (E)	Reference (F)
1 Ac-globulin	Plasma, serum, platelets	From plasma, Ware and Seegers	1	Rate of formation of thrombin from purified prothrombin	1
		Lewis and Ware	2		
		From platelets, Ware, Fahey and Seegers	3	Use of aged oxalated plasma	2
2 Antihemophilic factor	Fresh plasma	Adsorption on kaolin	4	Thromboplastin generation test	29
		Fractional precipitation	5,6	Ability to promote prothrombin consumption in hemophilic plasma	6
3 Antithrombin	Plasma, in trace amounts, in the α -globulin fraction	Adsorption by aluminum hydroxide	7,8	Incubation with excess thrombin and calculation of thrombin remaining	7,27
4 Factor VII	Serum	Adsorption by BaSO_4 and elution	9,10	Acceleration of thrombin formation	10
5 Fibrinogen	Plasma, 120-470 mg %	Cold ethyl alcohol fractionation, salt fractionation, centrifugation of plasma at 0°C	11	Purity based on clotability (i.e. % clottable protein)	30
6 Heparin	Connective tissue generally; high level in liver of dog, ox, pig, and sheep, in lung of ox and skin of rat, associated with distribution of mast cells	Charles and Scott	12	On bases of anticoagulant potency	
		Bell and Jaques	13	Whole blood	31,32
		Monkhouse	14,15	Recalcified blood	33
		Cass, Riley and West	16	Antithrombin potency	32
		Snellman, Jensen and Sylven	17	Metachromatic potency	34,35
				Combining power:	
				With protamine	36
				With toluidine blue	37
7 Heparin-cofactor	Plasma in the α -globulin fraction (may be identical to antithrombin)	Adsorption by aluminum hydroxide	7,8	Inhibits clotting of fibrinogen by thrombin in presence of heparin	7
8 Lipid anticoagulant	Blood, plasma	Cold methanol extraction	18,19	Inhibition of clotting time of recalcified plasma	38
				Activation of decalcified plasma by brain thromboplastin	19
9 Lipid antithromboplastin	Brain, spinal cord	Purified serine phosphate fractions	20	Inhibits action of tissue thromboplastin, blocks formation of plasma thromboplastin	39
10 Plasma thromboplastin component (PTC)	Plasma and serum	Adsorption and elution	21	With PTC deficient blood	21
				Thromboplastin generation test	40
11 Plasminogen	Plasma	Ammonium sulfate precipitation	22	Lysis of standard clot	23
		Isoelectric precipitation and adsorption by kaolin	23	Digestion of casein	
12 Platelets and platelet fractions	Whole blood or lightly centrifuged plasma	Fractional centrifugation	24,25	For whole platelets, microscopic observations and counting; for fractions, specific for each material. (Not yet completely worked out.)	30
13 Prothrombin	Estimated 15 mg % in plasma	Acid precipitation from diluted plasma adsorbed by magnesium hydroxide and eluted with CO_2 under pressure	26	One-stage method	26,41
		Adsorption on barium citrate	2	Two-stage method	42
		Other methods	11		

Substance	Distribution	Preparation		Assay	
		Method	Reference	Method	Reference
(A)	(B)	(C)	(D)	(E)	(F)
14 Thrombin	From prothrombin	Conversion of prothrombin with thromboplastin and calcium	27	According to its ability to clot fibrinogen	43
		Conversion with 25% citrate	28	Hydrolysis of p-toluene-sulfonylarginine methyl ester	41

Contributors: (a) Monkhouse, Frank C., (b) Tocantins, L. M.

- References.** [1] Ware, A. G., and W. H. Seegers. 1948. *J. Biol. Chem.* 172:699. [2] Lewis, M. L., and A. G. Ware. 1953. *Proc. Soc. Exp. Biol., N. Y.* 84:636. [3] Ware, A. G., J. L. Fahey, and W. H. Seegers. 1948. *Am. J. Physiol.* 154:140. [4] Lorand, L., and K. Laki. 1954. *Biochim. biophys. acta*, *Amst.* 13:448. [5] Bidwell, E. 1955. *Brit. J. Haemat.* 1:35. [6] Spaet, T. H., and B. G. Kinsell. 1954. *Proc. Soc. Exp. Biol., N. Y.* 84:314. [7] Monkhouse, F. C., and D. W. Clark. 1957. *Canad. J. Biochem. Physiol.* 35:373. [8] Monkhouse, F. C., E. France, and W. H. Seegers. 1955. *Circul. Res., N. Y.* 3:397. [9] DeVries, A., B. Alexander, and R. Goldstein. 1949. *Blood*, *N. Y.* 4:247. [10] Duckert, F., F. Koller, and M. Matter. 1953. *Proc. Soc. Exp. Biol., N. Y.* 82:259. [11] Ware, A. G., and G. F. Lanchantin. 1954. *Physiol. Rev.* 34:714. [12] Charles, A. F., and D. A. Scott. 1936. *Biochem. J., Lond.* 30:1927. [13] Bell, Helen J., and L. B. Jaques. 1956. *Bull. Soc. chim. Belg.* 65:36. [14] Monkhouse, F. C. 1956. *Canad. J. Biochem. Physiol.* 34:757. [15] Monkhouse, F. C., and Ruth MacKneson. 1957. *Fed. Proc., Balt.* 16:89. [16] Cass, R., J. F. Riley, and G. B. West. 1954. *Nature*, *Lond.* 174:318. [17] Snellman, O., R. Jensen, and B. Sylven. 1948. *Ibid.* 161:639. [18] Goldsmith, D. P. J., and C. W. Mushett. 1954. *J. Biol. Chem.* 211:169. [19] Carroll, R. T., and L. M. Tocantins. 1949. *Tr. Conf. Blood Clotting*, *N. Y.* 2:11. [20] Silver, M. J., D. L. Turner, and L. M. Tocantins. 1957. *Am. J. Physiol.* 190:8. [21] Aggeler, P. M., S. G. White, Mary B. Glendenning, E. W. Page, T. B. Leake, and G. Bates. 1952. *Proc. Soc. Exp. Biol., N. Y.* 79:692. [22] Loomis, E. C., C. George, Jr., and A. Ryder. 1947. *Arch. Biochem., N. Y.* 12:1. [23] Remmert, L. F., and P. P. Cohen. 1949. *J. Biol. Chem.* 181:431. [24] Schneider, C. L., E. B. Claxton, C. H. Hughes, and S. A. Johnson. 1954. *Am. J. Physiol.* 179:236. [25] Dillard, G. H. L., G. Brecher, and E. P. Cronkite. 1951. *Proc. Soc. Exp. Biol., N. Y.* 78:796. [26] Seegers, W. H., E. C. Loomis, and J. M. Vandenbelt. 1945. *Arch. Biochem., N. Y.* 6:85. [27] Seegers, W. H., K. D. Millar, E. B. Andrews, and R. C. Murphy. 1952. *Am. J. Physiol.* 169:700. [28] Seegers, W. H., R. I. McClaughry, and J. L. Fahey. 1950. *Blood*, *N. Y.* 5:421. [29] Biggs, R., J. Eveling, and G. Richards. 1955. *Brit. J. Haemat.* 1:20. [30] Tocantins, L. M. 1955. *The coagulation of blood: methods of study*. Grune and Stratton, New York. [31] Jorpes, J. E. 1955. *Acta pharm. tox., Kbh.* 11:367. [32] Jaques, L. B., and A. F. Charles. 1941. *Q. J. Pharm., Lond.* 14:1. [33] Kuizenga, M. H., J. W. Nelson, and G. F. Cartland. 1943. *Am. J. Physiol.* 139:612. [34] Jaques, L. B., M. Bruce-Mitford, and A. G. Ricker. 1947. *Rev. canad. biol.* 6:740. [35] Jaques, L. B., F. C. Monkhouse, and M. Stewart. 1949. *J. Physiol., Lond.* 109:41. [36] Allen, J. G., M. Sanderson, M. Milham, A. Kirschon, and L. O. Jacobson. 1948. *J. Exp. M.* 87:71. [37] Mackintosh, F. C. 1941. *Biochem. J., Lond.* 35:776. [38] Mushett, C. W., D. J. Goldsmith, and K. L. Kelly. 1954. *J. Biol. Chem.* 211:163. [39] Silver, M. J., D. L. Turner, and L. M. Tocantins. 1959. *Progr. Hemat.* 2:264. [40] Biggs, R., and A. S. Douglas. 1953. *J. Clin. Path., Lond.* 6:23. [41] Sherry, S., and W. Troll. 1954. *J. Biol. Chem.* 208:95. [42] Warner, E. D., K. M. Brinkhous, and H. P. Smith. 1936. *Am. J. Physiol.* 114:667. [43] Seegers, W. H., and H. P. Smith. 1942. *Ibid.* 137:348. [44] Owren, P. A. 1949. *Scand. J. Clin. & Invest., Oslo* 1:81. [45] Quick, A. J. 1947. *Am. J. M. Sc.* 214:272. [46] Riley, J. F., D. M. Shepherd, G. B. West, and S. W. Stroud. 1955. *Nature*, *Lond.* 176:1123. [47] Seegers, W. H., and D. A. McGinty. 1942. *J. Biol. Chem.* 146:511.

101. ORAL ANTICOAGULANTS: MAN

All subjects, except those treated with Warfarin and Coumatron Series II, had an initial basal prothrombin concentration of 70% or above. The major purpose for establishing this criterion was to create a uniform group of subjects with a normal, coagulative response for the proper evaluation of therapeutic effectiveness. The same therapeutic range, 10-25% prothrombin concentration (21-45 seconds), was used in all groups, except for Warfarin in which a range of 10-35% was used. Prothrombin values in all groups were determined by the method of Quick [7]. Values in parentheses are ranges, estimate "c" (cf. Introduction)

Drug	No. of Sub-jects	Dose mg.			Achievement of Therapeutic Effect da	Subjects Achieving Therapeutic Effect ² %			Average Recovery Time ³ da	Subjects with Prothrombin Escape		Subjects with Prothrombin Value <10% (M)	Incidence of Bleeding ⁴ (N)	Reference
		Effective Initial	Usual Maintenance	Cor-rec-tive (level)		Within 24 hr (G)	Within 48 hr (H)	Within 72 hr (I)		% (K)	Duration, da (J)			
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)	(K)	(L)	(M)	(N)	(O)
1 Dicumarol (3, 3'-Methylene-bis-4-hydroxycoumarin)	122	646	(25-125)	256	2.3	14	55	80	5	37	1.5	15	11% of 303 cases	1, b
2 Tromexan (3, 3'-Carboxy-methylene-bis-4-hydroxycoumarin)	50	(1500-1800)	(300-1200)	2100	2.1	12	82	96	2.5	90	1.8	20	9% of 100 cases	1, b
3 Warfarin (3-(6-Acetylbenzyl)-4-hydroxycoumarin)	100	(40-50)	9.7(6.25-12.5)			23	75	85					5% of 100 cases	2, b
4 Liqummar (3-(11-Phenylpropyl)-4-hydroxycoumarin)	100	(15-18)	(3-9)		2.0	25	90	92					5% of 100 cases	b
Coumatron Series I (3, 4-(12'-Methyl-2'-methoxy-4'-phenyl)pyranocoumarin)	10	380	77(25-175)	135	2.1	0	50	100	3.3	58	1.4	15	8% of 13 cases	3
Series II	100					5	85	96		4		22	5% of 100 cases	4
									1.5	81		27	11% of 70 cases	5
Sintrom (Nirophenyl acetyl ethyl-4-oxy-coumarin)	52	58	(6-16)		2.3	2	60					10	6% of 200 cases	1
Phenindione (2-Phenyl-1, 3-indanedione)	133	685	130(25-200)	430	1.3	18	95	100	1.5	17	1.6			
Dypaxin (2-Diphenylacetyl-1, 3-indanedione)	64	63	9(2.5-15)	27	2.3	5	67	89	4.1	18	1.7	10	11% of 104 cases	6

1/1 Mean dose required to return prothrombin below 10% in those subjects with "prothrombin escape." 2/2 Cumulative. 3/3 For 40% recovery or better.

4/4 All cases, regardless of initial prothrombin value.

Contributors: (a) Coon, William W., (b) Wright, Irving S. References: [1] Coon, W. W., I. F. Duff, P. E. Hodgson, and N. England J. A. 255, 491. [2] Duff, I. F. Unpublished. [3] Griffith, P. E., Hodgson, D. R., Griffith, J. A., W. S. Williams, P. W. Willis, J. R. Gamble, D. R. Gamble, and I. F. Duff, 1951, J. Laboret. Clin. M. 42:968. [4] Quick, A. J. 1951. Physiology and pathology of hemostasis. Lea and Febiger, Philadelphia.

5/5 All cases, regardless of initial prothrombin value. Contributors: (a) Coon, William W., (b) Wright, Irving S. References: [1] Coon, W. W., I. F. Duff, P. E. Hodgson, and N. England J. A. 255, 491. [2] Duff, I. F. Unpublished. [3] Griffith, P. E., Hodgson, D. R., Griffith, J. A., W. S. Williams, P. W. Willis, J. R. Gamble, D. R. Gamble, and I. F. Duff, 1951, J. Laboret. Clin. M. 42:968. [4] Quick, A. J. 1951. Physiology and pathology of hemostasis. Lea and Febiger, Philadelphia.

6/6 All cases, regardless of initial prothrombin value. Contributors: (a) Coon, William W., (b) Wright, Irving S. References: [1] Coon, W. W., I. F. Duff, P. E. Hodgson, and N. England J. A. 255, 491. [2] Duff, I. F. Unpublished. [3] Griffith, P. E., Hodgson, D. R., Griffith, J. A., W. S. Williams, P. W. Willis, J. R. Gamble, D. R. Gamble, and I. F. Duff, 1951, J. Laboret. Clin. M. 42:968. [4] Quick, A. J. 1951. Physiology and pathology of hemostasis. Lea and Febiger, Philadelphia.

7/7 All cases, regardless of initial prothrombin value. Contributors: (a) Coon, William W., (b) Wright, Irving S. References: [1] Coon, W. W., I. F. Duff, P. E. Hodgson, and N. England J. A. 255, 491. [2] Duff, I. F. Unpublished. [3] Griffith, P. E., Hodgson, D. R., Griffith, J. A., W. S. Williams, P. W. Willis, J. R. Gamble, D. R. Gamble, and I. F. Duff, 1951, J. Laboret. Clin. M. 42:968. [4] Quick, A. J. 1951. Physiology and pathology of hemostasis. Lea and Febiger, Philadelphia.

8/8 All cases, regardless of initial prothrombin value. Contributors: (a) Coon, William W., (b) Wright, Irving S. References: [1] Coon, W. W., I. F. Duff, P. E. Hodgson, and N. England J. A. 255, 491. [2] Duff, I. F. Unpublished. [3] Griffith, P. E., Hodgson, D. R., Griffith, J. A., W. S. Williams, P. W. Willis, J. R. Gamble, D. R. Gamble, and I. F. Duff, 1951, J. Laboret. Clin. M. 42:968. [4] Quick, A. J. 1951. Physiology and pathology of hemostasis. Lea and Febiger, Philadelphia.

9/9 All cases, regardless of initial prothrombin value. Contributors: (a) Coon, William W., (b) Wright, Irving S. References: [1] Coon, W. W., I. F. Duff, P. E. Hodgson, and N. England J. A. 255, 491. [2] Duff, I. F. Unpublished. [3] Griffith, P. E., Hodgson, D. R., Griffith, J. A., W. S. Williams, P. W. Willis, J. R. Gamble, D. R. Gamble, and I. F. Duff, 1951, J. Laboret. Clin. M. 42:968. [4] Quick, A. J. 1951. Physiology and pathology of hemostasis. Lea and Febiger, Philadelphia.

102. ANTICOAGULANTS IN VIVO: VERTEBRATES

Anticoagulant dosage is usually determined in vitro from a coagulation time--the clotting time for heparin and heparinoid compounds, and the prothrombin time for indirect anticoagulants [4, 26-28]. However, this may not always correspond to the desired action in vivo [29], as the dosage required to prevent coagulation may be many times that indicated by the test in vitro.

Direct Anticoagulants: Heparin and heparinoid compounds, injected intravenously, give a prolonged coagulation time; for some purposes, the high peak blood levels resulting from this mode of administration are essential [2]. Heparin is inactive orally [30], and intramuscular and subcutaneous injections are not as generally effective as those given intravenously [b]. However, the subcutaneous route is the commonly accepted method of choice in doses of 75-100 mg, given each 8-12 hours (depending on the clotting time [b]). It is desirable that the clotting time be approximately twice that of the normal control before the next dose is administered [b]. The effect on clotting time is increased and prolonged by caronamide [31] and phosphorylated heparidin [32]. The value for maximum clotting time can be estimated from the effect of clotting time in vitro [3]. The international unit of heparin = the weight of 1/130 mg of international standard; commercial heparin = 90-120 units/mg.

Indirect Anticoagulants: These drugs are given orally. Dosage is dependent on the technique used to detect the change in prothrombin time and on individual susceptibility to the drug. A considerable number of individuals in some species are refractory to one or more of these drugs [11]. The effect on prothrombin time, which is cumulative, can be greatly enhanced by administering an initial dose followed by smaller repeated doses. Such administration also avoids toxic side effects. Dicumarol must be dissolved with a small amount of 5N sodium hydroxide for intravenous use. It can be given intraperitoneally in suspension in propylene glycol (10-100 mg/ml). Warfarin and Tromexan are more soluble than Dicumarol.

Values in parentheses are ranges, estimate "b" (cf. Introduction).

Animal (A)	Dose (B)	Route (C)	Maximum Clotting Time or Prothrombin Time		Achievement of Therapeutic Effect (F)	Recovery Time (G)	Remarks (H)	Reference (I)
			Control (D)	Experimental (E)				
1 Man	1000 units (10 mg)	IV		<4.5 min				
2				4.5-7 min			28 hyporeactors.	1
3				>7 min			50 normal reactors.	
4	5000 units	IV	15 min				7 hyperreactors.	
5	150 mg	IV		115 min				
6	100 mg	IV		>80 min				
7	75 mg	IV		>80 min		4 hr	Incoagulable for 1 hr.	2
8	50 mg	IV		80 min		4 hr	Incoagulable for 1/2 hr.	
9	150 mg	IM	12 min	55 min		2 hr		
10	100 mg	IM	12 min	30 min		10 hr		
11	25, 50, 75 mg	IM	12 min	27 min		6 hr		
12 Dog	30 units/kg	IV	12 min	34 min	1 hr	3-4 hr		
13	100 units/kg	IV	12 min			32 min		
14	430 units/kg	IV		>2 hr				
15	2-3 units/kg/min for 2 hr	SC	20 sec	0.25-3.5 hr 5 min		2-13 hr	Prolonged by anesthetics.	3
16	65-290 units/kg	IV	20 sec	5 min		4 hr	Thrombin time used. Heparin appeared in lymph in 10-90 min.	4
17 Rabbit	10.5 mg/kg	IV	20 sec	5 min		200 min	Duration proportional to dose.	5
18	(50 units/kg)					100 min		6
19	2.5 mg/kg			>60 min		2 hr		
20 Rat	1.0 mg/kg	SC		40 min		1 hr		7
21	1950 units/kg	SC		>24 hr		6 hr		
22 Rabbit	5500 units/kg	SC		>24 hr		12 hr	Normal in 12 hr.	8
23	21 mg/kg			Dextran Sulfate				
24	17 mg/kg			>80 min		3 hr		
	7 mg/kg			>80 min		100 hr	Normal in 18 hr.	
				>80 min		50 min		7

25	Rabbit	30 mg/kg 15 mg/kg	65 min >60 min	Thrombocid (Xylan Polysulfuric Acid)	2.5 hr 1.5 hr	7
26						
27	Man	300 mg	30 sec	Picumarol	3-4 da	Varies
28		250 mg	63 sec			
29		500 mg	106 sec		6 da	Quick prothrombin time. Effective only with Vitamin K- deficient diet.
30		750 mg	37 (10 4-45.2) sec			Quick prothrombin time.
31	Dog	5 mg/kg	No effect		24 hr	Quick prothrombin time.
32	Hamster	5 mg/kg			10 da	Susceptibility decreased in puerperum.
33	Mouse	5 mg/kg	10 6(7 6-13 4) sec		5 da	
34	Rabbit	5 mg/kg	8 0(7 6-8 4) sec		4 da	
35		2 5 mg	15-16 sec		1 da	
36		10 mg/kg	65 min		2.5 da	
37		25 mg/kg	15-16 sec		3 da	
38		50 mg/kg	20 min		1 da	
39		100 mg/kg	28 sec		1-2/3 da	
40		0 37 mg	28 sec		2 da	
41		0 75 mg	28 sec		2 5 da	
42		1 5 mg	28 sec		3 da	
43		3 mg	28 sec		19 9(18 7-21 1) sec	
44	Rat	5 mg/kg				
45	Chicken	100 mg/kg				
46		300 mg/kg				
47				Tromexan (3, 3'-Carboxymethylene-bis-4-hydroxycoumarin)	30 hr	56 hr
48	Man	1200 mg	14 sec		30 hr	60 hr
49		1200 mg	14 sec		2 da	5 da
50	Dog	10 mg/kg	15 sec		3 da	5 da
51		40 mg/kg	15 sec		30 hr	48 hr
52	Rabbit	400 mg	12 sec			Minimum effective dose.
53	Chicken	100-500 mg/kg		Warfarin (3-(3-acetylbenzyl)-4-hydroxycoumarin)		5-10 mg/da for maintenance.
54	Man	45-50 mg initial	12 sec		18-24 hr	6 da
55		100 mg	40 sec		48 hr	
56	Rabbit	40 mg/kg	9 6 sec		36 7 sec	
57		5 mg/kg	7 9 sec		35 2 sec	
58	Rat	0 03 mg/kg/da	25-36 sec		No effect	
59		0 05 mg/kg/da	25-36 sec		55 sec	
60		0 10 mg/kg/da	25-36 sec		2 min	
				Liquamar (3-(11-Phenylpropyl)-4-hydroxycoumarin)		
61	Man	18 mg	20 9 sec		92 hr	144 hr
62		21 mg	20 9 sec		96 hr	168 hr
63	Rabbit	2 5 mg	14 sec		48 hr	96 hr
64		4 mg	14 sec		7 2 hr	160 hr
65	Chicken	50 mg/kg				Minimum effective dose.

/1/ Intra-arterial injection of 1 mg heparin/100 ml of blood, plus 1.2 mg protamine/100 ml of blood into venous outflow of organ (limb, kidney), gives satisfactory local heparinisation without general heparinisation in man and dog (24). /2/ To neutralize heparin in man and dog, slowly inject protamine in the amount of 1.2-2.5 x weight of heparin in blood, as determined by in vitro titration; excess of protamine will be anticoagulant (8, 24, 25). /3/ Crude heparin.

102. ANTICOAGULANTS IN VIVO: VERTEBRATES (Concluded)

Animal	Dose	Route	Maximum Clotting Time or Prothrombin Time		Achievement of Therapeutic Effect (F)	Recovery Time (G)	Remarks	Reference
			(D)	(E)				
65 Man	0.5 g				EDC [Ethylene-bis-4-hydroxycoumarin]			(1)
67 Rabbit	20 mg				Great individual variation		0.2 g maintenance dose.	21
68	30 mg	PO	7 sec		10 sec			
69 Dog	50 mg/kg	PO	7 sec		27 sec			21
70 Rabbit	50 mg/kg				Phenindione (2-Phenyl-1,3-indanedione)			
					9-13 sec			
					10-12 sec			
71 Man	20 mg				Dipaxin (2-Diphenylacetyl-1,3-indanedione)			
					25 hr	36 hr		22
					25 hr	40 hr		22
					1-2 da	6-10 da	2-4 mg maintenance dose.	23

Contributors (a) Jaques, Louis B. (b) Wright, Irving S.

- References. [1] De Takata, G. 1943. Surg. Gyn. Obst. 77:31. [2] Jorpes, J. E. 1950. Acta chir. scand. 99:476. [3] Jaques, L. B., and A. G. Ricker. 1948. Blood, N. Y. 3:1197. [4] Jaques, L. B. 1950. N. England J. M. 243:395. [5] Willis, P. W., J. A. MacCris, E. W. Dennis, P. E. Hodgson, W. W. Coon, and I. F. Duff. 1953. J. Laborat. Clin. M. 42:968. [6] Gross, P. 1929. Proc. Soc. Exp. Biol., N. Y. 26:383. [7] Astrup, T., H. H. K. Flyger. and J. Gormsen. 1955. Scand. J. Clin. & Invest., Oslo 7:204. [8] Jaques, L. B., A. F. Charles, and C. H. Best. 1938. Acta med. scand., Suppl. 90:190. [9] Allen, E. V. 1947. J. Am. M. Ass. 134:323. [10] Jansen, K. F. 1944. Dikumamol. Thesis. Copenhagen. [11] Jaques, L. B., E. Froese, R. O'Toole, and J. W. T. Spinks. 1957. Arch. Internat. pharm. dyn., Gand 11:478. [12] Link, K. P. 1943. Harvey Lect., Balt. 39:162. [13] Dam, H., T. Gell, E. Lund, and E. Sondergaard. 1954. Proc. First Internat. Conf. Thrombosis and Embolism. Basel. [14] Burke, G. E., and I. S. Wright. 1951. Circulation, N. Y. 3:164. [15] Gruber, C. M., K. S. Lee, Z. T. Lasziczenko, and C. M. Gruber, Jr. 1951. Arch. internat. pharm. dyn., Par. 87:402. [16] Clatsloff, D. V., and O. O. Meyer. 1956. Arch. Int. M. 97:753. [17] Freeman, D. J., and O. O. Meyer. 1956. Proc. Soc. Exp. Biol., N. Y. 92:52. [18] Hetsey, S. R., J. P. Saunders, and K. C. Olson. 1956. J. Agr. and Food Chem. 4:144. [19] Ashwin, J., and L. B. Jaques. Unpublished. [20] Bourgain, R., M. Todd, L. Herzog, and I. S. Wright. 1954. Circulation, N. Y. 10:680. [21] Fanth, P., and M. H. Nance. 1947. Med. J. Australia 34:133. [22] Jaques, L. B., E. Lepp, and E. Gordon. 1950. Tr. Third Conf. Blood Clot and Allied Probs. Josiah Macy, Jr., Foundation, New York. v. 11. [23] Field, J. B., M. S. Goldfarb, A. B. Ware, and G. C. Griffith. 1952. Proc. Soc. Exp. Biol., N. Y. 81:678. [24] Gordon, L. A., V. Richards, and H. A. Perkins. 1956. N. England J. M. 255:1026. [25] Chagaff, E., and K. B. Olson. 1937. J. Biol. Chem. 122:153. [26] Brown, A., and A. S. Douglas. 1952. Glasgow M. J. 33:225. [27] Jaques, L. B. 1955. Rev. hemat., Par. 10:379. [28] Schwabe, B. 1954. Proc. First Internat. Conf. Thrombosis and Embolism, Basel. [29] Solandt, D. Y., and C. H. Best. 1940. Lancet, Lond. 1:1042. [30] Fisher, A., and T. Astrup. 1939. Proc. Soc. Exp. Biol., N. Y. 42:81. [31] Sirak, H. D., R. S. McCleery, and C. P. Artz. 1948. Surgery 24:811. [32] Evans, J. M., I. Hau, and T. H. Korthals, 1955. Am. Surgeon 21:745.

104. BLOOD COAGULATION TESTS, NORMAL VALUES: MAN

Values in parentheses are ranges, estimate "c" (cf. Introduction).

Test (A)	No. of Observations (B)	Normal Value (C)	Reference (D)
1 Two-tube clotting time	50	11.35(7.45-15.0) min	1
2 Two-tube silicone clotting time	54	22.51(14.3-31.0) min	1
One-stage prothrombin test			
3 Quick	53	11.68(9.7-13.0) sec	2
4 Owren	48	29.06(19.5-35.0) sec	3
5 Ware	45	22.38(19.0-30.0) sec	4
6 Two-stage prothrombin test		356(290-425) units	5
7 Factor V	45	18.5(15.6-20.8) sec	6
Factor VII			
8 Serum	37	21.0(19.4-24.0) sec	7
9 Plasma	40	25.3(22.2-27.4) sec	7
10 Prothrombin consumption	50	96.5(74-100) %	8
11 Thromboplastin generation	50	17.02(9.0-21.2) sec	9
12 Fibrinogen	57	273.5(120-470) mg %	10

Contributor: Coon, William W.

References: [1] Duff, I. F., R. H. Landaburu, J. Conrad, and J. A. Polhemus. Unpublished. [2] Quick, A. J. 1951. Physiology and pathology of hemostasis. Lea and Febiger, Philadelphia. [3] Owren, P. A. 1949. Scand. J. Clin. Lab. Invest. 1:81. [4] Ware, A. G., and R. Stragnell. 1952. Am. J. Clin. Path. 22:791. [5] Ware, A. G., and W. H. Seegers. 1949. Ibid. 19:471. [6] Wolf, P. 1953. J. Clin. Path., Lond. 6:34. [7] Tocantins, L. M. 1955. The coagulation of blood, methods of study. Grune and Stratton, New York. p. 144. [8] Quick, A. J. 1949. Blood, N. Y. 4:1281. [9] Biggs, R., and A. S. Douglas. 1953. J. Clin. Path., Lond. 6:23. [10] Ratnoff, O. D., and C. Menzie. 1951. J. Laborat. Clin. M. 37:316.

105. EFFECT OF HEPARIN AND HEPARINOID COMPOUNDS ADMINISTERED INTRAVENOUSLY: MAN

Hemorrhagic complications are considered as an extension of the therapeutic effect of the agent, rather than as a side effect. Injection of heparin and other heparinoid compounds contributes to a clearing effect on lipemic blood in man, resulting in a fall in serum triglycerides and a rise in serum fatty acids. [3]

Drug (A)	Dose (B)	Effective Duration (C)	Side Effect (D)	Reference (E)
1 Heparin	50 mg	1 hr	Occasional transient alopecia.	1, 2
2	75 mg	2 hr		
3	100 mg	2-3 hr		
4	150 mg	4-5 hr		
5 Paritol ¹	200 mg	2-3 hr	Nausea, vomiting, paresthesia, headache, vasomotor collapse; late development of alopecia. Increased incidence of telangiectases to repeated doses	3, 4
6	300 mg	2-6 hr (highly variable)		
7				2, 5
8				
9				2, 6-8
sulfate			treatment prolonged beyond 3-5 da.	

1/11 major toxic reactions in 125 subjects receiving Paritol.

Contributor: Coon, William W.

References: [1] Duff, I. F., J. W. Linman, and R. Birch. 1951. Surg. Gyn. Obst. 93:343. [2] Hirschboeck, J. S., F. W. Madison, and A. V. Pisciotto. 1954. Am. J. M. Sc. 227:279. [3] Shore, B., A. V. Nichols, and N. K. Freeman. 1953. Proc. Soc. Exp. Biol., N. Y. 83:216. [4] Duff, I. F. Unpublished. [5] Hodgson, P. E., and W. W. Coon. Unpublished. [6] Ricketts, C. R., K. W. Walton, B. D. Van Leuven, A. Birbeck, A. Brown, A. C. Kennedy, and C. C. Burt. 1953. Lancet, Lond. 2:1004. [7] Griffith, D. R., R. H. Landaburu, J. A. Polhemus, P. W. Willis, and I. F. Duff. Unpublished. [8] Jeavons, S. M., K. W. Walton, and C. R. Ricketts. 1956. Brit. M. J. 2:1016.

106. EFFECT OF HEPARIN ADMINISTERED SUBCUTANEOUSLY AND INTRAMUSCULARLY: MAN

Heparin	Dose mg	Route	Achievement of Therapeutic Effect ¹	Time from Injection to Maximal Prolongation of Clotting Time	Time from Injection to Fall in Clotting Time below Therapeutic Level			Side Effects
					Minimal	Usual	Maximal	
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)
1 Aqueous concentration	200	SC	2-4 hr	5-6 hr	4 hr	7-12 hr	16 hr	Rarely complete resistance; local pain, tenderness, ecchymoses.
2 In Pitkin menstruum	400 ²	IM	3-8 hr	19 hr	Frequently complete resistance	24 hr	31 hr	Frequently complete resistance; local pain, tenderness, hematoma
3 In gelatin-dextrose ("Depo-heparin")	400 ³	IM	2-6 hr	8-12 hr	16 hr	24 hr	30 hr	Occasionally complete resistance; local pain, tenderness, hematoma.
4	400 ²	IM	2-9 hr	8-12 hr	16 hr	16 hr	21 hr	

/1/ Twice normal clotting time. /2/ 200 mg with, and 200 mg without, vasoconstrictors. /3/ Without vasoconstrictors.

Contributor: Coon, William W.

Reference. [1] Duff, I. F., J. W. Linman, and R. Birch. 1951. Surg. Gyn. Obst. 93 343.

107. EFFECT OF COAGULANTS AND ANTICOAGULANTS ON CIRCULATORY FUNCTION: VERTEBRATES

Drug (A)	Animal and Dose-- (B)			
1 Fibrin (thrombin and fibrinogen)	Dog, rabbit, intracisternally			
2 Fibrinogen	Cat	pulmonary vein		
3 Fibrinopeptide (from iodinated fibrinogen)	Frog	Removed by isolated heart.		2
		Stimulated isolated heart.		3
4 Thrombin	Dog, intravenous drip	Fall in systemic, rise in pulmonary, blood pressure.	15-30 sec apnea, followed by dyspnea or sporadic gasping.	4
5	Rabbit, 10-100 mg/min, intravenously	Change in QRS, ST, T; random myocardial damage; bradycardia; arterio-venous dissociation.	Death in convulsions.	5
6 Thromboplastin (present in crude tissue extracts)	Dog, 0.2-0.3 g/kg	Shock. Clot in right heart and portal system.		6
7	Dog, 0.4-0.5 g/kg	Clot in right heart and portal system. Instant death.		
8	Dog, unanesthetized	Rise in blood pressure.		
9	Dog, anesthetized	Marked, sustained fall in blood pressure.	Abolished by heparin, cord section.	7
10	Frog, brain trauma	Shock	Not abolished by cord section.	
11 Lung extract	Dog	Fall in systemic, marked rise in pulmonary, blood pressure. Failure of peripheral pulse. Extreme cardiac dilatation.	15-30 sec apnea, followed by dyspnea or sporadic gasping. Death prevented by artificial respiration.	8

Values in parentheses are ranges, estimate "c" (cf. Introduction).

Test (A)	No. of Observations (B)	Normal Value (C)	Reference (D)
1 Two-tube clotting time	50	11.35(7.45-15.0) min	1
2 Two-tube silicone clotting time	54	22.51(14.3-31.0) min	1
One-stage prothrombin test			
3 Quick	53	11.68(9.7-13.0) sec	2
4 Owren	48	29.06(19.5-35.0) sec	3
5 Ware	45	22.38(19.0-30.0) sec	4
6 Two-stage prothrombin test		356(290-425) units	5
7 Factor V	45	18.5(15.6-20.8) sec	6
Factor VII			
8 Serum	37	21.0(19.4-24.0) sec	7
9 Plasma	40	25.3(22.2-27.4) sec	7
10 Prothrombin consumption	50	96.5(74-100) %	8
11 Thromboplastin generation	50	17.02(9.0-21.2) sec	9
12 Fibrinogen	57	273.5(120-470) mg %	10

Contributor: Coon, William W.

References: [1] Duff, I. F., R. H. Landaburu, J. Conrad, and J. A. Polhemus. Unpublished. [2] Quick, A. J. 1951. Physiology and pathology of hemostasis. Lea and Febiger, Philadelphia. [3] Owren, P. A. 1949. Scand. J. Clin. Lab. Invest. 1:81. [4] Ware, A. G., and R. Stragrell. 1952. Am. J. Clin. Path. 22:791. [5] Ware, A. G., and W. H. Seegers. 1949. Ibid. 19:471. [6] Wolf, P. 1953. J. Clin. Path., Lond. 6:34. [7] Tocantins, L. M. 1955. The coagulation of blood, methods of study. Grune and Stratton, New York. p. 144. [8] Quick, A. J. 1949. Blood, N. Y. 4:1281. [9] Biggs, R., and A. S. Douglas. 1953. J. Clin. Path., Lond. 6:23. [10] Ratnoff, O. D., and C. Menzie. 1951. J. Laborat. Clin. M. 37:316.

105. EFFECT OF HEPARIN AND HEPARINOID COMPOUNDS ADMINISTERED INTRAVENOUSLY: MAN

Hemorrhagic complications are considered as an extension of the therapeutic effect of the agent, rather than as a side effect. Injection of heparin and other heparinoid compounds contributes to a clearing effect on lipemic blood in man, resulting in a fall in serum triglycerides and a rise in serum fatty acids. [3]

Drug (A)	Dose (B)	Effective Duration (C)	Side Effect (D)	Reference (E)
1 Heparin	50 mg	1 hr	Occasional transient alopecia.	1,2
2	75 mg	2 hr		
3	100 mg	2-3 hr		
4	150 mg	4-5 hr		
5 Paritol ¹	200 mg	2-3 hr	Nausea, vomiting, paresthesia, headache, vasomotor collapse, late development of alopecia. Increased individual tolerance to repeated doses.	3,4
6	300 mg	2-6 hr (highly variable)		
7 Treburon	200 mg	2 hr	Prolonged administration resulted in high incidence of diarrhea and transient alopecia.	2,5
8	300 mg	2-3 hr		
9 Dextran sulfate	5000 units	1-3 hr	Transient alopecia. Marked cumulative effect when treatment prolonged beyond 3-5 da.	2,6-8

/1/ 11 major toxic reactions in 125 subjects receiving Paritol.

Contributor: Coon, William W.

References: [1] Duff, I. F., J. W. Linman, and R. Birch. 1951. Surg. Gyn. Obst. 93:343. [2] Hirschboeck, J. S., F. W. Madison, and A. V. Pisciotto. 1954. Am. J. M. Sc. 227:279. [3] Shore, B., A. V. Nichols, and N. K. Freeman. 1953. Proc. Soc. Exp. Biol., N. Y. 83:216. [4] Duff, I. F. Unpublished. [5] Hodgson, P. E., and W. W. Coon. Unpublished. [6] Ricketts, C. R., K. W. Walton, B. D. Van Leuven, A. Birbeck, A. Brown, A. C. Kennedy, and C. C. Burt. 1953. Lancet, Lond. 2:1004. [7] Griffith, D. R., R. H. Landaburu, J. A. Polhemus, P. W. Willis, and I. F. Duff. Unpublished. [8] Jeavons, S. M., K. W. Walton, and C. R. Ricketts. 1956. Brit. M. J. 2:1016.

Heparin	Dose mg	Route	Achievement of Therapeutic Effect ¹	Time from Injection to Maximal Prolongation of Clotting Time	Time from Injection to Fall in Clotting Time below Therapeutic Level			Side Effects
					Minimal	Usual	Maximal	
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)
1 Aqueous concentration	200	SC	2-4 hr	5-6 hr	4 hr	7-12 hr	16 hr	Rarely complete resistance; local pain, tenderness, ecchymoses.
2 In Pitkin menstruum	400 ²	IM	3-8 hr	19 hr	Frequently complete resistance	24 hr	31 hr	Frequently complete resistance; local pain, tenderness, hematoma.
3 In gelatin-dextrose ("Depo-heparin")	400 ³	IM	2-6 hr	8-12 hr	16 hr	24 hr	30 hr	Occasionally complete resistance; local pain, tenderness, hematoma.
4	400 ²	IM	2-9 hr	8-12 hr	16 hr	16 hr	21 hr	

/1/ Twice normal clotting time. /2/ 200 mg with, and 200 mg without, vasoconstrictors /3/ Without vasoconstrictors.

Contributor: Coon, William W

Reference [1] Duff, I. F., J. W. Linman, and R. Birch. 1951. Surg. Gyn. Obst. 93:343.

107. EFFECT OF COAGULANTS AND ANTICOAGULANTS ON CIRCULATORY FUNCTION, VERTEBRATES

Drug	Animal and Dosage	Effect on Circulation	Remarks	Reference
(A)	(B)	(C)	(D)	(E)
1 Fibrin (thrombin and fibrinogen)	Dog, rabbit, intracisternally	Increased systemic and pulmonary blood pressure. Dilatation of left atrium and pulmonary vein.	Not blocked by vagotomy, sympathectomy; ganglionic blockade effective.	1
2 Fibrinogen	Cat	Removed by isolated heart.		2
3 Fibrinopeptide (from iodinated fibrinogen)	Frog	Stimulated isolated heart.		3
4 Thrombin	Dog, intravenous drip	Fall in systemic, rise in pulmonary, blood pressure.	15-30 sec apnea, followed by dyspnea or sporadic gasping.	4
5	Rabbit, 10-100 mg/min, intravenously	Change in QRS, ST, T, random myocardial damage; bradycardia; arterio-venous dissociation.	Death in convulsions	5
6 Thromboplastin (present in crude tissue extracts)	Dog, 0.2-0.3 g/kg	Shock. Clot in right heart and portal system.		6
7	Dog, 0.4-0.5 g/kg	Clot in right heart and portal system. Instant death		
8	Dog, unanesthetized	Rise in blood pressure.		
9	Dog, anesthetized	Marked, sustained fall in blood pressure.	Abolished by heparin, cord section.	7
10	Frog, brain trauma	Shock	Not abolished by cord section.	
11 Lung extract	Dog	Fall in systemic, marked rise in pulmonary, blood pressure. Failure of peripheral pulse. Extreme cardiac dilatation.	Not abolished by cord section. 15-30 sec apnea, followed by dyspnea or sporadic gasping. Death prevented by artificial respiration.	8

107. EFFECT OF COAGULANTS AND ANTICOAGULANTS
ON CIRCULATORY FUNCTION: VERTEBRATES (Continued)

	Drug (A)	Animal and Dosage (B)	Effect on Circulation (C)	Remarks (D)	Reference (E)
12	Thromboplastin (present in crude tissue extracts) (concluded) Kidney extract	Dog, bilaterally nephrectomized	Hemolytic necrosis of cere- bral vascular walls.		9
13	Cephalin	Dog	No effect on blood pressure.		10
14		Frog	Antagonized calcium in isolated heart.		11
15	Kaolin	Dog			
16	Agar	Dog			
17	Russel viper venom (MgATP)	Cat, rabbit, rat	Many vessels.	vagotomy. Tachypnea.	
18	Russel viper venom, asp venom, eel serum, cobra venom	Rabbit	Prevented fall in blood pres- sure.		13, 14
19	Plasmin (fibrino- lysin)	Dog	No effect on hypertensin, oxytocin; destroyed vaso- pressin.		15
20		Dog, guinea pig	Decreased blood pressure.	Due to another plasma protein in fibrinolysis preparations.	16
21		Turtle	Decreased output of isolated heart.		
22	Trypsin	Dog, 13,000 units/kg	Transient drop to 40 mm Hg in blood pressure. Caused intravascular clotting.		17
23		Dog, 25,000 units/kg	Caused intravascular clotting. Death in 15-45 min.		
24		Rabbit	Clotting and symptoms same as for thrombin and thromboplastin (see above).		18
25		Rabbit, 0.2-1.0 mg/min	Change in P, QRS, ST, T; arterio-venous dissociation, random myocardial damage.		5
26	Vitamin K, dietary deficiency of	Mouse, rat, on sulfaguanidine	Myocardial lesions. Hemor- rhage necrosis, leucocytic infiltration.		19
27	Menadiolone (K ₃)	Dog, 2-10 mg/kg	Decrease in blood pressure. No effect on blood vessels.	Arrested isolated heart, 1:10 ⁵ . Negative ino- tropic action, 1:10 ⁶ to 1:10 ⁷ . No effect on action of adrenalin, ephedrine, amphet- amine, acetylcholine.	20, 21
28		Dog, 50 mg/kg	Hypotension, then hyperten- sion and death. No effect on vessels.		
29	K ₃ diphosphate sodium	Dog, 2 mg/kg	No effect on blood pressure or vessels.		
30	K ₃ disuccinate	Dog, 2 mg/kg	No effect on blood pressure or vessels.		
31	K ₃ diacetate	Dog, 2 mg/kg	Lowered blood pressure. No effect on vessels.		
32	K ₃ bisulfite sodium	Dog, 2 mg/kg	Lowered blood pressure. No effect on vessels.		
33	Platelets, purified bovine	Dog	Marked increase in pulmonic and systemic blood pres- sure.		22
34	Antiplatelet serum	Dog	Hypotension, shock.	Histamine liberated.	23
35	Heparin (commer- cial solutions containing tri- cresol)	Man	No effect on coronary dilator. ECG changes in standard exercise test.		24
36		Cat, dog	No effect on retrograde blood flow in vessels.		25
37		Dog	No effect on blood pressure.	No respiratory effect.	13, 26
38		Dog, 10 mg/kg	Sodium salt increased coro- nary flow, barium salt had no effect on flow.		27
39		Hamster	Increased blood flow in venules.		28

107. EFFECT OF COAGULANTS AND ANTICOAGULANTS
ON CIRCULATORY FUNCTION- VERTEBRATES (Continued)

	Drug (A)	Animal and Dosage (B)	Effect on Circulation (C)	Remarks (D)	Reference (E)
40	Heparin (commercial solutions containing tri-cresol) (concluded)	Rabbit, 1:10,000	Decreased coronary flow of isolated heart.		29
41		Rabbit, <1-10,000	No effect on heart.		30
42		Frog	Initial inhibition of perfused isolated heart	Due to dialyzable materials in commercial heparin.	31
43	Heparin and other mucopolysaccharides	Rat	Arrest of vasomotion.	Reversed action of adrenalin.	32
44	Heparin	Man	No effect on claudication time.		33
45		Man	No effect on heart effort in angina.		34, 35
46		Man, hypertensive	Fall in blood pressure to normal levels.	Blood pressure fall accompanied by urinary excretion of histamine.	36
47		Man, atherosclerotic	Increased forearm arterio-venous O ₂ difference in 60% of subjects.		37
48		Dog	Heparin fixed to cement substance in blood vessels.		38
49		Dog, ischemic compression shock	No effect on blood pressure.		39
50		Dog, adrenal chloroform syncope	Prevented intracardiac thrombosis without preventing syncope		40-42
51		Rat, hypertensive, with DOCA, 100 units/kg, subcutaneously	Fall in blood pressure to normal levels.	Duration of effect shortened by pretreatment with 48/80, or water intraperitoneally to disrupt mast cells.	
52		Rat, hypertensive, by renal encapsulation	Fall in blood pressure to normal levels.	Effect prevented by anti-histamines.	43
53	Heparin, plus ergotamine tartrate	Rat	No effect on gangrene.		44
54	Xylan sulfuric acid ester	Rabbit, 2.5-50 mg/kg	Fall in blood pressure.	Respiratory effects secondary to blood pressure changes.	26, 45
55		Rabbit, 80 mg/kg	Death.		46
56	Cellulose sulfuric acid ester	Rabbit, 3 mg/kg	Death.		26, 45
57	Liquid (Sodium polyanethol sulfuric acid)	Cat	Transitory fall in blood pressure with increase in venous pressure, often failure to return to initial blood pressure.		26, 45
58		Rabbit	Intracapillary precipitates		26, 45
59	Novirudin	Cat	Decrease in systolic blood pressure. Decrease in minute volume.		26
60		Rabbit	Hypertension for 2-6 min.		26
61	Chlorazol fast pink	Cat, dog	No effect on blood pressure		26
62		Guinea Pig, rabbit	Marked fall in blood pressure		26
63	Protamine Salmine	Cat, 5 mg/kg	Transitory fall in blood pressure.		46
64		Dog, 5 mg/kg	Marked fall in blood pressure when injected rapidly. Vasodilator in muscles	Blood pressure effect blocked by removal of platelets and modified by cord section.	46
65		Guinea pig, 5 mg/kg	Slow fall in blood pressure, death.		47
66		Rabbit, 5 mg/kg	Slight rise in blood pressure.		46
67	Salmine, clupeine	Rat, 20-30 mg	Collapse		48

107. EFFECT OF COAGULANTS AND ANTICOAGULANTS
ON CIRCULATORY FUNCTION: VERTEBRATES (Continued)

	Drug (A)	Animal and Dosage (B)	Effect on Circulation (C)	Remarks (D)	Reference (E)
68	Protamines (con- cluded) Clupeine	Dog, 50-100 mg/kg	Marked fall in blood pressure when injected rapidly. Vasodilator in muscles.		47, 49
69		Guinea pig, rat, 100 mg/kg	Lethal occlusion of blood vessels.		50
70		Rabbit, 15 mg/kg	50% reduction in vascular flow in isolated ear.		51
71		Rabbit, 20 mg/kg	Reduced coronary flow of iso- lated heart 50%. Complete contraction of medial artery of ear.		
72		Frog, 0.5-5.0 µg	Increased action of acetyl- choline on heart. Increased capillary permeability.	Displaced platelets and serum proteins from capillary endothelium.	52, 53
73		Frog, 20 µg	Strengthened isolated hypo- dynamic heart. Increased capillary permeability.		
74		Frog, 50 µg	Increased capillary permea- bility. Stopped heart.		
75	Toluidine blue, azure A	Dog, rabbit, 1-10 mg/kg	Slight increase in blood pres- sure. Bradycardia. Increase in T wave and high take off of P. No effect on heart response to acetyl- choline or epinephrine. Vasoconstriction of blood vessels in hind leg of dog.	Higher dose required for cardiac effects after atropine. Increased respiratory rate and depth of breathing.	54
76		Rabbit, isolated heart	No effect on transmission, cytotoxic effect.		
77		Frog, isolated heart	Cholinergic blockade.		
78	Dicumarol	Man	No effect on normal or abnor- mal ECG.		56
79			No effect on ECG with standard exercise tests.		24
80		Man, arterio- sclerotic	Slight reduction in extremity blood flow (plethysmograph).		57
81		Dog	Increased coronary flow.		45
82		Rabbit, 5-20 mg/kg	Momentary slight rise in blood pressure. 0.25% caused inhibition of isolated heart. Perfusion of peripheral vein, contraction, large doses caused dilatation.	No effect on respiration.	58, 59
83		Rabbit, 10 mg/kg	Pronounced engorgement of great veins.		60
84	Dicumarol, plus ergotamine	Rat	No effect on gangrene.		43
85	Dicumarol (also 4 hydroxycoumarin and indanedione)	Cat, rabbit, 10-50 mg/kg	Usually decreased, sometimes increased, blood pressure.	Rapid rise in O ₂ con- sumption, respiratory rate and amplitude. Effects on O ₂ con- sumption: increased by vitamin K ₃ , not by K ₁ .	61
86	2-Isovaleryl-1,3- indanedione	Cat, 5-60 mg/kg	Decline in blood pressure. Rapid vasodilation (direct).		62

Contributor: Jaques, Louis B.

References [1] Sarnoff, S. J., and C. Sarnoff. 1952. Circulation, N. Y. 6:51. [2] Gorham, L. W., and A. W. Morrison. 1909. Am. J. Physiol. 23:419. [3] Laki, K. 1951. Science 114 435. [4] Schneider, C. L., and

107. EFFECT OF COAGULANTS AND ANTICOAGULANTS
ON CIRCULATORY FUNCTION VERTEBRATES (Concluded)

- R. M. Engstrom. 1954. *Am. J. Obst.* 68 691. [5] Leiberman, J. S., A. Taylor, and I. S. Wright. 1954. *Circulation*, N. Y. 10:338. [6] Bertrand, L., M. T. Gayet-Hallion, and D. Quivy. 1943. *C. rend. Acad. sc.* 216:316. [7] Mylon, E., M. Winternitz, R. Katzenstein, and G. De Suto-Nagy. 1942. *Am. J. Physiol.* 137:260. [8] Moreau, L. 1946. *Ibid* 155 92. [9] Winternitz, M. C., R. Katzenstein, E. Mylon, J. P. Murphy, and H. M. Zimmerman. 1941. *Yale J. Biol.* 13:579. [10] Croxatto, H., and L. de Lerma. 1947. *Bol. Soc. biol. Santiago* 4 57. [11] Scheiner, H. 1934. *C. rend. Soc. biol.* 115 232. [12] Heard, E. E. 1952. *J. Path. Bact., Lond.* 64:13. [13] Doxiades, L., and H. J. Fuchs. 1933. *Zachr. ges. exp. Med.* 88:793. [14] Grassetti, E., and D. E. Schwartz. 1955. *Bull. Schweiz. Akad. med. Wiss.* 11:10. [15] Croxatto, H., W. Badia, and R. Croxatto. 1948. *Proc. Soc. Exp. Biol., N. Y.* 69:422. [16] Guest, M. M., R. C. Murphy, S. R. Bodnar, A. C. Ware, and W. H. Seegers. 1947. *Am. J. Physiol.* 150:471. [17] Roach, H. D., and H. Laufman. 1954. *Surgery* 35 45. [18] Polcak, J., B. Schreiber, D. Sevelova, L. Votava, and J. Runstukova. 1955. *Vnitřn. Lék.* 1:809. [19] Dessau, F. L., L. Lipchuck, and S. Klein. 1956. *Proc. Soc. Exp. Biol., N. Y.* 87:522. [20] Cannava, A. 1948. *Boll. Soc. Ital. biol. sper.* 24:587. [21] Cannava, A. 1947. *Ibid* 23 68. [22] Schneider, C. L., E. B. Claxton, C. H. Hughes, and S. A. Johnson. 1954. *Am. J. Physiol.* 179 236. [23] Crus, W. O., A. Baumgarten, and A. C. Oliveira. 1954. *Ibid* 177:515. [24] Russek, H. I., B. L. Zohman, and V. J. Dorset. 1955. *Am. J. M. Sc.* 229:46. [25] Leiter, E. 1942. *Proc. Soc. Exp. Biol., N. Y.* 51:187. [26] Kahlson, G., and E. Landby. 1937. *Skand. Arch. Physiol., Berl.* 77:301. [27] Gilbert, H. C., and L. Nalefaki. 1949. *J. Laborat. Clin. M.* 34:797. [28] Sollivän, B. J., and W. E. Masterson. 1953. *Am. J. Physiol.* 175:56. [29] Halonen, P. I., O. Vartiainen, and E. V. Venho. 1951. *Ann. med. exp. biol. fenn.* 29:23. [30] Chast, B. 1955. *Physiol. zool.* 28 315. [31] Bonfils, S., J. P. Hardouin, R. Derache, and A. Lambling. 1956. *C. rend. Soc. biol.* 150:884. [32] Simon, E. P., and I. S. Wright. 1953. *J. Am. M. Ass.* 153:98. [33] Rinsler, S. H., J. Travell, H. Bakst, Z. H. Benjamin, R. L. Rosenthal, S. Rosenfeld, and B. B. Hirsch. 1953. *Am. J. Med.* 14 438. [34] Keller, R. 1954. *Praxis, Bern* 43 113. [35] Sartorius, H., A. Moench, and H. Sarre. 1954. *Verh. Deut. Ges. inn. Med.* 60 565. [36] Engelberg, H., and R. Kuhn. 1956. *Angiology* 7:73. [37] Samuels, P. B., and D. R. Webster. 1952. *Am. Surg.* 136:422. [38] Boss, J. R., and H. D. Green. 1947. *Am. J. Physiol.* 150:697. [39] Quivy, D. 1946. *C. rend. Soc. biol.* 142:1223. [40] Waser, P., and R. Keller. 1954. *Helvet. physiol. pharm. acta* 12:94. [41] Keller, R. 1956. *Arch. internat. pharm. dyn.*, Gand 107:382. [42] Hardegg, W., W. Maass, C. Mayer, H. Abel, and K. Huhnstock. 1955. *Klin. Wschr.* 33 812. [43] Anderson, L. H., and J. A. Wells. 1948. *Proc. Soc. Exp. Biol., N. Y.* 67:53. [44] Wiedersheim, M., W. Hertlein, E. Husemann, and R. Lotterle. 1953. *Arch. exp. Path., Lpz.* 217:107. [45] Kauffman, R., and P. M. Dreyfus. 1953. *Arch. Path., Chic.* 56:597. [46] Jaques, L. B. 1949. *Brit. J. Pharm.* 4 135. [47] De Waele, H. 1944. *Arch. internat. pharm. dyn.*, Gand 69 450. [48] Ahlstrom, L., and H. von Euler. 1944. *Acta. Chem. Min. Geol.* A23 1. [49] Thompson, W. H. 1900. *Zachr. physiol. Chem.* 29 1. [50] Shelley, W. B., and R. Tarall. 1943. *Proc. Soc. Exp. Biol., N. Y.* 52:215. [51] Solis, J. T., and H. E. Essex. 1951. *Am. J. Physiol.* 167:1. [52] Danielli, J. F. 1940. *J. Physiol., Lond.* 98 109. [53] Heim, F. 1943. *Arch. exp. Path., Lpz.* 202 228. [54] Haley, T. J., and J. L. Leitch. 1953. *Arch. internat. pharm. dyn.*, Gand 93 341. [55] Leitch, J. L., and T. J. Haley. 1952. *J. Pharm. Exp. Ther.* 104 416. [56] Balkin, S. S., and A. Gootnick. 1948. *J. Laborat. Clin. M.* 33 972. [57] Wakim, K. G., M. S. Klackner, Jr., N. W. Barker, and W. F. Kvale. 1950. *Fed. Proc., Balt.* 9 131. [58] Kubo, H. 1954. *Kumamoto M. J.* 7:73. [59] Lowenthal, J. 1956. *Bull. soc. chim. Belg.* 65 124. [60] Dale, D. U., and L. B. Jaques. 1942. *Canad. M. Ass. J.* 46 546. [61] Soderberg, U., and C. A. Wachtmelster. 1956. *J. Pharm. Exp. Ther.* 117:298. [62] Heisley, S. R., J. P. Saunders, and K. C. Olson. 1956. *Proc. Soc. Exp. Biol., N. Y.* 91:86.

**107. EFFECT OF COAGULANTS AND ANTICOAGULANTS
ON CIRCULATORY FUNCTION: VERTEBRATES (Continued)**

	Drug (A)	Animal and Dosage (B)	Effect on Circulation (C)	Remarks (D)	Reference (E)
68	Protamines (concluded) Clupeine	Dog, 50-100 mg/kg	Marked fall in blood pressure when injected rapidly. Vasodilator in muscles.		47, 49
69		Guinea pig, rat, 100 mg/kg	Lethal occlusion of blood vessels.		50
70		Rabbit, 15 mg/kg	50% reduction in vascular flow in isolated ear.		51
71		Rabbit, 20 mg/kg	Reduced coronary flow of isolated heart 50%. Complete constriction of medial artery of ear.		
72		Frog, 0.5-5.0 µg	Increased action of acetylcholine on heart. Increased capillary permeability.	Displaced platelets and serum proteins from capillary endothelium.	52, 53
73		Frog, 20 µg	Strengthened isolated hypodynamic heart. Increased capillary permeability.		
74		Frog, 50 µg	Increased capillary permeability. Stopped heart.		
75	Toluidine blue, azure A	Dog, rabbit, 1-10 mg/kg	Slight increase in blood pressure. Bradycardia. Increase in T wave and high take off of P. No effect on heart response to acetylcholine or epinephrine. Vasoconstriction of blood vessels in hind leg of dog.	Higher dose required for cardiac effects after atropine. Increased respiratory rate and depth of breathing.	54
76		Rabbit, isolated heart	No effect on transmission, cytotoxic effect.		
77		Frog, isolated heart	Cholinergic blockade.		55
78	Dicumarol	Man	No effect on normal or abnormal ECG.		56
79			No effect on ECG with standard exercise tests.		24
80		Man, arterio-sclerotic	Slight reduction in extremity blood flow (plethysmograph).		57
81		Dog	Increased coronary flow.		45
82		Rabbit, 5-20 mg/kg	Momentary slight rise in blood pressure. 0.25% caused inhibition of isolated heart. Perfusion of peripheral vein, contraction, large doses caused dilatation.	No effect on respiration.	58, 59
83		Rabbit, 10 mg/kg	Pronounced engorgement of great veins.		60
84	Dicumarol, plus ergotamine	Hat	No effect on gangrene.		43
85	Dicumarol (also 4 hydroxycoumarin and indanediones)	Cat, rabbit, 10-50 mg/kg	Usually decreased, sometimes increased, blood pressure.	Rapid rise in O ₂ consumption, respiratory rate and amplitude. Effects on O ₂ consumption; increased by vitamin K ₃ , not by K ₁ .	61
86	2-Isosaleryl-1, 3-Indanedione	Cat, 5-60 mg/kg	Decline in blood pressure. Rapid vasodilation (direct).	Increase in depth of breathing.	62

Contributor. Jaques, Louis B.

References. [1] Sarnoff, S. J., and C. Sarnoff. 1952. Circulation, N. Y. 6:51. [2] Gorham, L. W., and A. W. Morrison 1909. Am. J. Physiol. 25:419. [3] Laki, K. 1951. Science 114:435. [4] Schneider, C. L., and

109. DRUGS PRIMARILY USED FOR DISORDERS OF THE CIRCULATORY SYSTEM

Drugs listed are characterized by prominent hemodynamic actions; grouping is based on a common action or common source. Official generic names are from United States Pharmacopeia XV, National Formulary X, or from New and Nonofficial Remedies, 1955. Clinical Use: AA = antiarrhythmic drugs, for control of cardiac arrhythmia, prophylactic and curative; AH = antihypertensive drugs, for acute and chronic relief of essential and other forms of hypertension; CT = cardiotonic drugs, for management of congestive heart failure; HT = hypertensive drugs, for pressor action in correction of hypotensive states, and for local vasoconstrictor action in nasal congestion; VD = vasodilators, for improving blood flow in diseased limb and coronary vascular beds, and for relief of systemic hypertension. Route: IH = inhalation, IM = intramuscular; IV = intravenous, PO = oral, SC = subcutaneous, SL = sublingual; TOP = topical application.

pressor activity				vasodilators, for improving blood flow in		IV = intravenous		Effect		Hemodynamic Effects Found on Following Pages (G)
Drug (Synonym)		Clinical Use (B)	Route (C)	Average Adult Dose (D)	Desirable (E)	Undesirable (F)				
(A)										
Veratrum Alkaloids										
1	Alkavervir (Veriloid)	AH	PO	9-15 mg daily	Hypotension without tachycardia.	Severe bradycardia, nausea, vomiting, salivation, and sweating.	258			
2	Protoveratrine A and B ¹ (Veralba, Provell maleate)	AH	PO	0.5-1.0 mg daily			261, 268			
3	Veratrone	AH	PO	0.25-0.5 mg daily			253, 261, 264, 265, 268			
4	Vergitryl ²	AH	PO	3-15 units daily						
5	Vertavis ³	AH	PO	3-15 units daily						
Rauwolfia Alkaloids										
6	Alseroxylon (Rauwolfoid, Rautensin, Rau-Tab, Rauwistan)	AH	PO	2-4 mg daily	Hypotension without tachycardia.	Mental depression, nasal congestion, nausea, bradycardia, weakness and fatigue.				
7	Raudixin, Rauwal	AH	PO	50-500 mg daily						
8	Rescinnamine	AH	PO	2-4 mg daily			253, 258, 261, 271			
9	Reserpine (Serpassil, Serpiloid, Reserpidol, Raused serpate)	AH	PO	0.1-2 mg daily						
Miscellaneous Antihypertensives										
10	Hydralazine (Aprevaline)	AH	PO	100-400 mg daily	Hypotension	Palpitation, headache, lupus-like syndrome.	253, 257, 261, 265, 267			
11	Potassium or sodium thiocyanate	AH	PO	0.05-1.4 g daily		Drowsiness, vomiting.	253			
Ganglion Blocking Drugs										
12	Azamethonium dibromide (Pendiomide)	VD	IV	50-100 mg	Increased blood flow in limbs, lowering of arterial pressure if hypertension present.	Severe hypotension (unless intentional for hypertensive surgery), generalized ganglion blockade manifested as blurring of vision, dryness of mouth, tachycardia, constipation.	252, 258, 261, 266			
13	Chlorisondamine dimethochloride (Ecolid)	VD	PO	50-100 mg daily			261, 266			
14	Hexamethonium bromide or chloride (Bistrium, Methium, Esomid, Hexametol, Hiohex)	VD	PO	200-400 mg			252, 257, 258, 261, 263, 266			
15	Mecamylamine HCl (Inversine)	VD	PO	5-100 mg			261			
16	Pentolinium tartrate (Ansolyse, Penta-pyrrolidinium)	VD	PO	20-100 mg						
17	Pentamethonium chloride	VD	IV	100-150 mg			253, 267			
18	Tetraethylammonium bromide or chloride (Etamon)	VD	IV, IM	200-500 mg			253, 263, 267, 271			
19	Trimethaphan camphor-sulfate (Arfonad)	VD	IV	1 mg/min			253, 261			

1/1 Base, not the salt. 2/2 Standardized for hypotensive potency by effect on carotid sinus reflex in vagotomized dogs. 3/3 Standardized by its cardiac arrest in a crustacean, *Daphnia magna*.

Representative drugs are listed in tables 109 and 110 on the pages indicated in columns D and F.

Site of Action		Stimulant	Page	Depressant	Page	
General	Specific					
(A)	(B)					
Nervous Mechanisms Controlling Circulation						
1	Central nervous system (direct action)	Vasoconstrictor center	Analeptics. ¹	251	General anesthetics. ¹	250
2		Cardio-accelerator center				
3		Cardio-inhibitory center ²	General anesthetics ¹	250	Analeptics ¹	251
4		Other areas that modify vasomotor activity, so far unidentified	None recognized		Rauwolfia alkaloids. Hydralazine. Morphine.	247 247 250
5	Sensory receptors in heart, lungs, blood vessels	Stretch receptors in carotid sinuses and aortic arch ³	Local application of sympathomimetic amines.	248, 249	Local anesthetics. Receptors indirectly inactivated by hypotensive drugs.	250 248
			Veratrum alkaloids (esters of tertiary amines).	247		
			Activated indirectly by hypertensive drugs.	248, 249		
6		Chemoreceptors in carotid and aortic bodies ⁴	Ganglion stimulants. Inhibitors of oxidation, such as cyanide and sulfide. Papaverine. Nikethamide. Aminophylline.	252 252 248 251 248	Local application of ganglion blocking drugs depresses action of ganglion stimulants. O ₂ inhalation depresses action of cyanide.	247
7	Autonomic nervous system	Stretch receptors in heart, lung parenchyma, and blood vessels ⁵	Veratrum alkaloids (esters of tertiary amines).	247	Local anesthetics.	250
8		Receptors of various types in respiratory passages and joints ⁶	Veratrum alkaloids (esters of tertiary amines). Low concentration of volatile anesthetics.	247 250	High concentration of volatile anesthetics.	250
9		Ganglia, both sympathetic ⁷ and parasympathetic ⁸	Ganglion stimulants.	252	Ganglion blocking drugs. High concentration of nicotine.	247 252
10		Sympathetic ⁷	Sympathomimetics.	248-9	Adrenergic blocking drugs.	248
11		Parasympathetic ⁸	Parasympathomimetics.	248	Parasympathetic blocking drugs.	251
Muscular Components of Circulatory System						
12	Blood vessels	With specific antagonist	Methacholine. Histamine.	248 252	Atropine. Antihistamines.	251 251
13		Without specific antagonist	Musculotropic vasoconstrictors.	251, 252	Musculotropic vasodilators.	248
14	Heart	Contractility of heart muscle	Cardiac glycosides.	249, 250	Quinidine. General anesthetics. ¹	250 250
15		Ectopic automaticity of heart	Proarrhythmic drugs Chloroform. Cyclopropane.	250 250 250	Antifibrillatory drugs Quinidine.	250 250

/1/ None selective. /2/ Reciprocal innervation with cardio-accelerator center. /3/ Type 1 receptors, producing perfect inhibition of vasoconstrictor, cardio-accelerator and respiratory centers. /4/ Type 3 receptors, producing pure stimulation of vasoconstrictor, cardio-accelerator and respiratory centers. /5/ Type 2 receptors, producing imperfect inhibition of vasoconstrictor, cardio-accelerator and respiratory centers. /6/ Type 4 receptors, producing imperfect stimulation of vasoconstrictor, cardio-accelerator and respiratory centers. /7/ Sympathetic stimulation causes cardiac acceleration and constriction of most blood vessels. /8/ Parasympathetic stimulation causes cardiac slowing and dilatation of some blood vessels.

Contributors (a) Aviado, Domingo M., Jr., and Carl F. Schmidt, (b) Kray, Otto

Reference: Aviado, D. M., Jr., and C. F. Schmidt. 1955. *Physiol. Rev.* 35:247.

109. DRUGS PRIMARILY USED FOR DISORDERS OF THE CIRCULATORY SYSTEM (Continued)

109. DRUGS PRIMARILY USED FOR DISORDERS OF THE CIRCULATORY SYSTEM (continued)				Effect		Hemodynamic Effects Found on Following Pages: (G)
Drug (Synonym)	Clinical Use	Route	Average Adult Dose	Desirable (E)	Undesirable (F)	
(A)	(B)	(C)	(D)	Sympathomimetic Amines for Systemic Pressor Action (concluded)		(G)
49. Epinephrine, epinephrine bitartrate (Adrenalin, Suprarenin)	HT	SC IV	0.5 mg 5 µg/min	Rise in systemic arterial pressure.	Some amines induce cardiac arrhythmias and cerebral excitation.	254,257,259,262, 264,267,270
50. Hydroxylamphetamine hydrobromide (Paredrine)	HT	IM TOP	10 mg 0.25-1%			254,264
51. Levarterenol bitartrate (Levophed, Noradrenalin)	HT	IV	5 µg/min			254,257,259,262, 264,267,270
52. Mephentermine sulfate (Wyamine)	HT	IM, IV TOP	20 mg 0.5%			254,264,270
53. Methamphetamine HCl (Methedrine, Desoxyephedrine, Desoxin, Dexoval, Doxyfed, Efraxine, Norodin)	HT	PO SC, IM, IV TOP	5 mg 10-30 mg 0.5-1%			270
54. Methoxamine HCl (Vasoxyl)	HT	IM	5 mg			254,257,264,271
55. Orcethyl HCl	HT	IV IM	10-25 mg 75-100 mg			254,264,271
56. Phenylephrine HCl (Neosynephrine, Isophrin)	HT	SC, IM IV TOP	5 mg 0.5 mg 0.25%			
57. Synephrine (Sympatol, Synhephate)	HT	SC, IM	100 mg			
Sympathomimetic Amines for Topical Application Only						
58. Amphetamine sulfate (Benzedrine, Actedron)	HT	IM		For local nasal decongestion only; these amines are too toxic for parenteral use, with the exception of amphetamine sulfate and dextroamphetamine sulfate.		259,270
59. Dextroamphetamine sulfate (Dexedrine)	HT	TOP	1%			254,259,262,264
60. Metaraminol bitartrate (Aramine)	HT	TOP	0.25%			
61. Methylhexanamine (Fortane)	HT	IM				
62. Naphazoline HCl (Privine)	HT	TOP	0.1%			
63. Nordefrin HCl (Cobefrin)	HT	TOP	0.01%			
64. Phenylpropanolamine HCl (Propadrine)	HT	TOP	1%			270
65. Phenylpropylmethylamine (Vonedrine)	HT	IM, TOP	2.5%			270
66. Propylhexedrine (Benzedrex)	HT	IM				
67. Tuaminoheptane sulfate (Tuamine)	HT	IM, TOP	1%			270
Cardiac Glycosides ⁵						
68. Deslanoside (Cedilanid D)	CT	IV-1	1.6 mg	Improved ventricular function of heart in failure with and without atrial fibrillation; A-V block.	Anorexia, vomiting, headache, yellow vision, muscular weakness, ventricular arrhythmias, excessive A-V block.	254,259,264,270
69. Digifolin	CT	PO-1 M	10 units 1 unit			
70. Digiloid	CT	PO-1 M	3-6 mg 0.3-0.6 mg			
71. Digitalis, powdered	CT	PO-1 M	1-2 g 0.1-0.2 g			
72. Digitalis, tincture ⁶	CT	PO-M	1-2 ml			
73. Digitoxin	CT	PO-1	1.0 mg			
74. Digoxin	CT	PO-1 M	1.5 mg 0.5 mg			254,262,264
75. Gitalin (Gitaligin)	CT	PO-1	4-8 mg			
76. Lanatoside C (Cedilanid)	CT	PO-1 M	8 mg 1 mg			254,264

⁵/5/ I = initial dose, M = daily maintenance doses. ⁶/6/ 1 ml tincture = 0.1 g powdered digitalis

109. DRUGS PRIMARILY USED FOR DISORDERS OF THE CIRCULATORY SYSTEM (Continued)

Drug (Synonym)	Clinical Use	Route	Average Adult Dose	Effect		Hemodynamic Effects Found on Following Pages:
				Desirable	Undesirable	
(A)	(B)	(C)	(D)	(E)	(F)	(G)
Adrenergic Blocking Drugs						
20 Azapetine phosphate (Ildar)	VD	PQ	75-300 mg daily	Increased blood flow to limbs, hypotensive action useful in diagnosis of pheochromocytoma.	Drowsiness, nausea, vomit- ing, tachy- cardia, nasal congestion.	261,267
21 Dibenamine HCl	VD	IV	150 mg			253,263,269
22 Dihydroergocornine methanesulfonate	VD	IV, IM	0.5 mg/kg 0.25-1 mg			258,267
23 Dihydroergotamine methanesulfonate	VD	IV, IM	1-2 mg			253,263,269
24 Hydergine ¹	VD	SL, IV, IM	0.3-1.0 mg			253,261,263,267
25 Phenoxybenzamine HCl (Dibenzyline)	VD	IV	50-300 mg			258,261,267
		PO	10-60 mg daily			
26 Phentolamine HCl, or methanesulfonate (Regitine)	VD	PO	200-400 mg daily			258,267,269
		IV, IM	5 mg			
27 Piperoxan HCl (Benodaine)	VD	IV	0.25 mg/kg			
28 Ronicol tartrate	VD	PO	30-50 mg daily			267
29 Tolazoline HCl (Priscoline)	VD	PQ	200 mg daily			253,258,263,267, 269
		IV, IM, SC	15-25 mg			
Cholinergic Vasodilators						
(Mecholyl)					Peripheral vaso- dilatation accompanied by parasympatho-	257,265,270 253 254,267,271
33 Amyl nitrite	VD					
34 Erythritol tetranitrate	VD					
35 Glyceryl trinitrate (Nitroglycerin)	VD	SL	daily 0.4 mg		emia, develop- ment of tolerance.	254,257,258,269
36 Inositol hexanitrate	VD	PO	30-90 mg daily			
37 Mannitol hexanitrate	VD	PO	50-150 mg daily			
38 Pentaerythritol tetra- nitrate (Peritrate)	VD	PQ	30-60 mg daily			
39 Sodium nitrite	VD	PO	100-200 mg daily			254,264
40 Triethanolamine trinitrate biphosphate (Metamine)	VD	PO	2-6 mg daily			
41 Aminophylline (Theo- phylline ethylene- diamine)	VD	PO, SC, IM, IV	0.5-1.0 g 0.25-0.5 g		Variable effects on coronary and limb blood flow, incon- sistent clinical results.	253,257,258,261, 264,270
42 Dioxylone phosphate (Paveril)	VD	PO	0.6 g			254,257,259,261, 267
43 Ethyl alcohol	VD	PO	Variable			254,257,265
44 Khellin (Eske)	VD	PO	30-160 mg			
(Ephedrine)						
48 Ephedrine sulfate	HT	PO, SC, IM	15-50 mg	arterial pres- sure.	induce cardiac arrhythmias and cerebral excita- tion.	254,270

1/4) Equal parts of dihydroergocornine, dihydroergocristine, and dihydroergokryptine methanesulfonates.

109. DRUGS PRIMARILY USED FOR DISORDERS OF THE CIRCULATORY SYSTEM (Continued)

Drug (Synonym)	Clinical Use	Route	Average Adult Dose	Effect		Hemodynamic Effects Found on Following Pages ¹ (G)
				Desirable (E)	Undesirable (F)	
(A)						
49 Epinephrine, epineph bitartrate (Adrenalin, Suprarenin)	HT	IM	10 mg		an and cerebral excitation.	254,257,259,262, 264,267,270
50 Hydroxylamphetamine hydrobromide (Paredrine)	HT	TOP	0.25-1%			254,264
51 Levarterenol bitartrate (Levophed, Noradrena- lin)	HT	IV	5 µg/min			254,257,259,262, 264,267,270
52 Mephentermine sulfate (Wyamine)	HT	IM, IV	20 mg			254,264,270
53 Methamphetamine HCl (Methedrine, Desoxy- ephedrine, Desoxin, Desoval, Doxyfed, Etroxine, Norodin)	HT	PO SC, IM, IV TOP	5 mg 10-30 mg 0.5-1%			270
54 Methoxamine HCl (Vasoxyl)	HT	IM	5 mg			254,257,264,271
55 Oenethyl HCl	HT	IV	10-25 mg			
56 Phenylephrine HCl (Neosynephrine, Isophrin)	HT	IM IV TOP	75-100 mg 5 mg 0.5 mg 0.25%			254,264,271
57 Symphephrine (Sympatol, Synphenate)	HT	SC, IM	100 mg			
Sympathomimetic Amines for Topical Application Only						
58 Amphetamine sulfate (Benzedrine, Actedron)	HT	IM	1H	For local nasal decongestion only, these amines are too toxic for par- enternal use, with the excep- tion of amphet- amine sulfate and dextroampheta- mine sulfate.		259,270
59 Dextroamphetamine sulfate (Dexedrine)	HT	TOP	1%			254,259,262,264
60 Metaraminol bitartrate (Aramine)	HT	TOP	0.25%			
61 Methylhexanamine (Forthane)	HT	IM				
62 Naphazoline HCl (Privine)	HT	TOP	0.1%			
63 Nordefrin HCl (Cobefrin)	HT	TOP	0.01%			
64 Phenylpropanolamine HCl (Propadrine)	HT	TOP	1%			270
65 Phenylpropylmethylamine (Vonedrine)	HT	IM, TOP	2.8%			270
66 Propylhexedrine (Benzedrex)	HT	IM				
67 Tuaminoheptane sulfate (Tumaine)	HT	IM, TOP	1%			270
Cardiac Glycosides ²						
68 Deslanoside (Cedilanid D)	CT	IV, I	1.6 mg	Improved ventric- ular function of heart in failure with and without atrial fibrilla- tion; A-V block.	Anorexia, vomit- ing, headache, yellow vision, muscular weak- ness, ventricu- lar arrhythmias, excessive A-V block.	254,259,264,270
69 Digifolin	CT	PO I	10 units			
70 Digilanid	CT	PO I	1-6 mg			
71 Digitalis, powdered	CT	PO I	0.3-0.6 mg			
72 Digitalis, tincture ³	CT	PO I	0.8-1.5 mg			
73 Digitoxin	CT	PO I	1-2 g			
74 Digoxin	CT	PO I	0.1-0.2 g			
75 Gitalin (Gitaligin)	CT	PO I	1-2 ml			
76 Lanatoside C (Cedilanid)	CT	PO I	1.0 mg			
		PO I	1.5 mg			
		PO I	0.5 mg			
		IV: I	1.0 mg			
		PO I	4-8 mg			
		M	0.25-1.0 mg			
		PO I	8 mg			
		M	1 mg			

¹/5/ 1 = initial dose, M = daily maintenance doses. ²/6/ 1 ml tincture = 0.1 g powdered digitalis.

109. DRUGS PRIMARILY USED FOR DISORDERS OF THE CIRCULATORY SYSTEM (Continued)

Drug (Synonym)	Clinical Use	Route	Average Adult Dose	Effect		Hemodynamic Effects Found on Following Pages
				Desirable	Undesirable	
(A)	(B)	(C)	(D)	(E)	(F)	(G)
Cardiac Glycosides						
				heart in failure with and without atrial fibrillation; A-V block.	Yellow vision, muscular weakness, ventricular arrhythmia, excessive A-V block.	257
Cinchona Alkaloids						
82 Quinidine gluconate	AA	IM	0.4-0.8 g	Antifibrillatory action.	Cinchonism, vascular collapse following IV injection.	
83 Quinidine HCl	AA	IM, IV	0.45-0.65 g			
84 Quinidine sulfate	AA	PO	0.2 g			255, 269
Synthetic Antiarrhythmic Drugs						
85 Ambonestyl HCl	AA	IV	0.5 g	Depression of cardiac muscle contractility and hypotension.		269
86 Procaine amide HCl (Pronestyl)	AA	PO	6.0 g			255, 264, 269
		IV	1 g			

/5/ I = initial dose, M = daily maintenance doses.

Contributors: (a) Aviado, Domingo M., Jr., and Carl F. Schmidt, (b) Kraye, Otto, (c) Freis, Edward D.

110. DRUGS PRIMARILY USED FOR DISORDERS NOT LIMITED TO THE CIRCULATORY SYSTEM

Drugs listed are primarily used for disorders of systems other than the circulatory; grouping is based on a common action. Official generic names are from United States Pharmacopeia XV, National Formulary X, or from New and Nonofficial Remedies, 1955.

Drug	Use in Clinical Disorder of Circulatory System	Cardiovascular Action	Hemodynamic Effects Found on Following Pages
(A)	(B)	(C)	(D)
Hypnotics and Sedatives			
1 Amobarbital sodium	Sedative in hypertension and heart disease.	Large doses depressant to heart muscle contraction.	255
2 Pentobarbital sodium			267
3 Phenobarbital sodium			271
Analgesics			
4 Meperidine HCl	Sedative for acute myocardial infarction or acute congestive failure.	Large doses produce vasomotor depression and bradycardia.	269
5 Morphine sulfate or tartrate			262
6 Sodium salicylate	Antipyretic for rheumatic carditis.	Claimed to have direct effect on cardiac lesion.	255
General Anesthetics			
7 Chloroform	No direct use, cardiac patients may be subjected to general anesthetics if necessary.	Depressant to heart muscle contraction, in general, all general anesthetics act as dilators of blood vessels; chloroform and cyclopropane can cause dangerous cardiac arrhythmias.	271
8 Cyclopropane			255, 271
9 Ether			255, 271
10 Ethyl chloride			271
11 Nitrous oxide			255, 259, 271
12 Thiopental sodium			271
13 Trichloroethylene			271
14 Vinyl ether			271
Local or Spinal Anesthetic			
15 Procaine HCl	Applied locally to exposed heart during cardiac surgery to prevent arrhythmia.	Spinal anesthesia may reduce arterial blood pressure by sympathetic blockade and decreased thoracic movement.	255, 257, 259, 262, 264, 268, 269
Anticonvulsants			
16 Diphenylhydantoin sodium	None	Antiarrhythmic activity.	269
17 Trimethadione			

110. DRUGS PRIMARILY USED FOR DISORDERS NOT LIMITED TO THE CIRCULATORY SYSTEM (Continued)

Drug	Use in Clinical Disorder of Circulatory System	Cardiovascular Action	Hemodynamic Effects Found on Following Pages
(A)	(B)	(C)	(D)
Analeptics			
18. Caffeine and sodium benzoate	Stimulant to vasomotor and respiratory centers.	Direct contraction of heart muscle and dilatation of blood vessels.	258,270
19. Nikethamide		Direct effects on heart and coronaries uncertain.	257
Tranquilizing Drug			
20. Chlorpromazine	Prevention of nausea in uremia, eclampsia and veratrum therapy, and relief of mental aberrations	Essentially secondary to actions on nervous system.	259,271
Sympathomimetic Bronchodilators			
21. Isoproterenol HCl	Relief of bronchial asthma.	Hypotension.	254,264,267,270
22. Methoxyphenamine HCl			
Parasympathetic Blocking Drugs			
23. Atropine sulfate	Diagnosis and treatment of cardiac arrhythmia due to increased vagal tone; antidote for poisoning by anticholinesterase agents	Cardio-accelerator.	267,270
24. Methantheline bromide			
25. Scopolamine hydrobromide			
Neuromuscular Blocking Drugs			
26. Succinylcholine chloride	None.	Curare preparations cause hypotension by histamine liberation, ganglion blockade, and decreased muscle tonus	267
27. Tubocurarine chloride			
Antihistamines			
28. Diphenhydramine HCl	None.	Antifibrillatory action demonstrated in animals.	269
29. Pyridamine maleate			
30. Triphenylamine HCl			
Diuretics			
31. Acetazolamide (Diamox)	Mobilization of water and sodium ion in congestive heart failure	Decreased blood volume aids in treating congestive heart failure, large doses depressant to heart muscle.	
32. Meralluride sodium			
33. Mercurophylline sodium			
34. Mersalyl sodium			255
Anticoagulants			
35. Ethyldiisocoumarin	Prophylaxis against thrombo-embolic diseases	Prevention of thrombosis and emboli in coronary and other vascular beds.	254
36. Heparin			
Antibiotics			
37. Benzathine penicillin G	Control of subacute bacterial endocarditis	Secondary to arrest of infectious process	
38. Procaine penicillin G			260
Antimalarials			
39. Chloroquine diphosphate	Antifibrillatory agent	Decreased automaticity (ectopic) and contractility of heart muscle	269
40. Quinacrine HCl			269
Endocrine Preparations			
41. Corticotropin	None	None	
42. Cortisone acetate			259
43. Desoxycorticosterone acetate			255
44. Insulin			259
45. Pitressin			
46. Posterior pituitary extract		Vasoconstrictor.	259
47. Thiouracil	Treatment of hyperthyroid heart disease	None directly	255,265
Vitamins			
48. Cyanocobalamin (B ₁₂)	None	None	
49. Nicotinic acid	Treatment of peripheral vascular disease	Vasodilator	259
50. Thiamine HCl	Treatment of cardiac beriberi	None, unless deficiency symptoms exist.	254,259,267
111 Questionable			

110. DRUGS PRIMARILY USED FOR DISORDERS NOT LIMITED TO THE CIRCULATORY SYSTEM (Concluded)

Drug	Use in Clinical Disorder of Circulatory System	Cardiovascular Action	Hemodynamic Effects Found on Following Pages:
(A)	(B)	(C)	(D)
Plasma Expanders			
51 Dextran, 6% solution	Plasma expander in shock.	Increased cardiac output.	265
52 Gelatin, 6% solution			
53 Sodium chloride, 0.9% solution			260,262,265
Anions and Cations			
54 Ammonium chloride	Limited clinical trial as anti- arrhythmic drug or as stim- ulant to arrested ventricles.	Alterations in automaticity of heart.	259
55 Barium chloride			270
56 Calcium chloride			271
57 Magnesium sulfate			262,271
58 Potassium chloride			271
59 Sodium bicarbonate			260
60 Sodium lactate			270
Miscellaneous Drugs for Animal Experimental Use			
61 Aconitine	None because of toxicity.	Proarrhythmic.	270
62 Chloralose		General anesthetic.	
63 β -Dihydroisolasodine		Antiaccelerator.	270
64 DMPP (1,1-Dimeth- yl-4-phenyl- piperazinium iodide)		Ganglion stimulant.	
65 α -Fagarine		Antiarrhythmic.	269
66 Histamine phosphate or diphosphate		Vasodilator.	257,258,262,265, 267
67 Lobeline		Ganglion stimulant.	
68 Nicotine			
69 Serotonin (5-Hy- droxytryptamine)		Vasoconstrictor.	255,265
70 Sodium cyanide		Chemoreceptor stimulant.	
71 Sodium sulfide			
72 Urethane		General anesthetic.	271
73 Veratramine		Antiaccelerator.	271
74 Veratridine		Visceral receptor stimulant.	

Contributors: (a) Aviado, Domingo M., Jr., and Carl F. Schmidt, (b) Kraye, Otto

111. HEMODYNAMIC EFFECTS OF DRUGS: MAMMALS

Part I: MEAN ARTERIAL PRESSURE, CARDIAC OUTPUT, AND TOTAL VASCULAR RESISTANCE: MAN, DOG

Values are averages for reported number of cases and represent per cent change from control. Route: IH = inhalation, IM = intramuscular, IS = intraspinal, IV = intravenous; PO = oral, SC = subcutaneous, SL = sublingual. (+) = increase, (-) = decrease, (0) = no significant change.

Drug	Route	Dose	Subjects			Mean Arterial (Systemic) Pressure ¹	Cardiac Output ²	Total Vascular (Systemic) Resistance ³	Reference
			Animal	No.	Condition	% Change			
						(G)	(H)	(I)	
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)
Ganglion Blocking Drugs									
1 Azamethonium dibromide	IV	0.6-1.4 mg/kg	Man	10	Peripheral vascular disease	-8	+23 BCG		1
2 Hexamethonium bromide or chloride	IV	0.2-1 mg/kg	Man	10	Normotension	-20	+	-17	2
			Man	5	Hypertension	-25	-46		3
			Man	6	Hypertension, compensated	-24	-22	0	4
5			Man	4	Hypertension, decompensated	-31	+38	-47	4

/1/ Measured by mercury manometer or electrometer. /2/ Measured by Fick principle, unless otherwise indicated (BCG = ballistocardiograph, DYE = dye-dilution technique). /3/ Ratio of pressure to flow, or absolute values.

111. HEMODYNAMIC EFFECTS OF DRUGS. MAMMALS (Continued)

Part I. MEAN ARTERIAL PRESSURE, CARDIAC OUTPUT, AND TOTAL VASCULAR RESISTANCE: MAN, DOG (Continued)

Part I. MEAN ARTERIAL PRESSURE AND TOTAL VASCULAR RESISTANCE: MAN, DOG (Continued)

Drug	Route	Dose	Subjects		Mean Arterial (Systemic) Pressure ¹	Cardiac Output ²	Total Vascular (Systemic) Resistance ³	Reference	
			Animal	No.					Condition
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)
Ganglion Blocking Drugs (concluded)									
Hexamethonium bromide or chloride (concluded)	IV	0.2-1 mg/kg	Man	8	Pulmonary hypertension	-27	-16	-12	5
			Dog		Anesthetized	-35	-30		6
Pentamethonium bromide	IV	0.3-0.7 mg/kg	Man	14	Peripheral vascular disease	-11	+29 BCG		1
Tetraethylammonium bromide or chloride	IV	2-6 mg/kg	Man	14	Normotension	-10	0	-	7,8
			Man	4	Hypertension	-	-	-	9
			Man	5	Mitral stenosis	-5	+12	-27	10,11
			Man	5	Left heart failure	-21	0	-28	10
			Man	10	Cardiopulmonary disease	-	-32	-	7
			Man	11	Peripheral vascular disease	-5	+35		12
Trimethaphan camphorsulfate	IV	0.03-0.2 mg/kg	Man	2	Normotension, non-pregnant	-5	0 BCG	-8	13
			Man	5	Normotension, pregnant	-38	-19 BCG	-22	13
			Man	3	Hypertension, pregnant	-33	-18 BCG	-19	13
Adrenergic Blocking Drugs									
Dibenzamine HCl	IV	0.5-3.0 mg/kg	Man	9	Normotension	-6	0		8
			Man	4	Mitral stenosis	-6	+7		10
			Man	3	Congestive heart failure		+28		14
			Man	4	Left heart failure	-9	-6	-4	10
Dihydroergotamine methanesulfonate	IV	1 mg	Man	46	Normotension	+3	0	+45	8,10, 15
Hyderygine	IV	0.5 mg	Man	9	Normotension	-4	0	-5	15
Tolazoline HCl	IV	0.4-0.6 mg/kg	Man	16	Peripheral vascular disease	+8	+28 BCG		16
Antihypertensive Drugs									
Hydralazine	IV	0.2-5 mg/kg	Man	4	Normotension	-25	+110	-61	17
			Man	17	Hypertension	-17	+33	-31	18
.									
Reserpine	PO	1 mg daily for 6-35 wks	Man	54	Normotension, pregnancy	-	+ BCG		23
	IV		Dog	5	Anesthetized	0	0		24
Sodium thiocyanate	IV	5-100 mg/kg	Dog	15	Anesthetized	0	-18	+	25
Veratrine	IM	0.4-1.0 ml	Man	2	Hypertension, compensated	-11	0	-6	26
			Man	3	Hypertension, decompensated	-14	0	-21	26
Miscellaneous Vasodilators									
Aminophylline	IV, IM	0.5 g	Man	7	Hypertension, compensated	0	+6	-4	27
			Man	4	Hypertension, decompensated	+8	+33	-17	27
			Man	7	Normotension	+7	+25 BCG	-19	28
Carbachol	SC	0.5 mg	Man	4	Peripheral vascular disease	-7	+12 BCG	-17	28

/1/ Measured by mercury manometer or electrometer. /2/ Measured by Fick principle, unless otherwise indicated (BCG = ballistocardiograph, DYE = dye-dilution technique). /3/ Ratio of pressure to flow, or absolute values.

111. HEMODYNAMIC EFFECTS OF DRUGS: MAMMALS (Continued)

Part I: MEAN ARTERIAL PRESSURE, CARDIAC OUTPUT,
AND TOTAL VASCULAR RESISTANCE: MAN, DOG (Continued)

Drug	Route	Dose	Subjects			Mean Arterial (Systemic) Pressure ¹	Cardiac Output ²	Total Vascular (Systemic) Resistance ³	Reference
			Animal	No.	Condition				
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)
Miscellaneous Vasodilators (concluded)									
40 Ethyl alcohol	PO	60-150 ml, 40%	Man	5	Peripheral vascular disease	0	-5 BCG		12
41									
43			Man	5	Chronic pulmonary disease		0		31
44 Methacholine chloride	SC	3.7-5 mg	Man	3	Peripheral vascular disease	-7	+29 BCG		12
45 Nicotinic acid	PO	50-100 mg	Man	8	Peripheral vascular disease	0	-4 BCG		12
46 Sodium nitrite	IV	40-60 mg	Man	14	Various cardio-vascular diseases	-3	+21		8
47			Man	4	Mitral stenosis	-5	+11	-15	10
48			Man	5	Left heart failure	-5	+10	-13	10
49 Ephedrine sulfate	SC								
50 Epinephrine	SC, IM ⁴								
51									
52									
53									
54 Hydroxyamphetamine	IM								
62 Metaraminol bitartrate	IV	0.65 mg	Man	7	Normotension	+50	0	+55	40
63 Methoxamine HCl	IV	10 µg/kg/min	Dog	4	Anesthetized	+	-30	+	41
71			Man	5	Left ventricular failure	0	+25	-28	45
72			Man	3	Cor pulmonale, not in failure	+18	0	+7	46
73			Man	5	Cor pulmonale, in failure	+10	+15	+2	46
74			Man	12	Enlarged heart, not in failure	+6	+9		47
75			Man	2	Pulmonary hypertension	+11	+7	+5	46
76 Lanatoside C	IV	1.5 mg	Man	6	Mitral stenosis	+	+		48
77			Man	9	Chronic pulmonary disease		+		49

ple, unless otherwise indicated flow, or absolute values.

111. HEMODYNAMIC EFFECTS OF DRUGS. MAMMALS (Continued)

Part I. MEAN ARTERIAL PRESSURE, CARDIAC OUTPUT,
AND TOTAL VASCULAR RESISTANCE. MAN, DOG (Continued)

Drug	Route	Dose	Subjects			Mean Arterial (Systemic) Pressure ¹	Cardiac Output ²	Total Vascular (Systemic) Resistance ³	Reference
			Animal	No.	Condition				
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)
General Anesthetics and Sedatives									
78 Amobarbital sodium	PO	0.4 g	Man	5	Normotension	+8	-26 BCG	+44	56
79			Man	5	Hypertension	-13	-26	+26	50
80 Cyclopropane	IH		Dog	21	Blood level, 9-29 mg %	0	0 DYE	0	51, 52
81 Ether	IH		Dog	20	Blood level, 85-165 mg %	-5	+71 DYE	-44	51, 52
82	IH		Dog	9	Blood level, 31-190 mg %		+100		53
83 Thiopental sodium	IV		Man	15	Light surgical anesthesia	-19	-24 DYE	0	54
84			Man	15	Deep surgical anesthesia	-16	-20 DYE	+4	55
85			Man	14	Deep surgical anesthesia	-4	-25 DYE		56
Miscellaneous Drugs									
86 Coriisone acetate	PO	2-6 g daily for 17-29 da	Man	9	Normal	+	0		57
87 Mersalyl sodium	IV	3 ml	Man	11	Congestive heart failure		+19		58
88 Posterior pituitary extract	SC, IV	20 units	Man	7	Various diseases	-2	-2 BCG	-1	28
89			Dog	10	Anesthetized		-54		
90 Procaine amide HCl	IV	1 g	Man	10	Heart disease	-17	-11 DYE	+5	59
91 Procaine HCl	IV	0.25-0.5 g	Man	9	Normal	+2	-2 BCG		60
92			Man	9	Peripheral vascular disease	-7.6	+6 BCG		61
93	IS	0.25-0.5 g	Man	12	Pregnant, high spinal anesthesia	-	+23 BCG		62
94 Quinidine sulfate	PO	0.8-1.2 g	Man	11	Normal heart	-25	0	-	63
95			Man	8	Heart disease	0	-4 BCG	+4	28
96 Serotonin	IV	0.5 mg	Dog	25	Anesthetized	+	+30 DYE	-3	64
97 Sodium salicylate	IV	100 mg/kg	Dog	7	Anesthetized	+3	+14		65

/1/ Measured by mercury manometer or electrometer. /2/ Measured by Fick principle, unless otherwise indicated (BCG = ballistocardiograph. DYE = dye-dilution technique. /3/ Ratio of pressure to flow, or absolute values.
/6/ Decreased systolic but increased diastolic blood pressure

Contributors: Aviando, Domingo M., Jr., and Carl F. Schmidt

- References [1] Needleman, H. L., and O. Horwitz. 1953. *Am. J. M. Sc.* 226:164. [2] Rakita, L., and S. M. Saccetta. 1953. *Circul. Res.*, N. Y. 1:499. [3] Grob, D., W. R. Scarborough, A. A. Kattus, and H. G. Langford. 1953. *Circulation*, N. Y. 8:352. [4] Freis, E. D., J. C. Rose, E. A. Partenope, T. F. Higgins, R. T. Kelley, H. W. Schnaper, and R. L. Johnson. 1953. *J. Clin. Invest.* 32:1285. [5] Saccetta, S. M. 1955. *Am. Heart J.* 49:501. [6] Wakim, K. G. 1955. *Ibid.* 50:435. [7] Fowler, N. O., R. N. Westcott, V. D. Hauenstein, R. C. Scott, and J. McGuire. 1950. *J. Clin. Invest.* 29:1387. [8] Halmagyi, D., B. Felkai, J. Ivanyi, and G. Hetenyi. 1952. *Brit. Heart J.* 14:101. [9] Friak, A. R., S. Hammarsstrom, H. Lagerlöf, L. Werkö, G. Björkenheim, A. Holmgren, and Y. Larsson. 1948. *Am. J. Med.* 5:807. [10] Halmagyi, D., B. Felkai, J. Ivanyi, T. Zwoter, M. Tenyi, and Zs. Szűcs. 1953. *Brit. Heart J.* 15:15. [11] Scott, R. C., S. Kaplan, and W. J. Sillea. 1955. *Am. Heart J.* 50:720. [12] Horwitz, O., H. Montgomery, E. D. Longaker, and A. Sayen. 1949. *Am. J. M. Sc.* 218:669. [13] Assali, N. S., R. A. Douglas, and R. Suyemoto. 1953. *Circulation*, N. Y. 8:62. [14] Brod, J., and Z. Fejfar. 1951. *Sporn. k. Invest.*, Oslo 5:85. [15] Halmagyi, D., J. Ivanyi, B. Felkai, T. Zwoter, M. Tenyi, and Zs. Szűcs. 1953. *Scand. J. Clin. k. Invest.*, Oslo 5:85. [16] Rottenstein, H., O. Horwitz, H. Montgomery, A. Sayen, and L. L. Siems. 1953. *Am.*

- J. M. Sc. 221:661. [17] Wilkinson, E. L., H. Backmann, and H. H. Hecht. 1952. *J. Clin. Invest.* 31:872.
- [18] Rowe, G. G., J. H. Huston, G. M. Maxwell, A. P. Crosley, and C. W. Crumpton. 1955. *Ibid.* 34:115.
- [19] Rowe, G. G., J. H. Huston, G. M. Maxwell, A. B. Weinstein, H. Tuchman, and C. W. Crumpton. 1955. *Ibid.* 34:696. [20] Freis, E. D., J. C. Rose, T. F. Higgins, F. A. Finnerty, R. T. Kelley, and E. A. Partenope. 1953. *Circulation*, N. Y. 8:199. [21] Werkö, L., E. Varnauskas, J. Ek, H. Bucht, B. Thomasson, J. Bergstrom, and H. Eliasch. 1954. *Ibid.* 9:700. [22] Assali, N. S., S. Kaplan, S. Oigheinstein, and R. Suyemoto. 1953. *J. Clin. Invest.* 32:922. [23] Cotten, H. B., W. S. Herren, W. C. McDary, and M. S. Klapper. 1955. *Am. J. M. Sc.* 230:408. [24] Trapold, J. H., A. J. Plummer, and F. F. Yonkman. 1954. *J. Pharm. Exp. Ther.* 110:205.
- [25] Bacq, Z. M., R. Charlier, E. Philippot, and P. Fischer. 1949. *Brit. J. Pharm.* 4:162. [26] Freis, E. D., J. R. Stanton, J. W. Culbertson, J. Litter, M. H. Halperin, C. H. Burnett, and R. W. Wilkins. 1949. *J. Clin. Invest.* 28:353. [27] Werkö, L., and H. Lagerlöf. 1950. *Scand. J. Clin. & Invest.*, Oslo 2:181. [28] Starr, I., C. J. Gamble, A. Margolies, J. S. Donal, N. Joseph, and E. Eagle. 1937. *J. Clin. Invest.* 16:799. [29] Wegria, R., J. L. Nickerson, R. B. Case, and J. F. Holland. 1951. *Am. J. Med.* 10:414. [30] Cash, H. R., and H. A. Zimmerman. 1952. *Dis. Chest* 21:137. [31] Silber, E. N., G. L. Snider, H. Goldberg, L. N. Katz, and D. B. Radner. 1951. *J. Clin. Invest.* 30:1046. [32] Witham, A. C., and J. W. Fleming. 1951. *Ibid.* 30:707.
- [33] Barcroft, H., and I. Starr. 1951. *Clin. Sc., Lond.* 10:295. [34] Goldenberg, M., K. L. Pines, E. deF. Baldwin, D. G. Greene, and C. E. Roth. 1948. *Am. J. Med.* 5:792. [35] Kaufman, J., A. Iglaier, and G. K. Herwitz. 1951. *Ibid.* 11:442. [36] Fowler, N. O., R. N. Westcott, R. C. Scott, and J. McGuire. 1951. *J. Clin. Invest.* 30:517. [37] Delaunais, A. L., and C. W. Vandenberghe. 1954. *Arch. internat. pharm. dyn.* Gand 99:435. [38] Gilmore, J. P., C. M. Smythe, and S. W. Handford. 1954. *J. Clin. Invest.* 33:884.
- [39] Brofman, B. L., H. K. Hellerstein, and W. H. Caskey. 1952. *Am. Heart J.* 44:396. [40] Livesay, W. R., J. H. Moyer, and D. W. Chapman. 1954. *Ibid.* 47:745. [41] West, J. W. Unpublished. [42] Horvath, S. M., and D. W. Knapp. 1954. *Am. J. Physiol.* 178:387. [43] Eichna, L. W., et al. 1953. *Circulation*, N. Y. 7:674.
- [44] Eichna, L. W., et al. 1951. *J. Clin. Invest.* 30:1250. [45] Harvey, R. M., M. I. Ferrer, R. T. Cathcart, D. W. Richards, Jr., and A. Courmand. 1949. *Am. J. Med.* 7:439. [46] Ferrer, M. I., R. M. Harvey, R. T. Cathcart, C. A. Webster, D. W. Richards, Jr., and A. Courmand. 1950. *Circulation*, N. Y. 1:161.
- [47] Harvey, R. M., M. I. Ferrer, R. T. Cathcart, and J. K. Alexander. 1951. *Ibid.* 4:369. [48] Gray, F. D., and F. G. Gray. 1954. *Am. Heart J.* 47:282. [49] Gray, F. D., M. H. Williams, and F. G. Gray. 1952. *Ibid.* 44:517.
- [50] Winchell, P., L. H. Taylor, and C. B. Chapman. 1951. *Circulation*, N. Y. 4:229. [51] Grelshelmer, E. M., et al. 1953. *Am. J. Physiol.* 175:171. [52] Grelshelmer, E. M., et al. 1954. *Ibid.* 177:489. [53] Brewster, W. R., J. P. Isaacs, and T. Walno-Anderson. 1953. *Ibid.* 175:399. [54] Fieldman, E. J., R. W. Ridley, and E. H. Wood. 1955. *Anesthesiology* 16:473. [55] Elder, J. D., S. M. Nagano, D. W. Eastwood, and D. Harnagel. 1955. *Ibid.* 16:394. [56] Etsten, B., and T. H. Li. 1955. *J. Clin. Invest.* 34:500. [57] Albert, R. E., W. W. Smith, and L. W. Eichna. 1955. *Circulation*, N. Y. 12:1047. [58] Pugh, L. G., and C. L. Wyndham. 1949. *Clin. Sc., Lond.* 8:11. [59] Wakim, K. G., C. Denton, and H. E. Essex. 1954. *Am. Heart J.* 47:77. [60] McClendon, R. L., W. R. Hansen, and J. M. Kinsman. 1951. *Am. J. M. Sc.* 222:375. [61] Jesar, R. A., O. Horwitz, and H. Montgomery. 1952. *Ibid.* 224:300. [62] Assali, N. S., and H. Prystowski. 1950. *J. Clin. Invest.* 29:1367.
- [63] Ferrer, M. I., R. M. Harvey, L. Werkö, D. T. Dresdale, A. Courmand, and D. W. Richards, Jr. 1948. *Am. Heart J.* 36:816. [64] MacCannon, D. M., and S. M. Horvath. 1954. *Am. J. Physiol.* 179:131.
- [65] Tenney, S. M., and R. M. Miller. 1955. *Am. J. Med.* 19:498.

111. HEMODYNAMIC EFFECTS OF DRUGS: MAMMALS (Continued)

† II CORONARY BLOOD FLOW, VASCULAR RESISTANCE, AND CARDIAC O₂ UPTAKE: MAN, DOG, RABBIT
Values are averages for reported number of cases and represent per cent change from control. Route: CA = injected directly into coronary artery; IH = inhalation; IM = intramuscular, IS = intrasplenic, IV = intravenous. Method: J = nitrous oxide sampling [16], RTM = rotameter inserted into coronary artery; BFM = bubble flowmeter inserted into coronary artery. CSC = collection of coronary sinus outflow by Morawitz cannula, PIII = perfused late heart. (+) = increased, (-) = decreased, {0} = no significant change.

	Coronary	Cardiac
--	----------	---------

Table 1. Effect of various drugs on the heart rate of the heart.										
Drug	Route	Dose	Subjects			Coronary Blood Flow		Coronary Vascular Resistance	Cardiac O ₂ Uptake ²	Reference
			Animal (D)	No. (E)	Condition (F)	Method (G)	% Change (H)			
(A)	(B)	(C)						(I)	(J)	(K)
Ganglion Blocking Drugs										
Hexamethonium bromide or chloride	IV	1 mg/kg	Man	2	Hypertension	N ₂ O	-	-	-	1
			Dog	16	Anesthetized	N ₂ O	-15	+2	+2	2
			Dog	7	Renal hypertension	N ₂ O	-14	+1	-	2
Cardiac Glycosides										
Ouabain	IV	0.03 mg/kg	Dog	11	Anesthetized	N ₂ O	-11	+34	+4	3
		0.04-0.06 mg/kg	Dog	7	Anesthetized	CSC	+190	-70	-	4
K-Strophanthin	IV	0.65 mg	Man	7	Normal	N ₂ O	-1	-3	+5	5
			Man	5	Heart failure	N ₂ O	+9	-5	+24	5
Sympathomimetic Amines										
Epinephrine	CA	20 µg	Dog	9	Anesthetized	RTM	+128	+59		10
Papaverine HCl	IV	15-20 µg	Dog	6	Anesthetized	BFM	+60	-	-	1
		4-40 µg	Dog	2	Anesthetized	BFM	+20	-	-	2
Sympathomimetic Amines										
Epinephrine	CA	Hyperosmotic doses	Dog	4	Anesthetized	RTM	+77		+80	9
Levarterenol bitartrate	CA	0.2 µg	Dog	9	Anesthetized	BFM	+35	-	-	11
		10 µg	Dog	9	Anesthetized	RTM	+36	-30	-	10
Methoxamine HCl	IV	0.1 mg/kg	Dog	6	Anesthetized	CSC	+	-	-	12
Local Anesthetics										
Nikethamide	IV	1-1.25 g	Dog	4	Anesthetized	BFM	+45	-	+13	7
		0.125-0.75 g	Dog	9	Anesthetized	BFM	+3	-	-	7
Procaine HCl	IS	5 mg	Dog	3	Anesthetized	BFM	+10	-	-	7
			Man	6	Low apical anesthesia	N ₂ O	-52	+7	-53	15

/1/ Calculated from ratio of mean arterial blood pressure to flow. /2/ Calculated by multiplying blood flow and arterial-sinus O_2 (content) difference.

Contributors. (a) Aviado, Domingo M., Jr., and Carl F. Schmidt, (b) Gregg, Donald E.

References [1] Swan, H. J. C. 1951. *J. Physiol.*, Lond. 112 426. [2] Sensenbach, W., L. Madelon, and L. Ochs 1953 *J. Clin. Invest.* 32:226 [3] Page, R. G., E. L. Foltz, W. F. Sheldon, and H. Wendel. 1951. *J. Pharm. Exp. Ther.* 101:112. [4] Sherrod, T. R. 1952. *Ibid.* 104 445. [5] Bing, R. J., F. M. Marriot, J. F. Damman, A. Draper, R. Heimbecker, R. Daley, R. Gerard, and P. Calazel. 1950. *Circulation*, N. Y. 2 511. [6] Foltz, E. L., A. Rubin, W. A. Steiger, and P. C. Gaze. 1950 *Ibid.* 2:215. [7] Eckenhoff, J. E., and J. H. Haftenschiel. 1947. *J. Pharm. Exp. Ther.* 91:362. [8] Lasker, N., T. R. Sherrod, and K. F. Killam. 1955. *Ibid.* 113 414. [9] Eckstein, R. W., W. B. Newberry, J. A. McEachen, and G. Smith. 1951. *Circulation*, N. Y. 4 534.

111. HEMODYNAMIC EFFECTS OF DRUGS: MAMMALS (Continued)

Part I. MEAN ARTERIAL PRESSURE, CARDIAC OUTPUT, AND TOTAL VASCULAR RESISTANCE: MAN, DOG (Concluded)

- J. M. Sc. 221:661. [17] Wilkinson, E. L., H. Backmann, and H. H. Hecht. 1952. J. Clin. Invest. 31:872.
- [18] Rowe, G. G., J. H. Huston, G. M. Maxwell, A. P. Crosley, and C. W. Crumpton. 1955. *Ibid.* 34:115.
- [19] Rowe, G. G., J. H. Huston, G. M. Maxwell, A. B. Weinstein, H. Tuchman, and C. W. Crumpton. 1955. *Ibid.* 34:696. [20] Freis, E. D., J. C. Rose, T. F. Higgins, F. A. Finnerty, R. T. Kelley, and E. A. Partenope. 1953. *Circulation*, N. Y. 8:199. [21] Werkö, L., E. Varnauskas, J. Ek, H. Bucht, B. Thomasson, J. Bergstrom, and H. Ellasch. 1954. *Ibid.* 9:700. [22] Assali, N. S., S. Kaplan, S. Oghenstein, and R. Suyemoto. 1953. J. Clin. Invest. 32:122. [23] Cotten, H. B., W. S. Herren, W. C. McDary, and M. S. Klapper. 1955. *Am. J. M. Sc.* 230:408. [24] Trapold, J. H., A. J. Plummer, and F. F. Yonkman. 1954. J. Pharm. Exp. Ther. 110:205.
- [25] Bacq, Z. M., R. Charlier, E. Philippot, and P. Fischer. 1949. *Brit. J. Pharm.* 4:162. [26] Freis, E. D., J. R. Stanton, J. W. Culbertson, J. Litter, M. H. Halperin, C. H. Burnett, and R. W. Wilkins. 1949. J. Clin. Invest. 28:353. [27] Werkö, L., and H. Lagerlöf. 1950. *Scand. J. Clin. & Invest.*, Oslo 2:181. [28] Starr, L., C. J. Gamble, A. Margolies, J. S. Donal, N. Joseph, and E. Eagle. 1937. J. Clin. Invest. 16:799. [29] Wegria, J. L., Nickerson, R. B. Case, and J. F. Holland. 1951. *Am. J. Med.* 10:414. [30] Cash, H. R., and H. A. Zimmerman. 1952. *Dis. Chest* 21:137. [31] Silber, E. N., G. L. Snider, H. Goldberg, L. N. Katz, and D. B. Radner. 1951. J. Clin. Invest. 30:1046. [32] Witham, A. C., and J. W. Fleming. 1951. *Ibid.* 30:707.
- [33] Barcroft, H., and I. Starr. 1951. *Clin. Sc.*, Lond. 10:295. [34] Goldenberg, M., K. L. Pines, E. deF. Baldwin, D. G. Greene, and C. E. Roh. 1948. *Am. J. Med.* 5:792. [35] Kaufman, J., A. Iglauer, and G. K. Herwitz. 1951. *Ibid.* 11:442. [36] Fowler, N. O., R. N. Westcott, R. C. Scott, and J. McGuire. 1951. J. Clin. Invest. 30:517. [37] Delaunais, A. L., and C. Vandenberghen. 1954. *Arch. internat. pharm. dyn.*, Gand 99:435. [38] Gilmore, J. P., C. M. Smythe, and S. W. Handford. 1954. J. Clin. Invest. 33:884.
- [39] Brofman, B. L., H. K. Hellerstein, and W. H. Caskey. 1952. *Am. Heart J.* 44:396. [40] Livesay, W. R., J. H. Moyer, and D. W. Chapman. 1954. *Ibid.* 47:745. [41] West, J. W. Unpublished. [42] Horvath, S. M., and D. W. Knapp. 1954. *Am. J. Physiol.* 178:387. [43] Eichna, L. W., et al. 1953. *Circulation*, N. Y. 7:674.
- [44] Eichna, L. W., et al. 1951. J. Clin. Invest. 30:1250. [45] Harvey, R. M., M. I. Ferrer, R. T. Cathcart, D. W. Richards, Jr., and A. Courmand. 1949. *Am. J. Med.* 7:439. [46] Ferrer, M. I., R. M. Harvey, R. T. Cathcart, C. A. Webster, D. W. Richards, Jr., and A. Courmand. 1950. *Circulation*, N. Y. 1:161.
- [47] Harvey, R. M., M. I. Ferrer, R. T. Cathcart, and J. K. Alexander. 1951. *Ibid.* 4:369. [48] Gray, F. D., and F. G. Gray. 1954. *Am. Heart J.* 47:282. [49] Gray, F. D., M. H. Williams, and F. G. Gray. 1952. *Ibid.* 44:517.
- [50] Winchell, P., L. H. Taylor, and C. B. Chapman. 1951. *Circulation*, N. Y. 4:229. [51] Greisheimer, E. M., et al. 1953. *Am. J. Physiol.* 175:171. [52] Greisheimer, E. M., et al. 1954. *Ibid.* 177:489. [53] Brewster, W. J. P. Isaacs, and T. Walno-Anderson. 1953. *Ibid.* 175:399. [54] Fieldman, E. J., R. W. Ridley, and E. H. Wood. 1955. *Anesthesiology* 16:473. [55] Elder, J. D., S. M. Nagano, D. W. Eastwood, and D. Harnagel. 1955. *Ibid.* 16:394. [56] Etsten, B., and T. H. Li. 1955. J. Clin. Invest. 34:500. [57] Albert, R. E., W. W. Smith, and L. W. Eichna. 1955. *Circulation*, N. Y. 12:1047. [58] Pugh, L. G., and C. L. Wyndham. 1949. *Clin. Sc.*, Lond. 8:11. [59] Wakim, K. G., C. Denton, and H. E. Essex. 1954. *Am. Heart J.* 47:77. [60] McClendon, R. L., W. R. Hansen, and J. M. Kinsman. 1951. *Am. J. M. Sc.* 222:375. [61] Jessar, R. A., O. Horwitz, and H. Montgomery. 1952. *Ibid.* 224:300. [62] Assali, N. S., and H. Prystowski. 1950. J. Clin. Invest. 29:1367.
- [63] Ferrer, M. I., R. M. Harvey, L. Werkö, D. T. Dresdale, A. Courmand, and D. W. Richards, Jr. 1948. *Am. Heart J.* 36:816. [64] MacCannon, D. M., and S. M. Horvath. 1954. *Am. J. Physiol.* 179:131.
- [65] Tenney, S. M., and R. M. Miller. 1955. *Am. J. Med.* 19:498.

111. HEMODYNAMIC EFFECTS OF DRUGS. MAMMALS (Continued)

Part III. CEREBRAL BLOOD FLOW, VASCULAR RESISTANCE, AND O₂ UPTAKE: MAN (Continued)

Drug	Route	Dose	Subjects		Cerebral Blood Flow ¹		Cerebral Vascular Resistance		Cerebral O ₂ Uptake		Reference
			No.	Condition	Control ²	% Change	Control ³	% Change	Control ²	% Change	
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)	(K)	(L)
Sympathomimetic Amines											
29 Amphetamine sulfate	IV	10 mg	2	Brain tumor	40	-2	2.5	+4	2.4	+4	18
30 Epinephrine	IV ⁵	20-70 µg/min	7	Normal	50	+22	1.8	0	3.4	+23	22
31			15	Normal	66	-4	1.2	0	4.2	-3	23
32 Levarterenol bitartrate	IV ⁵	6-28 µg/min	3	Normal	61	-8	1.6	+37	3.7	-5	22
33			13	Normal	61	-21	1.3	-8	4.0	-5	23
34			4	Normal	62	-12	1.6	+53	4.6	-15	24
35 Metaraminol bitartrate	IV infusion		7	Normal	58	-9	1.6	+50	3.4	-6	24
Nervous System Depressants											
36 Chlorpromazine	IV	200 mg	7	Normal	41	-9	2.9	-24	2.4	+4	25
37 Ethyl alcohol	IV		8	Normal	52	+8	2.1	-14	2.9	-3	25
38		Plus chlorpromazine	6	Normal	49	-6	2.0	-25	2.2	-4	25
39	PO		10	Normal	68	-18	1.6	+12	3.8	-10	26
40		After antabus	10	Normal	54	+18	2.0	-35	3.2	+16	26
41 Insulin (hypoglycemia)			7	Schizophrenia	54	+11	1.6	-13	3.3	-21	27
42			7	Schizophrenia, in coma	54	+16	1.6	-6	3.3	-42	27
43 Procaine HCl	IS		17	Hypertension	52	-12	3.1	-16	3.3	-3	28
44	Stellate block		13	Normotension and hypertension	49	-2	2.5	+8	3.2	-3	29
45			23	Cerebral vascular disease	57	-6	2.2	+4	3.6	-6	30
46 Thiopental sodium	IV		8	Schizophrenia, under semimarcosis	54	-2			3.3	0	27
47			8	Normal, anesthetized	61	-25			3.6	-39	31
48			11	Normal, anesthetized	54	+11	1.6	-19	3.3	-36	32
Miscellaneous Drugs											
49 Ammonium chloride	IV	0.8% (0.3 L)	7	Normal	66	-21	1.3	+23	4.2	+5	33
50 Corticotropin	PO	160-400 mg daily for 4 da	4	Various diseases	52	0	2.2	0	2.9	-14	34
51 Cyanocobalamin, plus liver extract			7	Pericious anemia	91	-56	0.9	+177	2.2	+22	35
52 Deoxycorticosterone acetate	IV	50 mg	15	Normal	58	-5	1.5	+6	3.1	-10	36
53 Digitalis preparations, and diuretics			12	Congestive heart failure	52	-4	3.2	-9	3.1	+13	37
54 Heparin	IV	200 mg	11	Senile dementia	34	-3	3.4	0	2.6	+4	38
55 Nicotinic acid	IV	300-800 mg	20	Various diseases	54	+6	2.3	+9	3.4	-3	39

1/ Measured by nitrous oxide method of Kety and Schmidt [42]. 2/ ml/100 g/min. 3/ Ratio of mean arterial blood pressure to cerebral blood flow. 4/ Or IM, 1 mg

111. HEMODYNAMIC EFFECTS OF DRUGS MAMMALS (Continued)

Part II. CORONARY BLOOD FLOW, VASCULAR RESISTANCE, AND CARDIAC O₂ UPTAKE: MAN, DOG, RABBIT (Concluded)

[10] Winbury, M.M., and D. M. Green. 1952. *Am. J. Physiol.* 170:555. [11] Eckenhoff, J. E., J. H. Hafkenschiel, and C. M. Landmesser. 1947. *Ibid.* 148:582. [12] West, J. W. Unpublished. [13] Rowe, G. G., J. H. Huston, G. M. Maxwell, A. B. Weinstein, H. Tuchman, and C. W. Crumpton. 1955. *J. Clin. Invest.* 34:696. [14] Leusen, I. R., and H. E. Essex. 1953. *Am. J. Physiol.* 172:226. [15] Hackel, D. B., S. M. Sancetta, and J. Kleinerman. 1956. *Circulation*, N. Y. 13:92.

Part III: CEREBRAL BLOOD FLOW, VASCULAR RESISTANCE, AND O₂ UPTAKE. MAN

Values are per cent change from control. Route: IM = intramuscular, IS = intraspinal, IV = intravenous, PO = oral, SL = sublingual. (+) = increased, (-) = decreased, (0) = no significant change. For a comprehensive review, see reference 43.

Drug	Route	Dose	Subjects		Cerebral Blood Flow ¹		Cerebral Vascular Resistance		Cerebral O ₂ Uptake		Reference
			No.	Condition	Control ²	% Change	Control ³	% Change	Control ²	% Change	
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)	(K)	(L)
Ganglion Blocking Drugs											
1 Azamethonium dibromide	IV	1-6 mg/min	5	Normotension	51	-8	2.0	-35	2.8	+7	1
2			9	Normotension	53	+12	1.5	-20	3.2	+10	2
3 Hexamethonium bromide or chloride	IV ⁴	1-8 mg/min	9	Normotension	57	-30	1.9	-11	3.2	-12	3
4			8	Normotension	57	-26	1.8	-17	3.2	-9	1
5			7	Young, normotension	46	-33	1.9	-37	2.8	+4	4
6			10	Elderly, normotension	48	-39	1.7	-31	2.7	0	4
7			10	Head up 15°, normotension	53	-11	2.4	-45	3.3	+3	5
8			8	Essential hypertension	48	-39	2.4	-33	2.8	-4	4
9			7	Malignant hypertension	60	-42	3.0	-17	3.5	-11	4
10			13	Hypertension	55	-16	3.5	-28	3.6	-3	6
11			6	Hypertension	50	-4	3.2	-27	3.5	0	7
Adrenergic Blocking Drugs											
12 Dihydroergocornine methanesulfonate	IM	0.5 mg	5	Normotension	60	-8	1.7	-12	3.8	-8	8
13			12	Hypertension	58	0	2.9	-31	3.5	+3	9
14 Phenoxybenzamine HCl	IV	0.5-0.7 mg/kg	7	Hypertension	48	-12	3.2	-31	3.6	-14	10
15 Phentolamine HCl	IM	1 mg/kg	6	Hypertension	53	0	2.8	-21	3.2	0	11
16 Tolazoline HCl	IV	20-100 mg	24	Normotension	59	-12	1.7	+6	3.6	-3	12
17			6	Mitral stenosis	45	+22	2.0	-25			13
Other Hypotensive Drugs											
18 Alkaverin	IV	1 µg/kg/min	27	Hypertension	57	-12	3.2	-28	2.9	+17	14
19 Aminophylline	IV	0.5 g	10	Normal	59	-25	1.7	+24	3.8	0	15
20			16	Congestive heart failure	45	-33	2.6	+8	3.2	-18	16
21 Caffeine and sodium benzoate	IV	0.25-0.5 g	7	Hypertension	54	-13	2.7	+22	3.2	+9	17
22			9	Brain tumor	44	-13	2.1	+28	2.6	+8	18
23 Glyceryl trinitrate	SL	0.6 mg	6	Mitral stenosis	40	-8	2.5	+4	3.3	-3	13
24 Histamine	IV	0.7 mg	10	Normal	57	0	1.7	-30	3.2	+6	18
25 Papaverine	IV	0.06-2.0 g	18	Normotension	57	+12	1.7	-17	3.7	0	19
26			4	Brain tumor	37	+8	3.6	-22	2.3	+4	18
27 Parephylin	IV	0.5 g	7	Hypertension	45	-4	3.1	+3	2.6	+23	20
28 Reserpine	IV	2.5-5.5 mg	5	Hypertension	50	-2	3.5	-17	3.4	-3	21

/1/ Measured by nitrous oxide method of Kety and Schmidt [42]. /2/ ml/100 g/min. /3/ Ratio of mean arterial blood pressure to cerebral blood flow /4/ Or IM, 1 mg/kg

111. HEMODYNAMIC EFFECTS OF DRUGS. MAMMALS (Continued)

Part III. CEREBRAL BLOOD FLOW, VASCULAR RESISTANCE, AND O₂ UPTAKE MAN (Continued)

Drug	Route	Dose	Subjects		Cerebral Blood Flow ¹		Cerebral Vascular Resistance		Cerebral O ₂ Uptake		Reference	
			No	Condition	Control ²	% Change	Control ³	% Change	Control ²	% Change		
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)	(K)	(L)	
Sympathomimetic Amines												
29	Amphetamine sulfate	IV	10 mg	2	Brain tumor	40	-2	2.5	+4	2.4	+4	18
30	Epinephrine	IV ⁵	20-70 µg/min	7	Normal	50	+22	1.8	0	3.4	+23	22
31				15	Normal	66	-4	1.2	0	4.2	-3	23
32	Levartenerol bitartrate	IV ⁵	6-28 µg/min	3	Normal	61	-8	1.6	+37	3.7	-5	22
33				13	Normal	61	-21	1.3	-8	4.0	-5	23
34				4	Normal	62	-12	1.6	+53	4.6	-15	24
35	Metaraminol bitartrate	IV infusion		7	Normal	58	-9	1.6	+50	3.4	-6	24
Nervous System Depressants												
36	Chlorpromazine	IV	200 mg	7	Normal	41	-9	2.9	-24	2.4	+4	25
37	Ethyl alcohol	IV		8	Normal	52	+8	2.1	-14	2.9	-3	25
38			Plus chlorpromazine	6	Normal	49	-6	2.0	-25	2.2	-4	25
39		PO		10	Normal	68	-18	1.6	+12	3.8	-10	26
40			After antabus	10	Normal	54	+18	2.0	-35	3.2	+16	26
41	Insulin (hypoglycemia)			7	Schizophrenia	54	+11	1.6	-13	3.3	-21	27
42				7	Schizophrenia, in coma	54	+16	1.6	-6	3.3	-42	27
43	Procaine HCl	IS		17	Hypertension	52	-12	3.1	-16	3.3	-3	28
44		Stellate block		13	Normotension and hypertension	49	-2	2.5	+8	3.2	-3	29
45				23	Cerebral vascular disease	57	-6	2.2	+4	3.6	-6	30
46	Thiopental sodium	IV		8	Schizophrenia, under semimarcosis	54	-2			3.3	0	27
47				8	Normal, anesthetized	61	-25			3.6	-39	31
48				11	Normal, anesthetized	54	+11	1.6	-19	3.3	-36	32
Miscellaneous Drugs												
49	Ammonium chloride	IV	0.8% (0.3 L)	7	Normal	66	-21	1.3	+23	4.2	+5	33
50	Corticotropin	PO	260-400 mg daily for 4 da	4	Various diseases	52	0	2.2	0	2.9	-14	34
51	Cyanocobalamin, plus liver extract			7	Pernicious anemia	91	-56	0.9	+177	2.2	+22	35
52	Deoxycorticosterone acetate	IV	50 mg	15	Normal	58	-5	1.5	+6	3.3	-10	36
53	Digitalis preparations, and diuretics			12	Congestive heart failure	52	-4	3.2	-9	3.1	+13	37
54	Heparin	IV	200 mg	11	Senile dementia	34	-3	3.4	0	2.6	+4	38
55	Nicotinic acid	IV	300-800 mg	20	Various diseases	54	+6	2.3	+9	3.4	-3	39

^{1/1} Measured by nitrous oxide method of Kety and Schmidt (43).

1/1 Measured by nitrous oxide method of Kety and Schmidt [42]. 1/2 ml/100 g/min. 1/3 Ratio of mean arterial blood pressure to cerebral blood flow 1/5 Or IM, 1 mg

111. HEMODYNAMIC EFFECTS OF DRUGS. MAMMALS (Continued)

Part II: CORONARY BLOOD FLOW, VASCULAR RESISTANCE, AND CARDIAC O₂ UPTAKE: MAN, DOG, RABBIT (Concluded)

- [10] Winbury, M.M., and D. M. Green. 1952. Am. J. Physiol. 170:555. [11] Eckenhoff, J. E., J. H. Hafkenschiel, and C. M. Landmesser. 1947. Ibid. 148:582. [12] West, J. W. Unpublished. [13] Rowe, G. G., J. H. Huston, G. M. Maxwell, A. B. Weinstein, H. Tuchman, and C. W. Crumpton. 1955. J. Clin. Invest. 34:696. [14] Leusen, I. R., and H. E. Essex. 1953. Am. J. Physiol. 172:226. [15] Hackel, D. B., S. M. Sanctetta, and J. Kleinerman. 1956. Circulation, N. Y. 13:92.

Part III: CEREBRAL BLOOD FLOW, VASCULAR RESISTANCE, AND O₂ UPTAKE: MAN

Values are per cent change from control. Route: IM = intramuscular, IS = intraspinal, IV = intravenous, PO = oral, SL = sublingual. (+) = increased, (-) = decreased, (0) = no significant change. For a comprehensive review, see reference 43.

Drug	Route	Dose	Subjects		Cerebral Blood Flow ¹		Cerebral Vascular Resistance		Cerebral O ₂ Uptake		Reference	
			No.	Condition	Control ²	%			Control ²	%		
tension												
6			10	Elderly, normotension	48	-39	1.7	-31	2.7	0	4	
7			10	Head up 15°; normotension	53	-11	2.4	-45	3.3	+3	5	
8			8	Essential hypertension	48	-39	2.4	-33	2.8	-4	4	
9			7	Malignant hypertension	60	-42	3.0	-17	3.5	-11	4	
10			13	Hypertension	55	-16	3.5	-28	3.6	-3	6	
11			6	Hypertension	50	-4	3.2	-27	3.5	0	7	
Adrenergic Blocking Drugs												
12	Dihydroergocornine methanesulfonate	IM 0.5 mg	5	Normotension	60	-8	1.7	-12	3.8	-8	8	
13			12	Hypertension	58	0	2.9	-31	3.5	+3	9	
14	Phenoxybenzamine HCl	IV 0.5-0.7 mg/kg	7	Hypertension	48	-12	3.2	-31	3.6	-14	10	
15	Phentolamine HCl	IM 1 mg/kg	6	Hypertension	53	0	2.8	-21	3.2	0	11	
16	Tolazoline HCl	IV 20-100 mg	24	Normotension	59	-12	1.7	+6	3.6	-3	12	
17			6	Mitral stenosis	45	+22	2.0	-25			13	
Other Hypotensive Drugs												
18	Alkavervir	IV 1 µg/kg/min	27	Hypertension	57	-12	3.2	-28	2.9	+17	14	
19	Aminophylline	IV 0.5 g	10	Normal	59	-25	1.7	+24	3.8	0	15	
20			16	Congestive heart failure	45	-33	2.6	+8	3.2	-18	16	
21	Caffeine and sodium benzoate	IV 0.25-0.5 g	7	Hypertension	54	-13	2.7	+22	3.2	+9	17	
22			9	Brain tumor	44	-13	2.1	+28	2.6	+8	18	
23	Glyceryl trinitrate	SL 0.6 mg	6	Mitral stenosis	40	-8	2.5	+4	3.3	-3	13	
24	Histamine	IV 0.7 mg	10	Normal	57	0	1.7	-30	3.2	+6	18	
									2	2.3	+4	18
										2.6	+23	20
									7	3.4	-3	21

^{1/1} Measured by nitrous oxide method of Kety and Schmidt [42]. ^{2/2} ml/100 g/min. ^{3/3} Ratio of mean arterial blood pressure to cerebral blood flow ^{4/4} Or IM, 1 mg/kg.

III. HEMODYNAMIC EFFECTS OF DRUGS: SYMPATHETIC (Continued)

Part III: Cerebral Blood Flow, Vascular Resistance, and O₂ Uptake: MAN (Continued)

Drug	Route	Dose	Subjects		Cerebral Blood Flow ¹		Cerebral Vascular Resistance		Cerebral O ₂ Uptake		Reference
			No.	Condition	Control ²	% Change	Control ³	% Change	Control ³	% Change	
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)	(K)	(L)
Sympathomimetic Amines											
29 Amphetamine sulfate	IV	10 mg	2	Brain tumor	43	-2	2.5	+4	2.4	+4	18
30 Ephedrine	IV	25-75 µg/min	7	Normal	50	+22	1.8	0	3.4	+23	22
31			15	Normal	66	-4	1.2	0	4.2	-3	23
32 Levarterenol bitartrate	IV	6-23 µg/min	3	Normal	61	-8	1.6	+37	3.7	-5	22
33			13	Normal	61	-21	1.3	-8	4.0	-5	23
34			4	Normal	62	-12	1.6	+53	4.6	-15	24
35 Metaraminol bitartrate	IV infusion		7	Normal	58	-9	1.0	+50	3.4	-0	24
Nervous System Depressants											
36 Chlorpromazine	IV	250 mg	7	Normal	41	-9	2.9	-24	2.4	+4	25
37 Ethyl alcohol	IV		8	Normal	52	+6	2.1	-14	2.4	-3	25
38		Plus chlorpromazine	6	Normal	49	-6	2.0	-25	2.2	-4	25
39	PO		13	Normal	68	-18	1.0	+12	3.8	-10	26
40		After atabax	10	Normal	54	+18	2.0	-15	3.2	+10	26
41			7	Schizophrenia	54	+11	1.0	-13	3.3	-21	27
42			7	Schizophrenia, in coma	54	+10	1.0	-0	3.3	-42	27
43 Procaine HCl	IS		17	Hypertension	52	-12	3.1	-10	3.3	-3	28
44	Stellate block		13	Hypertension and hypertension	43	-2	2.5	+4	3.2	-3	28
45			23	Cerebral vascular disease	57	-0	2.2	+4	3.0	-0	30
46 Thiopental sodium	IV		8	Schizophrenia, under semistarvation	54	-2			3.3	0	27
47			8	Normal, anesthetized	61	-25			3.0	-17	31
48			10	Normal, anesthetized	54	+11	1.0	-19	3.3	-10	32
Miscellaneous Drugs											
49 Ammonium chloride	IV	0.3% (0.3 L)	7	Normal	60	-21	1.3	+21	4.2	+5	33
50 Carnitropin	PO	200-400 mg daily for 4 d	4	Various diseases	52	0	2.2	0	2.9	-14	34
51 Cyanocobalamin, plus liver extract			7	Pernicious anemia	91	-56	0.9	+177	2.2	+22	35
52 Desoxyepinephrine acetate	IV	50 mg	15	Normal	58	-5	1.5	+0	3.1	-13	36
53 Digitalis preparations, and diuretics			12	Congestive heart failure	52	-4	3.2	-9	3.1	+13	37
54 Dipyrin	IV	250 mg	11	Senile dementia	34	-3	3.4	0	2.0	+4	38
55 Salicylic acid	IV	300-800 mg	20	Various diseases	54	+0	2.3	+3	3.4	-3	39

1/ Measured by nitrous oxide method of Kety and Schmidt (42). 2/ ml/100 g/min. 3/ Ratio of mean arterial blood pressure to cerebral blood flow. 4/ Or 10, 1 mg

111. HEMODYNAMIC EFFECTS OF DRUGS: MAMMALS (Continued)

Part II: CORONARY BLOOD FLOW, VASCULAR RESISTANCE, AND CARDIAC O₂ UPTAKE: MAN, DOG, RABBIT (Concluded)

[10] Winbury, M.M., and D. M. Green. 1952. *Am. J. Physiol.* 170:555. [11] Eckenhoff, J. E., J. H. Hafkenschiel, and C. M. Landmesser. 1947. *Ibid.* 148:582. [12] West, J. W. Unpublished. [13] Rowe, G. G., J. H. Huston, G. M. Maxwell, A. B. Weinstein, H. Tuchman, and C. W. Crumpton. 1955. *J. Clin. Invest.* 34:696. [14] Leusen, I. R., and H. E. Essex. 1953. *Am. J. Physiol.* 172:226. [15] Hackel, D. B., S. M. Sancetta, and J. Kleinerman. 1956. *Circulation*, N. Y. 13:92.

Part III: CEREBRAL BLOOD FLOW, VASCULAR RESISTANCE, AND O₂ UPTAKE: MAN

Values are per cent change from control. Route: IM = intramuscular, IS = intraspinal, IV = intravenous, PO = oral, SL = sublingual. (+) = increased, (-) = decreased, (0) = no significant change. For a comprehensive review, see reference 43.

Drug	Route	Dose	Subjects		Cerebral Blood Flow ¹		Cerebral Vascular Resistance		Cerebral O ₂ Uptake		Reference
			No.	Condition	Control ²	% Change	Control ³	% Change	Control ³	% Change	
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)	(K)	(L)
Ganglion Blocking Drugs											
1 Azamethonium dibromide	IV	1-6 mg/min	5	Normotension	51	-8	2.0	-35	2.8	+7	1
2 Hexamethonium bromide or chloride	IV ⁴	1-6 mg/min	9	Normotension	53	+12	1.5	-20	3.2	+10	2
3			9	Normotension	57	-30	1.9	-11	3.2	-12	3
4			8	Normotension	57	-26	1.8	-17	3.2	-9	1
5			7	Young; normotension	46	-33	1.9	-37	2.8	+4	4
6			10	Elderly; normotension	48	-39	1.7	-31	2.7	0	4
7			10	Head up 15°, normotension	53	-11	2.4	-45	3.3	+3	5
8			8	Essential hypertension	48	-39	2.4	-33	2.8	-4	4
9			7	Malignant hypertension	60	-42	3.0	-17	3.5	-11	4
10			13	Hypertension	55	-16	3.5	-28	3.6	-3	6
11			6	Hypertension	50	-4	3.2	-27	3.5	0	7
Adrenergic Blocking Drugs											
12 Dihydroergocornine methanesulfonate	IM	0.5 mg	5	Normotension	60	-8	1.7	-12	3.8	-8	8
13			12	Hypertension	58	0	2.9	-31	3.5	+3	9
14 Phenoxylbenzamine HCl	IV	0.5-0.7 mg/kg	7	Hypertension	48	-12	3.2	-31	3.6	-14	10
15 Phentolamine HCl	IM	1 mg/kg	6	Hypertension	53	0	2.8	-21	3.2	0	11
16 Tolazoline HCl	IV	20-100 mg	24	Normotension	59	-12	1.7	+6	3.6	-3	12
17			6	Mitral stenosis	45	+22	2.0	-25			13
Other Hypotensive Drugs											
18 Alkavervir	IV	1 µg/kg/min	27	Hypertension	57	-12	3.2	-28	2.9	+17	14
19 Aminophylline	IV	0.5 g	10	Normal	59	-25	1.7	+24	3.8	0	15
20			16	Congestive heart failure	45	-33	2.6	+8	3.2	-18	16
21 Caffeine and sodium benzoate	IV	0.25-0.5 g	7	Hypertension	54	-13	2.7	+22	3.2	+9	17
22			9	Brain tumor	44	-13	2.1	+28	2.6	+8	18
23 Glyceryl trinitrate	SL	0.6 mg	6	Mitral stenosis	40	-8	2.5	+4	3.3	-3	13
24 Histamine	IV	0.7 mg	10	Normal	57	0	1.7	-30	3.2	+6	18
					29	+17	1.7	-17	3.7	0	19
								-22	2.3	+4	18
								+3	2.6	+23	20
								-17	3.4	-3	21

1/1 Measured by nitrous oxide method of Kety and Schmidt [42]. 1/2 ml/100 g/min. 1/3 Ratio of mean arterial blood pressure to cerebral blood flow 1/4 Or IM, 1 mg/kg.

- 11148 [40] Patterson, J. L., Jr., A. Heyman, and F. T. Nichols. 1950. J. Clin. Invest. 29:1327. [41] Sokoloff, L., R. L. Wechsler, R. Mangold, K. Balls, and S. S. Kety. 1953. Ibid. 32:202. [42] Kety, S. S., and C. F. Schmidt. 1948. Ibid. 27:484. [43] Sokoloff, L. 1959. Pharm. Rev., Balt. 1:11.

Part IV. RENAL AND HEPATIC BLOOD FLOW AND VASCULAR RESISTANCE: MAN, DOG

Values are averages for reported number of cases and represent per cent change from control. Route: IM = intramuscular, IS = intraspinal, IV = intravenous, PO = oral. (+) = increase, (-) = decrease, (0) = no significant change.

Drug	Route	Dose	Subjects		Blood Flow	Vascular Resistance ³		Reference		
			Animal	Condition		Renal ¹	Hepatic ²		Renal ¹	Hepatic ²
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)	(K)
Ganglion Blocking Drugs										
1 Azamethonium dibromide	IV	Hypotensive doses	Man	9	Normotension	-8		-12		1
2			Dog	8	Anesthetized	0		-23		2
3 Chlorisondamine dimethochloride	IV	0.3 mg/kg	Dog	16	Anesthetized	-4	-4	0	+	3
4 Hexamethonium bromide	IV, IM	Hypotensive doses	Man	9	Normotension	-21 ⁵		+10 ⁵		4
5			Man	9	Normotension	+17 ⁶		-55 ⁶		4
6			Man	17	Normotension		-33		+5	5
7			Man	6	Hypertension		-28			6
8			Man	5	Hypertension	-53				7
9			Man	8	Hypertension	-12				8
10			Man	7	Hypertension	+15		-28		9
11			Dog	20	Anesthetized	-11		-19		10
12			Dog		Anesthetized	-56 ⁴		+14		11
13 Mecamylamine HCl	IV	0.5-2 mg	Dog	17	Anesthetized	-21		0		12
14 Trimethaphan camphorsulfate	IV		Man	6	Normotension	-18		+33		13
15						4		+4		13
16								-36		2
17 Azapetine phosphate	IV									8
18 Hydergine	IV	0.6 mg	Man	6	Hypertension	-11				8
19			Dog	14	Anesthetized	+5		-8		14
20 Phenox benzamine HCl	IV	5-10 mg/kg	Dog	5	Anesthetized	+20		-50		15
21			Dog	8	Anesthetized	+57		-40		16
22			Dog	12	Anesthetized	+15		-32		17
Other Antihypertensive Drugs										
23 Hydralazine	IV	0.25-1 mg/kg	Man	12	Normotension	+38				18
24			Man	8	Hypertension	+37				8
25			Man	6	Hypertension		+34		-41	19
26			Man	7	Toxemia of pregnancy	+9				20
27			Dog	6	Anesthetized	-30				21
28 Protervatrine	IV	100-200 µg	Dog	17	Anesthetized		+58		-36	22
29			Man	15	Hypertension	-8				23
30			Man	3	Hypertension	-25				23
31			Man	8	Hypertension	-7				23
32 Reserpine	PO	2 mg daily for 3-5 mo	Man	15	Hypertension	0				24
33 Veratrine	IM	0.4-1.0 ml	Man	2	Hypertension	-	-	-	-	25
34			Dog	3	Anesthetized	-	-	-	-	26
Musculotropic Vasodilators										
35 Aminophylline	IV	0.5 g	Man	10	Normal	+30 ⁷				27
36			Man	6	Cardiac failure	+12 ⁷				27
37 Ethyl alcohol	IV	25-81 mg % blood level	Man	5	Peptic ulcer		+45			28

1/1 Measured by p-aminobiphenyl renal clearance and blood hematocrit [2]. 1/2 Measured by bromsulphalein hepatic clearance and blood hematocrit [22]. 1/3 Ratio of mean arterial blood pressure to blood flow.

1/4 Measured directly by insertion of rotameter in artery 1/5 Immediate. 1/6 3 hours later. 1/7 Temporary increase accompanied by prolonged diuretic action.

Part III: CEREBRAL BLOOD FLOW, VASCULAR RESISTANCE, AND O₂ UPTAKE: MAN (Continued)

Drug	Route	Dose	Subjects		Cerebral Blood Flow ¹		Cerebral Vascular Resistance		Cerebral O ₂ Uptake		Reference
			No.	Condition	Control ²	% Change	Control ²	% Change	Control ²	% Change	
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)	(K)	(L)
Miscellaneous Drugs (concluded)											
56 Penicillin, plus fever therapy			6	Meningovascular syphilis	39	+18	2.6	-11	2.5	+10	40
57 Sodium bicarbonate	IV	1.2% (1 L)	6	Normal	60	+30	1.6	-25	3.6	+8	33
58		3% (1 L)	10	Normal	52	+69	1.8	-39	3.6	-6	33
59 Sodium chloride	IV	0.9% (1 L)	6	Normal	60	0	1.7	-6	3.7	0	33
60		2% (1 L)	7	Normal	59	+12	1.5	-7	3.6	-8	33
61 Thiouracil iodine and surgery			11	Thyrotoxicosis	69	-13	1.6	+31	3.5	-14	41

/1/ Measured by nitrous oxide method of Kety and Schmidt [42]. /2/ ml/100 g/min. /3/ Ratio of mean arterial blood pressure to cerebral blood flow.

Contributors: Aviado, Domingo M., Jr., and Carl F. Schmidt

References: [1] Moyer, J. H., and G. Morris. 1954. J. Clin. Invest. 33:1081. [2] Bernsmeier, A., and K. Siemons. 1953. Schweiz. med. Wochr. 83:210. [3] Morris, G. C., J. H. Moyer, H. B. Snyder, and B. W. Haynes. 1953. Ann. Surg. 138:706. [4] Finnerty, F. A., L. Wilkin, and J. F. Fazekas. 1954. J. Clin. Invest. 33:1227. [5] Stone, H. H., T. N. MacKrell, and R. L. Wechsler. 1955. Anesthesiology 16:168. [6] Crumpton, C. W., G. G. Rowe, R. C. Capps, J. J. Whitmore, and Q. R. Murphy. 1955. Circulation, N. Y. 11:106. [7] Dewar, H. A., S. G. Owen, and A. R. Jenkins. 1953. Brit. M. J. 2:1017. [8] Hafkenschiel, J. H., C. W. Crumpton, and J. H. Moyer. 1950. J. Pharm. Exp. Ther. 98:144. [9] Hafkenschiel, J. H., C. W. Crumpton, J. H. Moyer, and W. A. Jeffers. 1950. J. Clin. Invest. 29:408. [10] Moyer, J. H., H. Snyder, and S. I. Miller. 1954. Am. J. M. Sc. 228:563. [11] Moyer, J. H., and C. Caplovitz. 1953. Am. Heart J. 45:602. [12] Scheinberg, P., I. Blackburn, and M. Rich. 1953. J. Clin. Invest. 32:125. [13] Dewar, H. A., S. G. Owen, and A. R. Jenkins. 1953. Lancet, Lond. 1:867. [14] Moyer, J. H., S. I. Miller, A. B. Tashnek, H. Snyder, and R. O. Bowman. 1953. Am. J. Med. 14:175. [15] Wechsler, R. L., L. M. Kleiss, and S. S. Kety. 1950. J. Clin. Invest. 29:28. [16] Moyer, J. H., S. I. Miller, A. B. Tashnek, and R. O. Bowman. 1952. Ibid 31:267. [17] Moyer, J. H., A. B. Tashnek, S. I. Miller, H. Snyder, and R. O. Bowman. 1952. Am. J. M. Sc. 24:377. [18] Shenkin, H. A. 1951. J. Appl. Physiol. 3:465. [19] Jayne, H. W., P. Scheinberg, M. Rich, and M. S. Belle. 1952. J. Clin. Invest. 31:111. [20] Moyer, J. H., and H. Snyder. 1954. Am. Heart J. 47:912. [21] Hafkenschiel, J. H., A. M. Sellers, G. A. King, and M. W. Thorner. 1955. Ann. N. York Acad. Sc. 61:78. [22] King, B. D., L. Sokoloff, and R. L. Wechsler. 1952. J. Clin. Invest. 31:273. [23] Sensenbach, W., L. Madison, and L. Ochs. 1953. Ibid 32:226. [24] Moyer, J. H., G. Morris, and H. Snyder. 1954. Circulation, N. Y. 10:265. [25] Fazekas, J. F., S. N. Albert, and R. W. Alman. 1955. Am. J. M. Sc. 230:128. [26] Hine, C. H., A. F. Schick, L. Margolis, T. M. Burbridge, and A. Simon. 1952. J. Pharm. Exp. Ther. 106:253. [27] Kety, S. S., R. B. Woodford, M. H. Harmel, F. A. Freyhan, K. E. Appel, and C. F. Schmidt. 1948. Am. J. Psychiat. 104:765. [28] Kety, S. S., B. D. King, S. M. Horvath, W. A. Jeffers, and J. H. Hafkenschiel. 1950. J. Clin. Invest. 29:402. [29] Harmel, M. H., J. H. Hafkenschiel, G. M. Austin, C. W. Crumpton, and S. S. Kety. 1949. Ibid 28:415. [30] Scheinberg, P. 1950. Am. J. Med. 8:139. [31] Schieve, J. F., and W. P. Wilson. 1953. Ibid 15:171. [32] Wechsler, R. L., R. B. Dripps, and S. S. Kety. 1951. Anesthesiology 12:308. [33] Schieve, J. F., and W. P. Wilson. 1953. J. Clin. Invest. 32:33. [34] Alman, R. W., and J. F. Fazekas. 1951. Arch. Neur. Psychiat., Chic. 65:680. [35] Scheinberg, P. 1951. Blood, N. Y. 6:213. [36] Bentinck, R. C., et al. 1951. J. Clin. Invest. 30:200. [37] Moyer, J. H., S. I. Miller, and H. Snyder. 1955. Ibid 34:121. [38] Talley, R. W. 1955. Am. J. M. Sc. 230:61. [39] Scheinberg, P. 1950. Circulation, N. Y.

III. HEMODYNAMIC EFFECTS OF DRUGS: MAMMALS (Continued)

Part III: CEREBRAL BLOOD FLOW, VASCULAR RESISTANCE, AND O₂ UPTAKE: MAN (Continued)

1148. [40] Patterson, J. L., Jr., A. Heyman, and F. T. Nichols. 1950. J. Clin. Invest. 29:1327. [41] Sokoloff, L., R. L. Wechsler, R. M. Magold, K. Balls, and S. S. Kety. 1953. Ibid. 32:202. [42] Kety, S. S., and C. F. Schmidt. 1948. Ibid. 27:434. [43] Sokoloff, L. 1959. Pharm. Rev., Balt. 11:1.

Part IV. RENAL AND HEPATIC BLOOD FLOW AND VASCULAR RESISTANCE. MAN. DOG

Values are averages for reported number of cases and represent per cent change from control. Route: IM = intramuscular, IS = intrasplenic, IV = intravenous, PO = oral. (+) = increase, (-) = decrease, (?) = no significant change

Drug	Route	Dose	Subjects		Blood Flow		Vascular Resistance		Reference
			Animal No.	Condition	Renal	Hepatic	Renal	Hepatic	
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(X)
Ganglion Blocking Drugs									
1. Azamethonium dibromide	IV	Hypertensive doses	Man 9	Normotension	-8		-12		1
2. Chlorisondamine dimethochloride	IV	0.3 mg/kg	Dog 18	Anesthetized	0		-25		2
3. Chlorisondamine dimethochloride	IV	0.3 mg/kg	Dog 10	Anesthetized	-4	-4	0	0	3
4. Hexamethonium bromide	IV, IM	Hypertensive doses	Man 9	Normotension	-21.5		+13.5		4
5. Hexamethonium bromide	IV, IM	Hypertensive doses	Man 9	Normotension	+17.5		-55.5		4
6. Hexamethonium bromide	IV, IM	Hypertensive doses	Man 17	Normotension		-23		+5	5
7. Hexamethonium bromide	IV, IM	Hypertensive doses	Man 8	Hypertension		-23			6
8. Hexamethonium bromide	IV, IM	Hypertensive doses	Man 3	Hypertension	-50				7
9. Hexamethonium bromide	IV, IM	Hypertensive doses	Man 8	Hypertension	-12				8
10. Hexamethonium bromide	IV, IM	Hypertensive doses	Man 7	Hypertension	+15		-2.8		9
11. Hexamethonium bromide	IV, IM	Hypertensive doses	Dog 20	Anesthetized	-11		-13		13
12. Hexamethonium bromide	IV, IM	Hypertensive doses	Dog 1	Anesthetized	-50.4		-10		1
13. Mecamylamine HCl	IV	0.3-2 mg	Dog 17	Anesthetized	-21		0		13
14. Trimethaphan camphorsulfate	IV	1-1.5 mg/min	Man 9	Normotension	-19		+12		13
15. Trimethaphan camphorsulfate	IV	1-1.5 mg/min	Man 18	Hypertension	-14		0		13
16. Trimethaphan camphorsulfate	IV	1-1.5 mg/min	Dog 19	Anesthetized	0		-10		1
Adrenergic Blocking Drugs									
17. Alapril phosphate	IV	3-10 mg	Man 8	Hypertension	-7				9
18. Dipyridine	IV	0.6 mg	Man 6	Hypertension	-11				8
19. Dipyridine	IV	0.6 mg	Dog 14	Anesthetized	+5		-9		14
20. Phenoxymethamine HCl	IV	5-10 mg/kg	Dog 5	Anesthetized	+20		-50		13
21. Phenoxymethamine HCl	IV	5-10 mg/kg	Dog 8	Anesthetized	+27		-40		10
22. Phenoxymethamine HCl	IV	5-10 mg/kg	Dog 12	Anesthetized	+15		-32		13
Other Antihypertensive Drugs									
23. Hydralazine	IV	0.25-1 mg/kg	Man 12	Normotension	+10				13
24. Hydralazine	IV	0.25-1 mg/kg	Man 8	Hypertension	+17				8
25. Hydralazine	IV	0.25-1 mg/kg	Man 6	Hypertension		+30		-41	15
26. Hydralazine	IV	0.25-1 mg/kg	Man 7	Toxemia of pregnancy	+3				23
27. Hydralazine	IV	0.25-1 mg/kg	Dog 6	Anesthetized	-10				1
28. Hydralazine	IV	0.25-1 mg/kg	Dog 17	Anesthetized		+1		+50	13
29. Proxeterazine	IV	100-200 µg	Man 15	Hypertension	-4				23
30. Proxeterazine	IV	100-200 µg	Man 13	Hypertension	-25				23
31. Reserpine	PO	2 mg daily for 3-5 mo	Man 8	Hypertension	-7				23
32. Reserpine	PO	2 mg daily for 3-5 mo	Man 13	Hypertension	0				24
33. Verapamil	IV	0.4-1.0 ml	Man 2	Hypertension	-	-	-	-	25
34. Verapamil	IV	0.4-1.0 ml	Dog 1	Anesthetized	-	-	-	-	25
Musculotropic Vasodilators									
35. Aminophylline	IV	0.5 g	Man 10	Normal	+10.1				27
36. Aminophylline	IV	0.5 g	Man 6	Cardiac failure	+27				27
37. Ethyl alcohol	IV	21-31 mg % blood level	Man 5	Peptic ulcer		+45			23

- (1) Measured by p-aminohippurate renal clearance and blood hematocrit. (2) Measured by transulphalain hepatic clearance and blood hematocrit. (23) Ratio of mean arterial blood pressure to blood flow. (4) Measured directly by insertion of cannulae in artery. (5) Immediate. (6) 1 hour later. (7) Temporary increase accompanied by prolonged diuretic action.

Part IV: RENAL AND HEPATIC BLOOD FLOW AND VASCULAR RESISTANCE: MAN, DOG (Continued)

Drug	Route	Dose	Subjects		Blood Flow	Vascular Resistance ³	Reference	
(A)	(B)							
38 Histamine	IV							
49					shock			
50								
51								
52					shock			
53 Metaraminol bitartrate	IV	15 mg	Man	8	Normal	-4	+56	36
59			Man	5	anesthesia High spinal anesthesia	-33	+7	44
60 Sodium chloride	IV	0.9% (1-3L)	Man	35	Normal	+4		45

/1/ Measured by p-aminohippurate renal clearance and blood hematocrit [2]. /2/ Measured by bromsulphalein hepatic clearance and blood hematocrit [22]. /3/ Ratio of mean arterial blood pressure to blood flow.

Contributors: Aviado, Domingo M., Jr., and Carl F. Schmidt

References: [1] Moyer, J. H., R. McConn, and G. C. Morris. 1955. *Anesthesiology* 16:355. [2] Moyer, J. H., and C. A. Handley. 1955. *J. Pharm. Exp. Ther.* 113:383. [3] Plummer, A. J., J. H. Trapold, J. A. Schneider, R. A. Maxwell, and A. E. Earl. 1955. *Ibid.* 115:172. [4] Moyer, J. H., G. Morris, and R. A. Seibert. 1955. *Surg. Gyn. Obst.* 100:27. [5] Reynolds, T. B., A. Paton, M. Freeman, F. Howard, and S. Sherlock. 1953. *J. Clin. Invest.* 32:793. [6] Freis, E. D., J. C. Rose, E. A. Partenope, T. F. Higgins, R. T. Kelley, H. W. Schnaper, and R. L. Johnson. 1953. *Ibid.* 32:1285. [7] Grob, D., W. R. Scarborough, A. A. Kattus, and H. G. Landford. 1953. *Circulation*, N. Y. 8:352. [8] Redisch, W., L. Wertheimer, C. Dellisle, and J. M. Steele. 1954. *Ibid.* 9:68. [9] Ford, R. V., J. H. Moyer, and C. L. Spurr. 1953. *J. Clin. Invest.* 32:1133. [10] Moyer, J. H., R. A. Huggins, C. A. Handley, and L. C. Mills. 1952. *J. Pharm. Exp. Ther.* 106:157. [11] Wakim, K. G. 1955. *Am. Heart J.* 50:435. [12] Moyer, J. H., R. Ford, E. Dennis, and C. A. Handley. 1955. *Proc. Soc. Exp. Biol.*, N. Y. 90:402. [13] Moyer, J. H., R. McConn, and R. A. Seibert. 1955. *Am. Heart J.* 49:360. [14] Moyer, J. H., C. A. Handley, and R. A. Seibert. 1955. *Am. J. Physiol.* 180:146. [15] Moyer, J. H., C. A. Handley, and R. A. Huggins. 1954. *Circul. Res.*, N. Y. 2:441. [16] Moyer, J. H., and C. A. Handley. 1952. *J. Pharm. Exp. Ther.* 104:329. [17] Handley, C. A., and J. H. Moyer. 1954. *Ibid.* 110:277. [18] Wilkinson, E. L., H. Backmann, and H. H. Hecht. 1952. *J. Clin. Invest.* 31:872. [19] Freis, E. D., J. C. Rose, T. F. Higgins, F. A. Finnerty, R. T. Kelley, and E. A. Partenope. 1953. *Circulation*, N. Y. 8:199. [20] Kaplan, S. A., and N. S. Assali. 1953. *Surg. Gyn. Obst.* 97:501. [21] Moyer, J. H., R. A. Huggins, and C. A. Handley. 1953. *J. Pharm. Exp. Ther.* 109:175.

- [22] Marks, P. A., P. C. Reynell, and S. E. Bradley. 1955. *Am. J. Physiol.* 183:144. [23] Mellman, E. 1953. *J. Clin. Invest.* 32:80. [24] Andreae, E., and F. E. Smith. 1955. *Am. J. M. Sc.* 230:45. [25] Freis, E. D., J. R. Stanton, J. W. Culbertson, J. Litter, M. H. Halperin, C. H. Burnett, and R. W. Wilkins. 1949. *J. Clin. Invest.* 28:353. [26] Mongessor, W. C., and E. P. Hatt. 1954. *J. Pharm. Exp. Ther.* 110:68. [27] Davis, J. O., and N. W. Shock. 1949. *J. Clin. Invest.* 28:1459. [28] Mendeloff, A. I. 1954. *Ibid.* 33:1298. [29] Blackmore, W. P., V. E. Wilson, and T. R. Sherrod. 1953. *J. Pharm. Exp. Ther.* 109:206. [30] Livesay, W. R., and J. H. Moyer. 1953. *Ibid.* 109:123. [31] Bearn, A. G., B. Billing, and S. Sherlock. 1951. *J. Physiol., Lond.* 115:430. [32] Jacobson, W. E., J. F. Hammersten, and B. I. Heller. 1951. *J. Clin. Invest.* 30:1503. [33] Moyer, J. H., and C. A. Handley. 1952. *Circulation, N. Y.* 5:91. [34] Kaplan, S. A., S. J. Foman, and S. Rappaport. 1952. *Am. J. Physiol.* 169:588. [35] Houck, C. R. 1951. *Ibid.* 166:649. [36] Livesay, W. R., J. H. Moyer, and D. W. Chapman. 1954. *Am. Heart J.* 47:745. [37] Moyer, J. H., G. Morris, and H. L. Beasley. 1955. *Circulation, N. Y.* 12:96. [38] Sampson, J. J., and A. Zipser. 1954. *Ibid.* 9:38. [39] Smythe, C. M., J. P. Gilmore, and S. W. Handford. 1954. *J. Pharm. Exp. Ther.* 110:398. [40] Farber, S. J., J. D. Alexander, E. D. Pellegrino, and D. P. Earle. 1951. *Circulation, N. Y.* 4:376. [41] Eichna, L. W., et al. 1951. *J. Clin. Invest.* 30:1250. [42] Heller, D. I., J. F. Hammersten, and F. L. Stutzman. 1953. *Ibid.* 32:858. [43] Smythe, C. M., and J. P. Gilmore. 1955. *J. Pharm. Exp. Ther.* 114:221. [44] Mueller, R. P., R. B. Lynn, and S. M. Sancetta. 1952. *Circulation, N. Y.* 6:694. [45] Crawford, B., and H. Ludemann. 1951. *J. Clin. Invest.* 30:1456.

Part V. PULMONARY ARTERIAL BLOOD PRESSURE, BLOOD FLOW, VASCULAR RESISTANCE, AND LOCAL EFFECT ON LUNG VESSELS: MAN, DOG

Values are averages for reported number of cases and represent per cent change from control. Route of administration is intravenous unless otherwise indicated. (+) = increase, (-) = decrease, (+) = increase or decrease, (0) = no significant change. Discrepancies between column (H) and column (I) are due to the fact that column (II) includes active changes in resistance caused by local action of drug plus passive changes caused by alteration in blood flow in man, whereas local effect on lung vessels was elicited from perfusion, at a constant flow, of dog or cat lung.

Drug	Dose	Subjects			Pulmonary Arterial Pressure ¹	Pulmonary Blood Flow ²	Pulmonary Vascular Resistance ³	Local Effect on Lung Vessels	Reference ⁴
		Animal	No.	Condition					
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)
Ganglion Blocking Drugs									
1 Hexamethonium bromide or chloride	15-50 mg	Man	10	Normal	-17	0	+21	None	1;2
2		Man	4	Mitral stenosis	-35	0	-35		3
3		Man	8	Pulmonary hypertension	-14	-36	+4		4
4 Tetraethylammonium chloride or bromide	2-6 mg/kg	Man	5	Normal	-3	0	-3	None	5,2
5		Man	6	Mitral stenosis	-26	0	0		6
6		Man	5	Mitral stenosis	-26	+32	-35		7
7		Man	5	Left heart failure	-20	0	-20		8
8		Man	10	Cardiopulmonary diseases	-19	-32	-14		5
Adrenergic Blocking Drugs									
9 Dibenzamine HCl	0.5-2 mg/kg	Man	4	Mitral stenosis	-27	+7	-34	Unknown	7
10		Man	4	Left heart failure	-21	-6	-17		7
11 Dihydroergotamine methanesulfonate	1 mg	Man	10	Normal	+28	0	+41	None	7,2
12		Man	8	Pulmonary diseases	+38	0	+45		9
13 Hydergine	0.3-0.5 mg	Man	9	Cardiopulmonary diseases	0	0	0	None	9;2
14 Tolazoline HCl	10-50 mg	Man	8	Pulmonary hypertension	-	-	-		10

1;2 also used by methamphetamine

...; semicolon pertains to

111. HEMODYNAMIC EFFECTS OF DRUGS: MAMMALS (Continued)

Part V: PULMONARY ARTERIAL BLOOD PRESSURE, BLOOD FLOW, VASCULAR RESISTANCE, AND LOCAL EFFECT ON LUNG VESSELS: MAN, DOG (Continued)

Drug	Dose	Subjects			Pulmonary Arterial Pressure ¹	Pulmonary Blood Flow ²	Pulmonary Vascular Resistance ³	Local Effect on Lung Vessels	Reference ⁴
		Animal	No.	Condition					
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)
Sympathomimetic Amines									
15 Epinephrine	0.5-0.7 mg ⁵	Man	13	Normal	+20	+30	+	Constrictor	11;12-16
16 Hydroxyamphetamine hydrobromide	2-20 mg	Dog	4	Unanesthetized	+			Variable	17;18
17 Isoproterenol HCl	0.25% ⁶	Man	18	Bronchospasm	-			Dilator	19;20
18 Levaterenol bitartrate	0.2-0.4 µg/kg/min	Man	13	Cardiopulmonary diseases	+33	-6	0	Constrictor	21;16, 22
19 Mephentermine sulfate	5-20 mg	Man	4	Normal	+	0	+	Dilator	23;18
20 Metaraminol bitartrate	0.65 mg	Man	7	Normal	+95	0	+	Constrictor	24;16
21 Methoxamine HCl	0.1 mg/kg	Dog	10	Anesthetized	-	-	-	Variable	25;16
22 Phenylephrine HCl	0.1-0.6 mg	Dog	14	Anesthetized	+70	-	+	Constrictor	26;16
Cardiac Glycosides									
23 Digitalis, various preparations		Man	6	Low output failure	-40	+46	-		27
24 Digoxin	1-1.5 mg	Man	2	Pulmonary hypertension	+4	+7		None	28;29
25		Man	5	Cor pulmonale, not in failure	0	+15			28
26		Man	3	Cor pulmonale, in failure	+7	0			28
27		Man	5	Left heart failure	-	+25			30
28		Man	5	Congestive heart failure	-12	+29			31
29		Man	13	Systemic hypertension, compensated	-12	+33	-37		34
32		Man	4	Systemic hypertension, decompensated	-12	+33	-37		34
33		Dog	3	Unanesthetized	+				17
34 Procaine amide HCl	1 g	Man	10	Heart disease	-	-11		None	36;2
35 Procaine HCl ⁷		Man	5	Low spinal anesthesia	-17	-16			37
36		Man	5	High spinal anesthesia	-34	-31			37
37 Sodium nitrite	40-60 mg	Man	4	Mitral stenosis	-10	+11	-19	Variable	7;2
38		Man	5	Left heart failure	-18	+10	-20		7
39 Veratrine	0.4-1.0 ml ⁵	Man	2	Systemic hypertension, compensated	-10			None	38;2

/1/ Measured by catheter registering via an electrical manometer or mercury manometer /2/ Measured by indirect methods, such as Fick principle or dye dilution. /3/ Ratio of pressure gradient in pulmonary circulation to blood flow; pressure gradient calculated by assuming pulmonary venous pressure unchanged at zero, or by using wedged arterial pressure as pulmonary venous pressure. /4/ Reference numbers following semicolon pertain to column (I) only. /5/ Intramuscular. /6/ Inhalation. /7/ Intraspinal.

III. HEMODYNAMIC EFFECTS OF DRUGS: MAMMALS (Continued)

PART V: PULMONARY ARTERIAL BLOOD PRESSURE, BLOOD FLOW, VASCULAR RESISTANCE, AND LOCAL EFFECT ON LUNG VESSELS. MAN, DOG (Continued)

Drug	Dose	Subjects		Pulmonary Arterial Pressure ¹	Blood Flow ²	Pulmonary Vascular Resistance ³	Local Effect on Lung Vessels	Reference ⁴
		Animal	No.	Condition	(F)	(G)	(H)	(I)
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)
Miscellaneous Drugs. Predominantly Pulmonary Hypertensive (concluded)								
40 Veratrine (concluded)	0.4-1.0 ml ⁵	Man	3	Systemic hypertension, decompensated	-29			34
Miscellaneous Drugs. Not Pulmonary Hypertensive								
41 Acetylcholine chloride	25-150 µg	Dog	3	Unanesthetized	+			Constructive 17, 39
42 Amyl nitrite		Dog	4	Unanesthetized	0			Variable 17, 39
43 Dextran	0.4 (500 ml in 20 min)	Man	5	Normal	+52	+36		42
44 Eserine phosphate	0.3-2 mg	Dog	3	Unanesthetized	+			Constructive 17, 41; 39
45 Erythrane	0.2-0.5 mg/kg	Man	4	Normal	+45	+115	+27	42
46		Man	12	Mitral stenosis	+62	+33	+3	43
47		Man	11	Mitral stenosis	+19	+25		44
48		Man	17	Systemic hypertension	-9	+11	-24	45
49		Man	8	Systemic hypertension	+	+		46
50		Man	11	Systemic hypertension	0	+	-	47
51		Man	3	Toxemia of pregnancy	0	+31	-62	48
52 Kevlon	100-200 mg ⁵	Man	5	Pulmonary hypertension	-4	-6		Note 49, 50
53		Man	6	Chronic pulmonary diseases	0	0	0	50
54 Prazosin HCl	32-64 mg	Dog	4	Unanesthetized	+			Variable 17, 40
55 Prazosin	0.1 unit/kg	Dog	4	Unanesthetized	+			Variable 17, 40
56 pituitary extract		Dog	12	Anesthetized	+	-54	+	51
57 Serpentin	0.5 mg	Dog	25	Anesthetized	+	+	+50	Constructive 52, 53
58 Sodium chloride	0.5% (1 L in 12 min)	Man	12	Normal	+10	+19		54

^{1/1} Measured by catheter registering via an electrical manometer or mercury manometer. ^{1/2} Measured by indirect methods, such as Fick principle or dye dilution. ^{1/3} Ratio of pressure gradient in pulmonary circulation to blood flow; pressure gradient calculated by assuming pulmonary venous pressure unchanged at zero, or by using wedged arterial pressure as pulmonary venous pressure. ^{1/4} Reference numbers following semicolon pertain to column (I) only. ^{1/5} Intramuscular.

Contributors: Ariado, Domingos M., Jr., and Carl F. Schmidt

- References: [1] Bakita, L., and S. M. Sarnetia. 1953. *Circul. Res.*, N. Y. 1:479. [2] Ariado, D. M., Jr. Unpublished. [3] Davies, L. G., J. F. Goodwin, and B. D. Van Leuven. 1954. *Brit. Heart J.* 16:447. [4] Sarnetia, S. M. 1955. *Am. Heart J.* 49:501. [5] Fowler, N. O., R. S. Westcott, T. D. Bawtstein, R. C. Scott, and J. McGure. 1950. *J. Clin. Invest.* 29:1337. [6] Scott, R. C., S. Kaplan, and W. J. Stiles. 1955. *Am. Heart J.* 50:722. [7] Halmagyi, D., B. Felhai, J. Tranyi, T. Zoster, M. Tenyl, and Zs. Szlos. 1953. *Brit. Heart J.* 15:15. [8] Ford, R. V., J. H. Moyer, and C. L. Spurr. 1953. *J. Clin. Invest.* 32:1131. [9] Halmagyi, D., J. Tranyi, B. Felhai, T. Zoster, M. Tenyl, and Zs. Szlos. 1953. *Scand. J. Clin. & Invest.*, Oslo 5:15. [10] Gardner, J. M. 1954. *Australas. Am. M.* 3:59. [11] Witham, A. C., and J. W. Fleming. 1951. *J. Clin. Invest.* 30:727. [12] Baxter, L. G., and J. W. Pearce. 1951. *J. Physiol., Lond.* 115:415. [13] Edwards, W. S. 1951. *Am. J. Physiol.* 157:756. [14] Goldham, J. H., and P. Bodz. 1933. *J. Physiol., Lond.* 77:137. [15] Miller, G. S. L. Wernberg, and A. Pick. 1952. *Circulation*, N. Y. 6:41. [16] Tyner, J. H. 1953. *Q. J. Exp. Physiol., Lond.* 38:164.

111. HEMODYNAMIC EFFECTS OF DRUGS: MAMMALS (Continued)

Part V: PULMONARY ARTERIAL BLOOD PRESSURE, BLOOD FLOW, VASCULAR RESISTANCE, AND LOCAL EFFECT ON LUNG VESSELS: MAN, DOG (Continued)

Drug	Dose	Subjects			Pulmonary Arterial Pressure ¹	Pulmonary Blood Flow ²	Pulmonary Vascular Resistance ³	Local Effect on Lung Vessels	Reference ⁴
		Animal	No.	Condition					
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)
Sympathomimetic Amines									
15 Epinephrine	0.5-0.7 mg ⁵	Man	13	Normal	+20	+30	±	Constrictor	11;12-16
16 Hydroxymphetamine hydrobromide	2-20 mg	Dog	4	Unanesthetized	+			Variable	17,18
17 Isoproterenol HCl	0.25% ⁶	Man	18	Bronchospasm	-			Dilator	19,20
18 Levarterenol bitartrate	0.2-0.4 µg/kg/min	Man	13	Cardiopulmonary diseases	+33	-6	0	Constrictor	21;16, 22
19 Mephentermine sulfate	5-20 mg	Man	4	Normal	+	0	+	Dilator	23;18
20 Metaraminol bitartrate	0.65 mg	Man	7	Normal	+95	0	+	Constrictor	24;18
21 Methoxamine HCl	0.1 mg/kg	Dog	10	Anesthetized	-	-	-	Variable	25;18
22 Phenylephrine HCl	0.1-0.6 mg	Dog	14	Anesthetized	+70	-	+	Constrictor	26;18
Cardiac Glycosides									
23 Digitalis, various preparations		Man	6	Low output failure	-40	+46	-		27
24 Digoxin	1-1.5 mg	Man	2	Pulmonary hypertension	+4	+7		None	28,29
25		Man	5	Cor pulmonale, not in failure	0	+15			28
26		Man	3	Cor pulmonale, in failure	+7	0			28
27		Man	5	Left heart failure	-	+25			30
28		Man	5	Congestive heart failure	-12	+29			31
29		Man	12	Enlarged heart	-7	+9			32
30 Lanatoside C	1.4 mg	Man	6	Mitral stenosis	±	+	-	None	33;29
Miscellaneous Drugs, Predominantly Pulmonary Hypotensive									
31 Aminophylline	0.5 g	Man	7	Systemic hypertension, compensated	-25	+6	+4	Dilator	34;35
32		Man	4	Systemic hypertension, decompensated	-12	+33	-37		34
33		Dog	3	Unanesthetized	+				17
34 Procaine amide HCl	1 g	Man	10	Heart disease	-	-11		None	36,2
35 Procaine HCl ⁷		Man	5	Low spinal anesthesia	-17	-16			37
36		Man	5	High spinal anesthesia	-34	-31			37
37 Sodium nitrite	40-60 mg	Man	4	Mitral stenosis	-10	+11	-19	Variable	7;2
38		Man	5	Left heart failure	-18	+10	-20		7
39 Veratrine	0.4-1.0 ml ⁵	Man	2	Systemic hypertension, compensated	-10			None	38;2

/1/ Measured by catheter registering via an electrical manometer or mercury manometer. /2/ Measured by indirect methods, such as Fick principle or dye dilution. /3/ Ratio of pressure gradient in pulmonary circulation to blood flow; pressure gradient calculated by assuming pulmonary venous pressure unchanged at zero, or by using wedged arterial pressure as pulmonary venous pressure. /4/ Reference numbers following semicolon pertain to column (I) only. /5/ Intramuscular. /6/ Inhalation. /7/ Intraspinal.

111. HEMODYNAMIC EFFECTS OF DRUGS: MAMMALS (Continued)

Part V. PULMONARY ARTERIAL BLOOD PRESSURE, BLOOD FLOW, VASCULAR RESISTANCE, AND LOCAL EFFECT ON LUNG VESSELS MAN, DOG (Continued)

Drug	Dose	Subjects			Pulmonary Arterial Pressure ¹	Pulmonary Blood Flow ²	Pulmonary Vascular Resistance ³	Local Effect on Lung Vessels	Reference ⁴
		Animal	No.	Condition					
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)
Miscellaneous Drugs. Predominantly Pulmonary Hypotensive (concluded)									
40 Veratropine (concluded)	0.4-1.0 ml ⁵	Man	3	Systemic hypertension, decompensated	-29				38
Miscellaneous Drugs, Not Pulmonary Hypotensive									
41 Acetylcholine chloride	25-250 µg	Dog	3	Unanesthetized	+			Constrictor	17,2, 39
42 Amyl nitrite		Dog	4	Unanesthetized	0			Variable	17,29
43 Dextran	6% (500 ml in 20 min)	Man	5	Normal	+50	+36			40
44 Histamine phosphate	0.3-2 mg	Dog	3	Unanesthetized	+			Constrictor	17,41, 39
45 Hydralazine	0.2-0.5 mg/kg	Man	4	Normal	+45	+110	+27		42
46		Man	10	Mitral stenosis	+40	+38	+3		43
47		Man	11	Mitral stenosis	+16	+25			44
48		Man	17	Systemic hypertension	-9	+33	-28		45
49		Man	8	Systemic hypertension	+	+			46
50		Man	11	Systemic hypertension	0	+	-		47
51		Man	3	Toxemia of pregnancy	0	+31	-60		48
52 Aethin	100-200 mg ⁵	Man	5	Pulmonary hypertension	-4	-6		None	49,2
53		Man	6	Chronic pulmonary disease	0	0	0		50
54 Papaverine HCl	32-64 mg	Dog	4	Unanesthetized	+			Variable	17,29
55 Posterior pituitary extract	0.1 unit/kg	Dog	4	Unanesthetized	+			Variable	17,2
56		Dog	10	Anesthetized	+	-54	+		51
57 Serotonin	0.5 mg	Dog	25	Anesthetized	+	+	+50	Constrictor	52,53
58 Sodium chloride 0.9% (1 L in 10 min)		Man	12	Normal	+66	+19			54

1/1 Measured by catheter registering via an electrical manometer or mercury manometer. 1/2 Measured by indirect methods, such as Fick principle or dye dilution. 1/3 Ratio of pressure gradient in pulmonary circulation to blood flow; pressure gradient calculated by assuming pulmonary venous pressure unchanged at zero, or by using wedged arterial pressure as pulmonary venous pressure. 1/4 Reference numbers following semicolon pertain to column (I) only. 1/5 Intramuscular

Contributors. Aviado, Domingo M., Jr. and Carl F. Schmidt

- References [1] Rakita, L., and S. M. Sancetta. 1953. *Circul. Res.*, N. Y. 1:499. [2] Aviado, D. M., Jr. Unpublished. [3] Davies, L. G., J. F. Goodwin, and B. D. Van Leuven. 1954. *Brit. Heart J.* 16:440. [4] Sancetta, S. M. 1955. *Am. Heart J.* 49:501. [5] Fowler, N. Q., R. N. Westcott, V. D. Hauenstein, R. C. Scott, and J. McGulre. 1950. *J. Clin. Invest.* 29:1387. [6] Scott, R. C., S. Kaplan, and W. J. Stiles. 1955. *Am. Heart J.* 50:720. [7] Halmagyi, G., H. Feik, J. Ivanyi, T. Zsoter, M. Tenyi, and Zs. Szűcs. 1953. *Brit. Heart J.* 15:15. [8] Ford, R. V., J. H. Moyer, and C. L. Spurr. 1953. *J. Clin. Invest.* 32:1133. [9] Halmagyi, D., J. Ivanyi, B. Felai, T. Zsoter, M. Tenyi, and Zs. Szűcs. 1953. *Scand. J. Clin. & Invest.*, Oslo 5:85. [10] Gardiner, J. M. 1954. *Australas. Ann. M.* 3:59. [11] Witham, A. C., and J. W. Fleming. 1951. *J. Clin. Invest.* 30:707. [12] Baxter, L. G., and J. W. Pearce. 1951. *J. Physiol.*, Lond. 115:410. [13] Edwards, W. S. 1951. *Am. J. Physiol.* 167:756. [14] Gaddum, J. H., and P. Holtz. 1933. *J. Physiol.*, Lond. 77:139. [15] Miller, G., S. L. Weinberg, and A. Pick. 1952. *Circulation*, N. Y. 6:41. [16] Tyrer, J. H. 1953. *Q. J. Exp. Physiol.*, Lond. 38:169.

- [1] Friedberg, E. D., L. N. Katz, and F. S. Steinitz. 1943. *J. Pharm. Exp. Ther.* 78:80. [18] Aviado, D. M., Jr., C. F. Schmidt. 1957. *Ibid.* 120:512. [19] Buhr, G. 1953. *Zschr. Kreislaufforsch.* 42:669. [20] Hebb, C. O., H. Konzett. 1949. *J. Pharm. Exp. Ther.* 96:228. [21] Forman, S., L. G. May, A. Bennett, M. Kobayashi, and Gregory. 1953. *Proc. Soc. Exp. Biol.*, N. Y. 83:847. [22] Konzett, H., and C. O. Hebb. 1949. *Arch. internat. pharm. dyn.*, Gand 78:210. [23] Brofman, B. L., H. K. Hellerstein, and W. H. Coskey. 1952. *Am. Heart J.* 44:396. [24] Livesay, W. R., J. H. Moyer, and D. W. Chapman. 1954. *Ibid.* 47:745. [25] Wnuck, L. Unpublished. [26] Horvath, S. M., and D. W. Knapp. 1954. *Am. J. Physiol.* 178:387. [27] Eichna, L. W., et al. 1953. *Circulation*, N. Y. 7:674. [28] Ferrer, M. I., R. M. Harvey, R. T. Cathcart, C. A. Webster, D. W. Richards, Jr., A. Courmand. 1950. *Ibid.* 1:161. [29] Matzen, R. Unpublished. [30] Harvey, R. M., M. I. Ferrer, R. T. Cathcart, D. W. Richards, Jr., and A. Courmand. 1949. *Am. J. Med.* 7:439. [31] Eichna, L. W., et al. 1951. *J. Clin. Invest.* 30:1250. [32] Harvey, R. M., M. I. Ferrer, R. T. Cathcart, and J. K. Alexander. 1951. *Circulation*, N. Y. 4:369. [33] Gray, F. D., and F. G. Gray. 1954. *Am. Heart J.* 47:282. [34] Werkö, L., and H. Lagerlöf. 1950. *Scand. J. Clin. & Invest.*, Oslo 2:181. [35] Quimby, C. W., Jr., D. M. Aviado, Jr., and C. F. Schmidt. 1958. *J. Pharm. Exp. Ther.* 122:396. [36] McClendon, R. L., W. R. Hansen, and J. M. Kinsman. 1951. *Am. J. Sc.* 222:375. [37] Pritchard, W. H., W. J. MacIntyre, W. C. Schmidt, B. L. Brofman, and D. J. Moore. 1952. *Circulation*, N. Y. 6:572. [38] Freis, E. D., J. R. Stanton, J. W. Culbertson, J. Litter, M. H. Halperin, C. H. Bennett, and R. W. Wilkins. 1949. *J. Clin. Invest.* 28:353. [39] Daly, J. deBurgh. 1933. *Physiol. Rev.* 13:149. [40] Witham, A. C., J. W. Fleming, and W. L. Bloom. 1951. *J. Clin. Invest.* 30:897. [41] Woodbury, R. A., and F. Hamilton. 1941. *J. Pharm. Exp. Ther.* 71:293. [42] Wilkinson, E. L., H. Backmann, and H. H. Hecht. 1952. *Clin. Invest.* 31:872. [43] Aitcheson, J. D., W. I. Cranston, and E. A. Priest. 1955. *Brit. Heart J.* 17:425. [44] Werkö, L., E. Varnauskas, J. Ek, H. Bucht, B. Thomasson, J. Bergstrom, and H. Eliasson. 1954. *Circulation*, N. Y. 9:700. [45] Rowe, G. G., J. H. Huston, G. M. Maxwell, A. P. Crosley, and C. W. Crumpton. 1955. *J. Clin. Invest.* 34:115. [46] Stein, D. H., and H. H. Hecht. 1955. *Ibid.* 34:867. [47] Rowe, G. G., J. H. Huston, G. M. Maxwell, A. B. Weinstein, H. Tuchman, and C. W. Crumpton. 1955. *Ibid.* 34:696. [48] Assali, S., S. Kaplan, S. Olghenstein, and R. Suyemoto. 1953. *Ibid.* 32:922. [49] Cash, H. R., and H. A. Zimmerman. 1952. *Dis. Chest* 21:137. [50] Silber, E. N., G. L. Snider, H. Goldberg, L. N. Katz, and D. B. Radner. 1951. *J. Clin. Invest.* 30:1046. [51] Wakim, K. G., G. Denton, and H. E. Essex. 1954. *Am. Heart J.* 47:77. [52] MacCannon, D. M., and S. M. Horvath. 1954. *Am. J. Physiol.* 179:131. [53] Ginzler, K. H., and S. R. Heggstad. 1953. *Q. J. Exp. Physiol.*, Lond 38:225. [54] Doyle, J. T., J. S. Wilson, E. H. Estes, and J. V. Brennan. 1951. *J. Clin. Invest.* 30:345.

Part VI: LIMB BLOOD FLOW AND LOCAL EFFECT ON LIMB VESSELS: MAN, DOG

Unless otherwise specified, data are for man. Route: IM = intramuscular, IS = intraspinal, IV = intravenous, PO = oral, SC = subcutaneous. Local effect on limb vessels elicited by direct arterial injection into intact limb or into perfused limb.

Drug	Route	Dose	Subjects		Blood Flow ¹	Local Effect on Limb Vessels	Reference
			No.	Condition			
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)
Ganglion Blocking Drugs							
Azamethonium dibromide	IV	0.6-1.3 mg/kg	10	Peripheral vascular disease	Increased	None	1
Chlorisondamine dimethochloride	PO	100 mg	4	Hypertension	Increased		2
Hexamethonium bromide or chloride	IV	5-100 mg	22	Hypertension	Increased	None	3-6
			10	Heart failure	Not increased		7

¹ Determined by skin temperature of fingers or toes, or by limb plethysmography.

111. HEMODYNAMIC EFFECTS OF DRUGS. MAMMALS (Continued) Part VI. LIMB BLOOD FLOW AND LOCAL EFFECT ON LIMB VESSELS MAN, DOG (Continued)

Drug	Route	Dose	Subjects		Blood Flow ¹	Local Effect on Limb Vessels	Reference
			No.	Condition			
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)
Ganglion Blocking Drugs (concluded)							
5 Pentamethonium chloride	IV	0.3-0.7 mg/kg	14	Peripheral vascular disease	Increased	None	8
6 Tetraethylammonium bromide or chloride	IV	200-500 mg	14	Normal	Increased	Variable	9
			11	Peripheral vascular disease	Increased		10
Adrenergic Blocking Drugs							
8 Azapetine phosphate	IV	5-10 mg	8	Hypertension	Increased		5
9 Dihydroergocornine methane-sulfonate	IV	0.25-1 mg	34	Normal	Increased	Weak constrictor	11, 12
			6	Hypertension	Increased		13
			20	Various diseases	Increased		14
12 Hydergine	IV	0.6 mg	8	Hypertension	Increased	Weak constrictor	5
13 Phenoxybenzamine HCl	PO	20-480 mg daily	16	Peripheral vascular disease	Increased		15, 16
14 Phentolamine HCl	IM	0.6-0.75 mg/kg	5	Normal	Increased	Dilator	17
			4	Peripheral vascular disease	Increased		17
			13	Peripheral organic occlusion	Increased		17
17 Ronidacil tartrate	IV	30-50 mg	30	Pregnant	Unchanged		18
18 Tolazoline HCl	IM, IV	30-50 mg	10	Normal	Increased	Dilator	17, 19, 20
			16	Peripheral vascular disease	Increased		21
Other Vasodilators							
20 Atropine sulfate	IM	2 mg	73	Normal	Increased	Dilator	22
21 Ethyl alcohol	PO		5	Peripheral vascular disease	Increased	Dilator	10
22 Histamine diphosphate	Femoral arterial injection		8	Peripheral vascular disease		Dilator	23
23 Hydralazine	IV	0.25-0.5 mg/kg	12	Normotension	Unchanged	Weak dilator	24
			14	Hypertension	Decreased		5, 25
25 Isoproterenol HCl	IV	2-20 µg/min	9	Normal	Increased	Dilator	26, 27
26 Methacholine chloride	SC	3-5 mg	3	Peripheral vascular disease	Unchanged	Dilator	10
27 Nicotinic acid	PO	50-100 mg	8	Peripheral vascular disease	Decreased	Dilator	10
28 Pentobarbital sodium	IV	35 mg/kg	8 ²	Anesthetized	Increased ³	Dilator	28
29 Tubocurarine chloride	Femoral arterial injection		28 ²	Anesthetized		Dilator	29, 30
Miscellaneous Drugs							
30 Epinephrine	IV	10 µg/min	22	Normal	Increased	Constrictor and dilator ⁴	31-34
31 Levatterenol bitartrate	IV	10-25 µg/min	8	Normal	Unchanged	Constrictor	26, 34

^{1/1} Determined by skin temperature of fingers or toes, or by limb plethysmography. ^{2/2} Dogs. ^{3/3} Measured by appropriate flowmeters attached to vessels of hind limbs. ^{4/4} Dependent on dosage and on nature of reactive vascular bed, i.e., skin or muscle.

Part VI: LIMB BLOOD FLOW AND LOCAL EFFECT ON LIMB VESSELS: MAN, DOG (Concluded)

Drug	Route	Dose	Subjects		Blood Flow ¹	Local Effect on Limb Vessels	Reference
			No.	Condition			
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)
Miscellaneous Drugs (concluded)							
32 Procaine HCl	IV	250-500 mg	9	Normal	Decreased	Dilator	35
33			9	Peripheral vascular disease	Variable		35
34	IS	250-500 mg	5	Low spinal anesthesia	Decreased ⁵		36
35			5	High spinal anesthesia	Increased ⁵		36
36 Protoveratrine	IV	100-170 µg	25	Hypertension	Increased	None	37
37			8	Hypertension	Unchanged		5
38 Veratrone	IM	0.4-1.0 ml	2	Hypertension	Decreased	None	38

/1/ Determined by skin temperature of fingers or toes, or by limb plethysmography. /5/ Hand blood flow.

Contributors: Aviado, Domingo M., Jr., and Carl F. Schmidt

References: [1] Moyer, J. H., and C. A. Handley. 1955. *J. Pharm. Exp. Ther.* 113:383. [2] Grimson, K. S., A. K. Tarazi, and J. W. Frazer. 1955. *Circulation*, N. Y. 11:733. [3] Freis, E. D., J. C. Rose, E. A. Partenope, T. F. Higgins, R. T. Kelley, H. W. Schnaper, and R. L. Johnson. 1953. *J. Clin. Invest.* 32:1285. [4] Restall, P. A., and F. H. Smirk. 1952. *Brit. Heart J.* 14:1. [5] Redisch, W., L. Wertheimer, C. Delisle, and J. M. Steele. 1954. *Circulation*, N. Y. 9:68. [6] Schnaper, H. W., R. L. Johnson, E. B. Tuohy, and E. D. Freis. 1951. *J. Clin. Invest.* 30:786. [7] Shuman, C. R., N. Learner, and J. H. Doane. 1954. *Am. Heart J.* 47:737. [8] Needleman, H. L., and O. Horwitz. 1953. *Am. J. M. Sc.* 226:164. [9] Hoobler, S. W., S. D. Malton, H. T. Ballantine, S. Cohen, R. B. Neligh, M. M. Peet, and R. H. Lyons. 1949. *J. Clin. Invest.* 28:638. [10] Horwitz, O., H. Montgomery, E. D. Longaker, and A. Sayen. 1949. *Am. J. M. Sc.* 218:669. [11] Barcroft, H., H. Konzett, and H. J. C. Swann. 1951. *J. Physiol., Lond.* 112:273. [12] Bluntschli, H. J., and R. H. Goetz. 1948. *Am. Heart J.* 35:873. [13] Freis, E. D., J. R. Stanton, J. Litter, J. W. Culbertson, M. H. Halperin, F. C. Moister, and R. W. Wilkins. 1949. *J. Clin. Invest.* 28:1387. [14] Hayes, D. W., K. G. Wakim, B. T. Horton, and G. A. Peters. 1949. *Ibid.* 28:615. [15] Wertheimer, L., W. Redisch, and J. M. Steele. 1954. *Circulation*, N. Y. 10:366. [16] Redisch, W., E. C. Tester, R. M. Howard, P. H. Stillman, and J. M. Steele. 1952. *Ibid.* 6:352. [17] Clarke, C. W., D. R. Hays, and T. B. Van Itallie. 1953. *Ibid.* 8:715. [18] Lynch, P. R., D. L. Webber, and M. J. Oppenheimer. 1955. *Anesthesiology* 16:632. [19] Green, H. D., W. K. Gobel, M. J. Moore, and T. C. Prince. 1952. *Circulation*, N. Y. 6:520. [20] Van Itallie, T. B., and C. W. Clarke. 1951. *Ibid.* 3:820. [21] Rottenstein, H., O. Horwitz, H. Montgomery, A. Sayen, and L. L. Siems. 1951. *Am. J. M. Sc.* 221:661. [22] Kaiser, M. H., C. W. Frye, and A. S. Gordon. 1954. *Circulation*, N. Y. 10:413. [23] Dixon, J. A., W. J. M. Scott, and M. A. Epstein. 1952. *Ibid.* 5:661. [24] Wilkinson, E. L., H. Backmann, and H. R. Hecht. 1952. *J. Clin. Invest.* 31:872. [25] Werkö, L., E. Varnauskas, J. Ek, H. Bucht, B. Thomasson, J. Bergstrom, and H. Ellasch. 1954. *Circulation*, N. Y. 9:700. [26] Barcroft, H., and H. Konzett. 1949. *J. Physiol., Lond.* 110:194. [27] Walters, P. A., T. W. Cooper, A. B. Denison, and H. D. Green. 1955. *J. Pharm. Exp. Ther.* 115:323. [28] Imig, C. J., F. Randall, and H. M. Hines. 1953. *Proc. Soc. Exp. Biol.*, N. Y. 82:9. [29] Bean, J. W., and L. H. Elwell. 1951. *Am. J. Physiol.* 164:734. [30] Elwell, L. H., and J. W. Bean. 1953. *Ibid.* 174:185. [31] Allen, W. J., H. Barcroft, and O. G. Edholm. 1946. *J. Physiol., Lond.* 105:255. [32] Lanier, J. T., H. D. Green, J. Hardaway, H. D. Johnson, and W. B. Donald. 1953. *Circul. Res.*, N. Y. 1:40. [33] Swan, H. J. C. 1951. *J. Physiol., Lond.* 112:426. [34] Youmans, P. L., H. D. Green, and A. B. Denison. 1955. *Circul. Res.*, N. Y. 3:171. [35] Jessar, R. A., O. Horwitz, and H. Montgomery. 1952. *Am. J. M. Sc.* 224:300. [36] Pritchard, W. H., W. J. MacIntyre, W. C. Schmidt, B. L. Brofman, and D. J. Moore. 1952. *Circulation*, N. Y. 6:572. [37] Currens, J. H., J. F. McGinty, R. B. Khambatta, and L. Gordon. 1955. *Ibid.* 11:440. [38] Freis, E. D., J. R. Stanton, J. W. Culbertson, J. Litter, M. H. Halperin, C. H. Burnett, and R. W. Wilkins. 1949. *J. Clin. Invest.* 28:353.

112. EFFECT OF DRUGS ON CARDIAC PROPERTIES

oses are not indicated and by consulting the diastolic rhythm result of mechanical, etc. required after effective contraction for the muscle. Measurements made either on animal or on human heart tissue. Conduction time (atrial or ventricular muscle): time required for an impulse to travel from one point to another. Conduction time (atrial or ventricular muscle strips): time required for an impulse to travel from one point to another. Rate (sino-atrial node): measured by ECG of intact animals. Contractility (total heart): measured by the bundle of His). A-V system = atrioventricular conduction system (atrial muscle, ventricular muscle).

Drug (A)	Location (B)	Property (C)	Effect (D)	Reference (E)
Antiarrhythmic or Antifibrillatory Drugs Clinically Useful				
1 Ambonestyl HCl	Ventricular muscle	Ectopic automaticity	Depressed	1
2		Refractory period	Prolonged	1
3 Procaine amide HCl	Total heart	Contractility	Decreased	2
4	Sino-atrial node	Rate	Accelerated	1
5	Atrial muscle	Ectopic automaticity	Depressed	2
6		Refractory period	Prolonged	3, 4
7	A-V system	Conduction time	Prolonged	2
8	Ventricular muscle	Automaticity	Depressed	1, 4
9		Ectopic automaticity	Depressed	2
10		Refractory period	Unchanged	3, 4
11		Conduction time	Prolonged	2
12 Procaine HCl	Total heart	Contractility	Decreased	2
13	Sino-atrial node	Rate	Accelerated	2
14	Atrial muscle	Ectopic automaticity	Depressed	5
15	Ventricular muscle	Ectopic automaticity	Depressed	6, 7
16		Contractility	Decreased	2
17 Quinidine sulfate	Total heart	Contractility	Accelerated	2
18	Sino-atrial node	Rate	Depressed	8
19	Atrial muscle	Ectopic automaticity	Depressed	8-13
20		Refractory period	Prolonged	12, 13
21		Conduction time	Prolonged	2
22	A-V system	Conduction time	Shortened	2
23	Ventricular muscle	Automaticity	Depressed	13, 14
24		Ectopic automaticity	Depressed	11
25		Refractory period	Prolonged	8, 15
26 Chloroquin				16, 17
27				17
28				18
29				17
30				16
31 Dibenzylamine				19
32 Dihydroethanol methanesulfonate				
33 Diphenhydramine HCl	Atrial muscle	Refractory period	Prolonged	20, 21
34	Ventricular muscle	Ectopic automaticity	Depressed	22
35	Ventricular muscle	Ectopic automaticity	Depressed	23, 24
36 Diphenylhydantoin sodium	Atrial muscle	Ectopic automaticity	Depressed	5
37 a-Fagarine		Refractory period	Prolonged	5, 9
38		Conduction time	Prolonged	5, 25
39 Glyceryl trinitrate	Total heart	Contractility	Increased	26
40	Ventricular muscle	Ectopic automaticity	Depressed	27
41 Meperidine HCl	Ventricular muscle	Ectopic automaticity	Depressed	28
42 Papaverine HCl	Atrial muscle	Refractory period	Prolonged	9
43	A-V system	Conduction time	Prolonged	5
44	Ventricular muscle	Ectopic automaticity	Depressed	14
45 Phentolamine HCl	Ventricular muscle	Ectopic automaticity	Depressed	29
46 Quinacrine HCl	Atrial muscle	Ectopic automaticity	Depressed	16, 30
47	Ventricular muscle	Ectopic automaticity	Depressed	31
48 Tolazoline HCl	Ventricular muscle	Ectopic automaticity	Depressed	28

1/1 Vagolytic 1/2 Vagolytic, plus direct action on heart muscle.

112. EFFECT OF DRUGS ON CARDIAC PROPERTIES (Continued)

Drug (A)	Heart Component (B)	Property (C)	Effect (D)	Referen (E)
Heart Muscle Stimulants				
49 Aminophylline	Total heart	Contractility	Increased	2
50	Sino-atrial node	Rate	Accelerated	2
51 Amphetamine sulfate	Total heart	Contractility	Increased	32
52	Sino-atrial node	Rate	Accelerated	2
53 Atropine sulfate	Sino-atrial node	Rate	Accelerated ¹	2
54	Atrial muscle	Ectopic automaticity	Depressed ¹	33
55		Refractory period	Prolonged ¹	2
56		Conduction time	Prolonged ¹	2
57	A-V system	Conduction time	Shortened	2
58	Ventricular muscle	Refractory period	Prolonged	11
59		Ectopic automaticity	Depressed	11
60 Caffeine citrate	Total heart	Contractility	Increased	34
61	Sino-atrial node	Rate	Accelerated	2
62 Cardiac glycosides	Total heart	Contractility	Increased	34-37
63	Sino-atrial node	Rate	Decelerated	2
64	Atrial muscle	Refractory period	Shortened or prolonged ³	2
65		Conduction time	Shortened or prolonged ³	2
66	A-V system	Refractory period	Prolonged ³	2
67		Conduction time	Prolonged ³	2
68	Ventricular muscle	Automaticity	Increased	38, 39
69		Ectopic automaticity	Enhanced or depressed	2
70		Refractory period	Shortened	2
71		Conduction time	Prolonged	2
72 Ephedrine sulfate	Total heart	Contractility	Increased or decreased	34
73	Sino-atrial node	Rate	Accelerated	2
74 Epinephrine	Total heart	Contractility	Increased	2
75	Sino-atrial node	Rate	Decelerated ⁴ or accelerated	2
76	Atrial muscle	Refractory period	Shortened	40
77	A-V system	Conduction time	Shortened or prolonged ⁴	2
78	Ventricular muscle	Automaticity	Increased	41
79		Ectopic automaticity	Depressed or enhanced	14
80		Refractory period	Shortened	11
81 Isoproterenol HCl	Sino-atrial node	Rate	Accelerated	42, 43
82	Ventricular muscle	Automaticity	Increased	42, 44
83		Ectopic automaticity	Unchanged	45
84 Levarterenol bitartrate	Sino-atrial node	Rate	Decelerated ⁴ or accelerated	32, 46-48
85	Ventricular muscle	Automaticity	Unchanged	44, 49
86				44, 49
87 Mephentermine sulfate	Total heart			32
88 Methamphetamine HCl	Total heart			32
89	Ventricula			50
90 Phenylpropanolamine HCl	Total heart			32
91 Phenylpropylmethylamine	Total heart			32
92 Sodium lactate ⁵	Total heart			51
93	Ventricular			52
94 Tuaminoheptane sulfate	Total heart			32
95	Ventricular			53
96 Acetylcholine chloride	Sino-atrial		Decelerated or accelerated ⁶	54
97	Atrial muscle	Ectopic automaticity	Enhanced	55
98		Refractory period	Shortened	40, 56
99		Conduction time	Shortened	2
100	A-V system	Conduction time		2
101	Ventricular			11
102				11
103 Aconitine	Atrial muscle			57
104 Barium chloride	Total heart			34
105	Ventricular			58, 59
106 β -Dihydrosolasodine ⁷	Sino-atrial r			60

/1/ Vagolytic. /3/ Direct action, plus increased vagal tone. /4/ Deceleration due to reflex increase in vagal tone arising from increase in arterial pressure. /5/ Molar solution. /6/ Acceleration due to reflex inhibition of vagal tone arising from fall in arterial pressure. /7/ Antagonizes cardio-accelerator action of epinephrine.

112. EFFECT OF DRUGS ON CARDIAC PROPERTIES (Continued)

112. EFFECT OF DRUGS ON CARDIAC PROPERTIES (Continued)					Reference (E)
Drug (A)	Heart Component (B)	Property (C)	Effect (D)		
Miscellaneous Drugs (concluded)					
107 108 109	Calcium chloride	Total heart	Contractility	Increased	34
		A-V system	Conduction time	Prolonged	61, 62
		Ventricular muscle	Ectopic automaticity	Enhanced	63
		Ventricular muscle	Ectopic automaticity	Depressed	64
110 111 112 113			Contractility	Depressed	65
				Decelerated or accelerated	66
				Enhanced	55
				Shortened	40, 56
114 115 116 117			Conduction time	Shortened	2
				Prolonged	2
				Enhanced	11
				Enhanced	11
118 119 120 121				Enhanced	66
				Enhanced	67
				Enhanced	32
				Enhanced	68
122 123 124 125				Enhanced or enhanced	61, 69
				Enhanced	15
				Enhanced	70
				Enhanced	71
126 127 128 129				Enhanced	72
				Enhanced	73
				Enhanced	74
				Enhanced	75
130 131 132 133	Chloroform ^a	Total heart	Contractility	Enhanced	58
		Ventricular muscle	Ectopic automaticity	Enhanced	76
		Total heart	Contractility	Decreased	58
		Ventricular muscle	Ectopic automaticity	Enhanced	75-77
134 135 136 137	Ether	Total heart	Contractility	Decreased	76, 79
		Ventricular muscle	Ectopic automaticity	Decreased	76, 80
		Total heart	Contractility	Decreased	2
		Ventricular muscle	Contractility	Decreased	23
138 139 140 141	Phenobarbital sodium	Total heart	Ectopic automaticity	Decreased	76
		Ventricular muscle	Contractility	Decreased	79
		Total heart	Ectopic automaticity	Unchanged	81
		Ventricular muscle	Ectopic automaticity	Enhanced	82
142 143 144 145	Thiopental sodium	Ventricular muscle	Ectopic automaticity	Enhanced	82
		Ventricular muscle	Ectopic automaticity	Enhanced	75
		Ventricular muscle	Ectopic automaticity	Enhanced	75
		Ventricular muscle	Ectopic automaticity	Enhanced	75

cardio-accelerator action of epinephrine; 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299, 300, 301, 302, 303, 304, 305, 306, 307, 308, 309, 310, 311, 312, 313, 314, 315, 316, 317, 318, 319, 320, 321, 322, 323, 324, 325, 326, 327, 328, 329, 330, 331, 332, 333, 334, 335, 336, 337, 338, 339, 340, 341, 342, 343, 344, 345, 346, 347, 348, 349, 350, 351, 352, 353, 354, 355, 356, 357, 358, 359, 360, 361, 362, 363, 364, 365, 366, 367, 368, 369, 370, 371, 372, 373, 374, 375, 376, 377, 378, 379, 380, 381, 382, 383, 384, 385, 386, 387, 388, 389, 390, 391, 392, 393, 394, 395, 396, 397, 398, 399, 400, 401, 402, 403, 404, 405, 406, 407, 408, 409, 410, 411, 412, 413, 414, 415, 416, 417, 418, 419, 420, 421, 422, 423, 424, 425, 426, 427, 428, 429, 430, 431, 432, 433, 434, 435, 436, 437, 438, 439, 440, 441, 442, 443, 444, 445, 446, 447, 448, 449, 450, 451, 452, 453, 454, 455, 456, 457, 458, 459, 460, 461, 462, 463, 464, 465, 466, 467, 468, 469, 470, 471, 472, 473, 474, 475, 476, 477, 478, 479, 480, 481, 482, 483, 484, 485, 486, 487, 488, 489, 490, 491, 492, 493, 494, 495, 496, 497, 498, 499, 500, 501, 502, 503, 504, 505, 506, 507, 508, 509, 510, 511, 512, 513, 514, 515, 516, 517, 518, 519, 520, 521, 522, 523, 524, 525, 526, 527, 528, 529, 530, 531, 532, 533, 534, 535, 536, 537, 538, 539, 540, 541, 542, 543, 544, 545, 546, 547, 548, 549, 550, 551, 552, 553, 554, 555, 556, 557, 558, 559, 560, 561, 562, 563, 564, 565, 566, 567, 568, 569, 570, 571, 572, 573, 574, 575, 576, 577, 578, 579, 580, 581, 582, 583, 584, 585, 586, 587, 588, 589, 590, 591, 592, 593, 594, 595, 596, 597, 598, 599, 600, 601, 602, 603, 604, 605, 606, 607, 608, 609, 610, 611, 612, 613, 614, 615, 616, 617, 618, 619, 620, 621, 622, 623, 624, 625, 626, 627, 628, 629, 630, 631, 632, 633, 634, 635, 636, 637, 638, 639, 640, 641, 642, 643, 644, 645, 646, 647, 648, 649, 650, 651, 652, 653, 654, 655, 656, 657, 658, 659, 660, 661, 662, 663, 664, 665, 666, 667, 668, 669, 670, 671, 672, 673, 674, 675, 676, 677, 678, 679, 680, 681, 682, 683, 684, 685, 686, 687, 688, 689, 690, 691, 692, 693, 694, 695, 696, 697, 698, 699, 700, 701, 702, 703, 704, 705, 706, 707, 708, 709, 710, 711, 712, 713, 714, 715, 716, 717, 718, 719, 720, 721, 722, 723, 724, 725, 726, 727, 728, 729, 730, 731, 732, 733, 734, 735, 736, 737, 738, 739, 740, 741, 742, 743, 744, 745, 746, 747, 748, 749, 750, 751, 752, 753, 754, 755, 756, 757, 758, 759, 760, 761, 762, 763, 764, 765, 766, 767, 768, 769, 770, 771, 772, 773, 774, 775, 776, 777, 778, 779, 780, 781, 782, 783, 784, 785, 786, 787, 788, 789, 790, 791, 792, 793, 794, 795, 796, 797, 798, 799, 800, 801, 802, 803, 804, 805, 806, 807, 808, 809, 810, 811, 812, 813, 814, 815, 816, 817, 818, 819, 820, 821, 822, 823, 824, 825, 826, 827, 828, 829, 830, 831, 832, 833, 834, 835, 836, 837, 838, 839, 840, 841, 842, 843, 844, 845, 846, 847, 848, 849, 850, 851, 852, 853, 854, 855, 856, 857, 858, 859, 860, 861, 862, 863, 864, 865, 866, 867, 868, 869, 870, 871, 872, 873, 874, 875, 876, 877, 878, 879, 880, 881, 882, 883, 884, 885, 886, 887, 888, 889, 890, 891, 892, 893, 894, 895, 896, 897, 898, 899, 900, 901, 902, 903, 904, 905, 906, 907, 908, 909, 910, 911, 912, 913, 914, 915, 916, 917, 918, 919, 920, 921, 922, 923, 924, 925, 926, 927, 928, 929, 930, 931, 932, 933, 934, 935, 936, 937, 938, 939, 940, 941, 942, 943, 944, 945, 946, 947, 948, 949, 950, 951, 952, 953, 954, 955, 956, 957, 958, 959, 960, 961, 962, 963, 964, 965, 966, 967, 968, 969, 970, 971, 972, 973, 974, 975, 976, 977, 978, 979, 980, 981, 982, 983, 984, 985, 986, 987, 988, 989, 990, 991, 992, 993, 994, 995, 996, 997, 998, 999, 1000.

Contributors Aviado, Domingo M., Jr., and Carl F. Schmidt

- References [1] Lanzoni, V., and B. B. Clark 1955. Circul. Res., N. Y. 3 335. [2] Goodman, L. S., and A. Gilman. 1955 The pharmacological basis of therapeutics Macmillan, New York [3] Miller, G. S. L. Weinberg, and A. Pick 1952 Circulation, N. Y. 6:41 [4] Woske, H., J. Belford, F. N. Fastier, and C. McBrooks. 1953. J. Pharm. Exp. Ther. 107 134 [5] DiPalma, J. R., and J. E. Schulta. 1950 Medicine, Hall, 29 123. [6] Burstein, C. 1946 Anesthesiology 7 113 [7] Wiggers, C. J., and R. Wegria. 1940 Am. J. Physiol. 131 296. [8] Lewis, T. 1921 Heart, Lond 9 55 [9] Alles, G. A., and C. H. Ellis 1946. J. Pharm. Exp. Ther. 94 416 [10] Dawes, G. S. 1946 Brit J Pharm. 1 90. [11] DiPalma, J. R., and A. V. Mascarello 1951. J. Pharm. Exp. Ther. 101 243 [12] Prinzmetal, M., E. Corday, R. W. Oblath, H. E. Kruger, I. C. Brill, J. Fields, S. R. Kenemer, J. A. Osborne, L. A. Smith, A. L. Sellers, W. Flieg, and E. Pinston. 1951. Am J. Med. 11:410. [13] Wedd, A. M., H. A. Blair, and R. E. Gosselin. 1942 J Pharm. Exp Ther 75 251. [14] Wegria, R., and N. D. Nickerson 1942 Ibid 75 50 [15] Garb, S., and M. B. Chenoweth. 1949. Am. J. Physiol. 156 27. [16] Burno, F., F. Burnstein, and J. R. DiPalma. 1954. Circul Res., N. Y. 2:414. [17] Hess, M. E. Unpublished. [18] Sanabria, A. 1955. Southwest. M. 36:474. [19] White, J. M., M. H. Noltenmeyer, and L. E. Morris. 1951. Arch. internat pharm. dyn., Gand 88:361. [20] Dews, P. B., and J. D. P. Graham. 1946 Brit. J. Pharm. 1:278.

- [21] Dutta, N. K. 1949. *Ibid.* 4:281. [22] Levitan, B. A., and H. J. Scott. 1949. *Canad. M. Ass. J.* 61:306.
- [23] Harris, A. S., and R. H. Kokernot. 1950. *Am. J. Physiol.* 163:505. [24] Mosey, L., and M. D. Tyler. 1954. *Circulation*, N. Y. 10:65. [25] Alman, R. W., and J. F. Fazekas. 1951. *Arch. Neur. Psychiat.*, Chic. 65 680.
- [26] Goldberg, L. I., and R. P. Walton. 1952. *Arch. internat. pharm. dyn.*, Gand 89:399. [27] Melville, K. I. 1948. *J. Pharm. Exp. Ther.* 94:136. [28] Troch, E. E. 1953. *Arch. internat. pharm. dyn.*, Gand 94:175.
- [29] Morris, L. E., C. S. Yein, B. Hald, and J. M. White. 1952. *J. Pharm. Exp. Ther.* 106:49. [30] Gertler, M. M., and D. Karp. 1947. *Proc. Soc. Exp. Biol.*, N. Y. 64:213. [31] Melville, K. I. 1946. *J. Pharm. Exp. Ther.* 87:350. [32] Goldberg, L. I., M. deV. Cotten, T. D. Darby, and E. V. Howell. 1953. *Ibid.* 108:177. [33] Resnik, W. H. 1924. *J. Clin. Invest.* 1:181. [34] Walton, R. P., and O. J. Brodie. 1947. *J. Pharm. Exp. Ther.* 90:26.
- [35] Gruhzt, C. C., and A. E. Farah. 1953. *Ibid.* 108 112. [36] Masuoko, D. T., and P. R. Saunders. 1950. *Proc. Soc. Exp. Biol.*, N. Y. 74:879. [37] Walker, J. M., E. M. Lourie, and J. H. Burn. 1950. *Brit. J. Pharm.* 5:306. [38] Farah, A., and T. A. Loomis. 1950. *Circulation*, N. Y. 2:742. [39] Mendez, R., and C. Mendez. 1953. *J. Pharm. Exp. Ther.* 107:24. [40] De Ellis, F. J. 1947. *Brit. J. Pharm.* 2:131. [41] Carlen, S. A., and L. N. Katz. 1939. *Am. J. Physiol.* 127:272. [42] Garb, S. 1950. *Proc. Soc. Exp. Biol.*, N. Y. 73:134. [43] Lands, A. M., and J. W. Howard. 1952. *J. Pharm. Exp. Ther.* 106:65. [44] Nathanson, M. H., and H. Miller. 1952. *Circulation*, N. Y. 6:238. [45] Garb, S., and M. B. Chenoweth. 1948. *J. Pharm. Exp. Ther.* 94:12. [46] Baker, J. B. E. 1953. *J. Physiol.*, Lond. 120:122. [47] Hilton, J. G. 1955. *Am. J. Physiol.* 180:371. [48] Serin, F. 1952. *Acta physiol. scand.* 26:291. [49] Nathanson, M. H., and H. Miller. 1950. *Am. Heart J.* 40:374.
- [50] Lynch, P. R., D. L. Webber, and M. J. Oppenheimer. 1955. *Anesthesiology* 16:632. [51] DeGuzman, S. Unpublished. [52] Bellet, S., R. Wasserman, and J. I. Brody. 1955. *Circulation*, N. Y. 11:685. [53] Murphy, Q. G. S. O'Brien, and W. J. Meek. 1950. *Anesthesiology* 11 437. [54] Burgen, A. S. V., and K. G. Terroux. 1953. *J. Physiol.*, Lond. 120:449. [55] Hoff, H. E., and L. A. Geddes. 1955. *J. Appl. Physiol.* 8:177. [56] West, T. C., L. D. Turner, and T. A. Loomis. 1954. *J. Pharm. Exp. Ther.* 111:475. [57] Scherf, D. 1947. *Proc. Soc. Exp. Biol.*, N. Y. 64:233. [58] Dawes, G. S. 1952. *Pharm. Rev.*, Balt. 4 43. [59] Rothberger, C. J. 1939. *Q. J. Exp. Physiol.*, Lond. 29:69. [60] Krayner, O., and L. H. Briggs. 1950. *Brit. J. Pharm.* 5:118. [61] Wiggers, C. J. 1930. *Am. J. Physiol.* 93:107. [62] Wolffe, J. B., and S. Bellet. 1930. *Ann. Int. M.* 4:795. [63] Kay, J. H., and A. Blalock. 1951. *Surg. Gyn. Obst.* 93:97. [64] Bourgeois-Gavardin, M., W. K. Nowill, G. Margolis, and C. R. Stephen. 1955. *Anesthesiology* 16:829. [65] Harris, A. S., A. Estandia, H. T. Smith, R. W. Olsen, T. J. Ford, and R. F. Tillotson. 1953. *Am. J. Physiol.* 172 251. [66] West, J. W. Unpublished. [67] Lahti, R. E., I. C. Brill, and E. L. McCawley. 1955. *J. Pharm. Exp. Ther.* 115:268. [68] Deterling, R. A., S. H. Ngai, J. H. Laragh, and E. M. Papper. 1954. *Anesthesiology* 15:11. [69] Enselberg, S. D., H. G. Simmons, and A. A. Mintz. 1950. *Am. Heart J.* 39:713. [70] Tripod, J., and R. Meier. 1954. *Arch. internat. pharm. dyn.*, Gand 97:251.
- [71] Trapold, J. H., A. J. Plummer, and F. F. Yonkman. 1954. *J. Pharm. Exp. Ther.* 110:205. [72] Yonkman, F. F., A. J. Plummer, and W. B. Reitze. 1952. *Arch. internat. pharm. dyn.*, Gand 88:401. [73] Wynne, N. A., and P. Szekely. 1952. *Acta med. scand.* 143:129. [74] Krayner, O., and P. Ourlisson. 1954. *J. Pharm. Exp. Ther.* 112:341. [75] Boniface, K. J., J. M. Brown, and P. S. Kronen. 1955. *Ibid.* 113:64. [76] Price, H. L., and M. Helrich. 1955. *Ibid.* 115:206. [77] Bennett, L. L., and C. W. Fisher. 1948. *Ibid.* 93:482. [78] Greishelmer, E. M., M. J. Oppenheimer, D. W. Ellis, P. R. Lynch, and L. Shapiro. 1954. *Anesthesiology* 15:51. [79] Mosey, L., and J. W. Stutzman. 1950. *Proc. Soc. Exp. Biol.*, N. Y. 75:34. [80] Moe, G. K., B. R. Rennick, W. A. Freyburger, and S. D. Malton. 1949. *Anesthesiology* 10:706. [81] Morris, L. E., M. H. Noltensmeyer, and J. M. White. 1953. *Ibid.* 14:153. [82] Wedd, A. M., and H. A. Blair. 1948. *Proc. Soc. Exp. Biol.*, N. Y. 68:304.

DATA ARE NOT IN MAN, unless dog is specified. Normal serum concentration of sodium (Na^+) = 137-145 mEq/L [42, 43], normal serum sodium levels antidiuretic, and decreased sodium levels enhance, the deleterious effects of potassium on the electrocardiogram [9, 36, 37]. The combination of hypokalaemia and hypocalcaemia is difficult to recognize in the electrocardiogram [46]. The administration of either potassium or calcium corrects only the electrocardiographic abnormality due to the deficiency of that particular electrolyte, without affecting the changes due to the deficiencies of the other electrolyte (1, 51). The electrocardiogram is not a reliable tool for the measurement of electrolyte disturbance [46-48].

Specification	Effect of Serum Concentration of Inorganic Ions						Reference
	Calcium (Ca^{++})	Magnesium (Mg^{++})	Potassium (K^+)	Increased ions ^a	Decreased ions ^b	Reference	
	Increased (B)	Increased (D)	Increased (E)	Increased (F)	Decreased (G)		(H)
1 Systolic interval	Prolonged.	None		Shortened.	Normal or shortened.	B, 1, 2, C, 3, D, 4, F, 2, G, 1	
2 Diastolic interval	Shortened	In man, slight prolongation may occur (not of diagnostic value clinically), prolonged in dog		Prolonged.	P wave decrease in height, increase in duration; PR interval may be prolonged.	B, C, F, 2	
3 P wave and PR interval	Flattening or inversion of P wave, prolonged PR interval (transient)				PR interval may be prolonged; increase in amplitude and width of P wave.	B, 3, 6, C, 7, D, 4, F, 8-11, G, 12	
4 QRS duration	Normal.	Normal			May be wide, bizarre, diphasic; intraventricular block.	B, C, F, 7, D, 4; F, 8-10, 13; G, 7	
5 QT interval	Shortened due to shortening of ST segment (not of diagnostic value clinically).		Shortened in dog.		May be shortened, unchanged, or lengthened, later QT is prolonged.	B, 3, 5, 14, C, 1, 3, 14-16; 19, F, 20, 21, G, 1, 7, 10, 22	
6 ST segment	Shortened or absent.		Sagging.		Depressed in precordial leads; elevated in cavity leads and occasionally elevated in precordial leads.	B, 3, 5, 14, C, 7, 14, 16, E, 19, F, 20, 21, G, 1, 7, 10, 22	
7 T wave	Normal, flat, or inverted	Inconstant minor variations.	Low in standard and left precordial leads.		Tent-shaped; tall, narrow, steep, pointed.	B, 3, C, 1, 15, 23, D, 24, E, 19, F, 8, 9, 11, 13, 16, 25, 26, G, 1, 7, 22, 23, 27	

1/1 Normal = 4.5-5.5 mEq/L (9-11 mg %). [42, 43] 2/2 Normal = 1.5-2.0 mEq/L (3.5-4 mg %). [44, 45] 3/3 Normal = 3.5-5.0 mEq/L. 4/4 Electrocardiographic changes are estimated to occur in 80-100% of subjects with blood levels of potassium above 6.8 mEq/L [23, 46]. Serum potassium may be elevated without definite electrocardiographic evidence of potassium intoxication [25, 47]. 5/5 Electrocardiographic changes are estimated to occur in 80% of subjects with blood levels of potassium below 3.5 mEq/L, and in 100% of subjects with potassium levels below 2.5 mEq/L [23, 46]. Potassium deficiency may occur without recognizable electrocardiographic changes, or the pattern of potassium intoxication may appear [23, 47-50]. 6/6 Pre-existent abnormalities of the ST segment and T wave may modify abnormalities induced by electrolyte disturbance [25, 51, 52].

113. EFFECT OF INORGANIC IONS ON THE HEART: MAN, DOG (Concluded)

Specification	Effect of Serum Concentration of Inorganic Ions					Reference
	Calcium (Ca ⁺⁺) ¹	Magnesium (Mg ⁺⁺) ²	Potassium (K ⁺) ³			
(A)	Increased Ions (B)	Decreased Ions (C)	Increased Ions (D)	Decreased Ions (E)	Increased Ions ⁴ (F)	Decreased Ions ⁵ (G)
8 U wave	Increased height.	Decreased height.		Prominent, but not large.	Decreased height.	Increased height; may become tall and pointed; increase in amplitude in left precordial leads.
9 Rate and rhythm	Bradycardia; later tachycardia, sinus arrhythmia, atrioventricular conduction defects, ventricular tachycardia or fibrillation may occur following calcium infusions.	Tachycardia.	Tachycardia initially and then bradycardia in dog; slowing may occur in man; occasional sinoatrial and atrioventricular block.		First degree atrioventricular block, idioventricular rhythm; premature ventricular beats; ventricular tachycardia; ventricular and atrial fibrillation; sinus arrest; ventricular standstill.	Supra-ventricular tachycardia may occur; various arrhythmias may occur in subjects with low potassium.
10 Remarks	When serum calcium exceeds 6.5 mEq/L (13 mg %), the QTc interval and ST segment are always shortened; when serum calcium is 3.0 mEq/L (6 mg %) or less, the QTc interval and ST segment are always prolonged.	Normal serum calcium levels antagonize, and decreased calcium levels enhance, the deleterious effects of excess potassium on the electrocardiogram. ⁷	Cardiac arrest rarely precedes respiratory arrest, changes not diagnostic in man; inconsistent, transient variations in QRS amplitudes.	In dog, no consistent electrocardiographic changes attributed to magnesium depletion.	Disturbed depolarization, terminal slur in QRS or S wave in lead I or V6 is frequent in early hyperkalemia. Progressive hyperkalemia causes peaking of T waves (at about 6.5 mEq/L), then widening of "S-ST" angle, QRS broadening with delay in activation of base of right ventricle, disturbance of rhythm, and the "sine" wave of ventricular fibrillation.	B.14;C.25, 35-39;D, 24,31;E, 17;F.4,10, 20,40,41; G.7,10,22

1/1 Normal = 4.5-5.5 mEq/L (9-11 mg %). [42, 43]. 2/2 Normal = 1.5-2.0 mEq/L (1.8-2.4 mg %). [44, 45] 3/3 Normal = 3.5-5.0 mEq/L. 4/4 Electrocardiographic changes are estimated to occur in 80-100% of subjects with blood levels of potassium above 6.8 mEq/L [23, 46]. 5/5 Electrocardiographic changes are estimated to occur in 80% of subjects with blood levels of potassium below 3.5 mEq/L, and in 100% of subjects with potassium levels below 2.3 mEq/L [23, 46]. Potassium deficiency may occur without recognizable electrocardiographic changes, or the pattern of potassium intoxication may appear [23, 47-50]. 7/7 These findings have not been confirmed in recent studies [41].

- References* [1] Surawicz, B., and E. Lepeschkin 1953. *Circulation*, N. Y. 8:801. [2] Best, C. H., and N. D. Taylor 1950. *The physiological basis of medical practice*. Ed. 5. Williams and Wilkins, Baltimore. [3] Barker, P. S., F. D. Johnston, and P. M. Wilson. 1937. *Am. Heart J.* 14:62. [4] Smith, P. K., A. W. Winkler, and H. Hoff 1939. *Am. J. Physiol.* 128:720. [5] Kellogg, F., and W. J. Kerr. 1936. *Am. Heart J.* 12:346. [6] Berliner, K. 1936. *Am. J. M. Sc.* 191:117. [7] Ernsteins, A. C., and W. L. Proudfoot. 1949. *Am. Heart J.* 38:240. [8] Merrill, J. P., H. D. Levine, W. Somerville, and S. Smith 1950. *Ann. Int. M.* 33:797. [9] Keith, N. M., and H. B. Burchell 1949. *Am. J. M. Sc.* 217:1. [10] Burchell, H. B. 1952. *Electrocardiographic changes related to disturbances in potassium metabolism*. A symposium on the metabolism of potassium, University of Minnesota. [11] Levine, H. D. 1954. *Mod. Concepts Card. Dis.* 23:246. [12] Weller, J. M., B. Lowy, B. V. Holgers, N. P. Wyatt, M. Cristofalio, J. P. Merrill, and S. A. Levine. 1953. *Circulation*, N. Y. 11:44. [13] Tazari, R. 1948. *Am. J. Med.* 5:828. [14] Yu, P. N. G. 1952. *Am. J. M. Sc.* 224:413. [15] Barker, J. M. 1952. *The unipolar electrocardiogram*. Appleton-Century-Crofts, New York. [16] Bechler, J. T., J. E. White, and E. H. Estes, Jr. 1956. *Circulation*, N. Y. 13:837. [17] Danzig, L. E., and W. J. Walker 1955. *Clin. Res. Proc.* 3:106. [18] Langendorf, R., and C. L. Pirani. 1947. *Am. Heart J.* 33:282. [19] Agna, J., and R. Scott. Unpublished. [20] Levine, H. D., J. P. Merrill, and W. Somerville. 1951. *Circulation*, N. Y. 3:889. [21] Levine, H. D., S. H. Wanser, and J. P. Merrill 1952. *Am. Heart J.* 43:634. [22] Bernstein, M., and S. Simkins. 1939. *J. Laborat. Clin. M.* 25:131. [23] Levine, H. D., J. P. Vaziradar, B. Lowy, and J. P. Merrill 1952. *Am. Heart J.* 43:437. [24] Finch, C. A., G. C. Sawyer, and J. M. Flynn. 1946. *Am. J. Med.* 1:337. [27] Sjostrand, T. 1953. *Acta med. scand.* 151:73. [28] Lepeschkin, E. 1955. *Arch. Int. M.* 96:600. [29] Clark, N. E. 1941. *Am. Heart J.* 22:367. [30] Myers, G. B. 1956. *The interpretation of the unipolar electrocardiogram*. Mosby Company, St. Louis. [31] Smith, P. K., A. W. Winkler, and H. Hoff. 1939. *Am. J. Physiol. vectocardiography*. Ed. 3. Lea and Febiger, Philadelphia. [34] Kolff, W. J., D. B. Effter, L. K. Graves, G. Peersboom, S. Aoyama, and F. M. Sones, Jr. 1936. *Cleveland Clin. Q.* 23:98. [35] Butcher, W. A., H. B. Burchell, K. G. Wakim, and H. E. Essex. 1951. *Am. J. Physiol.* 157:773. [36] Butcher, W. A., K. G. Wakim, H. E. Essex, R. D. Pruitt, and H. B. Burchell. 1952. *Am. Heart J.* 43:801. [37] Meroney, W. H., and R. F. Herndon. 1954. *J. Am. M. Ass.* 155:877. [38] Winkler, A. W., H. E. Hoff, and P. K. Smith. 1939. *Am. J. Physiol.* 127:430. [39] Braun, H. A., R. Van Horne, J. C. Bettinger, and S. Bellet 1955. *J. Laborat. Clin. M.* 46:544. [40] Herndon, R. F., W. H. Meroney, and C. M. Pearson. 1955. *Am. Heart J.* 50:188. [41] Braun, H. A., B. Surawicz, and S. Bellet. 1955. *Am. J. M. Sc.* 230:147. [42] Ham, T. H., ed. 1950. *A syllabus of laboratory examination in clinical diagnosis*. Harvard University Press, Cambridge. [43] Sunderman, F. W., and F. Boerner. 1949. *Normal values in clinical medicine*. W. B. Saunders, Philadelphia. [44] Flinn, E. B., P. L. Suttman, A. R. Anderson, T. Konig, and R. Fraser. 1954. *J. Laborat. Clin. M.* 43:169. [45] Agna, J. Unpublished. [46] Dreifus, L. S., and A. Pick. 1956. *Circulation*, N. Y. 14:815. [47] Scott, R. C. Unpublished. [48] Schwartz, W. B., H. D. Levine, and A. S. Reiman. 1954. *Am. J. Med.* 16:395. [49] Bellet, S., and D. Finkelstein. 1951. *Am. J. M. Sc.* 222:263. [50] Currena, J. H., and J. D. Crawford. 1950. *N. England J. M.* 243:843. [51] Bellet, S. 1955. *Arch. Int. M.* 96:618. [52] Dodge, H. T., R. P. Grant, and P. W. Seavey. 1953. *Am. Heart J.* 45:755.

114. EFFECT OF INORGANIC IONS ON CARDIAC PROPERTIES

(+) = increased, (-) = decreased, (0) = no effect or unknown.

Cardiac Property	Effect of:							
	Increased / Decreased		Increased / Decreased		Increased / Decreased		Increased / Decreased	
	Calcium (Ca^{++})		Magnesium (Mg^{++})		Potassium (K^{+})		Sodium (Na^{+})	
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)
1 Automaticity	-	0	-	0	-	0	0	+
2 Conductivity	0	0	-	0	-	0	0	0
3 Contractility	+	0	-	0	0	0	0	-
4 Excitability	+	-	-	0	-	0	0	0
5 Relaxation	-	0	+	0	+	0	0	+
6 Rhythm	+	0	-	0	-	0	0	+

Contributor: Gertler, Menard M.

115. ION RATIO REQUIRED TO MAINTAIN BEAT OF ISOLATED HEART: INSECTS

Ion Ratio	Insect	Stage	Optimal Ratio	Reference
(A)	(B)	(C)	(D)	(E)
1 $\text{Na}^{+}:\text{K}^{+}$	Calliphora	Larva	8:33.3	1
2	Chortophaga	Nymph and adult	3:34	2
3	Samia	Pupa	0.07.34	2
4	Galleria	Larva	8:16	3
5	Gryllus		8:16	4
6	Periplaneta	Nymph and adult	3:30	5
7	Tenebrio	Larva	1.4:32.4	6
8		Pupa	2.2:8.8	6
9		Adult	0.8:14.2	6
10 $\text{K}^{+}:\text{Ca}^{++}$	Chortophaga	Nymph and adult	1:3	2
11	Samia	Pupa	1:3	2
12	Periplaneta	Nymph and adult	0.9:3.5	5
13	Tenebrio	Larva	1:3	6
14		Pupa	1:3	6
15		Adult	1:3	6

Contributor: Ludwig, Daniel

References. [1] Levy, R. 1928. C. rend. Soc. biol. 99:1485. [2] Barasa, M. C. 1954. J. Gen. Physiol. 38:79. [3] Drieux, P. 1950. C. rend. Soc. biol. 144:803. [4] Bergerard, J. 1947. Ibid. 141:1079. [5] Ludwig, D., Sr., K. M. Tracey, and M. L. Burns. 1957. Ann. Entom. Soc. America 50:244. [6] Butz, A. 1956. Anat. Rec. 125:554.

116. EFFECT OF CHEMICAL SUBSTANCES ON THE HEART: INSECTS

Effect differs with time, temperature, pretreatment, relation to other chemicals, concentration, type of preparation, type of pacemaker tissue, stage of development, age, nutritional status. In many cases, specific concentrations were not available. Consult references for details.

Part I. INORGANIC IONS

Ion	Insect ¹	Stage	Salt Concentration in Perfusion Fluid ²	Effect (E)	Reference (F)
(A)	(B)	(C)	(D)	(E)	(F)
1 Calcium	Anopheles ³	Larva	Excessive	No effect or stimulates	1
2	Chortophaga	Nymph and adult	Excessive	Inhibits	2
3	Cossus	Larva	Excessive	Stimulates	3
4	Culex ³	Larva	Excessive	Stimulates	4
5	Samia	Pupa	Excessive	No effect or inhibits	2
6	Galleria	Larva	Excessive	Inhibits	3
7	Galleria	Larva	Reduced	Stimulates	5
8	Galleria	Larva	Absent	Stoppage in diastole	5
9 Magnesium	Anopheles ³	Larva	Excessive	Inhibits	1
10	Chortophaga	Nymph and adult	Isotonic MgCl ₂	Inhibits	2
11	Culex ³	Larva	Excessive	Inhibits	4
12	Samia	Pupa	Isotonic MgCl ₂	Inhibits	2
13 Potassium	Anopheles ³	Larva	Excessive	Stimulates	1
14	Calliphora	Larva	Excessive	Stimulates	6
15	Calliphora	Larva	Reduced	Inhibits	6
16	Calliphora	Larva	Absent	Inhibits	6
17	Chortophaga	Nymph and adult	Isotonic KCl	Inhibits	2
18	Cossus	Larva	Excessive	Inhibits	3
19	Culex ³	Larva	Excessive	Stimulates	4
20	Samia	Pupa	Isotonic KCl	No effect	2
21	Galleria	Larva	Excessive	Stops in systole	5
22	Galleria	Larva	Excessive	Inhibits	7
23	Galleria	Larva	Reduced	Inhibits	5
24	Galleria	Larva	Absent	Stoppage in diastole	5
25	Cryllus	Nymph and adult	Excessive	Stoppage in systole	8
26	Phormia	Larva	Excessive	Stimulates	6
27	Phormia	Larva	Reduced	Inhibits	6
28	Phormia	Larva	Absent	Inhibits	6
29	Tenebrio	Larva, pupa, adult	Excessive	No effect	9
30 Sodium	Anopheles ³	Larva	Excessive	No effect or stimulates	1
31	Chortophaga	Nymph and adult	Isotonic NaCl	Inhibits	2
32	Cossus	Larva	Reduced	Inhibits	3
33	Culex ³	Larva	Excessive	Stimulates	4
34	Samia	Pupa	Isotonic NaCl	No immediate effect	2
35	Tenebrio	Larva, pupa, adult	Excessive	No effect	9
36	Tenebrio	Larva, pupa, adult	Reduced	No effect	9

¹/ Unless otherwise indicated, heart exposed by dissection, then perfused ²/ "Excessive" varies from 2-10 x isotonic concentration ³/ Intact specimen totally immersed in solution.

Contributor Jones, Jack Colvard

References. [1] Jones, J. C. 1956. J. Exp. Zool. 133:125. [2] Barra, M. C. 1954. J. Gen. Physiol. 38:79. [3] Clark, A. J. 1927. Comparative physiology of the heart. Macmillan, New York. p. 157. [4] Wixforth, E. 1924. Arch. Naturgesch. 90(AK5):193. [5] Dreux, P. 1950. C. rend. Soc. Biol. 144:804. [6] Levy, R. 1928. Ibid. 99 1482 [7] Millman, N. 1938. Some aspects of the physiology of the heart of *Galleria mellonella* (bee moth). M. S. Thesis. Brown University, Providence. p.52. [8] Fisser, J. 1950. C. rend. Soc. Biol. 144 815. [9] Butz, A. 1956. Anat. Rec. 125 554.

Part II: DRUGS AND TOXIC COMPOUNDS

	Substance (A)	Insect (B)	Stage (C)	Condition (D)	Effect (E)	Refer- ence (F)
1	Acetylcholine	Anopheles	Adult		No effect.	1, 2
2		Apis			Acceleration.	3
3		Blatta			Acceleration.	3
4		Galleria			No effect.	4
5		Melanoplus			Acceleration.	5
6		Periplaneta			Acceleration.	6, 7
7		Stenopelmatus			Acceleration.	8
8	Adrenalin	Anopheles	Larva		Slight stimulation.	1
9		Anopheles	Adult	Intact	No effect.	1
10		Cossus		Intact	Acceleration.	9
11		Culex		Intact	Acceleration.	10
12		Periplaneta			Acceleration.	6, 7
13		Stenopelmatus			Depression.	8
14	Aldrin	Periplaneta	Adult		Rate increased.	11
15	Anabasine sulfate	Blatta orientalis	Larva	Intact	Transient effects.	12
16		Pieris brassicae	Larva	Dissected	Instant arrest.	12
17		Pteronux	Larva	Intact	Submersion caused transient increase in rate, normal in 30 min. Rate increased fourfold when compound applied with brush; recovery after 4-hr paralysis.	12
18		Periplaneta	Adult	Intact	Marked decrease in amplitude, diastolic arrest.	13
19	Atropine	Anopheles	Larva		Depression.	2
20		Anopheles	Adult	Intact	No effect; did not block nicotinic effects.	1
21						
22						
23						
24						
25						
26	BHC (Benzene hexachloride)	Allomyrina dichotomus			nicotine effects.	15
27		Locusta migratoria danica	Larva	Dissected	No marked effect after prolonged perfusion.	16
28						
29						
30					diastolic arrest.	
31						18
32						0
33						
34						
35						
36		Stenopelmatus			effects. No marked effect, did not modify acetylcholine effects.	8
37	DDT (Dichlorodiphenyl-trichloroethane)	Anopheles	Larva	Intact	No effect prior to convulsions at concentration of 10^{-5} , eserine and atropine had no effect on rate.	18
38		Corethra	Larva	Partially dissected	Rate decreased with doses of 1:1000 to 1:10,000 in 1.5% NaCl emulsions.	19
39		Drosophila	Larva	Intact	Rate increased when DDT first injected, then returned to normal.	20

116. EFFECT OF CHEMICAL SUBSTANCES ON THE HEART: INSECTS (Continued)

Part II: DRUGS AND TOXIC COMPOUNDS (Continued)

	Substance	Insect	Stage	Condition	Effect	Reference
	(A)	(B)	(C)	(D)	(E)	(F)
40	DDT (Dichlorodiphenyl-trichloroethane) (concluded)	<i>Locusta migratoria danica</i>	Larva	Dissected	Rate greatly decreased and irregular with spasms of body and alary muscles; heart not stimulated by low concentration of acetylcholine after prolonged perfusions.	16
41		<i>Periplaneta</i>	Adult	Intact	Heart beat after extremities stopped twitching.	21
42		<i>Periplaneta</i>	Adult	Intact	No immediate effect; diastolic arrest.	17
43		<i>Periplaneta</i>	Adult	Intact	Rate decreased slightly.	22
44		<i>Periplaneta</i>	Adult	Dissected	No effect for 5-7 min at concentration of 10^{-5} , irregular beat later, previous atropine treatment had no effect	7
45		<i>Stenopelmatus</i>	Adult	Dissected	No effect in 5% emulsion, heart beat at reduced rate for hours in 100 ppm.	8
46	Dieldrin	<i>Anopheles</i>	Larva	Intact	No marked primary effects.	18
47		<i>Periplaneta</i>	Adult		Rate increased.	11
48	Digitalis	<i>Anopheles</i>	Adult			
49		<i>Culex</i>				
50		<i>Periplaneta</i>				
51	Diphtheria toxin	<i>Hyalophora</i>	Pupa			
52	DNOC (4,6-Dinitro-o-cresol), DNOCHP (Dinitro-o-cyclohexylphenol)	<i>Periplaneta</i>	Adult	Intact	massive dose. Rate increased significantly; when substance first injected, posterior heart showed temporary stoppage in diastole.	17
53	DNP (Dinitrophenylphthalate)	<i>Anopheles</i>	Larva	Intact	Rate increased greatly at concentrations of 10^{-2} to 10^{-3} ; no effect at concentrations of 10^{-4} to 10^{-5} .	18
54	EPN (Ethyl p-nitrophenyl thionobenzenephosphate)	<i>Anopheles</i>	Larva	Intact	Rate increased with larval agitation.	18
55	Ergotamine	<i>Periplaneta</i>			Stimulation, then depression.	7
56	Eserine	<i>Anopheles</i>			No marked effect.	1
57		<i>Culex</i>	Adult	Intact	Stimulation.	10
58		<i>Oncopeltus</i>			No effect.	24
59		<i>Periplaneta</i>			Stimulation.	7
60		<i>Stenopelmatus</i>			No marked effect, potentiated acetylcholine effects.	8
61	Hexamethonium	<i>Periplaneta</i>			No effect.	7
62	Lethane 60 (mixed aliphatic thiocyanates)	<i>Periplaneta</i>	Adult	Intact	Injection caused slight rate increase, followed by gradual decrease; diastolic arrest.	17
63	Lethane 384 (mixed aliphatic thiocyanates)	<i>Periplaneta</i>	Adult	Intact	3% dilution caused sharp decrease, irregular beat, death. 1.25% resulted in sharp decrease, rise above normal, recovery.	25
64	Lindane (1,2,3,4,5,6-Hexachlorocyclohexane)	<i>Periplaneta</i>			Irregular beat.	17
65	Lobeline	<i>Periplaneta</i>				
66	Mecholyl	<i>Anopheles</i>			Stimulation with low concentrations, depression with high concentrations.	6
67	Methoxychlor	<i>Anopheles</i>	Larva	Intact	No effect.	2
					No marked primary effects.	18

PART II: DRUGS AND TOXIC COMPOUNDS (Continued)

	Substance	Insect	Stage	Condition	Effect	Reference
	(A)	(B)	(C)	(D)	(E)	(F)
68	Morphine	Anopheles	Adult		Depression.	1
69		Periplaneta			Stimulation with low concentrations, depression with high concentrations.	6
70	Muscarine	Stenopelmatus			No effect.	8
71	Nicotine	Anopheles			No marked effect.	1
72		Culex		Intact	Stimulation with low concentrations, depression with high concentrations.	10
73		Culex		Intact	Depression.	26
74		Melanoplus			Rate temporarily increased, then decreased.	5
75		Periplaneta			Irregular decline.	25
76		Periplaneta			Stimulation with low concentrations, depression with high concentrations.	6, 7, 27
77		Prodenia			Initial stimulation followed by partial depression, extent depending on concentration.	27
78		Stenopelmatus			Acceleration.	8
79	Para-oxon	Periplaneta	Adult	Dissected	No effect at concentration of 3×10^{-6} , rate increased immediately then declined at concentration of 15×10^{-7} ; blocked by atropine.	7
80	Parathion	Corethra	Larva	Partially dissected	Rate increased.	19
81						17
82						17
83					Increased steadily with irradiated parathion; blocked atropine; potentiated acetylcholine effects.	7
84	Phenothiazine	Anopheles	Larva	Intact	No marked primary effects.	18
85		Corethra	Larva	Intact	Beats continued for 3 wk.	28
86	Pilocarpine	Anopheles			No marked effect.	1, 2
						6
						29
						23
						29
						30
					more than 5 da.	
92		Corethra	Larva	Intact	Rate decreased	28
93		Corethra	Larva	Intact	Normal during initial convulsions; rate decreased slightly after 1 da, decreased markedly after several da	31
94		Galleria	Larva	Dissected	Anterior heart stopped first in diastole	32
95		Lymantria	Larva	Intact	Weak beats continued more than 5 da	30
96		Periplaneta	Adult	Intact	Sublethal dose caused rate increase, lethal dose caused continuous decrease.	25
97		Periplaneta	Adult	Dissected	Prolonged rate increase at concentration of 10^{-6} , atropine abolished effect.	7

116. EFFECT OF CHEMICAL SUBSTANCES ON THE HEART: INSECTS (Continued)

Part II. DRUGS AND TOXIC COMPOUNDS (Continued)

	Substance	Insect	Stage	Condition	Effect	Reference
	(A)	(B)	(C)	(D)	(E)	(F)
98	Pyrethrum (concluded)	Vanessa	Larva	Intact	Weak beats continued for more than 5 da	30
99	Rotenone	Allomyrina dichotomus			Temporary increase in rate and decrease in amplitude; subsequent rate decrease and irregularities.	15
100		Anopheles	Larva	Intact	No marked primary effects.	18
101		Bombyx	Larva	Intact	Rate decreased 10-15 min before other symptoms appeared	33
102		Culex	Larva	Intact	Rate decreased.	33
103		Dendrolimus	Larva	Intact	Beats continued 6 da.	30
104		Periplaneta	Nymph and adult	Intact	Injection or contact caused steady rate decrease; anterior heart stopped first in diastole.	17, 33
105		Periplaneta	Nymph and adult	Intact	Heart stopped in 26-42 min.	22
106		Periplaneta	Nymph and adult	Dissected	Rate decreased to cessation in diastole.	7, 34
107		Plutella	Larva	Intact	Rate decreased.	33
108		Tomato cutworm	Larva	Intact	Rate decreased	33
109		Vanessa	Larva	Intact	Beats continued for 6 da.	30
110	Sodium arsenite and arsenate	Bombyx	Larva	Intact	Injection caused transient rate decrease; contact caused slow decrease.	35
111	Sodium fluoride	Anopheles	Larva	Intact	Transient rate increase, then decrease.	36
112	Sodium metarsenite	Anopheles	Larva	Intact	Rate decreased.	18
113	Strychnine	Culex		Intact	Stimulation.	10
114		Periplaneta			Stimulation with low concentrations, depression with high concentrations.	6
115	TEPP (Tetraethyl pyrophosphate)	Periplaneta	Adult	Intact	Beats occurred in prostrate roach.	36
116		Periplaneta	Adult	Dissected	0.02% dilution caused decreased rate, systolic arrest, concentration of 8×10^{-8} resulted in sharply increased rate and amplitude.	34
117	Thiocyanate	Blattia orientalis	Adult	Dissected	Dilatation from increased myotonus of alary muscles, rate decreased.	37
118	Toxaphene	Anopheles	Larva	Intact	No effect in 30 min at concentration of 10^{-4} .	18
119		Periplaneta	Adult	Intact	Rate irregular; stoppage in diastole.	17
120	Veratrine	Corethra			Slightly stimulating effects.	14

Contributors: (a) Jones, Jack Colvard, (b) Ludwig, Daniel, (c) Beard, Raimon L., (d) Dresden, D., and A. Glenn Richards, (e) Brown, A. W. A.

References: [1] Jones, J. C. 1956. J. Exp. Zool. 133:573. [2] Jones, J. C. 1954. J. Morph. 94:71. [3] Prosser, C. L. 1942. Biol. Bull. 83:145. [4] Millman, N. 1938. Some aspects of the physiology of the heart of Galleria mellonella (bee moth). M. S. Thesis. Brown University, Providence. p. 52. [5] Hamilton, H. L. 1939. J. Cellul. Physiol. 13:91. [6] Krijgsman, B. J., and N. E. Krijgsman-Berger. 1951. Bull. Entom. Res., Lond. 42:143. [7] Naidu, M. B. 1955. Ibid. 46:205. [8] Davenport, D. 1949. Physiol. Zool. 22:35. [9] Clark, A. J. 1927. Comparative physiology of the heart. Macmillan, New York. p. 157. [10] Wixforth, E. 1924. Arch. Naturgesch. 90(A):193. [11] Giannotti, O., R. L. Metcalf, and R. B. March. 1956. Ann. Entom. Soc. America 49:588. [12] Tarasova, K. L. 1936. Izv. Kuzn. prikl. Zool. 6:15. [13] Beck, S. D. 1950. J. Econom. Entom. 43:103. [14] Dogiel, J. 1877. Mém. Acad. Imp. Sci., S. Petersb. Ser. 7, 24:1. [15] Yamasaki, T., and T. Ishii. 1951. Oyo-Kontyu 7:58. [16] Yamasaki, T., and T. Ishii. 1950. Ibid. 5:155. [17] Orser, W. B., and A. W. A.

Part II: DRUGS AND TOXIC COMPOUNDS (Concluded)

Brown. 1951. *Canad. J. Zool.* 29:54. [18] Jones, J. C. 1957. *Mosquito News* 17:214. [19] Florey, E. 1951. *Pflüger* 7:81. [20] Bodenstein, D. 1946. *Biol. Bull.* 90:148. [21] Tobias, J. M., and J. J. Kollros. 1946. *Ibid.* 91:247. [22] Bellemare, E. R., and J. Belcourt. 1955. *Canad. J. Zool.* 33:175. [23] Pappenheimer, A. M., Jr., and C. M. Williams. 1952. *J. Gen. Physiol.* 35:727. [24] Beard, R. L. 1953. *Circulation*. In K. D. Roeder, ed. *Insect physiology*. J. Wiley and Sons, New York. [25] Coon, B. F. 1944. *J. Econom. Entom.* 37:785. [26] Richardson, C. H., and H. H. Shepard. 1930. *J. Agr. Res.* 41:337. [27] Yeager, J. F., and J. B. Gahan. 1937. *Ibid.* 55:1. [28] Deonier, C. C., and A. W. Lindquist. 1942. *J. Econom. Entom.* 35:188. [29] Harvey, W. R., and C. M. Williams. 1953. *Anat. Rec.* 117:544. [30] Klinger, H. 1936. *Arb. Physiol. angew. Entom.* 3:49. [31] Krüger, F. 1931. *Zachr. angew. Entom.* 18:344. [32] Bellevue, G. 1938. *Ann. physiol.*, Par. 14 717. [33] Tischler, N. 1935. *J. Econom. Entom.* 28:215. [34] Krilgsmann, B. J., D. Dresden, and N. E. Krilgsmann-Berger. 1950. *Bull. Entom. Res.*, Lond. 41:141. [35] Campbell, F. L. 1926. *J. Pharm. Exp. Ther.* 26:277. [36] Roan, C. C., H. E. Fernando, and C. W. Kearns. 1950. *J. Econom. Entom.* 43:319. [37] Yeager, J. F., A. Hager, and J. M. Straley. 1935. *Ann. Entom. Soc. America* 28:256.

Part III GASES AND VAPORS

All specimens intact.

Gas or Vapor	Insect	Stage	Effect	Reference
(A)	(B)	(C)	(D)	(E)
1 Acetone	<i>Macrosiphum tulipae</i>		Heart stopped in 10 min.	1
2 Benzene	<i>Macrosiphum tulipae</i>		Heart stopped in 8 min.	1
3 Carbon dioxide	<i>Anopheles</i>	Larva	100% concentration caused diastolic arrest in 30 sec-3 min; normal rate following anesthesia.	2
4	<i>Anopheles</i>	Larva	5% concentration stimulated heart rate.	2
5	<i>Corethra</i>	Larva	Irregularities.	3
6	<i>Graesshopper</i>		Beat continued for 6 hr; isolated heart stopped in 20-60 sec.	4
7	<i>Oncopeltus</i>		Heart stopped rapidly.	5
8 Carbon disulfide	<i>Macrosiphum tulipae</i>		Heart stopped in 4-7 min.	1
9 Carbon monoxide	<i>Corethra</i>	Larva	Brief stimulation.	3
10	<i>Graesshopper</i>		No recovery after 48-hr exposure.	4
11 Carbon tetrachloride	<i>Macrosiphum tulipae</i>		Heart stopped in 13 min.	1
12 Chloroform	<i>Macrosiphum tulipae</i>		Heart stopped in 5-10 min.	1
13 Cyanide	<i>Calliphora</i>	Larva	Heart rate declined rapidly following 10-min exposure.	6
14	<i>Periplaneta</i>		Heart rate declined sharply following 1-hr exposure, then rose above normal.	7
15 Ether	<i>Anopheles</i>	Larva	No effect after 20-min exposure to 1/2-saturated aqueous solution.	2
16	<i>Graesshopper</i>		No effect during 3-hr exposure; isolated heart stopped in 1 min.	4
17 Ethyl acetate plus chloroform (2:1 ratio)	<i>Macrosiphum tulipae</i>		Heart stopped in 11 min.	1
18 Formaldehyde, 40%	<i>Macrosiphum tulipae</i>		Heart stopped in 57 min.	1
19 Hydrogen	<i>Graesshopper</i>		Beat continued for 5 da, isolated heart beat up to 4 da.	4
20 Nitrobenzene	<i>Macrosiphum tulipae</i>		Heart stopped in 50 min.	1
21 Nitrogen	<i>Anopheles</i>	Larva	99% concentration caused rapid decline in 1 hr.	2
22 Oxygen	<i>Anopheles</i>	Larva	1-100% concentration had no effect or caused only slight stimulation.	2

Contributors (a) Jones, Jack Colvard, (b) Ludwig, Daniel

References. [1] Kirschner, R. 1932. *Zachr. angew. Entom.* 19:544. [2] Jones, J. C. 1956. *J. Exp. Zool.* 131:257. [3] Dogiel, J. 1877. *Mém. Acad. Imp. Sc.*, S. Petersb., Ser. 7, 24:1. [4] Walling, E. V. 1906. *J. Exp. Zool.* 3:621. [5] Beard, R. L. 1953. *Circulation*. In K. D. Roeder, ed. *Insect physiology*. J. Wiley and Sons, New York. [6] Myers, J. G. 1927. *Bull. Entom. Res.*, Lond. 17:219. [7] Coon, B. F. 1944. *J. Econom. Entom.* 37:785.

117. TRANSLOCATION OF ORGANIC SUBSTANCES PLANTS

Data are for substances translocated predominantly in the phloem. Translocation of antibiotics in plants has not been included, as much of the data indicates acropetal movement chiefly within the transpiration stream or xylem. Translocation rates, where given, do not necessarily represent maximum or even mean values for the conditions of the experiment, for in many cases they are simply based on the time interval between treatment and harvest or assay of specific plant parts. In most instances, where the rate is based on time of absorption from cuticular surface to phloem, the true translocation rate within the phloem will be greater than the figures given. It is known, of course, that many organic substances applied to a plant are broken down into different metabolic products within the plant. Recent research (made possible largely by chromatography) indicates, however, that at least a significant portion of most organic materials is translocated within the plant in the exact molecular form in which applied. For simplification of tabulation, the organic materials appearing in the substance column are listed as they were applied to the plant, i.e., before metabolic change. Direction pertains only to the given experiment, and may vary according to growth stage, concentration of translocated substance, and other conditions; A = acropetal, B = basipetal.

Part I ASSIMILATES AND CARBOHYDRATES

Plant	Part	Substance	Translocation		Remarks	Reference
			Rate cm/hr	Direction		
(A)	(B)	(C)	(D)	(E)	(F)	(G)
1 Ash, white (<i>Fraxinus americana</i>)	Trunk	Mannitol, raffinose, stachyose, sucrose, verbascose		B	Transport of specific substances determined by chromatography of sieve tube exudate obtained from cut in bark. Concentration gradient of individual sugars varied with height in tree, but overall molar gradient was zero after leaf abscission.	1, 2
2 Bean (<i>Phaseolus vulgaris</i>)	Leaf, stem	Carbohydrates		A ¹ , B ¹ , 2	Application of 2, 4-D to various loci along primary leaf midveins reduced carbohydrate and dry matter accumulation in aerial parts, as determined by analysis 9-29 da later. Application to leaf tips less effective than to leaf center or pulvinus.	3
3	Leaf	Starch, sugars		B	Starch and sugar analysis of leaf blades, with petiole maintained at 1-25°C (maximum translocation at 25°C). Encasing petioles with N ₂ inhibited translocation.	4
4	Leaf	Sugars		B	Increased translocation of sugars resulting from application of boron may be due to element's tendency to decrease enzymatic conversion of glucose-1-phosphate to starch. Increased synthesis of sucrose or other hexose phosphates from resultant glucose-1-phosphate may cause the increased translocation.	5
5	Leaf, stem	Fructose, glucose, sucrose	10?	A ¹ , B ¹ , 2	C ¹⁴ activity in various constituents of stem determined by autoradiography after application of C ¹⁴ -labelled CO ₂ to 1 leaflet. P ³² and THO, applied simultaneously with C ¹⁴ , were translocated at a slower rate. Concentration gradient of THO within stem was different from that of C ¹⁴ and P ³² .	6, 7
6	Leaf, stem	Sucrose		A ¹ , B ³	Solution applied to 1 primary leaf, with petiole maintained at 5-40°C (maximum translocation at 20-30°C). Subsequent apical growth in dark measured. Retarding effect of low temperatures decreased with time.	8

1/1 Stem 1/2 Leaf 1/3 Petiole.

Part I: ASSIMILATES AND CARBOHYDRATES (Continued)

Plant	Part	Substance	Translocation		Remarks	Reference
			Rate cm/hr	Direction		
(A)	(B)	(C)	(D)	(E)	(F)	(G)
7 Bean (Phaseolus vulgaris) (concluded)	Leaf, stem	Sucrose		A ¹ , B ³	Solution applied to 1 primary leaf, with petiole maintained at fluctuating temperatures in 12-hr cycles: 25° and 50°, 40° and 25°, and 25°C continuous (maximum translocation). Stem elongation, during 81-hr period in dark, measured.	9
8	Leaf, stem	Photosynthate		A, B	One primary leaf exposed to C ¹³ -labelled CO ₂ ; active photosynthate determined in various plant parts.	10
9	Leaf, stem	Photosynthate, sucrose		A ¹ , B ¹ , 3	Hypocotyls maintained at 7-40°C (maximum translocation at 33°). Plants placed either in light, or in dark with 1 primary leaf in solution. Apical growth measured over period of 115 hr.	11
10 Bean (Phaseolus vulgaris) Beet, sugar (Beta vulgaris) Cotton (Gossypium sp) Plantain (Plantago sp) Pumpkin (Cucurbita pepo)	Leaf, stem, root, fruit	Sucrose, hexose phosphates	30-100	A ¹ , B ¹ , 2	Autoradiography and radiochromatography of isolated vascular bundles, after application of C ¹⁴ -labelled sugar or CO ₂ to beet leaves, showed sucrose to be the dominant sugar. Presence of phosphorylase and phosphatase, and of sugar-phosphate esters in other plants, suggested esters as being an important translocatory form along with sucrose. In pumpkin, translocation rate greatest in young plants, and greater in basipetal direction than in acropetal.	12-14
11 Bean (Phaseolus vulgaris) Milkweed (Asclepias syriaca) Tomato (Lycopersicon esculentum)	Leaf	Soluble sugars, starch		B	Change in carbohydrate content and dry weight of matched pairs of leaves, both excised and intact, during 13-hr dark period Temp. = 4-40°C (maximum translocation at 20-30°)	15
12 Bean (Phaseolus vulgaris) Tomato (Lycopersicon esculentum)	Leaf, stem, root	Sucrose		A, B	C ¹⁴ activity in stem tip and roots determined 4-24 hr after submersion of lower leaf in C ¹⁴ -labelled solution. Addition of 10 ppm boron to solution increased activity in tip by 550% in tomato.	16
13 Beet, sugar (Beta vulgaris)	Leaf, stem, root	Fructose, glucose, sucrose		A, B	Chemical analysis of plant parts. Translocation only out of mature leaves, but could take place into young leaves	17
14	Leaf, stem, root	Glucose, sucrose	2-3	A, B	Application of C ¹⁴ -labelled glucose or sucrose to 4 basal leaves. Movement to youngest leaf and roots determined by autoradiogram: activity in specific sugars noted by radiochromatogram. Temp = 26°C.	18

/1/ Stem. /2/ Leaf. /3/ Petiole.

117. TRANSLOCATION OF ORGANIC SUBSTANCES. PLANTS (Continued)
Part I. ASSIMILATES AND CARBOHYDRATES (Continued)

Plant	Part	Substance	Translocation		Remarks	Reference
			Rate cm/hr	Direction		
(A)	(B)	(C)	(D)	(E)	(F)	(G)
15 Beet, sugar (<i>Beta vulgaris</i>) (concluded)	Leaf, stalk, root	Sucrose, mono- saccharides	70-100	B	Sucrose was first free sugar formed during photosynthesis. It penetrated vascular bundles and moved to lower stalk and root more rapidly than mono- saccharides, as determined by radiochromatography.	19,20
16	Leaf, stem, root	Reducing sugars, sucrose		B	Chemical analysis of different sugars in various plant parts and tissues. Reducing sugars apparently polymerized into sucrose in leaf phloem.	21
17 Cotton (<i>Gossypium</i> sp)	Stem	Total sugars	2-3	B	Chemical analysis of tissue in iso- lated and semi-isolated flaps of stem phloem under various con- ditions. Temp. = 26°C.	22
18 Cucumber (<i>Cucumis</i> <i>sativus</i>) Pumpkin (<i>Cucurbita pepo</i>)	Peduncle	Assimilates	14	Into fruit	Rate of movement based on calcu- lations of sap flow required to develop full-size cucumber (based on exudation rate of cut peduncles), and from cross- sectional area of phloem. Temp. = 2-24°C.	23
19 Geranium (<i>Pelargonium</i> <i>hortorum</i>)	Leaf, stem	Photosynthate		A, B	C ¹⁴ -labelled CO ₂ applied to leaf above stem section with phloem separated from xylem, and K ₂ H ₂ P ₃ O ₄ applied to leaf below section. Simultaneous move- ment of C ¹⁴ and P ³² in opposite directions through separated phloem. Temp. = 28-29°C.	24
20 Grape, concord (<i>Vitis labrus-</i> <i>cana</i>)	Leaf, cane	Sucrose	60	A ¹ , B ^{1,2}	Radiochemical analysis at various locl along cane, after application of C ¹⁴ -labelled CO ₂ to single leaf, indicated sucrose to be only translocated sugar. The fructose and glucose found were apparently hydrolytic products of the translocatory sucrose.	25
21 Hornbeam (<i>Carpinus</i> <i>betulus</i>) Locust, black (<i>Robinia pseudo-</i> <i>acacia</i>) Wax plant (<i>Hoya carnosa</i>)	Leaf	Sucrose			Sucrose, but no monosaccharides or sugar phosphates, identified in phloem exudate by chroma- tography. Presence of phos- phorylated compounds in bundle sheath suggested splitting and synthesis into sucrose before entering phloem.	26
22 Kelp, giant (<i>Macrocystis</i> <i>pyraifera</i>)	Stipe	Photosynthate		A, B	Translocation into youngest leaves and basal holdfast determined from measurements of photo- synthesis, respiration, and dry weight change of blades at various levels. Temp. = 19- 21°C.	27
23 Maize (<i>Zea mays</i>)	Leaf, stalk	Primarily sucrose		A, B	Translocation from leaves to grain during period of development shown by chemical analysis of various plant parts.	28

1/1 Stem 1/2 Leaf.

117. TRANSLOCATION OF ORGANIC SUBSTANCES: PLANTS (Continued)

Part I ASSIMILATES AND CARBOHYDRATES (Continued)

	Plant	Part	Substance	Translocation		Remarks	Reference
				Rate cm/hr	Direction		
	(A)	(B)	(C)	(D)	(E)	(F)	(G)
7	Bean (Phaseolus vulgaris) (concluded)	Leaf, stem	Sucrose		A ¹ , B ³	Solution applied to 1 primary leaf, with petiole maintained at fluctuating temperatures in 12-hr cycles: 25° and 50°, 40° and 25°, and 25°C continuous (maximum translocation). Stem elongation, during 81-hr period in dark, measured.	9
8		Leaf, stem	Photosynthate		A, B	One primary leaf exposed to C ¹³ -labelled CO ₂ ; active photosynthate determined in various plant parts.	10
9		Leaf, stem	Photosynthate, sucrose		A ¹ , B ¹ , 3	Hypocotyls maintained at 7-40°C (maximum translocation at 33°). Plants placed either in light, or in dark with 1 primary leaf in solution. Apical growth measured over period of 115 hr.	11
10	Bean (Phaseolus vulgaris) Beet, sugar (Beta vulgaris) Cotton (Gossypium sp) Plantain (Plantago sp) Pumpkin (Cucurbita pepo)	Leaf, stem, root, fruit	Sucrose, hexose phosphates	30-100	A ¹ , B ¹ , 2	Autoradiography and radiochromatography of isolated vascular bundles, after application of C ¹⁴ -labelled sugar or CO ₂ to beet leaves, showed sucrose to be the dominant sugar. Presence of phosphorylase and phosphatase, and of sugar-phosphate esters in other plants, suggested esters as being an important translocatory form along with sucrose. In pumpkin, translocation rate greatest in young plants, and greater in basipetal direction than in acropetal.	12-14
11	Bean (Phaseolus vulgaris) Milkweed (Asclepias syriaca) Tomato (Lycopersicon esculentum)	Leaf	Soluble sugars, starch		B	Change in carbohydrate content and dry weight of matched pairs of leaves, both excised and intact, during 13-hr dark period Temp. = 4-40°C (maximum translocation at 20-30°).	15
12	Bean (Phaseolus vulgaris) Tomato (Lycopersicon esculentum)	Leaf, stem, root	Sucrose		A, B	C ¹⁴ activity in stem tip and roots determined 4-24 hr after submersion of lower leaf in C ¹⁴ -labelled solution. Addition of 10 ppm boron to solution increased activity in tip by 550% in tomato.	16
13	Beet, sugar (Beta vulgaris)	Leaf, stem, root	Fructose, glucose, sucrose		A, B	Chemical analysis of plant parts. Translocation only out of mature leaves, but could take place into young leaves.	17
14		Leaf, stem, root	Glucose, sucrose	2-3	A, B	Application of C ¹⁴ -labelled glucose or sucrose to 4 basal leaves. Movement to youngest leaf and roots determined by autoradiogram. activity in specific sugars noted by radiochromatogram. Temp. = 26°C.	18

/1/ Stem. /2/ Leaf. /3/ Petiole.

117. TRANSLOCATION OF ORGANIC SUBSTANCES. PLANTS (Continued)

Part I: ASSIMILATES AND CARBOHYDRATES (Continued)

Plant	Part	Substance	Translocation		Remarks	Reference
			Rate cm/hr	Direction		
(A)	(B)	(C)	(D)	(E)	(F)	(G)
34 Soybean (<i>Glycine max</i>)	Leaf, stem	Fructose, glucose, sucrose	84	A, B	C ¹⁴ -labelled CO ₂ applied to 1 leaf during photosynthesis, recovered 20 min later in sucrose and invert sugars of apical portion of stem. Treatment of single leaflet did not result in appreciable movement to other leaflets. Temp. = 26-30°C.	40, 41
35	Leaf, stem, root	Fructose, glucose, photosyn- thate, sucrose	90	A ¹ , B ¹ , 2	C ¹⁴ -labelled sucrose or glucose, applied to primary leaf, was translocated more in dark than in light. Sugar uptake was increased by boron and sodium dodecyl sulfate.	42
36	Leaf, stem, root	Fructose, glucose, photosyn- thate, sucrose	100 ⁴ , 100 ⁵	A ¹ , B ¹ , 2	Translocation of C ¹⁴ -labelled sugars applied to leaf was increased by boron and surfactant. With hexoses, activity in stem decreased logarithmically from introduction point, with sucrose, it decreased linearly. Hexoses passed killed stem sections, but were impeded at nodes, sucrose did not pass killed sections, and was not impeded at nodes.	43, 44
37	Leaf, stem, root, nodule	Photosynthate		B	Chromatographic analysis of nodules and root tissue 1 da, or 1 da and 1 night, after application of C ¹⁴ -labelled CO ₂ to foliage. Distribution of C ¹⁴ in amino acids, organic acids, and carbohydrates different in nodules and root tissue, and different in the 2 harvest periods.	45
38	Cotyledon	Reducing sugars, total sugars		From coty- ledon	Chemical analysis of cotyledons showed sugars to have nearly disappeared by emergence (10 da). K and P were also exported rapidly, followed by ether-extractables and proteins.	46
39 Squash (<i>Cucurbita pepo</i>)	Peduncle	Assimilates	8-177	Into fruit	Phloem exudate collected hourly over 24-hr period. Rates, generally decreasing with time, based on exudate collected and observed cross-sectional area of sieve tubes or phloem. Later data indicated that N ₂ content of phloem exudate was higher than N ₂ content of fruit, relative to C content. Phloem exudate may, therefore, not be precisely identical to true assimilate stream. Temp. = 20-32°C.	47, 48
40 Sugar cane (<i>Saccharum officinarum</i>)	Leaf, stem, root	Photosynthate	43	B	C ¹⁴ -labelled CO ₂ applied to 1 leaf. Activity detected in roots after 6 hr, and in sister shoot of same plant after 20 hr.	49

1/1 Stem 2/ Leaf. 4/ Old seedlings. 5/ Young seedlings.

Part I. ASSIMILATES AND CARBOHYDRATES (Continued)

Plant	Part	Substance	Translocation		Remarks	Reference
			Rate cm/hr	Direction		
(A)	(B)	(C)	(D)	(E)	(F)	(G)
Maize (Zea mays) (concluded)	Shoot, root	Photosynthate		B	Activity of various substances in shoot and root determined after 24 hr of photosynthesis in atmosphere of C^{14} -labelled CO_2 . Maximum activity in carbohydrates, organic and amino acids, with less activity in pigments, fat, protein, starch, and cellulose.	29
Molds, black, blue, green (Aspergillus flavus, A. niger, A. oryzae, Penicillium notatum, Rhizopus nigricans, R. oryzae)	Mycelium	Glucose, sucrose (also nitrogen, phosphorus, fluorescein)		Unidirectional	Translocation determined by ability of a mold to grow from a full-nutrient agar into a deficient agar. Positive ability (<i>Rhizopus</i> only) associated with protoplasmic streaming. Visual observation corroborated by chemical analysis.	30
Oak, red (Quercus rubra) Other broad-leaved trees	Trunk	Assimilates, sugars	10-200	B	Determinations of sugar content of sap at heights between 1.5 and 12.5 m at various time intervals of day and night, and measurement of exudation rates.	31,32
Palm, date (Phoenix dactylifera)	Peduncle, trunk	Assimilates		Into fruit	Increment in pericarp dry weight, measured at 6 a.m. and 6 p.m. on consecutive days, indicated approximately 10 times more assimilate translocated into fruit during night than during day.	33
Pea (Pisum sativum)	Leaf	Assimilates		B	Change in weight of matched pairs of leaflets, both excised and intact, during growth in light or a 15-hr dark period. Translocation at $10^{\circ}C$ as great as, or greater than, that at $20^{\circ}C$.	34
Pea (Pisum sativum)	Leaf	Photosynthate		B	Measurement of photosynthesis, respiration, and dry weight change of pea leaves indicated maximum export during afternoon. Potato showed maximum export during night.	35
Potato (Solanum tuberosum)	Rhizome	Soluble carbohydrates	40	Into tuber	Calculated rate based on assumed movement of 10% solution in phloem, cross-sectional area of which was measured.	36
Pumpkin, Connecticut field Squash, early prolific straightneck	Peduncle	Assimilates	55-160	Into fruit	Translocation rate calculated from increment in dry weight of fruit at various stages of development, from observed cross-sectional area of peduncle sieve tube lumina, and from 10-20% concentration of assimilates therein	37
Raspberry (Rubus idaeus)	Stem	Reducing sugars, starch, sucrose		A, B	Both reducing sugars and sucrose translocated, but sucrose accumulated to a greater degree above stem girdle.	38
Sausage tree (Kigelia africana)	Peduncle	Sugars		Into fruit	The 1.75% sugar content of flowing sap is seemingly too low a concentration to account for growth rate of fruit on assumption of a mass flow hypothesis.	39

117. TRANSLOCATION OF ORGANIC SUBSTANCES: PLANTS (Continued)

Part I. ASSIMILATES AND CARBOHYDRATES (Concluded)

- References** [1] Zimmerman, M. H. 1957. *Plant Physiol.* 32:288. [2] Zimmerman, M. H. 1957. *Ibid.* 32:399. [3] Garren, R., L. P. Reimert, and N. L. Lawrence. 1953. *Bot. Gaz.* 115:105. [4] Curtis, O. F. 1929. *Am. J. Bot.* 16:154. [5] Dugger, W. M., Jr., T. E. Humphreys, and B. Calhoun. 1957. *Plant Physiol.* 32:364. [6] Biddulph, O., and R. Cory. 1957. *Ibid.* 32:608. [7] Biddulph, S., O. Biddulph, and R. Cory. 1958. *Am. J. Bot.* 45:648. [8] Swanson, C. A., and R. H. Böhnig. 1951. *Plant Physiol.* 26:557. [9] Kendall, W. A. 1952. *Ibid.* 27:631. [10] Rabideau, G. S., and G. O. Burr. 1949. *Am. J. Bot.* 32:349. [11] Böhnig, R. H., C. A. Swanson, and A. J. Linck. 1952. *Plant Physiol.* 27:417. [12] Kursanov, A. L. 1956. *Proc. Internat. Conf. Peaceful Uses Atomic Energy*, Geneva 12:165. United Nations, New York. [13] Kursanov, A. L., M. V. Turkina, and I. M. Dubinina. 1953. *Doklady acad. nauk. S. S. S. R.* 93:1115. [14] Kursanov, A. L., and E. I. Vyskrebentseva. 1954. *Fiziol. Rastenii* 1:156. [15] Hewitt, S. P., and O. F. Curtis. 1948. *Am. J. Bot.* 35:746. [16] Gauch, H. G., and W. M. Dugger, Jr. 1953. *Plant Physiol.* 28:457. [17] Leonard, O. A. 1939. *Ibid.* 14:55. [18] Hall, H. M. 1952. *Am. J. Bot.* 39:661. [19] Kursanov, A. L. 1954. *Bot. Zhur. S. S. S. R.* 39:482. [20] Turkina, M. V. 1954. *Biokhimiya*, Moskva 19:357. [21] William, A. 1945. *Inst. Amelior Betterave* 13:256. [22] Mason, T. G., and E. J. Maskell. 1928. *Ann. Bot., Lond.* 42:189. [23] Crafts, A. S. 1932. *Plant Physiol.* 7:183. [24] Chen, S. L. 1953. *Ibid.* 38:203. [25] Swanson, C. A., and E. D. H. El-Shishiny. 1958. *Plant Physiol.* 33:33. [26] Wanner, H. 1952. *Planta* 41:190. [27] Sargent, M. C., and L. W. Lantrip. 1952. *Am. J. Bot.* 39:99. [28] Loomis, W. E. 1945. *Science* 101:398. [29] Rabideau, G. S., and L. W. Mericle. 1953. *Plant Physiol.* 28:329. [30] Schütte, K. H. 1946. *New Phytol.* 55:164. [31] Huber, B., E. Schmidt, and H. Jahnel. 1937. *Tharandt. Gest. Jb.* 88:1017. [32] Huber, B. 1941. *Ber. deut. bot. Ges.* 59:181. [33] Curtis, O. F., Jr. 1947. *Am. J. Bot.* 34:388. [34] Monsellise, S. P., and F. W. Went. 1958. *Plant Physiol.* 33:372. [35] Tchesnokov, V., and K. Bazyrina. 1930. *Planta* 11:473. [36] Dixon, H. H., and N. G. Ball. 1922. *Nature*, Lond. 109:236. [37] Crafts, A. S., and O. A. Lorenz. 1944. *Plant Physiol.* 19:131. [38] Engard, C. J. 1939. *Bot. Gaz.* 100:439. [39] Clements, H. F. 1940. *Plant Physiol.* 15:689. [40] Vernon, L. P., and S. Aronoff. 1952. *Arch. Biochem.*, N. Y. 36:383. [41] Aronoff, S. 1955. *Plant Physiol.* 30:184. [42] Nelson, C. D. 1956. *Plant Physiol., Suppl.* 31:33. [43] Nelson, C. D., and P. R. Gorham. 1957. *Canad. J. Bot.* 35:359. [44] Nelson, C. D., and P. R. Gorham. 1957. *Ibid.* 35:703. [45] Bach, M. K., W. E. Magee, and R. H. Burris. 1958. *Plant Physiol.* 33:118. [46] McAllister, D. F., and O. A. Krober. 1951. *Plant Physiol., Suppl.* 26:525. [47] Crafts, A. S. 1936. *Plant Physiol.* 11:63. [48] Crafts, A. S., and O. A. Lorenz. 1944. *Ibid.* 19:326. [49] Harth, Constance E., and G. O. Burr. 1950. *Proc. Internat. Bot. Congr.* 7:748. [50] Burr, G. O., T. Yamamoto, C. E. Harth, A. Forbes, G. Sadaoka, F. M. Ashton, J. H. Payne, J. A. Silva, and G. E. Sloane. 1956. *Proc. Internat. Conf. Peaceful Uses Atomic Energy*, Geneva 12:177. United Nations, New York. [51] Skok, J. 1957. *Plant Physiol.* 32:308. [52] Goddall, D. W. 1946. *Ann. Bot., Lond.* 10:305. [53] Went, F. W. 1944. *Am. J. Bot.* 31:597. [54] Went, F. W., and H. M. Hall. 1949. *Plant Physiol.* 24:505. [55] Böhnig, R. J., W. A. Kendall, and A. J. Linck. 1953. *Am. J. Bot.* 40:150. [56] Slater, E. C., W. M. Dugger, Jr., and H. G. Gauch. 1956. *Plant Physiol.* 31:11. [57] Zholkevich, V. 1954. *Doklady acad. nauk. S. S. S. R.* 94:653. [58] McDonough, W. T. 1958. *Plant Physiol., Suppl.* 33:42.

117. TRANSLOCATION OF ORGANIC SUBSTANCES: PLANTS (Continued)

Part 1: ASSIMILATES AND CARBOHYDRATES (Continued)

	Plant	Part	Substance	Translocation		Remarks	Reference
				Rate cm/hr	Direction		
	(A)	(B)	(C)	(D)	(E)	(F)	(G)
41	Sugar cane (<i>Saccharum officinarum</i>) (concluded)	Leaf, stalk, root	Photosynthate	43	A ¹ , 2, B ¹ , 2	Translocation of C ¹⁴ -labelled CO ₂ recovered from entire plant 18 hr after application to 5th leaf, and respired labelled CO ₂ recovered from roots 6 hr after application. C ¹⁴ -labelled CO ₂ recovered from 16 stalks of a single stool 44 hr after application to a leaf of 1 stalk.	50
42	Sunflower, mammoth Russian (<i>Helianthus</i> sp)	Cotyledon, stem	Sucrose		A ¹ , B ¹ , 6	Translocation of C ¹⁴ -labelled sucrose from cotyledon to stem was not influenced by addition of boron or germanium. This and other data suggest boron does not affect sugar transport through formation of a boron-sugar complex.	51
43	Tomato (<i>Lycopersicon esculentum</i>)	Cotyledon, leaf, stem	Assimilates		A, B	Dry weight changes in individual leaves, both intact and separated. Greatest translocation during day, with movement generally from cotyledons and older leaves to younger leaves.	52
44		Leaf, stem	Sucrose		B	Chemical analysis of leaves and accumulation of sucrose above stem girdle Temp. = 18-26°C (maximum translocation at 18°).	53
45		Leaf, stem, root	Sucrose	3-4	B	Translocation of foliar-applied sucrose into roots inferred by increased bleeding rate from cut stump, occurred 8-12 hr after sucrose application. Stems held at 1-23°C (maximum translocation at 10°).	54
46		Leaf, stem	Sucrose		A ¹ , B ³	Elongation of stem and apical leaf during 110 hr in dark while basal leaf was immersed in sucrose solution. Petioles or stems maintained at 12-30°C (maximum translocation at 24°).	55
47		Leaf, stem	Photosynthate, sucrose		A ¹ , B ¹ , 2	C ¹⁴ activity decreased logarithmically up and down stem from node of leaf supplied with C ¹⁴ -labelled CO ₂ , transport being reduced in boron-deficient plants. Boron added to C ¹⁴ -labelled sucrose increased its translocation	56
48	Wheat (<i>Triticum</i> sp)	Leaf, stem, seed	Photosynthate		A ¹ , B ¹ , 2 into seed	Analysis of seed at intervals, after application of C ¹⁴ -labelled CO ₂ to leaves, indicated more rapid translocation in irrigated areas than in water-deficient areas. Transport from awns accounted for 12% of total kernel dry weight accumulation from 2nd through 4th week after heading	57,58

/1/ Stem. /2/ Leaf. /3/ Petiole. /6/ Cotyledon.

Contributor: Hull, Herbert M

117. TRANLOCATION OF ORGANIC SUBSTANCES. PLANTS (Continued)

Part II: DYES (Concluded)

Plant	Part	Substance	Translocation		Remarks	Reference
			Rate cm/hr	Direction		
(A)	(B)	(C)	(D)	(E)	(F)	(G)
9 <i>Geranium</i> (<i>Pelargonium</i> sp.)	Leaf, stem	Fluorescein	1-30	A ² , B ¹ , 2	Applied in gelatin block to scarified leaf or stem surface, subsequent movement through sieve tube protoplasm microscopically observed with ultraviolet light. Most rapid movement in young leaves. Measurements at 1-30°C (maximum translocation at 20-30°C).	10
10 (<i>P. zonale</i>)	Leaf	Sodium fluorescein, rhodamine B and other dyes	0-2	A, B	Applied to incision on ventral petiole surface, 1 cm below blade. Movement through sieve tube observed microscopically for 3 hr in dark, humid chamber, with petiole maintained at temperatures of 2-21°C (maximum translocation at 21°). Variation from other findings perhaps due to shorter time of movement and length of translocation path, different pH and osmotic pressure of solutions, delay in entrance from wound to sieve tube.	11
11	Leaf	Potassium fluorescein (also N and P compounds)			Translocation of dye and N compounds in petiole sieve tubes not affected by N ₂ , H ₂ , or CO, but reversibly inhibited by HCN. Inhibition less intense with P compounds, suggesting independent movement. Transport apparently dependent on a respiratory process.	12
12 Pumpkin (<i>Cucurbita</i> sp.)	Leaf	Potassium fluorescein		A, B	Artificially reversing (by osmosis) polar streaming in sieve tube vacuoles caused simultaneous bi-directional movement in external and internal phloem. Transport apparently passive in streaming solution of the vacuoles, and only secondarily drawn into the cytoplasm.	13

11/ Petiole. 12/ Stem.

Contributor: Hull, Herbert M.

- References: [1] Döpp, W. 1939. Jb. wiss. Bot. 87:679. [2] Crafts, A. S. 1939. Bot. Rev. 5:471. [3] Palmquist, E. M. 1938. Am. J. Bot. 25:97. [4] Palmquist, E. M. 1939. Ibid. 26:665. [5] Parker, J., and R. G. Bowmer. 1956. Plant Physiol., Suppl. 31:16. [6] Schumacher, W. 1948. Planta 35:642. [7] Bauer, L. 1949. Ibid. 37:221. [8] Schlafke, E. 1958. Ibid. 50:388. [9] Schumacher, W. 1947. Naturwissenschaften 6:176. [10] Schumacher, W. 1933. Jb. wiss. Bot. 77:685. [11] Bauer, L. 1943. Planta 42:367. [12] Willenbrink, J. 1957. Ibid. 48:269. [13] Rouschal, E. 1941. Flora 35:135.

Part II: DYES

Plant	Part	Substance	Translocation		Remarks	Reference
			Rate cm/hr	Direction		
(A)	(B)	(C)	(D)	(E)	(F)	(G)
1 Apple (<i>Malus sylvestris</i>) Pumpkin (<i>Cucurbita pepo</i>) Quince (<i>Cydonia oblonga</i>)	Peduncle	Cyanol (also assimilates)		Into fruit	From observation of fruit growth and dye movement after severing peduncle xylem vessels, conclusion reached that water required by fruit could not be supplied through cortex. No protoplasmic streaming occurred in sieve tubes, but assimilates were transported without simultaneous movement of water.	1
2 Bean (<i>Phaseolus vulgaris</i>)	Leaf, stem	Fluorescein	2-10	B ¹	Immersion of intact leaf in 0.1% solution; subsequent movement observed with microscope.	2
3	Leaf, stem	Sodium fluorescein (uranin)		A, B	Terminal leaflet submerged in solution; substance detected in petioles of lateral leaflets after 15 hr in dark. During this period, lateral leaflets lost dry weight, suggesting simultaneous movement of dye and assimilates in opposite directions. Scalding rachises inhibited movement of dye.	3, 4
4 Bean (<i>Phaseolus vulgaris</i>) Cotton (<i>Gossypium</i> sp)	Leaf, stem, root	Sodium fluorescein (also P32)	7	A, B	Basipetal stem transport slower in cotton than in bean. P32, applied to leaf higher than dye-treated leaf, overtook dye in movement down stem. Cotton: P32 moved faster than dye in petiole, and moved up from roots through phloem by a point on the stem where xylem had been removed, while dye was simultaneously moving down.	5
5 Bryony (<i>Bryonia dioica</i>)	Leaf, stem	Sodium fluorescein	15-65	A, B	Solution applied to small epidermal wounds of stems, petioles, and leaves of intact plants. Translocation rates in stem greatest in basal portion.	6
6 Bryony (<i>Bryonia dioica</i>) Elodea (<i>Anacharis densa</i>)	Leaf	Berberine sulfate, fluorescein, primuline, quinine chloride, rhodamine B and 6G		A, B	Relative distribution of dye between sieve tube cytoplasm and vacuole dependent on concentration and pH. Microscopic examination of the preceding dye front suggested mass flow through vacuole, possibly by electro-osmotic phenomena.	7
7 Chickweed (<i>Stellaria media</i>) Elodea (<i>Anacharis densa</i>)	Leaf	Potassium fluorescein, other acid fluorescent dyes	0-30		In elodea, movement in parenchyma cell walls slower than diffusion in gelatine and not affected by narcotics. Volume of tissue into which dye moved was independent of its concentration and loci of application. In <i>Stellaria</i> , different dyes moved at different rates, a humid atmosphere causing a reduction of rate. Maximum indicated rate in midvein.	8
8 Dodder (<i>Cuscuta</i> sp)	Leaf, stem	Fluorescein	2-50	A ² , B ¹ , 2	Substance moved basally to stem when applied to leaf veins, and then became bi-directional. Movement apparently from cell to cell through total intercellular surface and not through plasmodesmata.	9

/1/ Petiole. /2/ Stem.

117. TRANSLOCATION OF ORGANIC SUBSTANCES. PLANTS (Continued)

Part III. HERBICIDES AND GROWTH REGULATORS (Continued)

Plant	Part	Substance	Translocation		Remarks	Reference
			Rate cm/hr	Direction		
(A)	(B)	(C)	(D)	(E)	(F)	(G)
Barley (<i>Hordeum vulgare</i>) Bean (<i>Phaseolus vulgaris</i>) Cucumber (<i>Cucumis sativus</i>) Maize (<i>Zea mays</i>) Sunflower (<i>Helianthus</i> sp) Tomato (<i>Lycopersicon esculentum</i>)	Leaf, stem	α -Methoxyphenyl- acetic acid		A, B	Applied to stem (dicots) or leaf (monocots), compound caused epinasty or repres- sion of terminal growth. Although phenylacetic acid produced gall at site of application in bean, it did not produce a secondary gall at internode above, as did α -methoxyphenyl- acetic acid.	6
Barley (<i>Hordeum vulgare</i>) Grass, Bermuda (<i>Cynodon dactylon</i>) Wandering Jew (<i>Tradescantia fluminensis</i> , <i>Zebrina pendula</i>)	Leaf, stem, root	3-Amino-1, 2, 4- triazole; 2, 4-D, 1, 2-dihydro- pyridazine-3, 6-dione ¹ ; 3-(p- chlorophenyl)- 1, 1-dimethyl- urea ² , 2, 4, 5-T, urea	0-30	A, B	Translocation determined by autoradiography at various intervals after application of C ¹⁴ -labelled compound to 1 or several leaves, or to stolons. Movement of 2, 4-D restricted due to absorption in phloem, but absorption minimized in rapidly growing plants. Other compounds translo- cated throughout plants to a greater extent, some mov- ing in the phloem, others also entering xylem or moving predominantly in xylem. None moved out of chlorotic leaves.	7-9
8 Bean (<i>Phaseolus vulgaris</i>)	Petiole, stem, root	3-Indoleacetic acid	0.17	B	Donor, containing 3-indole- acetic acid, and receptor agar blocks applied to ends of 5 mm section. Assay of block by <i>Avena</i> curvature test. Infusion of 2, 4-D or 2, 3, 5-triiodobenzoic acid into section (directly or via foliage before cutting) markedly inhibited ability to translocate 3-indole- acetic acid	10
9	Hypocotyl	3-Indoleacetic acid	0.17	B	Method of determination same as No. 8 above. Only basi- petal translocation observ- ed; maximum intensity at top of older (8 da) hypo- cotyls. Temp. = 24°C.	11
10	Leaf, stem	3-Indoleacetic acid, 2, 4-D (morpholine salt), β -hydroxyethyl- 4-chlorophen- oxyacetate; p-chlorophenyl-2, 4, 5-trichlorophenoxyacetate; 2-bromo-3, 5-dichloroben- zoic acid, (2, 4-dichloro- phenoxyacetyl)-urea		A, B	Substances all translocated, as detected by stem bend- ing after application to 1 primary leaf. Sucrose, glucose, fructose, maltose, lactose, and galactose each caused 2, 4-D to translocate out of pre-darkened leaf when applied simultane- ously with 2, 4-D.	12

11/ Maleic hydrazide. 12/ Monuron.

117. TRANSLOCATION OF ORGANIC SUBSTANCES: PLANTS (Continued)

Part III: HERBICIDES AND GROWTH REGULATORS

	Plant	Part	Substance	Translocation		Remarks	Reference
				Rate cm/hr	Direction		
	(A)	(B)	(C)	(D)	(E)	(F)	(G)
1	Alligator weed (<i>Alternanthera phloxeroides</i>)	Leaf, stem	2, 4-D	4.2-4.3	A, B	C ¹⁴ -labelled 2, 4-D applied to 1 pair leaves. Stems sectioned and extracted for activity determinations 8 or more hr after treatment. Acropetal movement in stem more extensive and slightly more rapid than basipetal.	1
2	Alta fescue (<i>Festuca arundinacea</i>) Barley (<i>Hordeum vulgare</i>) Grass, rye (<i>Lolium</i> sp) Oat (<i>Avena</i> sp)	Leaf	Lactic acid N- phenylcarbamate, lactic acid N-3- chlorophenyl- carbamate		A, B	Application to first leaf, with measurement of subsequent repression of young leaf growth. Greater transloca- tion of acid compounds than of corresponding iso- propyl compounds, the lactic acid molecular con- figuration apparently being advantageous to transloca- tion.	2
3	Apple (<i>Malus</i> sp) Pear (<i>Pyrus</i> sp)	Twig	Native auxin, 3-indoleacetic acid	0.3-0.4	B	Translocation determined in 5-10 mm sections by agar block method. Auxin in receptor block analyzed by <i>Avena</i> test. Native auxin moved only basipetally in active twigs. Applied 3-indoleacetic acid moved in similar manner, but acropetally to a very slight extent in dormant twigs.	3
4	Apple, crab (<i>Malus</i> sp)	Stem	Native auxin	0.5-0.7	B	Stem sections placed with base on agar block. Auxin diffusing into block meas- ured by <i>Avena</i> curvature test. O ₂ tension, ranging from zero to that in air, gave straight line correla- tion with translocation rate. Within temperature range of 0-42°C, maxi- mum transport occurred at 27-30°C and was nearly zero at extremes.	4
5	Autumn-crocus (<i>Colchicum autumnale</i>) Buttercup (<i>Ranunculus</i> sp) Chickweed (<i>Stellaria media</i>) Dandelion (<i>Taraxacum</i> sp) Mustard, white (<i>Brassica hirta</i>) Orache (<i>Atriplex patula</i>) Pigweed (<i>Chenopodium album</i>)	Leaf, stem	2, 4-D		A, B	Transport in orache and pig- weed initially basipetal and later acropetal. Sensi- tivity to a second 2, 4-D application was variable among different plants and depended on length of time after first application.	5

III. TRANSLLOCATION OF ORGANIC SUBSTANCES: PLANTS (Continued)

Part III: REMEDIES AND GROWTH REGULATORS (Continued)

Plant	Part	Substance	Translocation		Remarks	Reference
			Rate cm/hr	Direction		
(A)	(B)	(C)	(D)	(E)	(F)	(G)
18. Bean (<i>Phaseolus vulgaris</i>) (continued)	Leaf, stem, root	2,4-D		A ¹ , B ¹ , 4	Bioassay of stem and roots, after application of C ¹⁴ -labelled 2,4-D to primary leaf, showed activity in 6 hr. with transport chiefly basipetal. After 1 wk. approximately 1/3 of C ¹⁴ in stem still in 2,4-D. much of remainder being in an unknown water-soluble acid.	23,24
19	Leaf, stem	2,4-D			Pretreatment with gibberellic acid 48 hr before 2,4-D application increased amount of 2,4-D in untreated plant parts after several da. Gibberellic acid apparently affects translocation rather than absorption or degradation of 2,4-D.	22
20	Leaf, stem, root	2,4-D			Applied to leaves and determined in various parts by bioassay. No transport from leaves of plants kept in dark 18-24 hr prior to and during treatment. Marmitol did not facilitate movement as did sucrose. Low temperatures reduced transport.	23
21	Entire plant	2,4-D		A, B	Primary leaves severed 24 hr after application and entire remainder of plant extracted. Presence of unaltered 2,4-D determined both by bioassay of extract and by chemical characterization.	24
22	Leaf, stem	2,4-D		A, B	Application directly to apical bud caused growth repression independent of temperatures from 22-34°C. Application via primary leaf resulted in maximum repression of apical bud and leaf expansion at lower temperature.	25
23	Leaf, stem	2,4-D	13-106	B	Application to primary leaf caused epicotyl curvature to appear sooner at 35°C than at 25°, but maximum angle of curvature was 60% greater at lower temperature.	26
24	Leaf, stem	2,4-D (also water)		B	The fact that labelled 2,4-D did not move rapidly downward through steam-girdled petioles (as did labelled water), along with the ability of water to move out of carbohydrate-depleted leaves (2,4-D does not), renders improbable a mass movement of 2,4-D and water in phloem.	27

13/ Stem. 14/ Leaf.

117. TRANSLOCATION OF ORGANIC SUBSTANCES: PLANTS (Continued)

Part III: HERBICIDES AND GROWTH REGULATORS (Continued)

Plant	Part	Substance	Translocation		Remarks	Reference
			Rate cm/hr	Direction		
(A)	(B)	(C)	(D)	(E)	(F)	(G)
11 Bean (Phaseolus vulgaris) (continued)	Leaf, stem, root	3-(p-Chloro- phenyl)-1, 1-di- methylurea		A, B	1 leaf treated with Cl^{14} - labelled compound; plant sectioned for assay of activity 2-72 hr after treat- ment. Translocation throughout plant, maximum concentration in non-treated leaves occurring only by 4th da.	13
12	Leaf, stem, hypocotyl	2-Iodo-3-nitro- benzoic acid		A, B	Accumulation in apical bud and hypocotyl indicated by radioactivity.	14
13	Leaf, stem, root	2, 4-Dichloro-5- iodophenoxy- acetic acid, and 14 derivatives		B	^{131}I -labelled compounds applied to leaves, stems, and roots. Movement from leaf to stem (determined by curvature of stem) occurred only if plant kept in light. As spray, ^{131}I - labelled 2, 4-D-5-iodo- phenoxyacetic acid less	15
14	Leaf, stem, root	2, 4-Dichloro-5- iodophenoxy- acetic acid (free acid and morpholine salt)		A, B	to 1 primary leaf. Concen- tration in stem significant- ly greater with acid than with morpholine salt, and markedly increased by addition of wetting agent.	
15	Leaf, stem	Various butyl, octyl, and octadecyl esters of 2, 4-D and 2, 4, 5-T		B	Application to 1 primary leaf, with resultant stem curva- ture measured after 4 hr. Only plants kept in light developed curvature Temp. = 21-24°C.	17
16	Leaf, stem, root	2, 4-D		A, B	Methylene Cl^{14} -labelled 2, 4-D applied to midrib of 1 pri- mary leaf. Plants section- ed 2-144 hr after applica- tion and activity deter- mined. Activity in apical bud after 8 hr, but no movement into leaf oppo- site treated leaf.	18
17	Leaf, stem	2, 4-D		A ³ , B ⁴	Content of 2, 4-D in various plant parts determined by bioassay after application to primary leaf. Entered epicotyl in 6 hr, but did not move below center of hypo- cotyl. Transport inhibited in case of 2nd application or too high a concentration, due to phloem destruction. Sucrose, but not mannitol and urea, enhanced trans- port in dark.	19

117. TRANSLOCATION OF ORGANIC SUBSTANCES PLANTS (Continued)
Part III. HERBICIDES AND GROWTH REGULATORS (Continued)

Plant	Part	Substance	Translocation		Remarks	Reference
			Rate cm/hr	Direction		
(A)	(B)	(C)	(D)	(E)	(F)	(G)
Bean (<i>Phaseolus vulgaris</i>) (continued)	Leaf, stem, root	2,4-D		A ³ , B ³ , 4	Radioassay of stem and roots, after application of C ¹⁴ -labelled 2,4-D to primary leaf. showed activity in 6 hr, with transport chiefly basipetal. After 1 wk. approximately 1/3 of C ¹⁴ in stem still in 2,4-D, much of remainder being in an unknown water-soluble acid.	20,21
19	Leaf, stem	2,4-D			Pretreatment with gibberellic acid 48 hr before 2,4-D application increased amount of 2,4-D in untreated plant parts after several da. Gibberellic acid apparently affects translocation rather than absorption or degradation of 2,4-D.	22
20	Leaf, stem, root	2,4-D			Applied to leaves and determined in various parts by bioassay. No transport from leaves of plants kept in dark 18-24 hr prior to and during treatment. Mannitol did not facilitate movement as did sucrose. Low temperatures reduced transport.	23
21	Entire plant	2,4-D		A, B	Primary leaves severed 24 hr after application and entire remainder of plant extracted. Presence of unaltered 2,4-D determined both by bioassay of extract and by chemical characterization.	24
22	Leaf, stem	2,4-D		A, B	Application directly to apical bud caused growth repression independent of temperatures from 22-34°C. Application via primary leaf resulted in maximum repression of apical bud and leaf expansion at lower temperature.	25
23	Leaf, stem	2,4-D	13-106	B	Application to primary leaf caused epicotyl curvature to appear sooner at 35°C than at 25°, but maximum angle of curvature was 60% greater at lower temperature.	26
24	Leaf, stem	2,4-D (also water)		B	The fact that labelled 2,4-D did not move rapidly downward through steam-girdled petioles (as did labelled water), along with the ability of water to move out of carbohydrate-depleted leaves (2,4-D does not), renders improbable a mass movement of 2,4-D and water in phloem.	27

13/ Stem. 14/ Leaf.

117. TRANSLOCATION OF ORGANIC SUBSTANCES: PLANTS (Continued)

Part II: HERBICIDES AND GROWTH REGULATORS (Continued)

	Plant	Part	Substance	Translocation		Remarks	Reference
				Rate cm/hr	Direction		
	(A)	(B)	(C)	(D)	(E)	(F)	(G)
25	Bean (Phaseolus vulgaris) (continued)	Leaf, stem, root	2, 4-D and derivatives	4	A, B	Translocation determined by autoradiogram, stem curvature, and apical leaf growth repression after application to 1 primary leaf. Kerosene served as good carrier in destarched plants, and appeared to move through intercellular spaces of phloem parenchyma.	28
26		Leaf, stem	2, 4-D; 2, 4-D complex		B	C ¹⁴ -labelled 2, 4-D applied to veins of 1 leaf of etiolated plants; activity determined in sections made 4 da later. Addition of glucose or sucrose to 2, 4-D increased movement; addition of glucose-1-phosphate did not. Temp. = 16-24°C.	29
27		Leaf, stem	2, 4-D (ammonium salt)		A, B	Primary leaf excised 4-72 hr after application, and fresh weight of apical leaves determined after 9 da. Increased translocation rate from leaf at high light intensities.	30
28		Leaf, stem	2, 4-D (ammonium salt); 2, 4, 5-T; α -naphthalene- acetic acid		B	Movement from primary leaf to stem determined by C ¹⁴ activity in stem (using C ¹⁴ -labelled 2, 4-D), or by stem curvature. Boron applied to tip of treated leaf failed to affect translocation, as did sugar. Boron added to sugar markedly increased translocation of growth substances.	31
29		Leaf, stem, root	2, 4-D (sodium salt); 2, 4, 5-T (sodium salt)	50-100	B	Applied to 1 primary leaf in a carrier of pH 8, 4 hr required for epicotyl to bend. 2 hr required at pH 2.	32
30		Leaf, stem, root	2, 4-D, 3-indole- acetic acid		A ³ , B ³ , 4	Activity determined in various plant parts after application of C ¹⁴ -labelled 3-indoleacetic acid to primary leaf. 10-14% of activity transported out of treated leaf in 14 da, but activity in petiole and lower stem reached maximum after 1 da. Terminal bud reached maximum after 7 da. 2, 4-D moved out of treated leaf more rapidly.	33
31		Leaf, stem, root	Benzoic acid, 2, 4-D (trieth- anolamine salts)		B	Transport, determined by autoradiography and counting, was maximum at high temperature (30° over 20°C) and high humidity (72% over 36%) Maximum effect at high humidity correlated with large stomatal openings.	34

/3/ Stem. /4/ Leaf.

117. TRANSLOCATION OF ORGANIC SUBSTANCES PLANTS (Continued)

Part III. HERBICIDES AND GROWTH REGULATORS (Continued)

Plant	Part	Substance	Translocation		Remarks	Reference
			Rate cm/hr	Direction		
(A)	(B)	(C)	(D)	(E)	(F)	(G)
Bean (<i>Phaseolus vulgaris</i>) (continued)	Hypocotyl	3-Indoleacetic acid		B	Transport in sections, determined by bioassay of receptor blocks, was inhibited by 2, 4-D; 2, 4-dichlorophenoxypropionic acid; 2, 4, 6-trichlorophenoxyacetic acid, 3-indolepropionic acid, and 2, 3, 5-triiodobenzoic acid. Other compounds did not inhibit transport.	35
	Leaf, stem, root	α -Methoxyphenylacetic acid		B	Radioassay of nutrient culture solution, after application of C^{14} -labelled compound to primary leaves, showed translocation through roots into nutrient, reaching maximum after 126 hr. Decreased nutrient aeration, or severing part of roots, decreased exudation into nutrient, but exudation later increased after severed roots renewed growth.	36
	Leaf, stem, root	2, 3, 6-Trichlorobenzoic acid, 2, 3, 5, 6-trichlorobenzoic acid		A, B	Application to first internode, cotyledon, or primary leaf, resulted in movement to root, through soil, and into adjacent plants, as determined by growth modification. Transport from leaf to stem took 4 hr. Other chlorinated benzoic acids did not affect adjacent plants.	37
	Stem	Gibberellic acid			C^{14} -labelled compound applied to first internodes of young plants; translocation determined by autoradiography after 24-48 hr.	38
	Leaf, stem	Gibberellic acid		A	Application by foliage spray or to stem base or apex caused an increase in stem length of 0-300% in various plant species, indicating translocation.	39
	Leaf, stem	2, 4-Dichlorobenzylpicotinium chloride; 3, 4-dichlorobenzylpicotinium chloride; p-chlorobenzylpicotinium chloride, other nicotinium compounds		A ³ , B ⁴	Translocation, indicated by per cent reduction in stem elongation 10 da following treatment of specific parts, as follows: first internode, 34%; hypocotyl, 23%; leaves, 7%; cotyledon, no reduction. 2, 4-compound most effective, and young plants most sensitive.	40

/3/ Stem. /4/ Leaf.

117. TRANSLOCATION OF ORGANIC SUBSTANCES: PLANTS (Continued)

Part II: HERBICIDES AND GROWTH REGULATORS (Continued)

Plant	Part	Substance	Translocation		Remarks	Reference
			Rate cm/hr	Direction		
(A)	(B)	(C)	(D)	(E)	(F)	(G)
38 Bean (<i>Phaseolus vulgaris</i>) (concluded)	Leaf, stem, root	3-Amino-1,2,4-triazole		A ³ , 4, B ³ , 4	C ¹⁴ -labelled compound applied to primary leaves moved basipetally in stem during first 24 hr, but radiochromatography of plant sections after 5 da showed C ¹⁴ concentrated in terminal bud and aminotriazole converted to 2 unknowns. Only 7% of total activity remained as aminotriazole by 5th da, and about half of activity had moved out of treated leaves.	41
39 Bean (<i>Phaseolus vulgaris</i>) Cotton (<i>Gossypium</i> sp) Cucumber (<i>Cucumis sativus</i>) Morning-glory (<i>Ipomoea</i> sp) Pea, black-eyed (<i>Vigna sinensis</i>)	Leaf, stem, root	2,4-D (acid, amine, ester, and sodium salt formulations)	0-70	A, B	Transport determined by autoradiography at intervals of 15 min-144 hr after application to specific cotyledons or leaves. 2,4-D reached bean roots in 3 hr and terminal bud in 6 hr, and moved out of cotyledons or basal leaves more rapidly than out of young leaves. It is transported with assimilates, and may move in water-stressed plants. Effect enhanced by low pH and surfactants.	42,43
40 Bean (<i>Phaseolus vulgaris</i>) Cotton (<i>Gossypium</i> sp) Maize (<i>Zea mays</i>) Millet (<i>Pennisetum</i> sp) Rice (<i>Oryza sativa</i>) Soybean (<i>Glycine max</i>)	Leaf, stem, root	2,4-D; other halogenated phenoxyacetic acids		A, B	Application of labelled compound to 1 leaf and harvest of plant at intervals for C ¹⁴ assay demonstrated that translocation, but not penetration or metabolic inactivation, was correlated with varietal susceptibility. Translocation of phenoxy acids in several dicots was 5-70 times greater than in monocots, in the following decreasing order: 4-fluoro-; 2,4-di-fluoro-; 4-chloro-, 2,4-dichloro-, 2,4,5-trichloro-.	44
41 Bean (<i>Phaseolus vulgaris</i>) Cotton (<i>Gossypium</i> sp) Soybean (<i>Glycine max</i>) Tomato (<i>Lycopersicon esculentum</i>)	Leaf, stem	2,4-D (ammonium salt)		A, B	Growth reduction as affected by application to individual leaves of various ages. Maximum reduction achieved with application to apical leaves or stem. No basipetal movement occurred through dead stem segments.	45
42 Bean (<i>Phaseolus vulgaris</i>) Maize (<i>Zea mays</i>)	Leaf, stem	Fungal metabolites		A ³ , B ⁴	Application of filtrate of <i>Penicillium thomii</i> or <i>Arachnoidiscus trisporus</i> culture to 1 leaf resulted in growth repression and abnormal development of subsequent leaves.	46

117. TRANSLLOCATION OF ORGANIC SUBSTANCES: PLANTS (Continued)

Part III HERBICIDES AND GROWTH REGULATORS (Continued)

Plant	Part	Substance	Translocation		Remarks	Reference
			Rate cm/hr	Direction		
(A)	(B)	(C)	(D)	(E)	(F)	(G)
43 Bean (Phaseolus vulgaris) Maize (Zea mays) Oat (Avena sp)	Leaf, stem	2-Iodo-3-nitro- benzoic acid		A ³ , 4, B ⁴	Activity of ¹³¹ I determined in young leaves 6 da after application of labelled com- pound to old leaves. Accumulation greatest in bean, with activity concen- trated in meristematic regions.	47
44 Bean (Phaseolus vulgaris) Maize (Zea mays) Sunflower (Helianthus sp)	Leaf, stem, root	2, 4-Dichloro-5- iodophenoxy- acetic acid (morpholine salt)		A, B	Determination of activity in various plant parts and time intervals after appli- cation of ¹³¹ I-labelled compound to 1 leaf. Activ- ity in bean hypocotyl appeared 6 hr after appli- cation. Movement slower from maize, oat, and wheat leaves.	48
45 Bean (Phaseolus vulgaris) Oat (Avena sp)	Leaf, stem, root	2-Iodo-4-chloro- phenoxyacetic acid	1.9-3.8 ⁵	A, B	¹³¹ I-labelled material applied to 1 leaf; followed at var- ious intervals by autoradi- ogram. Bean movement throughout plant in 27 hr.	49
46 Bean (Phaseolus vulgaris) Sugar cane (Saccharum officinarium)	Leaf, stem	2, 4-D		A ³ , B ³ , 6	Cl ⁴ -labelled 2, 4-D applied to a lower leaf (activity deter- mined by subsequent form- ative effect and radioassay) was readily absorbed by both bean and sugar cane. However, about 10 times more Cl ⁴ was translocated to the growing point of the bean than to the cane.	50
47 Bean, broad (Vicia faba) Maize (Zea mays)	Root	1-Indoleacetic acid, native auxin		A, B	Polarity of movement of externally applied or native auxin in root sections determined by bioassay of receptor agar block. Movement within 1-2 mm of tip was predominantly downward, but within zone 2-4 mm from tip movement was largely upward.	51-53
48 Bean, Florida velvet (Stizolobium deeringianum)	Leaf, stem	3-(p-Chloro- phenyl)-1, 1- dimethylurea		A	Short midstem section (intact plants) immersed in solu- tion of substance several da. Only leaves acropetal to treated section showed typical injury after 6 da.	54
49 Chilicothe (Echinocystia macrocarpa) Maize (Zea mays)	Stem	Gibberellic acid	0.5-1.0		Bioassay with mutant dwarf maize demonstrated longi- tudinal movement in inter- node sections to be double the rate of lateral move- ment.	55
50 Cocklebur (Xanthium sp) Maize (Zea mays) Peanut (Arachis hypogaea) Tomato (Lycopersicon esulentum)	Leaf, stem, root	1, 2-Dihydro- pyridazine- 3, 6-dione ¹		A ³ , B ³ , 6	Repeated application to single leaf or by spraying resulted in translocation of sub- stance to regions of meristematic activity, as indicated by formative effect.	56

(1) Maleic hydrazide. (3) Stem. (4) Leaf. (5) For bean; rate slower in oat. (6) Petiole.

117. TRANSLOCATION OF ORGANIC SUBSTANCES- PLANTS (Continued)

Part III: HERBICIDES AND GROWTH REGULATORS (Continued)

Plant (A)	Part (B)	Substance (C)	Translocation		Remarks (F)	Reference (G)
			Rate cm/hr (D)	Direction (E)		
51 Coleus (<i>Coleus blumei</i>)	Stem, root	Native auxin, 3-Indoleacetic acid	0.2	A 7, 8, B 3, 8	Donor, containing 3-indoleacetic acid, and receptor agar blocks placed on ends of sections; activity determined by pea root test. Acropetal movement of native auxin in intact plants shown by darkening stems; only flowering stems responded phototropically after treatment.	57
52	Stem	3-Indoleacetic acid		A, B	Transport in section from 2nd internode from apex determined by donor-receptor block technique, using <i>Avena</i> bioassay. Basipetal movement 3 times that of acropetal movement.	58
53 Cotton (<i>Gossypium</i> sp)	Leaf, stem	3-Amino-1,2,4- triazole		A, B	Apical or basal leaves submerged in solution. Injury appearing in non-treated leaves indicated bi-directional movement in stem, but only acropetal movement could traverse stem girdle.	59
54	Cotyledon, stem, root	3-Amino-1,2,4- triazole; 2,4-D, urea		B	Radioanalysis of culture solution a few days after application of C ¹⁴ -labelled compounds to cotyledons showed only 2,4-D exuded from roots, although all compounds were translocated to roots in 2 hr. Because of absorption in tissue, distribution of 2,4-D in plants was less than with other compounds.	60
55 Cotton (<i>Gossypium</i> sp) Flax (<i>Linum</i> sp) Milo (<i>Sorghum vulgare</i>) Rice (<i>Oryza sativa</i>) Watermelon (<i>Citrullus vulgaris</i>)	Cotyledon, leaf, stem	1,2-Dihydro- pyridazine- 3,6-dione ¹		A, B	Application as a spray, or only to cotyledons and stem, resulted in translocation to meristematic regions, as evidenced by stunting and development of abnormal leaves.	61
56 Cucumber (<i>Cucumis sativus</i>)	Cotyledon, hypocotyl	3-Indoleacetic acid	18.5 (15.1-21.9) ⁹		Difference in time for hypocotyl curvature to appear at 20°C, when substance is applied to cotyledon tip rather than in direct hypocotyl application. Indicates translocation rate in cotyledon.	62
57 Grass, canary (<i>Phalaris canariensis</i>)	Coleoptile	Native auxin		B	Phototropic response below tip of seedling receiving unilateral illumination. Curvature resulted from exposing seedlings to lateral light.	63

[1] Maleic hydrazide. [3] Stem. [7] Root tip. [8] Flowering stem. [9] Estimate "b" (cf. Introduction).

117. TRANSLOCATION OF ORGANIC SUBSTANCES. PLANTS (Continued)
Part III: HERBICIDES AND GROWTH REGULATORS (Continued)

Plant	Part	Substance	Translocation		Remarks	Reference
			Rate cm/hr	Direction		
(A)	(B)	(C)	(D)	(E)	(F)	(G)
Grass, Johnson (Sorghum balepense) Grass, nut (Cyperus rotundus)	Leaf, stem, rhizome	3-Amino-1,2,4- triazole	0.03-0.05	Horizontal through rhizomes A, B	30 da after foliage spray, rhizomes were separated and planted. Shoots arising were stunted and chlorotic. Translocation increased by addition of wetting agents.	64
Grass, nut (Cyperus rotundus)	Leaf, stem, root, tuber	3-Amino-1,2,4- triazole, 2,4-D, 2,4,5-T, 1,2- dihydropyrid- azine-3,6- dione ¹	20-25	A, 3, 4, B, 3, 4	Transport determined by autoradiography after application to single leaf. 3-Amino-1,2,4-triazole reached root tip in 2 hr; 2,4-D and 2,4,5-T were slower. No movement out of treated leaf if kept in dark.	65
Hickory (Carya tomentosa) Privet (Ligustrum vulgare)	Leaf, stem	Ammonium sulfamate	0.6-2.5	A, B	³⁵ S-labelled material applied to stem frill, followed by autoradiogram. Acropetal movement traversed stem girdle, but basipetal move- ment did not.	66
Lemon (Citrus limon)	Stem	3-Indoleacetic acid		B	Substance applied to upper edge or cut surface of cut- ting increased root forma- tion near base. Basipetal movement inhibited by cooling portion of cutting at 1-8°C, and also by girdling.	67
Maize (Zea mays)	Root	Native auxin		A	Excised root sections were inhibited in growth and regained geotropic response only when root tips were placed on apical end and not when placed on morphological upper end.	68
Maize (Zea mays) Oat (Avena sativa)	Leaf, stem, root	Isopropyl N- phenylcarba- mate		A, B	¹⁴ C-labelled compound applied to cut surface of oat leaf, translocation of approximately 1% or less to stem or root within 7 da. Absorption and movement from cut or intact root even smaller. No transport in maize.	69
Maize (Zea mays) Oat (Avena sativa) Papaya (Carica papaya)	Coleoptile, stem	3-Indoleacetic acid, indole- butyric acid, anthracene- acetic acid, naphthalene- acetic acid	0.2-1.2	B	Sections interposed between donor agar block and 1 side of cut surface of decapitated Avena coleoptile. Increased time for curva- ture to appear indicated time for transport through section. All substances used for oat, only 3-indole- acetic acid for maize and papaya.	70
Maize (Zea mays) Pea (Pisum sati- vum)	Leaf, stem, root	Trichloroacetic acid (sodium salt)		A, B	Outer half of 3rd leaf dipped into solution of ¹⁴ C-la- belled substance. After 24 hr, measurements and autoradiograms indicated movement into roots and other leaves. No apparent metabolism to other products.	71

1/1 Maleic hydrazide 3/3 Stem. 14/ Leaf

Part III: HERBICIDES AND GROWTH REGULATORS (Continued)

	Plant	Part	Substance	Translocation		Remarks	Reference
				Rate cm/hr	Direction		
	(A)	(B)	(C)	(D)	(E)	(F)	(G)
66	Maize (Zea mays) Wheat (Triticum sp)	Leaf, stem, root	2, 4-D		A, B	2, 4-D and 3 of its metabolic products recovered throughout plants 1-4 da after application of C ¹⁴ -labelled 2, 4-D to older leaf. Determinations by radiochromatography. Most activity occurred in young leaves, some in stem and sheath, and least in roots.	72
67	Marabu (Dichrostachys nutans)	Shoot	2, 4, 5-T (butoxy- ethanol ester)		A	Substance, applied in oil or water at different levels on surface of shoot, was translocated acropetally, as evidenced by bioassay of tissues and ultimate death of leaves and stems above treated area. Slight basipetal movement occurred with bark application, as indicated by inhibition of basal sprouting.	73
68		Stem	2, 4-D (butoxy- ethanol ester)		A, B	Positive recovery (determined by bioassay) 19 cm from top of 2-ft sucker, 6 hr after apical application. No recovery after 30 hr suggests rapid transport and/or metabolism. Transport also shown by effect on basal buds which were inhibited in sprouting even when shoot was excised 6 hr after application. Acropetal transport occurred through stem girdle; basipetal did not, and was therefore extra-axillary.	74
69	Mesquite (Prosopis juliflora)	Leaf	2, 4, 5-T (various formulations)		A ³ , B ³ , 4	Maximum translocation out of branch tips 50-90 da after leaf emergence in spring, when root sugars undergo rapid buildup. Little translocation if total root sugars are decreasing and reducing sugars are abundant. Anatomical studies showed phloem proliferation, and resultant transport inhibition could occur from too high concentration of either 2, 4, 5-T or surfactant.	75, 76
70		Leaf, stem, root	2, 4-Dichloro-5- iodophenoxy- acetic acid		B	Activity determined in more basal tissue 4 da after spraying upper half of plant with solution of ¹³¹ I-labelled compound. Use of emulsifying agent increased proportion of activity in root and hypocotyl over that in stem, but less than 3% moved from treated foliage.	77

117. TRANSLOCATION OF ORGANIC SUBSTANCES: PLANTS (Continued)

Part III HERBICIDES AND GROWTH REGULATORS (Continued)

Plant	Part	Substance	Translocation		Remarks	Reference
			Rate cm/hr	Direction		
(A)	(B)	(C)	(D)	(E)	(F)	(G)
71 Mesquite (Prosopis juliflora) (concluded)	Leaf, stem	2, 5-D (butoxy-ethoxypropanol ester), 3, 4-D (butoxyethanol ester), 2, 2-dichloropropionic acid		A, B	Apical epinasty and injury to young leaves, also injury to basal leaves determined at intervals after application to central leaves. Acropetal movement far more intense than basipetal.	78
72 Oak (Quercus douglasii, Q wislizenii) Ceanothus (Ceanothus cuneatus) Coyote brush (Baccharis pilularis) Manzanita (Arctostaphylos manzanita) Toyon (Photinia arbutifolia) Willow (Salix lasiolepis)	Leaf, stem	2, 4-D, urea		A ¹ , B ¹ , 4	Autoradiography of bark samples collected 1-21 da after treatment of single leaf with C ¹⁴ -labelled compound, at distances up to 150 cm from leaf. Acropetal transport less pronounced than basipetal, and dependent on plant species and season. Transport active many months in evergreen species, but only for short period in deciduous. Contact injury inhibited, and soil mixture enhanced, translocation.	79
73 Oat (Avena sativa)	Coleoptile	3-Indoleacetic acid		Lateral	After 1 hr of symmetrical application to cut surface, direct electrical current was applied across coleoptile for 2 min. Subsequent curvature indicated lateral transport.	80
74	Coleoptile	3-Indoleacetic acid	0.8-1.5	B	Donor agar blocks, containing 3-indoleacetic acid, and receptor blocks applied to ends of 2 mm sections; amount translocated determined by bioassay. Velocity of transport not wholly independent of temperature. Quantity increased from 0-40°C.	81
75	Coleoptile	3-Indoleacetic acid	0.4-0.8	B	Basal translocation, as determined by donor-receptor block method in 5 mm sections, inhibited by potassium cyanide and dinitrophenol.	82
76	Coleoptile	3-Indoleacetic acid		B	By using donor blocks (containing C ¹⁴ -labelled 3-indoleacetic acid), and analyzing receptor blocks (radio- and bioassay) after 1-3 hr transport through sections, it was shown that donor activity increased as original donor concentration increased up to 1.6 mg/L, whereas receptor activity reached maximum at a lower original donor concentration. Such activity indicates saturation of transport capacity at low concentrations.	83

1/3/ Stem, 1/4/ Leaf.

Part III. HERBICIDES AND GROWTH REGULATORS (Continued)

Plant	Part	Substance	Translocation		Remarks	Reference
			Rate cm/hr	Direction		
(A)	(B)	(C)	(D)	(E)	(F)	(G)
77 Oat (<i>Avena sativa</i>) (concluded)	Coleoptile	Native auxin	1.0-1.5	B	Calculation of transport rate based on various phototropic and geotropic responses, plus curvatures induced by applied auxin.	84
78 Pea (<i>Pisum sativum</i>)	Stem	Gibberellic acid	0.2	A, B	Donor-receptor block technique, utilizing 5-mm etiolated internode sections in conjunction with both photometric- and bioassay. No polarity. Acropetal and basipetal transport approximately equal.	85
79	Epicotyl	Gibberellic acid	60-600	A	Application of gibberellic acid to basal end of 100-mm etiolated section, subsequently followed by 3-indoleacetic acid application to a 5-mm section excised from apical end. Growth of section greater than that of controls receiving no gibberellic acid. Data resulting from adjustment of original section length and time interval before excision of apical section suggest rapid gibberellic acid transport and synergistic effect with 3-indoleacetic acid.	86
80 Pea (<i>Pisum sativum</i>) Tomato (<i>Lycopersicon esculentum</i>)	Leaf, stem, root	2, 4-D		A ³ , B ³ , 4	Basipetal movement more extensive than acropetal, as determined by radioassay at various intervals after application of C ¹⁴ -labelled compound to lower leaf. Of 2 active metabolic products of 2, 4-D, 1 was more common in pea and the other in tomato.	87
81 Peanut (<i>Arachis hypogaea</i>)	Gynophore	3-Indoleacetic acid	0.9	B	Short sections (3-5 mm) excised from gynophore just beneath ovule. Translocation determined by donor-receptor agar block method at 25°C, activity in blocks being determined by <i>Avena</i> test.	88
82 Potato (<i>Solanum tuberosum</i>)	Leaf, stem, root, tuber	Gibberellic acid		B	Foliar application 4, 2 or 1 wk before harvest of tubers resulted in their premature sprouting, thus suggesting rapid translocation in plants.	89
83 Soybean (<i>Glycine max</i>)	Leaf, stem, root	2, 4-D (sodium salt); also P ³²			Transport (determined by growth repression after severing basal leaf at various intervals following application of substance) was reduced by low leaf carbohydrate content. Temperature coefficient for transport of 2, 4-D and P ³² (determined by autoradiography) showed Q ₁₀ values of approximately 2.	90

/3/ Stem. /4/ Leaf.

117. TRANSLOCATION OF ORGANIC SUBSTANCES. PLANTS (Continued)
Part III: HERBICIDES AND GROWTH REGULATORS (Continued)

Plant	Part	Substance	Translocation		Remarks	Reference
			Rate cm/hr	Direction		
(A)	(B)	(C)	(D)	(E)	(F)	(G)
84 Spider flower, giant (<i>Cleome spinosa</i>)	Stem	2, 4-D; 3-Indole- acetic acid; naphthalene- acetic acid		A, B	Application to stem in form of ring. Translocation determined by degree of rooting of cuttings taken above or below application point. Transport in both directions across stem girdle indicated principal movement in xylem.	91
85 Squash (<i>Cucurbita maxima</i>) Tomato (<i>Lycopersicon esculentum</i>)	Stem	Native auxin, 3-Indoleacetic acid	20	B	Squash: analysis (<i>Avena</i> test) of 3-Indoleacetic acid con- tent above and below girdled stem sections at specific distances from point of application. Tomato: both substances moved only basipetally in stem sections, as deter- mined by agar block tech- nique. Temp. = 24-26°C.	92
86 Strawberry (<i>Fragaria</i> sp)	Leaf, runner	1, 2-Dihydro- pyridazine-3, 6-dione (diethanol- amine salt)		B ¹	Activity detected by auto- radiography in runner tips 30 hr after application of C14-labelled compound to mature leaf. Substance could move through several crowns to concentrate in a runner tip, but did not move against growth gradi- ent back toward formed crown of plant.	93
87 Sugar cane (<i>Saccharum officinatum</i>)	Leaf, stalk, root	2, 4-D		A, B	Radioassay, after application of C14-labelled 2, 4-D to single leaf, showed very little translocation to growing tips of shoot and root and to young leaves, as compared to transloca- tion in bean.	94
88 Sunflower (<i>Helianthus</i> sp)	Hypocotyl	3-Indoleacetic acid		B	Donor-receptor agar block technique demonstrated only basipetal transport in hypocotyl sections. Inhibi- tion of chlorinated phenoxy- acetic acids, upon polar transport of 3-Indoleacetic acid (IAA), studied by applying IAA at apical end and inhibitor at basal end IAA content in receptor block determined by <i>Avena</i> test. Increased inhibition correlated with increased chlorination of phenoxy- acetic acids.	95,96
89 Sunflower (<i>Helianthus</i> sp) Tobacco (<i>Nicotiana tabacum</i>)	Stem	3-Indoleacetic acid		B	Transport measured in sun- flower epicotyl sections by donor-receptor block technique, after infusion of various substances. Polar movement inhibited by respiratory inhibitors (e.g., KCN) and SH-inactivat- ing compounds (e.g., iodoacetate). Tobacco sections, cultured in nutrient with tritiodibenzole acid, con- tained equal distribution of 3-Indoleacetic acid (IAA), whereas controls accumulated IAA in basal end, thus indicating nullification of polarity.	97,98

1/1 Maleic hydrazide. /4/ Leaf.

117. TRANSLOCATION OF ORGANIC SUBSTANCES: PLANTS (Continued)

Part III: HERBICIDES AND GROWTH REGULATORS (Continued)

Plant	Part	Substance	Translocation		Remarks	Reference
			Rate cm/hr	Direction		
(A)	(B)	(C)	(D)	(E)	(F)	(G)
90 Sweet potato (Ipomoea batatas)	Leaf, petiole	3-Indoleacetic acid		B	Application of 3-Indoleacetic acid to lamina; transport determined by anomalous phototropic and geotropic response of petiole. Triiodobenzoic acid, placed between application point and responsive area, negated response, indicating inhibition of transport.	99
91 Thistle, Canada (Cirsium arvense) Grass, Johnson (Sorghum halepense) Soybean (Glycine max)	Leaf, stem, root	3-Amino-1, 2, 4-triazole		A, B	Autoradiography and chromatography of plant parts after application of C ¹⁴ -labelled compound to leaf, or leaf axil, showed movement throughout plant in 30 hr. Metabolism rate of 3-amino-1, 2, 4-triazole variable in different species, as determined by appearance of a labelled unknown.	100
92 Tobacco (Nicotiana sp)	Stem	1, 2-Dihydro- pyridazine- 3, 6-dione ¹		B	Application to cut surface after topping resulted in basipetal transport, as determined by inhibition of adventitious buds and increased formation of carbohydrates in leaves developing subsequent to treatment.	101
93	Stem	3-Indoleacetic acid, native auxin		B	2, 6-Dichlorobenzoic acid and 2, 5-dibromobenzoic acid (and related compounds) inhibited polar auxin transport, as determined by distribution of callus on stem segments, pith cell enlargement, and by distribution of C ¹⁴ -labelled 3-indoleacetic acid.	102
94 Tomato (Lycopersicon esculentum)	Leaf	3-[p-Chloro- phenyl]-1, 1- dimethylurea		A ¹⁰ , B ¹¹	Application of C ¹⁴ -ring-labelled compound to terminal leaflet; subsequent movement followed by autoradiogram. Very little basipetal movement in leaves. Acropetal movement only when compound given via root system.	103
95	Leaf, stem, root	3-Indoleacetic acid, indole- butyric acid, naphthalene- acetic acid, phenylacetic acid, phenyl- propionic acid		A ¹⁰ , B ¹¹	Application of substance in lanolin to lower stem. Epinasty of stem and leaves developed more slowly in older tissue.	104
96	Leaf, stem, root	2, 4-D		A ³ , B ³ , 4	C ¹⁴ -labelled 2, 4-D applied to first true leaf of P-deficient plants. Translocation from leaf determined by stem curvature and autoradiography. Addition of P to nutrient 8 hr or 3 da before application increased translocation, particularly with 3-da pretreatment.	105

/1/ Maleic hydrazide. /3/ Stem. /4/ Leaf. /10/ Movement principally in xylem. /11/ Slight movement.

117. TRANSLOCATION OF ORGANIC SUBSTANCES. PLANTS (Continued)

PART III: HERBICIDES AND GROWTH REGULATORS (Continued)

	Plant	Part	Substance	Translocation		Remarks	Reference
				Rate cm/hr	Direction		
	(A)	(B)	(C)	(D)	(E)	(F)	(G)
97	Tomato (<i>Lycopersicon esculentum</i>) (concluded)	Leaf, stem, root	2, 4-D		B	K-deficient plants given K via roots 3 da before, or simultaneously with, 2, 4-D application to 1 leaflet showed greater transport than plants given K 24 hr after 2, 4-D, as evidenced by growth repression, formative effect, and autoradiography.	106
98		Leaf, stem, root	2, 4-D (sodium salt)		A, B	Movement throughout plant determined by photometric chemical analysis 24-48 hr after application to 2nd basal leaf or to soil.	107
99		Leaf, stem, root	Gibberellic acid		A, B	Application to soil, individual leaves, stem apices, or flower peduncles, caused increase in parthenocarpic fruit set, suggesting translocation in either phloem or xylem.	108

Contributor: Hull, Herbert M

- References** [1] Earle, T. T., K. Rieas, and J. Hidalgo. 1951. *Science* 114:695. [2] Mitchell, J. W., P. C. Marth, and W. H. Preston, Jr. 1954. *Ibid.* 120:263. [3] Oserkowsky, J. 1942. *Am. J. Bot.* 29:858. [4] Gregory, F. G., and C. R. Hancock. 1955. *Ann. Bot.*, Lond. 19:451. [5] Linden, G. 1954. *Beitr. Biol. Pflanz.* 30:343. [6] Mitchell, J. W., and W. H. Preston, Jr. 1953. *Science* 118:518. [7] Crafts, A. S., and S. Yamaguchi. 1958. *Hilgardia* 27:421. [8] Crafts, A. S., H. B. Currier, and H. R. Drever. 1958. *Ibid.* 27:723. [9] Yamaguchi, S., and A. S. Crafts. 1957. *Plant Physiol.*, Suppl. 32:42. [10] Hay, J. R. 1956. *Plant Physiol.* 31:118. [11] Jacobs, W. P. 1950. *Am. J. Bot.* 37:248. [12] Weintraub, R. L., and J. W. Brown. 1950. *Plant Physiol.* 25:140. [13] Fang, S. C., V. H. Freed, R. H. Johnson, and D. R. Coffee. 1955. *J. Agr. Food Chem.* 3:400. [14] Wood, J. W., J. W. Mitchell, and G. W. Irving, Jr. 1947. *Science* 105:337. [15] Linder, P. J., J. W. Mitchell, and J. W. Wood. 1950. *Ibid.* 111:518. [16] Mitchell, J. W., and P. J. Linder. 1950. *Ibid.* 112:54. [17] Linder, P. J., J. W. Brown, and J. W. Mitchell. 1949. *Bot. Gaz.* 110:628. [18] Fang, S. C., E. G. Jaworski, A. V. Logan, V. H. Freed, and J. S. Butts. 1951. *Arch. Biochem.*, N. Y. 32:249. [19] Hay, J. R., and K. V. Thimann. 1956. *Plant Physiol.* 31:446. [20] Holley, R. W. 1952. *Arch. Biochem.*, N. Y. 35:171. [21] Holley, R. W., F. P. Boyle, and D. B. Hand. 1950. *Ibid.* 27:143. [22] Ashton, F. M. 1958. *Plant Physiol.*, Suppl. 33:39. [23] Hay, J. R. 1955. *Ibid.* 30:5. [24] Weintraub, R. L., J. W. Brown, and J. N. Yeatman. 1950. *Science* 111:493. [25] Weintraub, R. L., and J. W. Brown. 1950. *Am. J. Bot.* 37:682. [26] Day, B. E. 1952. *Plant Physiol.* 27:143. [27] Burrows, V. D., and J. Bonner. 1958. *Plant Physiol.*, Suppl. 33:21. [28] Rice, E. L., and L. M. Rohrbaugh. 1953. *Bot. Gaz.* 115:76. [29] Jaworski, E. G., S. C. Fang, and V. H. Freed. 1955. *Plant Physiol.* 30:272. [30] Rice, E. L. 1948. *Bot. Gaz.* 109:301. [31] Mitchell, J. W., W. M. Dugger, Jr., and H. G. Gauch. 1953. *Science* 118:354. [32] Crafts, A. S. 1953. *J. Agr. Food Chem.* 1:51. [33] Fang, S. C., and J. S. Butts. 1957. *Plant Physiol.* 32:253. [34] Pallas, J. E. 1958. *Plant Physiol.*, Suppl. 33:21. [35] Zwar, J. A., and A. H. G. C. Rijven. 1956. *Austral. J. Biol. Sc.* 9:528. [36] Linder, P. J., J. C. Craig, Jr., and T. R. Walton. 1957. *Plant Physiol.* 32:572. [37] Linder, P. J., J. C. Craig, Jr., F. E. Cooper, and J. W. Mitchell. 1958. *J. Agr. Food Chem.* 6:356. [38] Watanabe, R., and N. J. Scully. 1957. *Plant Physiol.*, Suppl. 32:56. [39] Marth, P. C., W. V. Audia, and J. W. Mitchell. 1956. *Bot. Gaz.* 118:106. [40] Mitchell, J. W., J. W. Wirwille, and L. Weil. 1949. *Science* 110:252. [41] Racusen, D. 1958

- Arch. Biochem., N. Y. 74:106. [42] Crafts, A. S. 1956. *Hilgardia* 26:287. [43] Crafts, A. S. 1956. *Ibid.* 26:335. [44] Weintraub, R. L., J. H. Reinhart, and R. A. Scherff. 1956. A conference on radioactive isotopes in agriculture. Atomic Energy Commission. Rpt. No. TID-7512. pp. 203-08. [45] Weaver, R. J., and H. R. De Rose. 1946. *Bot. Gaz.* 107:509. [46] Curtis, R. W. 1957. *Plant Physiol.* 32:56. [47] Mitchell, J. W., J. W. Wood, W. C. Wolfe, and G. W. Irving, Jr. 1947. *Science* 106:395. [48] Gallup, A. H., and F. G. Gustafson. 1952. *Plant Physiol.* 27:603. [49] Doxey, D. 1953. *J. Exp. Bot.* 4:53. [50] Ashton, F. M. 1958. *Weeds* 6:257. [51] Amlong, H. U. 1939. *Jb. wiss. Bot.* 88:421. [52] Gorter, C. J. 1936. *Biol. Jaarboek, Antwerp* 3:205. [53] Syre, H. Z. 1938. *Zschr. Bot.* 33:129. [54] Muzik, T. J., H. J. Cruzado, and A. J. Loustalot. 1954. *Bot. Gaz.* 116:65. [55] Neely, P. M., and B. O. Phinney. 1957. *Plant Physiol., Suppl.* 32:31. [56] Naylor, A. W., and E. A. Davis. 1950. *Bot. Gaz.* 112:112. [57] Leopold, A. C., and F. S. Guernsey. 1953. *Ibid.* 115:147. [58] Jacobs, W. P. 1954. *Am. Natur.* 88:327. [59] Hall, W. C., S. P. Johnson, and C. L. Leinweber. 1954. *Texas Agr. Exp. Sta. Bull.* 789, College Sta. [60] Clor, M. A., and A. S. Crafts. 1957. *Plant Physiol., Suppl.* 32:43. [61] Crafts, A. S., H. B. Currier, and B. E. Day. 1950. *Hilgardia* 20:57. [62] Laibach, F., and O. Fischnich. 1937. *Planta* 26:81. [63] Darwin, C. 1880. The power of movement in plants. John Murray, London. p. 592. [64] Hauser, E. W., and J. Thompson. 1954. *J. Agr. Food Chem.* 2:680. [65] Anderson, O. 1958. *Weeds* 6:370. [66] Carvell, K. L. 1955. *Forest Sc.* 1:41. [67] Cooper, W. C. 1936. *Plant Physiol.* 11:779. [68] Cholodny, N. 1934. *Planta* 21:517. [69] Baldwin, R. E., V. H. Freed, and S. C. Fang. 1954. *J. Agr. Food Chem.* 2:428. [70] Went, F. W., and R. White. 1939. *Bot. Gaz.* 100:465. [71] Blanchard, F. A. 1954. *Weeds* 3:274. [72] Fang, S. C., and J. S. Butts. 1954. *Plant Physiol.* 29:56. [73] Hay, J. R. 1956. *Weeds* 4:218. [74] Hay, J. R. 1956. *Ibid.* 4:349. [75] Fisher, C. E., C. H. Meadors, and R. Behrens. 1956. *Ibid.* 4:139. [76] Hull, H. M. 1957. *Plant Physiol., Suppl.* 32:43. [77] Blair, B. O., and W. H. Fuller. 1952. *Bot. Gaz.* 113:368. [78] Hull, H. M. 1956. *Weeds* 4:22. [79] Leonard, O. A., and A. S. Crafts. 1956. *Hilgardia* 26:366. [80] Webster, W. W., Jr., and R. A. Schrank. 1953. *Arch. Biochem., N. Y.* 47:107. [81] Van der Wel, H. G. 1932. *Rec. Tr. bot. néerl.* 29:379. [82] DuBuy, H. G., and R. A. Olson. 1940. *Am. J. Bot.* 27:401. [83] Goldsmith, M. H. M. 1958. *Plant Physiol., Suppl.* 33:39. [84] DuBuy, H. G., and E. Nuernbergk. 1932. *Erg. Biol.* 9:358. [85] Kato, J. 1958. *Science* 128:1008. [86] Galston, A. W. 1958. *Plant Physiol., Suppl.* 33:39. [87] Fang, S. C. 1958. *Weeds* 6:179. [88] Jacobs, W. P. 1951. *Am. J. Bot.* 38:307. [89] Lippert, L. F., L. Rappaport, and H. Timm. 1958. *Plant Physiol.* 33:132. [90] Barrier, G. E., and W. E. Loomis. 1957. *Ibid.* 32:225. [91] Ferri, M. G. 1945. *Contr. Boyce Thompson Inst.* 14:51. [92] Skoog, F. 1938. *Am. J. Bot.* 25:361. [93] Brown, M. S., and C. W. Hitz. 1957. *Proc. Am. Soc. Horticul. Sc.* 70:131. [94] Burr, G. O., T. Tanimoto, C. E. Hartt, A. Forbes, G. Sadaoka, F. M. Ashton, J. H. Payne, J. A. Silva, and G. E. Sloane. 1956. *Proc. Internat. Conf. Peaceful Uses Atomic Energy, Geneva* 12:177. *United Nations, New York.* [95] Von Guttenberg, H., and I. Eifler. 1952-53. *Ber. deut. bot. Ges.* 65:387. [96] Leopold, A. C., and E. Niedergang-Kamlen. 1958. *Plant Physiol., Suppl.* 33:21. [97] Niedergang-Kamlen, E., and A. C. Leopold. 1957. *Physiol. Plant* 10:29. [98] Niedergang-Kamlen, E., and F. Skoog. 1956. *Ibid.* 9:60. [99] Kuse, G. 1953. *Mem. Coll. Sc. Univ. Kyoto* SB20:207. [100] Rogers, B. J. 1957. *Weeds* 5:5. [101] Arnaud, J., P. Barge, M. Richez, and R. J. Gautheret. 1956. *C. rend. Acad. agr. France* 42:168. [102] Kelitt, G. W., Jr., and F. Skoog. 1957. *Plant Physiol., Suppl.* 32:20. [103] Haun, J. R., and J. H. Peterson. 1954. *Weeds* 3:177. [104] Hitchcock, A. E., and P. W. Zimmerman. 1935. *Contr. Boyce Thompson Inst.* 7:447. [105] Rohrbaugh, L. M., and E. L. Rice. 1956. *Plant Physiol.* 31:196. [106] Rice, E. L., and L. M. Rohrbaugh. 1958. *Ibid.* 33:300. [107] Dhillon, A. S., and E. H. Lucas. 1950. *Bot. Gaz.* 112:198. [108] Persson, A., and L. Rappaport. 1958. *Science* 127:816.

117. TRANSLOCATION OF ORGANIC SUBSTANCES PLANTS (Continued)
Part IV: VIRUSES

Plant	Part	Substance	Translocation		Remarks	Reference
			Rate cm/hr	Direction		
(A)	(B)	(C)	(D)	(E)	(F)	(G)
1 Bean (Phaseolus vulgaris)	Leaf, stem	Virus causing southern bean mosaic			Virus moved from inoculated leaf down stem, across approach graft, and up stem of 2nd plant through a steamed section (as evidenced by necrotic areas above steamed region), thus suggesting movement in tracheal elements.	1
2 Bean (Phaseolus vulgaris) Tobacco (Nicotiana tabacum)	Leaf, stem	Virus causing tobacco mosaic, tobacco ring spot, cucumber mosaic; potato virus X		A, B	Virus moved distally out of darkened (starch free), inoculated leaves as sugars moved in (indicated by accumulation of starch in such leaves, and by recovery of virus and inoculation to appropriate indicator plants).	2
3 Beet, sugar (Beta vulgaris)	Leaf	Virus causing sugar beet curlytop	9-36	B	1 leaf inoculated by leafhopper (Cerculifer tenellus), plant infectivity determined by excising petiole base at various intervals after inoculation Temp. = 34-40°C.	3
4	Leaf	Virus causing Argentine curlytop	15	B	Infectivity of plant determined after viruliferous leafhoppers (Agalliana ensigera) had fed 2 hr on leaf tip, leaf severed 15 cm below point of feeding. Temp. = 38°C.	4
5 Beet, sugar (Beta vulgaris) Tobacco (Nicotiana tabacum)	Leaf, cotyledon, stem	Virus causing sugar beet curlytop	Beet leaf: 25-152 Beet cotyledon: 30 at 16°C 51 at 29°C 76 at 43°C 76 at 57°C Tobacco 1.3	B	Tobacco. Viruliferous leafhoppers (Cerculifer tenellus) allowed to feed on youngest leaves. Stem later segmented, and segments rooted and observed for disease. No movement from apex during first 24 hr after feeding. Sugar beet. Leafhoppers allowed to feed on leaf or cotyledon tips, these organs then severed at specific times and measured at intervals below point of feeding.	5
6 Dodder (Cuscuta aubinculosa)	Stem	Virus causing cucumber mosaic	0.2-10.2	A	When dodder stem was connected in line to 3 tobacco plants, only center plant infected; movement of virus almost entirely to healthy plant at apical end of dodder. More rapid movement shown by leafhopper feeding experiments.	6
7 Globe amaranth (Gomphrena globosa) Potato (Solanum tuberosum) Tobacco (Nicotiana tabacum)	Leaf, stem	Potato viruses X and Y	0.2-0.71 0.004-0.012		Application over leaf veins resulted in more rapid movement along a 6-mm path than with application over intercostal areas. In stem, Y virus moved faster than X, possibly due to inactivation of X virus in phloem.	7
8 Maize (Zea mays)	Leaf	Virus causing maize streak	20	B	Inoculation at distal end of 1 leaf with viruliferous leafhopper (Cicadulina mbila), after which leaf was severed at known distance and plant retained for observation of infectivity. Temp. = 35°C.	8

1/1 Stem 1/2 Leaf

Part IV: VIRUSES (Continued)

Plant	Part	Substance	Translocation		Remarks	Reference
			Rate cm/hr	Direction		
(A)	(B)	(C)	(D)	(E)	(F)	(G)
9 Peach (<i>Prunus persica</i>)	Stem	Virus causing yellow-red or X disease of peach		A	Translocation of virus, from diseased bud placed on stem to apical portion of plant, could be increased by darkening upper half of plant for 2 wk, as shown by symptomatic appearance.	9
10 Potato (<i>Solanum tuberosum</i>)	Leaf, stem, rhizome,	Virus causing leaf roll		B	Per cent of infected tubers at various intervals after inoculation of leaf: 6 da, 0%; 10 da, 100%	10
11 obconica)		necrosis			did not cause full systemic infection as in certain other plants (only isolated islets of infected tissue appeared). In some cases presence could be detected only by transmission and serological tests. Temp. * 16-27°C.	
12 Raspberry (<i>Rubus</i> sp)	Cane	Virus causing raspberry leaf curl		A, B	An inoculated cane did not rapidly transmit virus to other canes on same plant unless canes were cut back. Assimilates essential for regrowth (secured from the infected cane) during their movement transported virus to new canes or suckers.	12
13 Tobacco (<i>Nicotiana tabacum</i>)	Normal callus tissue	Virus causing tobacco mosaic	0.0005		Transport rate determined by inoculating 1 end of strips cut from tissue cultures of normal callus. At intervals up to 32 da, strips cut into 4 pieces and each piece sub-cultured and tested for infectivity on <i>Nicotiana glutinosa</i> leaves.	13
14	Leaf, stem, root	Virus causing tobacco mosaic	0.02 ³ 0.04 ⁴ 0.8 ⁵		Virus inoculated into leaf moved at rates shown, and finally much faster in stem phloem, as deter- mined by cytological abnormali- ties. Virus may be transported not as complete entity, but as smaller, non-infective, precursory particles which are not capable of combining in sieve elements, but only in adjacent parenchyma.	14
15 Tobacco (<i>Nicotiana sylvestris</i>)	Leaf	Virus causing tobacco mosaic	0.0007-0.0009		Inoculation to upper leaf surface resulted in movement to lower surface in 36-40 hr at rate shown.	15
16 Tobacco (<i>Nicotiana sp</i>)	Leaf	Virus causing tobacco mosaic	0.1-0.3	B	Leaf severed at specific distance and time after inoculation of virus, plant observed for devel- opment of disease symptoms.	16
17 Tomato (<i>Lycopersicon esulentum</i>)	Leaf, stem, root	Virus causing tobacco mosaic	0.2-0.5	A, B	Tobacco inoculum applied to single basal quarter of central leaf. Movement to apical leaves occurred in 3 da, and to roots in 5 da. Spread of virus to other quarters of inoculated leaf took 2-3 wk at 24°C. Tomato Movement to extreme ends of plant occurred 6 da after inoculation of central leaf. Sectioning of stem, and subsequent germination of cuttings, indicated lack of virus concentration in any specific stem area.	17

1/ Stem. 2/ Leaf. 3/ Intercostal epidermis. 4/ Elongated epidermal cells along veins. 5/ Elongated bundle sheath cells.

Part IV: VIRUSES (Continued)

Plant	Part	Substance	Translocation		Remarks	Reference
			Rate cm/hr (D)	Direction (E)		
(A)	(B)	(C)	(D)	(E)	(F)	(G)
18 Tobacco (Nicotiana sp) Tomato (Lycopersicon esculentum) (concluded)	Leaf, stem, root	Virus causing tobacco mosaic, cucumber mosaic, tobacco streak, severe etch (tobacco), tobacco mosaic, cucumber mosaic, potato ring spot (tomato)	0.4	A, B	Application of inoculum to surface of central portion of root resulted essentially in movement toward root tip, as determined by later maceration of tissue and inoculation to similar host. Inoculation at stem base resulted, however, in movement to both aerial plant parts and roots.	18
19	Leaf, stem	Virus causing tomato auburn mosaic, tobacco mosaic (tomato, tobacco)	0.3	A, B	Virus inoculated on either apical or basal leaf did not pass to opposite end of plant if central internode had been steamed, suggesting movement in phloem. Tomato plants, inoculated on 1 leaf near center of stem, were cut into sections after 3 da and planted as cuttings. All cuttings germinated, but those apical to inoculated leaf showed greatest infectivity.	19
20 Tomato (Lycopersicon esculentum)	Stem	Virus causing tomato mosaic	0.1-0.2	A, B	Plants, having lateral branches layered and rooted in separate pots, were inoculated at apex of main stem. Secondary plants severed from inoculated parent plant after 3-19 da and observed for development of disease. Half of plants received virus if allowed to remain attached to parent 10 da.	20
21	Leaf, stem, root	Virus causing tomato mosaic	18	A, B	44 hr was minimum time for movement of virus from inoculated leaf throughout stem. Upon reaching stem, virus generally translocated bi-directionally, but occasionally moved only up or down. Presence was determined by inoculating tissue extracts on tobacco test plants.	21
22	Leaf, stem, root	Virus causing tobacco mosaic	1-4	A, B	Virus moved out of central inoculated leaf after 3 da, then to root in less than 12 hr, and finally rapidly to top. Virus moved to fruit 1-2 da after leaving inoculated leaf. Translocation determined by sectioning stem at various intervals, extracting sections, and inoculating test plant with extract.	22
23	Leaf, stem	Potato virus X; virus causing tomato auburn mosaic, tobacco mosaic	7-8	A, B	3-5 da required before virus exported from inoculated leaf, but moved bi-directionally in stem within 12 hr of entry. Movement determined by progress of disease symptoms. Temp. = 11-31°C.	23

Part IV: VIRUSES (Concluded)

Contributor: Hull, Herbert M.

References: [1] Schneider, I. R., and J. F. Worley. 1958. *Science* 127:1050. [2] Roberts, D. A. 1952. *Phytopathology* 42:381. [3] Severin, H. H. P. 1924. *Ibid.* 14:80. [4] Bennett, C. W., E. Carsner, G. H. Coons, and E. W. Brandes. 1946. *J. Agr. Res.* 72:19. [5] Bennett, C. W. 1934. *Ibid.* 48:665. [6] Bennett, C. W. 1944. *Phytopathology* 34:905. [7] Köhler, E. 1956. *Phytopathol. Zschr.* 26:147. [8] Storey, H. H. 1928. *Ann. Appl. Biol., Lond.* 15:1. [9] Hildebrand, E. M., and O. F. Curtis. 1942. *Science* 95:390. [10] De Montgrier, H. A. 1954. *Extr. Soc. Emul. Abbeville* 17:11. [11] Bawden, F. C., and B. Kassanis. 1947. *Ann. Appl. Biol., Lond.* 34:127. [12] Bennett, C. W. 1927. *Michigan Agr. Exp. Sta. Tech. Bull.* 80:26. [13] Kassanis, B., T. W. Tinsley, and F. Quak. 1958. *Ann. Appl. Biol., Lond.* 46:11. [14] Zech, H. 1952. *Planta* 40:461. [15] Uppal, B. N. 1934. *Ind. J. Agr. Sc.* 4:865. [16] Böning, K. 1928. *Zschr. Pflanzenerk.* 1:198. [17] Holmes, F. O. 1930. *Am. J. Bot.* 17:789. [18] Fulton, R. W. 1941. *Phytopathology* 31:575. [19] Caldwell, J. 1931. *Ann. Appl. Biol., Lond.* 18:279. [20] McCubbin, W. A., and F. F. Smith. 1927. *Science* 66:486. [21] Kunkel, L. O. 1939. *Phytopathology* 29:684. [22] Samuel, G. 1934. *Ann. Appl. Biol., Lond.* 21:90. [23] Caspoor, S. P. 1949. *Ibid.* 36:307.

Part V: FLORIGEN

Plant	Part	Substance	Translocation		Remarks	Reference
			Rate cm/hr	Direction		
(A)	(B)	(C)	(D)	(E)	(F)	(G)
1 Beet, sugar (<i>Beta vulgaris</i>)	Leaf, stem	Florigen		A, B	Separate shoots of single plant subjected to different day lengths, those receiving short days did not flower. Shoots kept in dark flowered when others on same plant were exposed to long days, thus suggesting movement of florigen into light-starved parts along with assimilates.	1
2 Cocklebur (<i>Xanthium pennsylvanicum</i>)	Leaf, stem	Florigen		A, B	Movement basipetally from receptor leaves subjected to short photoperiod, then transportation acropetally to plant parts maintained on long photoperiod, as indicated by induction of floral primordia. Florigen traversed approach graft, even when plants were separated by lens paper.	2
3	Leaf, stem	Florigen		A, B	Failure to translocate out of photoperiodically-induced donor branch of 2-branch plant, if either branch was girdled immediately above point of branching, thus demonstrating acropetal and basipetal movement in phloem (Floral primordia was induced on opposite branch in case of non-girdled plants.)	3
4	Leaf, stem	Florigen		A ¹ , B ²	Plants defoliated to 1 leaf and exposed to 1 short day; remaining leaf detached at various time intervals after end of 15-hr dark period. Maximum response in young leaves appeared to occur if treated leaf was retained on plant for 16 hr or more after termination of long dark period.	4
5	Leaf, stem	Florigen		A ¹ , B ²	Movement out of leaves, during 13 hr following inductive dark period, was considerably depressed if leaves were not exposed to a high intensity light. Depressing effect of light absence could be overcome by supplying sucrose to leaves.	5

[1] Stem. [2] Leaf.

117. TRANSLOCATION OF ORGANIC SUBSTANCES- PLANTS (Continued)

	Plant	Part	Substance	Translocation		Remarks	Reference
				Rate cm/hr	Direction		
	(A)	(B)	(C)	(D)	(E)	(F)	(G)
6	Cocklebur (Xanthium saccharatum)	Leaf, stem	Florigen		A ¹ , B ²	Although single short-day induction period produced floral primordia, light given after long dark period increased movement from leaf to apical meristem. Without light, induction was achieved only after 8 da of continuous darkness.	6
7	Cocklebur (Xanthium commune) Soybean (Glycine max)	Leaf, stem	Florigen (also assimilates)	0.33	A ¹ , B ²	Cooling stem segments to 3-5°C, or encasing with N ₂ , did not inhibit rate of florigen transport, although amount of assimilates was inhibited, and possibly the amount of florigen.	7
8	Soybean (Glycine max)	Leaf, stem	Florigen		A, B	Induction of floral primordia, as result of translocation from photoperiodically-induced leaf, was inhibited by scalding petiole of such leaf, suggesting movement in phloem.	8
9		Leaf, stem	Florigen		A, B	Chilling petiole of photoperiodically-induced leaf to 3°C inhibited outward movement of florigen, as determined by subsequent induction of floral primordia. Chilling to 5-6°C gave less inhibition, and at 10°C translocation was almost equal to that of controls at 25°C.	9
10	Su-Tsa (Perilla ocymoides)	Leaf	Florigen		B	Florigen formed in leaves exposed to short photoperiods and passed to shoots both via leaf veins and leaf parenchyma. Cutting leaf vein stopped outflow of nutrients, but not florigen, from leaf.	10

Contributor: Hall, Herbert M

Part VI. NITROGENOUS SUBSTANCES AND VITAMINS

Part IV: VIRUSES (Concluded)

Contributor: Hull, Herbert M.

- References. [1] Schneider, I. R., and J. F. Worley. 1958. Science 127:1050. [2] Roberts, D. A. 1952. Phytopathology 42:381. [3] Severin, H. H. P. 1924. Ibid. 14:80. [4] Bennett, C. W., E. Carsner, G. H. Coons, and E. W. Brandes. 1946. J. Agr. Res. 72:19. [5] Bennett, C. W. 1934. Ibid. 48:665. [6] Bennett, C. W. 1944. Phytopathology 34:905. [7] Köhler, E. 1956. Phytopathol. Zschr. 26:147. [8] Storey, H. H. 1928. Ann. Appl. Biol., Lond. 15:1. [9] Hildebrand, E. M., and O. F. Curtis. 1942. Science 95:390. [10] De Montgrentier, H. A. 1954. Extr. Soc. Emul. Abbeville 17:11. [11] Bawden, F. C., and B. Kassanis. 1947. Ann. Appl. Biol., Lond. 4:127. [12] Bennett, C. W. 1927. Michigan Agr. Exp. Sta. Tech. Bull. 80:26. [13] Kassanis, B., T. W. Tinsley, and F. Quak. 1958. Ann. Appl. Biol., Lond. 46:11. [14] Zech, H. 1952. Planta 40:461. [15] Uppal, B. N. 1934. J. Agr. Sc. 4:865. [16] Böning, K. 1928. Zschr. Parasitenk. 1:198. [17] Holmes, F. O. 1930. Am. J. Bot. 17:789. [18] Fulton, R. W. 1941. Phytopathology 31:575. [19] Caldwell, J. 1931. Ann. Appl. Biol., Lond. 18:279. [20] McCubbin, W. A., and F. F. Smith. 1927. Science 66:486. [21] Kunkel, L. O. 1939. Phytopathology 29:684. [22] Samuel, G. 1934. Ann. Appl. Biol., Lond. 21:90. [23] Capoor, S. P. 1949. Ibid. 36:307.

Part V: FLORIGEN

	Plant	Part	Substance	Translocation		Remarks	Reference
				Rate cm/hr	Direction		
	(A)	(B)	(C)	(D)	(E)	(F)	(G)
1	Beet, sugar (Beta vulgaris)	Leaf, stem	Florigen		A, B	Separate shoots of single plant subjected to different day lengths; those receiving short days did not flower. Shoots kept in dark flowered when others on same plant were exposed to long days, thus suggesting movement of florigen into light-starved parts along with assimilates.	1
2	Cocklebur (Xanthium pennsylvanicum)	Leaf, stem	Florigen		A, B	Movement basipetally from receptor leaves subjected to short photoperiod, then transportation acropetally to plant parts maintained on long photoperiod, as indicated by induction of floral primordia. Florigen traversed approach graft, even when plants were separated by lens paper.	2
3		Leaf, stem	Florigen		A, B	Failure to translocate out of photoperiodically-induced donor branch of 2-branch plant, if either branch was girdled immediately above point of branching, thus demonstrating acropetal and basipetal movement in phloem. (Floral primordia was induced on opposite branch in case of non-girdled plants.)	3
4		Leaf, stem	Florigen		A ¹ , B ²	Plants defoliated to 1 leaf and exposed to 1 short day; remaining leaf detached at various time intervals after end of 15-hr dark period. Maximum response in young leaves appeared to occur if treated leaf was retained on plant for 16 hr or more after termination of long dark period.	4
5		Leaf, stem	Florigen		A ¹ , B ²	Movement out of leaves, during 13 hr following inductive dark period, was considerably depressed if leaves were not exposed to a high intensity light. Depressing effect of light absence could be overcome by supplying sucrose to leaves.	5

1/ Stem. 2/ Leaf.

117. TRANSLOCATION OF ORGANIC SUBSTANCES. PLANTS (Continued)
Part VI: NITROGENOUS SUBSTANCES AND VITAMINS (Concluded)

Plant	Part	Substance	Translocation		Remarks	Reference
			Rate cm/hr	Direction		
(A)	(B)	(C)	(D)	(E)	(F)	(G)
11 Tobacco (<i>Nicotiana tabacum</i>)	Leaf, stem, root	Urea		A2, 3, B2, 3	Radioassay after application of N ¹⁵ . labelled urea to single leaf showed half of absorbed N ¹⁵ transported from leaf in 24 hr. Activity present after 6 hr in leaf opposite to treated leaf, as well as in stem and root, may indicate transfer from phloem to xylem.	11
12 Tomato (<i>Lycopersicon esculentum</i>)	Leaf, stem	Total nitrogen, non-protein nitrogen, sucrose, thiamine, pyridoxine, pantothenic acid, ribo- flavin		B	Compounds accumulated above steamed section of stem at 2nd node in 1-4 da. Chemical analysis of bioassay for vitamins indicated that sucrose accumulated mostly above steamed section, with other compounds accumu- lating in varying degrees.	12
13	Leaf, stem, root	Thiamine		A, B	Bioassay of various plant parts, 1-6 da after girdling different stem loci or petioles by steam, indicated older leaves have low thiamine concentration, transport greatest to young leaves and roots.	13
14 Wheat (<i>Triticum</i> sp)	Stem	Asparagine		A	Excised stem sections (10-35 cm long) placed with upper or lower ends in solu- tion. After 20 min, analysis of 5-cm pieces cut from each end showed N accumulation in morphological upper end regardless of which end was immersed, suggesting polar transport.	14

[2] Stem. [3] Leaf.

Contributor: Hall, Herbert M

References. [1] Bond, G. 1956. J. Exp. Bot. 7:387. [2] Ziegler, H. 1956. Planta 47:447. [3] Kazarian, V. O., and E. S. Avundzhyan. 1955. Doklady akad. nauk. S.S.S.R. 101:181. [4] Kazarian, V. O., and N. V. Balagerian. 1955. Ibid 103:337. [5] Nothes, K., and L. Engelbrecht. 1954. Flora 141:356. [6] Maskell, E. J., and T. G. Mason. 1929. Ann. Bot., Lond 43:205. [7] Hay, R. E., E. B. Easley, and E. E. de Turk. 1953. Plant Physiol. 28:606. [8] Kursanov, A. L. 1956. Proc. Internat. Conf. Peaceful Uses Atomic Energy, Geneva 12:165. United Nations, New York. [9] Nelson, C. D., and P. R. Gorham. 1958. Plant Physiol., Suppl. 33:21. [10] Arisz, W. H. 1953. Acta bot. neri 2:74. [11] Volk, R., and C. McAuliffe. 1954. Proc. Soil Sc. Soc. America 18:308. [12] Bonner, J. 1944. Am. J. Bot. 31:551. [13] Bonner, J. 1942. Ibid. 29:136. [14] Kursanov, A. L., and M. N. Zaprometov. 1949. Doklady akad. nauk. S.S.S.R. 69:89.

117. TRANSLOCATION OF ORGANIC SUBSTANCES. PLANTS (Continued)

Part VI. NITROGENOUS SUBSTANCES AND VITAMINS (Continued)

Plant	Part	Substance	Translocation		Remarks	Reference
			Rate cm/hr	Direction		
(A)	(B)	(C)	(D)	(E)	(F)	(G)
3 Ash (<i>Fraxinus</i> sp) Honeysuckle (<i>Lonicera</i> sp) Lilac (<i>Syringa</i> sp) Loxa (<i>Cinchona</i> sp)	Leaf, twig	Chlorophyll precursors		A	Darkening twig tips during spring development of leaves did not completely inhibit formation of chlorophyll, even when twigs were previously cut off. Transport of chlorophyll precursors into darkened leaves was apparently from phellogen cells, which simultaneously decreased in chlorophyll content.	3
4 Chrysanthemum (<i>Chrysanthemum</i> sp) Grass, pepper (<i>Lepidium</i> sp)	Stem, root	Glycocoll		A, B	By topping plants at various levels, applying glycocoll at cut surfaces, and following resultant absorption of N, polarity of movement was found to vary with developmental stage of plant and fruit.	4
5 Comfrey (<i>Symphytum officinale</i>)	Leaf, stem, root	Nitrogenous		A, B	Acropetal movement found to continue until fruiting; after fruiting, leaf proteins translocated basipetally and stored in roots largely in the form of allantoin. Root proteins did not translocate, being "protoplasmic proteins."	5
6 Cotton (<i>Gossypium</i> sp)	Leaf, stem	Nitrogenous		B ¹	Increase in total N in leaves during day and decrease during night, combined with accumulation of N above stem girdle and in semi-isolated bark flaps, suggested transport into leaves during day via xylem and from leaves both day and night via phloem.	6
7 Maize (<i>Zea mays</i>)	Leaf, stalk, grain	Nitrogenous		Into grain	Chemical analysis of various plant parts, during period of pollination to maturity, indicated plant lost 57% of total N to grain. Data suggested incorporation of nitrates with protein, or protein-like substances, and subsequent translocation to grain.	7
8 Pumpkin (<i>Cucurbita pepo</i>)	Leaf, stem	Amino acids (also sugars)		A ² , B ² , 3	1 leaf treated with known mixture of C ¹⁴ -labelled amino acids and sugars resulted in bi-directional movement on entry into stem. Relative movement of acids and sugars into fruit dependent on stage of development.	8
9 Soybean (<i>Glycine max</i>)	Petiole, stem	Alanine, arginine, asparagine, aspartic acid, glutamic acid, glutamine, glycine, serine, urea	350-1400	B	Transport measured by introducing C ¹⁴ -labelled substance through cut petiole, then sectioning plant after 5 min for assay. Rate different for every compound, but C ¹⁴ concentration always decreased logarithmically from point of introduction. Basipetal transport not inhibited by a 1-7 cm steam-killed stem section, but was by 16 cm section.	9
10 Sundew (<i>Drosera capensis</i>)	Tentacle	Amino acids (including asparagine, glutamine, urea, thiourea, caffeine)			Amino acids and glutamine absorbed and translocated through parenchyma of tentacles only in presence of O ₂ . Urea, thiourea, and caffeine translocated in absence of O ₂ .	10

/1/ Phloem. /2/ Stem. /3/ Leaf.

117. TRANSLOCATION OF ORGANIC SUBSTANCES: PLANTS (Continued)

Part VI. NITROGENOUS SUBSTANCES AND VITAMINS (Concluded)

Plant	Part	Substance	Translocation		Remarks	Reference
			Rate cm/hr	Direction		
(A)	(B)	(C)	(D)	(E)	(F)	(G)
11 Tobacco (<i>Nicotiana tabacum</i>)	Leaf, stem, root	Urea		A2, 3, B2, 3	Radioassay after application of N^{15} -labelled urea to single leaf showed half of absorbed N^{15} transported from leaf in 24 hr. Activity present after 6 hr in leaf opposite to treated leaf, as well as in stem and root, may indicate transfer from phloem to xylem.	11
12 Tomato (<i>Lycopersicon esculentum</i>)	Leaf, stem	Total nitrogen, non-protein nitrogen, sucrose, thiamine, pyridoxine, pantothenic acid, riboflavin		B	Compounds accumulated above steamed section of stem at 2nd node in 1-4 da. Chemical analysis of bioassay for vitamins indicated that sucrose accumulated mostly above steamed section, with other compounds accumulating in varying degrees.	12
13	Leaf, stem, root	Thiamine		A, B	Bioassay of various plant parts, 1-6 da after girdling different stem loci or petioles by steam, indicated older leaves have low thiamine concentration; transport greatest to young leaves and roots.	13
14 Wheat (<i>Triticum</i> sp.)	Stem	Asparagine		A	Excised stem sections (30-35 cm long) placed with upper or lower ends in solution. After 20 min, analysis of 5-cm pieces cut from each end showed N accumulation in morphological upper end regardless of which end was immersed, suggesting polar transport.	14

12/ Stem 13/ Leaf.

Contributor: Hull, Herbert M.

- References [1] Bond, G. 1956. J. Exp. Bot. 7:387. [2] Ziegler, H. 1956. Planta 47:447. [3] Kazarian, V. O., and E. S. Avundzhyan. 1955. Doklady akad. nauk. S. S. S. R. 101:181. [4] Kazarian, V. O., and N. V. Balagerian. 1955. Ibid. 103:337. [5] Mothes, K., and L. Engelbrecht. 1954. Flora 143:356. [6] Maskell, E. J., and T. G. Mason. 1929. Ann. Bot., Lond. 43:205. [7] Hay, R. E., E. B. Lacey, and E. E. de Turk. 1953. Plant Physiol. 28:606. [8] Kursanov, A. L. 1956. Proc. Internat. Conf. Peaceful Uses Atomic Energy, Geneva 12:165. United Nations, New York. [9] Nelson, C. D., and P. R. Gorham. 1958. Plant Physiol., Suppl. 33:21. [10] Arisz, W. H. 1953. Acta bot. nederl. 2:74. [11] Volk, R., and C. McAuliffe. 1954. Proc. Soil Sc. Soc. America 18:308. [12] Bonner, J. 1944. Am. J. Bot. 31:591. [13] Bonner, J. 1942. Ibid. 29:136. [14] Kursanov, A. L., and M. N. Zaprometov. 1949. Doklady akad. nauk. S. S. S. R. 69:89.

Part VII: ALKALOIDS AND INSECTICIDES

Plant	Part	Substance	Translocation		Remarks	Reference
			Rate cm/hr	Direction		
(A)	(B)	(C)	(D)	(E)	(F)	(G)
Alkaloids						
1 Alkekengi, or winter cherry (Physalis alkekengi) Belladonna (Atropa belladonna) Potato (Solanum tuberosum) Tomato (Lycopersicon esculentum)	Leaf, stem, root, tuber	Atropine, hyoscyne, hyoscyamine		A, B	Approach grafts of belladonna to alkekengi, potato, or tomato, allowed to remain 3-4 mo. Alkaloids, normally elaborated in belladonna roots, translocated into basal stem, root, and tuber portions of graft partners. Such transport occurred apparently in phloem with other assimilates.	1
2 Tobacco (Nicotiana tabacum, N. rustica) Tomato (Lycopersicon esculentum)	Leaf, stem	Nicotine		A, B	Tobacco scions, grafted to tomato rootstocks and topgrafted with tomato scions, produced nicotine in basal portion of tobacco scion. Nicotine moved acropetally through upper graft, but did not pass basal graft and enter rootstock, suggesting xylem and phloem translocation, respectively. Relative contribution of tobacco root and stem to leaf nicotine varied widely.	2, 3
Insecticides						
3 Apple (Malus sp) Bean, broad (Vicia faba) Chrysanthemum (Chrysanthemum sp) Coleus (Coleus sp)	Leaf, stem, root	Octamethyl pyrophosphoramidate		A, B	Radioassay of various plant parts, after application of P ³² -labelled compound to central leaves. Indicated movement to other leaves in 8 hr and to root in 72 hr (apple). Although light had pronounced and variable effect on different plants, temperature had less effect. In beans, translocation at 10-15°C approximately equivalent to that at 27°C.	4
4 Bean (Phaseolus vulgaris) Lemon (Citrus limon)	Leaf, stem	Octamethyl pyrophosphoramidate; O, O-diethyl-O-mercaptoethyl thiophosphate	10 2.5	A B	P ³² or S ³⁵ -labelled compounds applied to parts of foliage or bark; translocation in stem determined by bioassay (insects feeding on various stem sections).	5

Contributor: Hull, Herbert M.

References: [1] Wilson, P. M. W. 1952. New Phytol. 51:301. [2] Dawson, R. F., and M. L. Solt. 1957. Plant Physiol., Suppl. 32:36. [3] Solt, M. L., and R. F. Dawson. 1958. Plant Physiol. 33:375. [4] Thomas, W. D. E., and S. H. Bennett. 1954. Ann. Appl. Biol., Lond. 41:501. [5] Wedding, R. T. 1953. J. Agr. Food Chem. 1:832.

For a comprehensive review of the subject, see reference 4. Values in parentheses are ranges, estimate "c" unless otherwise indicated (cf. Introduction).

Specification	Normal Value	Trauma without Shock		Slight Shock ¹		Severe Shock ²		Reference
		No. of Subjects	Value	No. of Subjects	Value	No. of Subjects	Value	
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)
1 Pulse rate, beats/min	74 (51-96)	8	80 (70-98)	4	120 (100-140)	4	137 (112-205)	1
2 Arterial blood pressure, mm Hg								
Systolic	137 (122-149)	8	137 (102-167)	4	77 (48-113)	4	63 (46-80)	1
3 Diastolic	74 (68-83)	8	69 (44-87)	4	40 (28-50)	4	32 (15-42)	1
4 Mean	97 (88-106)	8	94 (67-113)	4	54 (36-67)	4	44 (22-57)	1
5 Venous blood pressure, mm H ₂ O								
Peripheral	74	8	86 (79-92)	4	77 (62-105)	4	90 (50-121)	1
6 Atrial	33	8	45 (12-70)	4	26 (10-47)	4	-8 (0 to -11)	1
7 Cardiac output								
Cardiac index, L/min/sq m	3.4 (2.1-4.3)	8	3.6 (2.2-4.3)	4	2.5 (2.2-2.7)	4	2.1 (1.3-3.0)	1
8 Stroke volume, ml/beat	83 (63-109)	8	78 (52-100)	4	36 (33-39)	4	26 (21-35)	1
9 Total peripheral vascular resistance, dynes/cm ⁵	1290	8	1281 (35)	4	1019 (738-1452)	4	1021 (567-1468)	1
10 " "					77 (58-82)	4	84 (60-100)	1
11 " "	1600	8	1597 (1130-2070)	4	1532 (1376-1675)	4	1225 (1033-1310)	1
12 Whole blood	2900	8	2791	4	2330	4	1670	1
13 RBC packed volume (hematocrit), %	47 (42-52)							
14 Hemoglobin, g/100 ml	16.0 (14.0-17.5)		(8.5-15.7) ^b		(8.1-14.5) ^b		8.8 (7.0-10.6) ^b	
15 Arterial blood								
O ₂ content, vol %	18.6	8	15.6 (13.1-18.5)	4	12.3 (9.8-14.4)	4	11.8 (10.2-14.6)	1
16 CO ₂ content, vol %	49.1	8	44.1 (42.7-50.6)	4	35.8 (32.7-40.1)	4	28.4 (16.5-39.4)	1
17 CO ₂ pressure, mm Hg	38.8	8	37.4 (30.4-42.8)	4	30.5 (27.0-33.4)	4	28.4 (17.4-39.4)	1
18 O ₂ saturation, %	96.0	8	94.2 (91.0-97.4)	4	90.8 (85.7-95.0)	4	94.6 (83.0-99.0)	1
19 pH	7.42	8	7.39 (7.36-7.43)	4	7.38 (7.32-7.41)	4	7.28 (7.26-7.29)	1
20 Mixed venous blood								
CO ₂ pressure, mm Hg	44.7	8	43.9 (38.3-48.2)	4	36.9 (32.0-39.5)	4	40.9 (31.4-46.7)	1
21 O ₂ saturation, %	67.7	8	66.7 (58.0-76.3)	4	38.4 (35.0-42.9)	4	28.2 (14.0-49.6)	1
22 pH	7.39	8	7.36 (7.33-7.39)	4	7.32 (7.29-7.35)	4	7.22 (7.19-7.26)	1
23 Plasma								
Protein, g/100 ml	6.5 (6.0-7.0)	15	6.6 (6.4-6.8) ^b	26	6.4 (6.2-6.6) ^b	25	6.0 (5.8-6.2) ^b	2
24 Non-protein nitrogen, mg/100 ml	33 (23-40)	18	34.0 (31.0-37.0) ^b	24	35.0 (31.4-38.6) ^b	24	44.0 (39.6-48.2) ^b	2

1/1 Characterized by decreased blood pressure of 20% or less, cool skin temperature, pale skin color, definite slowing of circulation (response to pressure, blanching), and clear but distressed mental state. 1/2 Characterized by weak-to-imperceptible pulse, decreased blood pressure of 40% to non-recordable, cold skin temperature, ashen-to-cyanotic (mottled) skin color, very sluggish circulation (response to pressure, blanching), severe thirst, apathetic-to-comatose mental state.

118. EFFECT OF SHOCK ON CIRCULATORY FUNCTION AND BLOOD COMPOSITION. MAN (Concluded)

	Specification	Normal Value	Trauma without Shock		Slight Shock ¹		Severe Shock ²		Reference
			No. of Sub-jects	Value	No. of Sub-jects	Value	No. of Sub-jects	Value	
	(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)
25	Plasma (concluded) Creatinine, mg/100 ml	1.0 (0.5-1.2)	18	1.0 (0.9-1.1) ^b	24	1.3 (1.1-1.5) ^b	23	2.1 (1.8-2.4) ^b	2
26	Uric acid, mg/100 ml	3.0 (1.5-3.5)	10	4.0 (2.8-5.2) ^b	16	4.5 (3.7-5.3) ^b	16	5.2 (4.1-6.3) ^b	2
27	Phosphorus, mg/100 ml	3.0 (1.6-3.5)	14	3.3 (2.8-3.8) ^b	20	3.8 (3.5-4.1) ^b	21	5.8 (4.8-6.8) ^b	2
28	Magnesium, mg/100 ml	1.8 (1.0-3.0)	11	1.7 (1.6-1.8) ^b	17	1.9 (1.8-2.0) ^b	16	2.5 (2.2-2.8) ^b	2
29	Bilirubin, mg/100 ml	(0.1-0.5)	11	0.43 (0.27-0.59) ^b	23	0.68 (0.48-0.88) ^b	21	0.47 (0.46-0.48) ^b	2
30	Glucose, mg/100 ml	(80-90)	9	134 (116-152) ^b	14	149 (131-167) ^b	13	202 (150-254) ^b	2
31	Chlorides, mEq/L	100 (97.5-104.0)	13	102 (96-108) ^b	24	101 (99.6-102.4) ^b	24	99.5 (97.7-101.3) ^b	2,3
32	CO ₂ combining power, mEq/L	27 (24-31)	13	27.9 (25.9-29.9) ^b	22	26.1 (24.7-27.5) ^b	25	22.1 (20.3-23.9) ^b	2
33	Serum sodium, mEq/L	143 (139-149)			6	142.1	5	144.7	2,3

/1/ Characterized by decreased blood pressure of 20% or less, cool skin temperature, pale skin color, definite slowing of circulation (response to pressure, blanching), and clear but distressed mental state. /2/ Characterized by weak-to-imperceptible pulse, decreased blood pressure of 40% to non-recordable, cold skin temperature, ashen-to-cyanotic (mottled) skin color, very sluggish circulation (response to pressure, blanching), severe thirst, apathetic-to-comatose mental state.

Contributors. (a) Wiggers, Carl J., (b) Engel, Frank L., (c) Page, Irvine H., (d) Fine, Jacob, Howard A. Frank, and Henry Korman

References: [1] Cournand, A., et al. 1943. Surgery 13 964. [2] The Board for the Study of the Severely Wounded. 1952. The physiologic effects of wounds. Office of the Surgeon General, Department of the Army, Washington. [3] Beecher, H. K., et al. 1947. Surgery 22.672. [4] Wiggers, C. J. 1950. Physiology of shock. Harvard University Press, Cambridge.

119. EFFECT OF SHOCK ON BLOOD COMPOSITION: LABORATORY ANIMALS

For a comprehensive review of the subject, see reference 27.

Specification	Animal	Control	Shock						Reference
			Gravity	Tourniquet	Noble-Uncond	Collip Cond.	Traumatic	Hemorrhagic	
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)
1 Glucose, mg/100 ml	Dog	85.9					86.9		1-3
2	Rat	87-95	67-196	80-150	100-190	100-125	70-195	70-195-40	4
3 Chlorides, mEq/L	Dog	107.4-115.3					115.7-116.2	108-111.4	5
4	Rabbit	107-114	95-110						6
5 Sodium, mEq/L	Dog	135-145					135-145	135-145	1
6	Dog	149-159					155-158	148-162	5
7 Calcium, mEq/L	Dog	5.1-6.6					5.1-5.4	5.4-5.8	5
8 Potassium, mEq/L	Dog	4.4-6.2					5.1-6.5	7.6-7.7	5
9	Dog	4.9					5-12		7
10	Rabbit	4.5-5.1							8,9
11 Magnesium, mEq/L	Dog	1.4-2.2					2.6-3.2	2.7-2.9	5
12 Pyruvic acid, mg/100 ml	Rat	1.4		1.2-2.2	1.5-5.0	1.5-4.0		1.4-5.0	2,4
13 Pyruvate, mEq/L	Dog	0.1-0.2					0.4-0.5	0.4-0.5	1,5
14 Lactate, mEq/L	Dog	2.1-3.9					8.0-18.8	6.0-14.6	5
15 Lactic acid, mg/100 ml	Dog	12.8					14.5		1
16	Rat	16-18		20-43	20-150	20-70			2
17 Uric acid, mg/100 ml	Rat	0.8			1.5				10
18 Plasma uric acid, mg/100 ml	Dog	0.5						4.0-5.0	11,12
19	Rat	0.6						8.7	13
20 Plasma amino acid nitrogen, mg/100 ml	Cat	4.2-7.6						4.6-11.8	14
21	Dog	2.6-5.0					5.0	5.0-7.5	1,15
22	Rat	7.0						7-40	4
23	Rat	5.0		6-12					2
24 RBC amino acid nitrogen, mg/100 ml	Rat	21						21-24	16
25 Whole blood amino acid nitrogen, mg/100 ml	Rat	14		10-30	13-23	11-17		14-40	2,4
26 Urea nitrogen, mg/100 ml	Cat	26-32					25-49		17
27	Rat	15		14.6-48.4					18
28 Creatine nitrogen, mg/100 ml	Cat	1.2-1.9					1.3-6.0		17
29	Rat	1.9			2.5				10
30 Non-protein nitrogen, mg/100 ml	Dog	15.1					19.7		1
31 HPO_4^{2-} and H_2PO_4^- , mEq/L	Dog	1.2-2.6					4.0-6.5	4.3-9.7	5
32 Plasma inorganic phosphorus, mg/100 ml	Dog	4.4		4.6			4.6	3.6	1
33	Rabbit	3.3-7.2	5.2-24.6						6
34	Rat	6.92	20.0						18
35	Rat	7.5		7.5-12	6-13	6-10			2
36 RBC inorganic phosphorus, mg/100 ml	Rabbit	2.2-2.7	2.6-13.0						6
37	Rat	2.35	7.70						18
38 Plasma organic phosphorus, mg/100 ml	Rabbit	0.1-0.8	0.4-8.6						6
39 RBC organic phosphorus, mg/100 ml	Rabbit	70-89	68-98						6
40 Acid-soluble organic phosphorus, mg/100 ml	Rat	9	86.9						18

119. EFFECT OF SHOCK ON BLOOD COMPOSITION: LABORATORY ANIMALS (Continued)

Specification	Animal	Control	Shock						Reference
			Gravity	Tourniquet	Noble-Collip Uncond.	Cond.	Traumatic	Hemorrhagic	
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)
41 Plasma acid-soluble organic phosphorus, mg/100 ml	Rat	1.20	2.35						18
42 Whole blood acid-insoluble phosphorus, mg/100 ml	Rat	20.2	15.0						18
43 Plasma iron, μ g/100 ml	Dog	73						275	19
44 Plasma iron-binding capacity, μ g/100 ml	Dog	170						10	19
45 Fibrinogen, %	Dog	100						70-100	20
46 Prothrombin, %	Dog	100						2.5-80	20
47 Plasma proteins, %	Dog	100						68-100	20
48 Total proteins, mg/100 ml	Cat	4.7-5.5					4.7-7.4		17
49 Ketones, mg/100 ml as acetone	Rat	3-6						6-0	21
50 Plasma amino-peptidase (glycyl-glycyl-glycine) μ M/0.1 ml plasma/hr	Rat	2.0						4-11	22
51 Serum neutral fat glycerol, mM/L	Rabbit	1		2.5-9					23
52 Serum "free glycerol," mM/L	Rabbit	1		2.5-3.5					23
53 Serum phospholipid, mg/100 ml	Rabbit	90-100		180-350					23
54 Ammonia, μ g/ml	Dog	0.7-1.2						1.8-4.1	24
55 Creatinine, mg/100 ml	Dog	0.97					1.03		1
56 Plasma pentose, mg/100 ml	Rat	7.0		6-18	4-17	4-8			2
57 Plasma H ₂ O, g/L	Dog	912-946					911-919	917-957	5
58 HCO ₃ , mEq/L	Dog	17.9-25.6					6.3-13.5	3.5-13.4	5
59 Base bound by protein, mEq/L	Dog	12.8-20.7					12.2-16.3	11.0-14.9	5
60 Total base, mEq/L	Dog	158.6-168.6					167.8-174.8	162.5-168.7	5
61 Sum of anions, mEq/L	Dog	146.6-160.7					154.3-162.0	148.6-151.8	5
62 Sum of cations, mEq/L	Dog	160.9-172.1					169.2-170.8	164.5-177.9	5
63 Cl/RBC plasma	Dog	0.45-0.61						0.52-0.64	5
64 CO ₂ pressure, mm Hg	Dog	30.0-43.7					14.7-31.8	6.8-27.9	5
65 CO ₂ content, vol %	Cat	36.5					22.3		17
66 Arterial O ₂ saturation, %	Cat	82-100						85-99	14
67 Venous O ₂ saturation, %	Cat	69-89						5-43	14
68 Average O ₂ content, right heart, vol %	Dog	12.88					5.23	4.88	25

119. EFFECT OF SHOCK ON BLOOD COMPOSITION: LABORATORY ANIMALS (Concluded)

Specification	Animal	Control	Shock				Traumatic	Hemorrhagic	Reference
			Gravity	Tourniquet	Noble-Collip				
					Uncond.	Cond.			
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)
69 Average O ₂ content, portal vein, vol %	Dog	12.32					7.50	4.99	25
70 Average O ₂ content, femoral vein, vol %	Dog	10.68					4.15	4.18	25
71 Average O ₂ content, jugular vein, vol %	Dog	13.86					4.68	6.0	25
72 Average O ₂ content, renal vein, vol %	Dog	14.54					10.83	7.62	25
73 Average O ₂ content, femoral artery, vol %	Dog	17.14					16.31	13.84	25
74 pH, arterial blood	Dog	7.28-7.42					7.19-7.33	6.92-7.32	5
75	Rat	7.34	6.83						18
76 RBC packed volume (hematocrit), %	Rat	45.7		45.8-67				25-30	26
77 Lymphocytes, %	Rat	72		2.43					18

Contributors. (a) Fine, Jacob, Howard A. Frank, and Henry Korman. (b) Engel, Frank L., (c) Baez, Silvio, (d) Page, Irvine H., (e) Wiggers, Carl J.

References: [1] Davis, H. A. 1948. Shock and allied forms of failure of the circulation. Grune and Stratton, New York. [2] McShan, W. H., et al. 1945-46. Am. J. Physiol. 145:93. [3] Brand, E. D., et al. 1951. Ibid. 165:539. [4] Engel, F. L., et al. 1943. J. Exp. M. 77:397. [5] Root, W. S., et al. 1947. Am. J. Physiol. 149:52. [6] Allison, J. B., and W. H. Cole. 1944. Proc. Soc. Exp. Biol., N. Y. 57:21. [7] Manery, J. F. 1943. Am. J. Physiol. 138:499. [8] Cole, W. H. 1943. Proc. Soc. Exp. Biol., N. Y. 54:215. [9] Clarke, A. P. 1942. Endocrinology 31:597. [10] Neufeld, A. K., C. G. Toby, and R. L. Noble. 1943. Proc. Soc. Exp. Biol., N. Y. 54:249. [11] Zweifel, B. W., et al. 1945. Surgery 18:48. [12] Van Slyke, D. D. 1948. Ann. N. York Acad. Sc. 49:593. [13] Green, S., and A. J. Mazur. 1957. J. Biol. Chem. 227:653. [14] Engel, F. L. 1944. J. Exp. M. 79:9. [15] Kline, D. L. 1946. Am. J. Physiol. 146:854. [16] Sayers, M. A., et al. 1945. Proc. Soc. Exp. Biol., N. Y. 60:20. [17] Aub, J. C. 1920. Am. J. Physiol. 54:416. [18] Wiedeman, Mary P., and C. R. Lewis. 1949. Proc. Soc. Exp. Biol., N. Y. 71:467. [19] Mazur, A., S. Baez, and E. Shorr. 1953. J. Biol. Chem. 213:1, 147. [20] Frank, E. D., et al. 1951. Am. J. Physiol. 167:499. [21] Engel, F. L., and K. Hewson. 1953. Proc. Soc. Exp. Biol., N. Y. 83:608. [22] Schwartz, T. H., and F. L. Engel. 1953. Ibid. 83:568. [23] Johnson, S. R., and L. B. Waldstrom. 1936. Scand J Clin. & Invest., Oslo 8:323. [24] Nelson, R. M., and D. Seligson. 1953. Surgery 34:1. [25] Bialock, A., and H. Bradburn. 1930. Arch. Surg. 20:26. [26] Gray, J. L., et al. 1950. Proc. Soc. Exp. Biol., N. Y. 75:189. [27] Wiggers, C. J. 1950. Physiology of shock. Harvard University Press, Cambridge.

Values in parentheses are ranges, estimate "c" (cf. Introduction).

Part I: BLOOD PRESSURES

Condition or Agent	No. of Subjects	Pressure, mm Hg					Right Ventricle (Diastolic)	Right Atrium (Mean)	Ventricular Work Against Pressure kg·m/min/sq m		Reference
		Arterial (Systemic) (Mean)	Pulmonary Capillary (Mean)	Pulmonary Artery (Mean)	(F)	(G)	(H)	(I)			
1 Pulmonary emphysema	12	(A)	(D)	(E)							(J)
2 Mild	18		8(1-14)	35(14-57)							1
3 Mild	6	89(74-100)	8(1 to 12)	23(9-49)							2
4 Severe	6	95(84-118)	25(23-26)	15(12-19)	2(1-4)						3
5 Cor pulmonale	11	88(75-106)	25(15-46)	2(0-6)							
6 Silicosis	5	82(65-90)	20(18-21)	2(0-4)							3
7 Severe	3	87(80-93)	23(18-36)	3(0-5)							
8 Diffuse	9	87(71-100)	2(15-34)	1(1 to 12)							
9 Mitral stenosis	12		(18-37)								
10	22	87(65-123)	29(17-54)	51(23-94)	8(-3 to +33)						4
11	15	(18-56)	(15-34)	(18-56)	(0-15)						5
12	17	(63-102)	(24-81)	(1-23)							6
13	36	(63-148)	27(11-36)	(7-39)							7
14	43		(15-28)	(15-86)	(-2 to +20)						8
15 Mild			25	45							9
16 Moderate ²			38	65							10
17 Severe ³											
18 Tricuspid stenosis	12		(14-35/20) ⁴	(15-5-65/30) ⁴	(15-7 to 65/-5) ⁴						11
19	5		(12-32)	(18-49)	(29/11-80/74) ⁴						12
20			(19-20)	(27/15-48/37) ⁴	34/54, 6						
21 Rest	3	(120/65-124/90) ⁴	(19-20)	(29-51)	(40/10-78/15) ⁴						13
22			(24-25)	(35-65)	53/20 ⁴						
23 Exercise	3	130/72 ⁴	(56/27-86/55) ⁴								
24 Tricuspid insufficiency	60										14
25 Rest	60										
26 Exercise	60										
27 Aortic valvular disease	9	90(77-118)	(6-17)	(11-25)							15
28 Aortic stenosis	11	102(72/85/65-130/100) ⁴	(5-25)	(14-50)							
29 Aortic stenosis, plus aortic insufficiency	5	99(75-115)	(8-25)	(14-47)							
30 Aortic insufficiency	5	174/55(140/50-240/60) ⁴									
31											
32											
33											
34											

35	Primary pulmonary hypertension	10	(100/76-153/110) ⁴	(3-13)	(56-105) (62/47-144/72) ⁴	(5-0-26/15) ⁴	16
36					(66-74)	(-7 to +1)	17
37		4	(90-94)	(3-5)	(61/31-112/30) ⁴	(11-17)	18
38			(107/70-128/88) ⁴	(2-11)	(70/38-143/47) ⁴	(1.50-4)	19
39		6	(178-188)		(34/213/4-79/0) ⁴	(13-15)	20
40			(104/54-139/92) ⁴		(2-2 to +5)	(57-4 to +10)	21
41	Patent ductus arteriosus	16			(6/916/-2 to 29/11) ⁴	(2-9)	22
42					(145-26)	(16/2-140/8) ⁴	23
43	Pulmonary stenosis	12			(23/914/0-33/24) ⁴	(37/11-37/31) ⁴	24
44		24			(18-104)	(135-25)	25
45	Interventricular septal defect	23	(81/32-150/95) ⁴	(6-12)	(135/6-153/79) ⁴	(11-22)	26
46		50	(60-62)	(10-28)	(24/135-45/20) ⁴	(110-21)	27
47	Constrictive pericarditis	13		(11-21)	(33-43) ⁴	(-15 to +15)	28
48		5	(100/85-136/82) ⁴	(13-21)	(31/18-44/23) ⁴	(46-5)	29
49	Congestive heart failure	13	(65/41-74)			(16/-4 to +23)	30
50	Acute hemorrhagic pericarditis	7	(73/32-104)				31
51							32
52	Acute hypoxia	2844	100	10	19		
53	Before	2848	100	20	29	5.0	
54	After	815	96	5	10	5.0	
55	Before	817	93	5	15	4.7	
56	After	1218	98	8	24	5.6	
57	Before	1218	102	9	37(10-73)		
58	After	2419	95(73-130)	(18/7-34)	34(17-84)		
59	100% oxygen	2519	91(58-122)	(9/6-35)	43		
60	Before			5	25		
61	After			4	20		
62	Norepinephrine	817	90	5	13		
63	Before	817	140	13	23		
64	After	220	144	8	17		
65	Before	510	108	20	28		
66	After	72	100	9	16		
67	Before	12	125	15	21		
68	After						

11/ With pulmonary artery systolic pressure <50 mm Hg. 12/ With pulmonary artery systolic pressure >50 mm Hg. 13/ With right ventricular failure. 14/ Systolic/diastolic. 15/ A wave. 16/ V wave. 17/ Rest. 18/ Exercise. 19/ Systolic. 110/ Early diastolic. 111/ Late diastolic. 112/ Systolic. 113/ Pulmonary. 114/ Breathing 100% O₂. 115/ Diagnostic criteria: (1) M- or W-shaped pattern of right atrial pressure, (2) an early diastolic dip followed by a rapid rise of right ventricular pressure to form a plateau, (3) ratio of right ventricular diastolic pressure to right ventricular systolic pressure of 1:1 to 40%. 116/ 22 normal subjects, 6 patients. 117/ Normal subjects. 118/ Patients with mitral stenosis. 119/ Hypertensive patients.

Part II. BLOOD FLOW AND RESISTANCE

Condition or Agent (A)	No of Subjects (B)	Blood Flow			Resistance, dynes sec./cm. ⁵			Reference
		Cardiac Index L./min./sq m (C)	Stroke Index ml/beat/sq m (D)	A-V O ₂ Difference ml/L (E)	Total Arterial (F)	Pulmonary (G)	Pulmonary Vascular (H)	
1 Pulmonary empty-	12	3.0 (1.9-4.9)		41 (28-63)				
2 Aortic stenosis	18	3.5 (2.3-6.6)				590 (145-1788)	450 (73-1322)	1
3 Mild	6	3.2 (2.7-3.6)				590 (145-1788)	450 (73-1322)	2
4 Severe	6	3.6 (2.4-4.5)				590 (145-1788)	450 (73-1322)	3
5 Cor pulmonale	11	3.4 (2.4-4.6)				590 (145-1788)	450 (73-1322)	
6 Silicosis	5	2.8 (2.4-3.2)						3
7 Mild	3	2.8 (2.4-3.2)						
8 Severe	9	4.0 (3.2-4.9)						
9 Diffuse	9	1.4 (1.3-1.6)	30 (9-54)	57 (30-86)	1799 (775-3760)	111 (212-2372)	49 (151-1480)	4
10 Mitral stenosis	2	2.6 (1.4-5.2)		(49-123)				5
11	15	1.1 (1.3-2.3)		(26-89)	1734 (4460)	157 (2805)	64 (1443)	6
12	17	2.0 (1.4-3.5)	(11-24)	48		514	231	7
13	36	2.0 (1.3-3.5)		48		1627	775	8
14	43	2.78		63				9
15 Mild	3	3.07				1320 (950)	110 (330)	10
16 Moderate	2	2.24						11
17 Severe	12	1.2 (0.6-8.1)	(28-39)	177 (104)				12
18 Tricuspid stenosis	5	(2.10-3.40)		(103-162)				13
19	3	(0.8-2.2)						
20 Rest	3	(2.2-2.7)						14
21 Exercise	60	1.94 (1.15-3.11)				643 (110-2109)		
22 Tricuspid insufficiency	60	2.25 (1.16-6.25)						15
23 Rest	60							
24 Exercise	60							
25 Aortic valve disease	9	(2.0-5.1)	(19-54)	(33-62)	1280 (4550)		73 (532)	
26 Aortic stenosis, plus	11	(2.6-5.2)	(26-54)	(41-83)	1310 (3002)		71 (590)	
27 Aortic insufficiency	5	(2.2-3.9)						16
28 Aortic insufficiency	10	(0.9-2.3) ²						17
29 Primary pulmonary hypertension	4	(1.6-2.8)						18
30	4	(1.08-2.33)						19
31 Patent ductus arteriosus	16	3.9 (1.7-5.5) ⁴						20
32 Pulmonary stenosis	12	3.4 (1.8-5.1)						21
33	24	(1.2-12.0)						22
34 Interventricular septal defect	23	(1.5-26.9) ⁵						

11/ Cardiac output, L/min. 12/ Pulmonary blood flow, L/min/sq m = 1.2-4.2 breathing room air, 1.3-4.7 breathing 100% O₂. 13/ Breathing 100% O₂. 14/ Systemic. 15/ Pulmonary. 16/ Blood O₂ content = 14.6-31.4 vol %, blood O₂ saturation = 69.2-101%. 17/ Blood O₂ content = 11.6-22.6 vol %, blood O₂ saturation = 51.5-90%.

Part I: BLOOD PRESSURES (Concluded)

Condition or Agent	No. of Subjects	Pressure, mm Hg				Ventricular Work Against Pressure		Reference
		Arterial (Systemic) (Mean)	Pulmonary Capillary (Mean)	Pulmonary Artery (Mean)	Right Ventricle (Diastolic)	Right Atrium (Mean)	Left Right	
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H) (I)	(J)
Saline infusion, Intravenous								
Before	12		5	12				
After	12		13	20				37
76								
77								
78		(76-110)						
79		(106/60-155/80)4				(-2 to +4)		38
After	10	(78-109)				(2-8)		
80		(107/64-158/80)4						
5% albumin infusion, Intravenous								
Before	7	(66-113)						
After	7	(107/47-163/86)4				(-1 to +5)		38
81		(67-84)				(3-14)		
82		(107/69-166/113)4						
83								
84								

/4) Systolic/diastolic.

Contributor: Yu, Paul N

- References [1] Dexter, L., J. L. Whittenberger, R. Gorlin, B. M. Lewis, F. W. Haynes, and R. J. Spiegel. 1951. *Tr. Ass. Am. Physicians* 65:226.
- [2] Yu, P. N., F. W. Lovejoy, Jr., H. A. Joos, R. E. Nye, Jr., and W. S. McCann. 1953. *J. Clin. Invest.* 32:130. [3] Harvey, R. M., M. L. Ferrer, D. W. Richards, Jr., and A. Courand. 1951. *Am. J. Med.* 10:719. [4] Dexter, L. 1952. *Bull. N. York Acad. Sci.* 28:90. [5] Gorlin, R., F. W. Haynes, W. T. Goodale, C. G. Sawyer, J. W. Dow, and L. Dexter. 1951. *Am. Heart J.* 41:1. [6] Curtis, P. C., G. Cohen, B. Castelman, J. G. Scannell, A. L. Friedlitch, and G. S. Myers. 1953. *Circulation*, N. Y. 8:893. [7] Ferrer, M. L., R. M. Harvey, R. T. Cathcart, A. Courand, and D. W. Richards, Jr. 1952. *Ibid.* 6:688. [8] Arufo, J., and D. S. Lukas. 1952. *J. Clin. Invest.* 31:1082. [9] Yu, P. N., H. H. Joos, and R. E. Nye, Jr. 1954. 23:499. [10] Cosby, R. S., G. C. Griffith, D. C. Levinson, W. J. Zimm, R. W. Olathe, S. F. Dumitroff, and L. M. Herman. 1953. *Dis. Chest.* 23:499. [11] Gibson, R., and P. Wood. 1955. *Brit. Heart J.* 17:552. [12] Yu, P. N., D. E. Harken, F. W. Lovejoy, Jr., R. E. Nye, Jr., and E. B. Mahoney. 1956. *Circulation*, N. Y. 13:680. [13] McCord, M. C., H. J. Swan, and S. G. Blount. 1954. *Am. Heart J.* 48:405. [14] Sepulveda, G., and D. S. Haynes. 1955. *Circulation*, N. Y. 11:552. [15] Gorlin, R., J. K. R. McMullan, W. E. Nedd, M. B. Mathews, and R. Dalley. 1955. *Am. J. Med.* 18:855. [16] Shepherd, J. T., J. E. Edwards, H. B. Burchell, H. J. C. Swan, and E. H. Wood. 1957. *Brit. Heart J.* 19:70. [17] Kuida, H., G. J. Dammin, F. W. McCord. 1955. *Ibid.* 18:871. [23] Hubbard, T. F., W. D. Angle, and J. H. Simpson. 1954. *Ann. Int. M.* 47:138. [19] Storslein, O., S. Humerfelt, O. Muller, and H. Raemussen. 1952. *Acta med. scand.* 141:419. [20] Sobin, S. S., M. J. Carson, J. L. Johnson, and C. R. Baker. 1954. *Am. Heart J.* 48:416. [21] Joos, H. A., P. N. Yu, F. W. Lovejoy, Jr., R. E. Nye, Jr., and B. J. Koszewski. 1957. *Am. Heart J.* 54:210. [24] Blount, S. G., H. W. Hoshel, C. Sadoff, and M. E. Dempsey. 1954. *Surgery* 42:184. [25] McKusick, V. A. 1952. *Bull. Johns Hopkins Hosp.* 90:3. [26] Wilson, R. H., W. Hoshel, C. Sadoff, and A. L. Friedlitch. 1954. *Am. Heart J.* 48:671. [27] Stead, E. A., Jr., J. V. Warren, and E. S. Brannon. 1946. *Ibid.* 31:407. [29] Merrill, A. J., J. V. Warren, E. A. Stead, Jr., and E. S. Brannon. 1946. *Ibid.* 31:413. [30] Warren, J. V., E. S. Brannon, E. A. Stead, Jr., and A. J. Merrill. 1946. *Ibid.* 31:418. [31] Westcott, R. N., N. O. Fowler, R. C. Scott, V. D. Hauenstein, and J. McGuire. 1951. *J. Clin. Invest.* 30:957. [32] Doyle, J. T., J. S. Wilson, and J. V. Warren. 1952. *Circulation*, N. Y. 5:263. [33] Yu, P. N., D. C. Bagby, F. W. Lovejoy, Jr., R. E. Nye, Jr., and H. A. Joos. 1956. *Am. Heart J.* 52:683. [34] Wilson, R. H., W. Hoshel, and M. E. Dempsey. 1955. *Ann. Int. M.* 42:629. [35] Nelson, R. A., L. G. May, A. Bennett, M. Kabayaski, and R. Gregory. 1955. *Am. Heart J.* 50:172. [36] Fowler, N. O., R. N. Westcott, R. C. Scott, and J. McGuire. 1951. *J. Clin. Invest.* 30:517. [37] Doyle, J. T., J. S. Wilson, E. H. Estes, and J. V. Warren. 1951. *Ibid.* 30:345. [38] Warren, J. V., E. S. Brannon, H. S. Weens, and E. A. Stead, Jr. 1948. *Am. J. Med.* 4:193.

- Richards, Jr., and A. Courmand 1931. *Am. J. Med* 10:719. [4] Dexter, L. 1932. *Bull. N. York Acad. Sci.* 28:90. [5] Corlin, R., F. W. Haynes, W. T. Goodale, C. G. Sawyer, J. W. Dow, and L. Dexter. 1931. *Am. Heart J.* 4:111. [6] Curti, P. C., G. Cohen, D. Castleman, J. G. Scannell, A. L. Friedlich, and G. S. Myers 1953. *Circulation*, N. Y. 8:993. [7] Farret, M. J., R. M. Harvey, R. T. Calhoun, A. Courmand, and D. W. Richards. *Jr* 1952. *Ibid* 6:689 [8] Araujo, J., and D. S. Lukas. 1952. *J. Clin. Invest.* 31:1082. [9] Yu, P. N., J. H. Simpson, F. W. Lovejoy, Jr., H. A. Joss, and R. E. Nye, Jr. 1954. *Am Heart J* 47:330 [10] Cosby, R. S., G. C. Griffith, D. C. Levinson, W. J. Zimm, R. W. Obleath, S. P. Dimitroff, and L. M. Herman 1953. *Dis. Chest* 23:499. [11] Gibson, R., and P. Wood. 1955. *Brit. Heart J.* 17:552. [12] Yu, P. N., D. E. Harker, F. W. Lovejoy, Jr., R. E. Nye, Jr., and E. B Mahoney 1956. *Circulation*, N. Y. 13:680. [13] McCord, M. C., H. J. C. Swan, and S. G. Blount. 1954. *Am. Heart J.* 48:405. [14] Sepulveda, G., and D. S. Lukas 1955. *Circulation*, N. Y. 11:552. [15] Corlin, R., J. K. R. McMillan, W. E. Medd, M. B. Matthews, and R. Deley. 1955. *Am. J. Med.* 18:855. [16] Shepherd, J. T., J. E. Edwards, H. B. Burchell, H. J. C. Swan, and E. H. Wood. 1957. *Brit. Heart J.* 19:70. [17] Kulda, H., G. J. Dammann, F. W. Haynes, E. Rappaport, and L. Dexter. 1957. *Am. J. Med* 23:166. [18] Yu, P. N. 1956. *Ann. Int. M.* 49:1138. [19] Storzstein, O., S. Humerfeldt, O. Muller, and H. Rasmussen. 1952. *Acta med scand* 141:419 [20] Sobin, S. S., M. J. Carson, J. L. Johnson, and C. R. Baker. 1954. *Am. Heart J.* 48:416 [21] Joss, H. A., P. N. Yu, F. W. Lovejoy, Jr., R. E. Nye, Jr., and J. H. Simpson. 1954. *Am. J. Med.* 17:6. [22] Blount, S. G., H. Mueller, and M. C. McCord 1955. *Ibid* 18:871 [23] Scannell, J. G., G. S. Myers, and A. L. Friedlich. 1952. *Surgery* 32:164. [24] McKusick, V. A. 1952. *Bull. Johns Hopkins Hosp* 90:3 [25] Wilson, R. H., W. Hoseth, C. Sadoff, and M. E. Dempsey. 1954. *Am. Heart J.* 48:671. [26] Stead, E. A., Jr., J. V. Warren, and E. S. Brannon. 1948. *Ibid* 35:529. [27] Brannon, E. S., E. A. Stead, Jr., J. V. Warren, and A. J. Merrill. 1946. *Ibid* 31:407. [28] Merrill, A. J., J. V. Warren, E. A. Stead, Jr., and E. S. Brannon. 1946. *Ibid* 31:413. [29] Warren, J. V., E. S. Brannon, E. A. Stead, Jr., and A. J. Merrill. 1946. *Ibid* 31:418 [30] Westcott, R. N., N. O. Fowler, R. C. Scott, V. D. Hausenstein, and J. McGuire. 1951. *J. Clin. Invest.* 30:957. [31] Doyle, J. T., J. S. Wilson, and J. V. Warren. 1952. *Circulation*, N. Y. 5:263. [32] Yu, P. N., D. C. Beatty, F. W. Lovejoy, Jr., R. E. Nye, Jr., and H. A. Joss. 1956. *Am Heart J.* 52:683 [33] Wilson, R. H., W. Hoseth, and M. E. Dempsey. 1955. *Ann. Int. M.* 42:679. [34] Fowler, N. O., R. N. Westcott, R. C. Scott, and J. McGuire. 1951. *J. Clin. Invest.* 30:517. [35] Doyle, J. T., J. S. Wilson, E. H. Esler, and J. V. Warren. 1951. *Ibid* 30:345. [36] Warren, J. V., E. S. Brannon, H. S. Weens, and E. A. Stead, Jr. 1948. *Am. J. Med.* 4:193.

Part II. BLOOD FLOW AND RESISTANCE (Concluded)

Condition or Agent (A)	No. of Subjects (B)	Blood Flow		Resistance, dynes sec/(cm ⁵)			Reference
		Cardiac Index L/min/sq m (C)	Stroke Index ml/beat/sq m (D)	A-V O ₂ Difference ml/L (E)	Total Arterial (Systemic) (F)	Pulmonary Total (G)	Pulmonary Vascular (H)
37 Constructive	13	(1.27-3.91)					
38 pericarditis	5	(1.7-3.3)					
39	10	(1.18-4.57)	(43-104) ⁸				
40 Congestive heart failure	48	2.0(1.3-4.0)		68(32-108)			
41 Acute hemorrhage	13	2.5(1.6-5.0)		54(26-105)	1300(700-2427)		
42 Chest injury	13	3.8(1.8-5.0)		48(35-63)	1000(490-1326)		
43 Pericardial tamponade	7	2.7(1.5-4.0)		61(52-84)	1250(930-1960)		
44 Acute hypoxia	28	3.0		46			133
45 Before	28	3.1		43			202
46 After	8	3.4			1214		66
47 Before	8	4.4			962		119
48 After	12	3.2			1494		262
49 Before	12	3.8			1450		416
50 After	24	3.1(1.42-5.22)	39(19-72)	46.7(28.5-70.4)	1540(690-3900)	610(120-1500)	27(40-880)
51 100% oxygen	24	3.63(1.81-10.75)	39(20-101)	41.9(13.1-61.3)	1370(315-3140)	690(160-1550)	418(80-980)
52 Before		5.61				417	
53 After		4.81				366	
Norepinephrine							
54 Before	12	3.0		40	1500		109
55 After	12	2.9		41	2090		118
Saline infusion, intravenous							
56 Before	12	3.6		43			
57 After	12	4.3		38			
58 Before	10	(2.3-5.0)		(23-46)			
59 After	10	(2.4-6.3)		(22-49)			
5% albumin infusion, intravenous							
60 Before	7	(2.5-4.6)		(26-56)			
61 After	7	(2.6-5.9)		(22-49)			
1/1 Cardiac output, L/min. 1/8 Stroke volume, ml/beat.							

Contributor: Yu, Paul N.

References. [1] Dexter, L., J. L. Whittenberger, R. Gorlin, B. M. Lewis, F. W. Haynes, and R. J. Spiegel. 1951. *Tr. Ass. Am. Physicians* 65:226.

[2] Yu, P. N., F. W. Lovejoy, Jr., H. A. Joos, R. E. Nye, Jr., and W. S. McCann. 1953. *J. Clin. Invest.* 32:130. [3] Harvey, R. M., M. I. Ferrer, D. W.

121. CHRONIC CONSTRUCTIVE PERICARDITIS: MAN (Concluded)
Part II INCIDENCE OF ACCOMPANYING CONDITIONS (Concluded)

Condition	No. of Subjects	Incidence %	Reference
(A)	(B)	(C)	(D)
	104	39	2-5
9 Cyanosis	59	90	4, 6, 7
10 Pulsus paradoxus	206	27	1-4, 6, 8
11 Atrial fibrillation or flutter	98	42	5, 8, 9
12 Protodiastolic sound	26	96	3, 9, 10
13 Right ventricular diastolic dip	78	22	2, 5
14 Albuminuria	267	52	2, 4, 8
15 Pericardial calcification ¹	215	50	1-4, 8
16 Enlarged cardiopericardial silhouette ¹	192	93	1-4, 6, 8
17 Diminished cardiac pulsations ¹	240	61	1-6, 8
18 Pleural effusion ¹	22	95	2
19 Low voltage ²	22	77	2
20 Low, inverted T waves ²			

1/1 Roentgenographic determination. 1/2 Electrocardiographic determination

Contributors: McGuire, Johnson, and Robert A. Helm

References [1] Chambliss, J. R., E. J. Jaruszewski, B. L. Drofman, J. F. Martin, and H. Feil. 1951. Circulation, N. Y. 4:816. [2] Deterling, R. A., and G. H. Humphries. 1955. Ibid. 12:30. [3] Harrington, S. W. 1944. Ann. Surg. 120:468. [4] Stewart, H. J., J. R. Carty, and J. R. Seal. 1943. Am. J. Roentg. 49:349. [5] Mortensen, V., and E. Warburg. 1948. Acta med. scand. 131:203. [6] Blalock, A., and C. S. Burwell. 1941. Surg. Gyn. Obst. 73:433. [7] Dalton, J. C., R. J. Pearson, Jr., and F. D. White. 1956. Ann. Int. M. 45:445. [8] Paul, O., B. Castleman, and P. D. White. 1948. Am. J. M. Sc. 216:361. [9] McKusick, V. A. 1952. Bull. Johns Hopkins Hosp. 90:3. [10] Wilson, R. H., W. Hoseth, C. Sadoff, and M. E. Dempsey. 1954. Am. Heart J. 48:671

122. HEMODYNAMICS OF ACUTE CARDIAC TAMPONADE. MAN

Values in parentheses are ranges, estimate "c" (cf Introduction).

Variable	No. of Subjects	Value	Reference
(A)	(B)	(C)	(D)
1 Heart rate, beats/min	20	116(76-150)	1, 2
2 Respiratory rate, breaths/min	13	27(22-35)	1
Arterial blood pressure, mm Hg			1-3
3 Systolic	21	86	
4 Diastolic	21	57	
5 Respiratory variation in arterial systolic pressure, mm Hg	7	26(16-49)	2
6 Venous pressure, mm saline	13	240	1
7 Arterial O ₂ saturation, %	3	94(92-97)	3
8 Cardiac index (Fick), L/min/sq m	7	2.5(1.5-4.0)	2, 3
Pulmonary artery pressure, mm Hg			3
9 Systolic	3	29(20-39)	
10 Diastolic	3	25(11-25)	
Right ventricular pressure, mm Hg			3
11 Systolic	3	29(20-39)	
12 Diastolic	3	15(8-17)	
13 Right ventricular pressure ratio (diastolic/systolic), %	3	46(38-61)	3
14 Right atrial pressure (mean), mm Hg	10	12.3(8-23)	2, 3
15 Superior vena caval pressure (mean), mm Hg	3	15.7(8-23)	3

Contributors: McGuire, Johnson, and Robert A. Helm

References [1] McGuire, J., and E. J. McGrath. 1941. Tr. Ass. Am. Physicians 56:194. [2] Warren, J. V., E. S. Brannon, E. A. Stead, Jr., and A. J. Merrill. 1946. Am. Heart J. 31:418. [3] Wilson, R. H., W. Hoseth, C. Sadoff, and M. E. Dempsey. 1954. Ibid. 48:671.

121. CHRONIC CONSTRUCTIVE PERICARDITIS. MAN

Part I: HEMODYNAMICS

Values in parentheses are ranges, estimate "c" (cf. Introduction).

	Variable (A)	No. of Subjects ¹ (B)	Value (C)	Reference (D)
1	Age, yr	125	47.4(6-77)	1-8
2	Heart rate, beats/min	89	96(60-150)	1, 2
3	Arterial blood pressure, mm Hg			2, 5-7, 9, 10
4	Systolic	98	110(80-210)	
5	Diastolic	98	78(50-120)	
6	Pulse pressure, mm Hg	61	33(10-70)	2
7	Circulation time (arm to tongue), sec	56	26(10-88)	2, 3, 8
8	Vital capacity, ml	49	1885(575-4000)	2
9	Venous pressure, cm H ₂ O	107	25.8(13-41)	1-5, 8-12
10	White blood cell count	61	6,870(4,300-13,300)	2
11	Blood urea nitrogen, mg %	61	12.2(6.0-22.5)	2
12	Serum protein, g %	59 ² , 15 ³	6.17 ² (3.8-8.8) ³	2, 3 ² , 3 ³
13	Serum albumin, g %	39 ² , 15 ³	3.46 ² (2.5-5.5) ³	2 ² , 3 ³
14	Serum globulin, g %	39 ² , 15 ³	2.61 ² (1.6-3.8) ³	2 ² , 3 ³
15	Blood volume ⁴ , ml/kg	19	93.5(71-112) ⁵	2
16	Arterial O ₂ saturation, %	27	93.9(85-100)	5, 7, 13
17	Cardiac index (Fick), L/min/sq m	30	2.39(1.2-3.9)	5, 7, 13
18	Pulmonary artery "wedge" pressure (mean), mm Hg	14	18.1(8-28)	5, 7, 13, 14
19	Pulmonary artery pressure, mm Hg			5, 7, 13, 14
20	Systolic	29	34.5(18-45)	
21	Diastolic	29	18.5(8-31)	
22	Right ventricular pressure, mm Hg			5, 7, 8, 13, 14
23	Systolic	35	37.4(18-57)	
24	Diastolic	35	17.2(8-32)	
25	Right atrial pressure (mean), mm Hg	18	17.9(10-25)	5, 7, 13, 14
26	Superior vena caval pressure (mean), mm Hg	9	18.0(15-22)	7
27	Glomerular filtration rate, ml/min	7	107.7(80-141)	13
28	Renal plasma flow, ml/min	7	350.1(171-595)	13
29	Filtration fraction	7	0.312(0.242-0.454)	13

/1/ Distribution among the sexes in 257 cases of chronic constrictive pericarditis: 75% males, 25% females [1-5, 8, 9]. /2/ and /3/ Matching footnotes in column D indicate source of data. /4/ Vital-red technique, /5/ Normal range, 70-100 ml/kg.

Contributors: McGuire, Johnson, and Robert A. Helm

References: [1] Blalock, A., and C. S. Burwell. 1941. Surg. Gyn. Obst. 73:433. [2] Chambliss, J. R., E. J. Jaruszewski, B. L. Brofman, J. F. Martin, and H. Feil. 1951. Circulation, N. Y. 4:816. [3] Deterling, R. A., and G. H. Humphries. 1955. Ibid. 12:30. [4] Harrington, S. W. 1944. Ann. Surg. 120:468. [5] Sawyer, C. G., C. S. Burwell, L. Dexter, E. C. Eppinger, W. T. Goodale, R. Gorlin, D. E. Harken, and F. W. Haynes. 1952. Am. Heart J. 44:207. [6] Stewart, H. J., J. R. Carty, and J. R. Seal. 1943. Am. J. Roentg. 49:349. [7] Wilson, R. H., W. Hoseth, C. Sadoff, and M. E. Dempsey. 1954. Am. Heart J. 48:671. [8] Yu, P. N. G., F. W. Lovejoy, Jr., H. A. Joos, R. E. Nye, Jr., and E. B. Mahoney. 1953. Circulation, N. Y. 7:102. [9] Mortensen, V., and E. Warburg. 1948. Acta med. scand. 131:203. [10] Cushing, E. H., and H. S. Feil. 1936. Am. J. M. Sc. 192:327. [11] Burwell, C. S., and A. Blalock. 1938. J. Am. M. Ass. 110:265. [12] Burwell, C. S., and D. Flickinger. 1935. Arch. Int. M. 56:250. [13] McKusick, V. A. 1952. Bull. Johns Hopkins Hosp. 90:3. [14] Lin, T. K., and M. Anache. 1956. Am. Heart J. 51:340.

Part II INCIDENCE OF ACCOMPANYING CONDITIONS

	Condition (A)	No of Subjects (B)	Incidence % (C)	Reference (D)
1	Dyspnea	273	76	1-5
2	Orthopnea	109	34	1-3
3	Cough	46	41	2, 5
4	Precordial pain	45	31	2, 5
5	Edema	295	73	1-6
6	Ascites	247	80	1-6
7	Hepatomegaly	93	99	1-6
8	Distended cervical veins	125	86	1, 3-5

Part I. ANEMIA, ARTERIO-VENOUS COMMUNICATION,
BERIBERI, PAGET'S DISEASE, THYROTOXICOSIS (Concluded)

Contributor. Helm, Robert A.

References: [1] Bishop, J. M., K. W. Donald, and O. L. Wade. 1955. *Clin. Sc.*, Lond. 14:329. [2] Brannon, E. S., A. J. Merrill, J. W. Warren, and E. A. Stead, Jr. 1945. *J. Clin. Invest.* 24:332. [3] Sharpey-Schafer, E. P. 1944. *Clin. Sc.*, Lond. 5:125. [4] Sharpey-Schafer, E. P. 1945. *Lancet*, Lond. 2:296. [5] Cohen, S. M., O. G. Enholm, S. Howarth, J. McMichael, and E. P. Sharpey-Schafer. 1948. *Clin. Sc.*, Lond. 7:36. [6] Burwell, C. S., and L. Dexter. 1947. *Tr. Ass. Am. Physicians* 60:59. [7] Fowler, N. O. 1953. *Ann. Int. M.* 38:478. [8] Lahay, W. J., D. B. Arst, M. Silver, C. R. Kleeman, and P. Kunkel. 1953. *Am. J. Med.* 14:248. [9] Porter, R. R., and R. S. Downs. 1942. *Ann. Int. M.* 17:645. [10] Campbell, J. A., L. A. Silverstone, and D. L. Donovan. 1951. *J. Clin. Invest.* 30:632. [11] Howarth, S. 1953. *Clin. Sc.*, Lond. 12:171. [12] Rappaport, E., H. Kusida, L. Dexter, P. H. Henneaman, and F. Albright. 1957. *Am. J. Med.* 22:252. [13] Myers, J. D., E. S. Brannon, and B. C. Holland. 1950. *J. Clin. Invest.* 29:1669.

Part II. CHRONIC PULMONARY EMPHYSEMA, WITH AND WITHOUT COR PULMONALE

For additional information on the subject, see references 4-9.

Variable	Emphysema						Reference
	Without Cor Pulmonale		With Chronic Cor Pulmonale				
	Compensated		Compensated		Decompensated		
(A)	No. of Sub-jects (B)	Value (C)	No. of Sub-jects (D)	Value (E)	No. of Sub-jects (F)	Value (G)	(H)
1 Ratio of residual volume to total capacity, %	28	62(40-86)	18	56(23-83)	5	56(42-73)	B-E, 1, 2, F, G, 2
2 Arterial O ₂ saturation, %	28	86(70-98)	22	79(62-90)	12	67(51-86)	B, C, 1, 2, D, E, 1-3, F, G, 2, 3
3 Maximal breathing capacity, % predicted	11	39(22-58)	11	43(21-77)	5	26(13-51)	B-G, 2
4 CO ₂ pressure, mm Hg	12	46(37-55)	11	50(38-58)	6	64(53-72)	
5 Hematocrit, %	27	44(34-51)	22	52(47-57)	12	63(54-70)	B, C, 1, 2, D, E, 1-3, F, G, 2, 3
6 Total blood volume, ml/kg m	28	3024(180-3740)	22	3630(2909-4035)	12	4893(3820-5070)	
7 Pulmonary artery pressure (systolic/diastolic), mm Hg	27	32/17 (23-71/7-37)	22	44/22 (26-76/8-37)	12	72/54 (47-110/8-24)	

Contributors (a) Helm, Robert A. (b) Fowler, Noble O.

References [1] Borden, C. W., R. H. Wilson, R. V. Ebert, and H. S. Wells. 1950. *Am. J. Med.* 8 701. [2] Harvey, R. M., M. I. Ferrer, D. W. Richards, Jr., and A. Courmand. 1951. *Ibid.* 10:719. [3] Fowler, N. O., R. N. Westcott, R. C. Scott, and E. Heas. 1952. *Circulation*, N. Y. 6:888. [4] Dexter, L., et al. 1951. *Tr. Ass. Am. Physicians* 44 22b. [5] Yu, P. N., et al. 1953. *J. Clin. Invest.* 32 130. [6] Hecht, H. H. 1956. *Circulation*, N. Y. 14,265. [7] Whitaker, W. 1954. *Q. J. Med.*, Oxf. 47:57. [8] Mounsey, P., et al. 1952. *Brit. Heart J.* 14 153. [9] Denolin, H., and J. Leguizans. 1951. *Arch. mal coeur* 5 391.

123. HEMODYNAMICS IN CONDITIONS OF HIGH CARDIAC OUTPUT: MAN

Cardiac output determined by catheterization. Values in parentheses are ranges, estimate "c" (cf. Introduction).

Part I ANEMIA, ARTERIO-VENOUS COMMUNICATION, BERIBERI, PAGET'S DISEASE, THYROTOXICOSIS

Variable	No. of Subjects	Value	Reference
(A)	(B)	(C)	(D)
Anemia			
1 Age, yr	16♂, 34♀	45(17-53)	1-4
2 Hemoglobin, g/100 ml	49	5.5(1.9-11.2)	
3 Heart rate, beats/min	48	91(63-124)	
4 Arterial blood pressure (systolic/diastolic), mm Hg	47	140/59(94-185/40-99)	
5 O ₂ consumption, ml/min	46	249(167-348)	1-3
6 Arterio-venous O ₂ difference, vol %	46	3.0(1.8-5.1)	
7 Cardiac output, L/min	50	8.6(3.9-14.3)	1-4
8 Cardiac index, L/min/sq m	28	5.26(2.5-8.8)	1,2
9 Cardiac output, L/min (Hb <6 g/100 ml)	34	9.07(6.9-13.6)	1-4
Arterio-venous Communication			
10 Age, yr	13♂	28(20-51)	1,5
11 Heart rate, beats/min	13	80(55-90)	
12 Arterial blood pressure (systolic/diastolic), mm Hg	12	120/63(98-134/55-78)	
13 O ₂ consumption, ml/min	13	308(243-367)	
14 Arterio-venous O ₂ difference, vol %	13	2.8(2.0-4.2)	
15 Cardiac output, L/min	13	11.65(6.4-16.1)	
Beriberi			
16 Age, yr	4♂, 1♀	29(23-35)	6-9
17 Heart rate, beats/min	5	104(89-124)	
18 Arterial blood pressure (systolic/diastolic), mm Hg	5	128/69(100-150/55-84)	
19 Pulmonary artery pressure (systolic/diastolic), mm Hg	3	44-25(34-64/15-36)	6-8
20 Venous pressure, cm H ₂ O	4	26.6(19.5-31.0)	6,8,9
21 O ₂ consumption, ml/min	5	114(9-14)	6-9
22 Cardiac output, L/min	5	310(259-355)	
23 O ₂ difference, vol %	5	3.4(2.2-6.2)	
24 Cardiac index, L/min/sq m	5	10.47(4.7-16.1)	
25 Cardiac output, L/min	5	5.67(2.9-7.8)	7-10
Paget's Disease¹			
26 Age, yr	7♂, 11♀	63(41-77)	11,12
27 Heart rate, beats/min	18	79(60-88)	
28 Arterial blood pressure (systolic/diastolic), mm Hg	17	140/73(116-160/55-90)	
29 O ₂ consumption, ml/min	18	268(147-328)	
30 O ₂ difference, vol %	18	3.5(2.0-5.5)	
31 Cardiac output, L/min	18	7.66(3.8-13.3)	
32 Cardiac index, L/min/sq m	12	33(10-64)	11
33 Cardiac output, L/min (skeletal involvement <30%)	12	57(11.5-120.5)	
34 Cardiac output, L/min (skeletal involvement >30%)	6	56(17-114)	12
35 Cardiac output, L/min (alkaline phosphatase >45 K-A units ²)	11	9.21(7.1-13.3)	11,12
36 Cardiac output, L/min (alkaline phosphatase <45 K-A units ²)	7	5.21(3.8-6.3)	
37 Cardiac output, L/min (alkaline phosphatase >45 K-A units ²)	15	8.16(3.8-13.3)	
38 Cardiac output, L/min (alkaline phosphatase <45 K-A units ²)	3	5.13(4.2-5.7)	11
Thyrototoxicosis			
39 Age, yr	10♂, 19♀	35(12-62)	1,13
40 Heart rate, beats/min	26	104(64-150)	
41 Arterial blood pressure (systolic/diastolic), mm Hg	25	142/75(106-183/50-96)	
42 Pulmonary artery pressure (systolic/diastolic), mm Hg	14	20.6(13-27)	
43 Cardiac output, L/min	20	36.5(23-54)	13
44 Cardiac index, L/min/sq m	20	2.5(-1 to 4.6)	
45 Cardiac output, L/min	24	35.8(27.5-42.1)	
46 Cardiac output, L/min	29	32.5(23.4-47.6)	1,13
47 Cardiac output, L/min	29	3.7(2.2-4.9)	
48 Cardiac output, L/min	29	5.65(3.3-7.3)	
49 Cardiac output, L/min	29	46.1(+23 to +111)	

1/1 Osteitis deformans generalized 1/2 Amount of phosphatase which, when allowed to act upon an excess of disodium phenylphosphate at pH 9 for 30 minutes (37.5°C), will liberate 1 mg of phenol. 1/3 Amount of phosphatase required to liberate 1 mg of phosphorus in the form of the phosphorus ion, in 1 hour of incubation with a substrate of sodium β -glycerophosphate.

Values in parentheses are ranges, estimate ¹⁰ (cf. Introduction).

Condition	No. of Subjects	Body Surface Area, sq m	Cardiac Index, L/min/sq m	Stroke Volume, ml/min/sq m	O ₂ Consumption, ml/min/sq m	Arterio-venous O ₂ Difference, vol %	Arterial O ₂ Concentration, vol %	Arterial CO ₂ Pressure, mm Hg	Pulmonary Artery Pressure, mm Hg			Reference				
									Systolic	Diastolic	Mean					
	(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)	(I)	(J)	(K)	(L)	(M)	(N)	(O)	(P)
1 Normal																
2 Obstructive pulmonary emphysema	11	1.56	(2.7-3.5)	70	130	4.2	19	(94-98)	40	15	30	5	22	1		
3	11	1.56	(2.7-3.5)	70	130	4.2	19	(94-98)	40	15	30	5	22	1		
4	11	1.56	(2.7-3.5)	70	130	4.2	19	(94-98)	40	15	30	5	22	1		
5	11	1.56	(2.7-3.5)	70	130	4.2	19	(94-98)	40	15	30	5	22	1		
6 Alveolar-capillary diffusion block	12	1.56	(2.6-4.9)	63	151	4.1	(3.1-5.6)	(78-98)	38	14	23					4
7 Primary pulmonary hypertension	3	1.53	1.9	33	126	6.8	18.2	98	81	43	62	80	132	5		
8	10	1.5	1.6	33	126	6.8	20.0	92	111	59	75			6		
9	3	1.5	1.6	33	126	6.8	20.0	92	111	59	75			6		
10	3	1.5	1.6	33	126	6.8	20.0	92	111	59	75			6		
						</										

1/ Recently in cardiac failure. 2/ End diastolic pressure. 3/ In reversible cardiac failure. 4/ In cardiac failure at time of observation. 5/ Adults of various ages, distribution among the sexes approximately equal. 6/ Majority of observations in young patients, most of whom were females. 7/ Advanced cases, representing extremes in this condition.

Contributor: Samet, Philip

References: [1] Harvey, R. M., M. J. Ferrer, D. W. Richards, Jr., and A. Cournaud. 1951. *Am J Med* 10:719. [2] Lewis, C. S., A. J. Samuels, M. C. Dafoe, and H. H. Hecht. 1952. *Circulation* N Y 6:874. [3] Fowler, N. O., R. N. Westcott, R. C. Scott, and E. H. Hays. 1952. *Ibid* 6:888. [4] Austrian, R., J. H. McClement, A. D. Hemmelt, Jr., K. W. Donald, R. L. Riley, and A. Cournaud. 1951. *Am J Med* 11:567. [5] Dresdale, D. T., N. Schutt, and R. J. Michtom. 1951. *Ibid* 11:686. [6] Shepherd, J. I., J. E. Edwards, H. B. Burchell, R. J. C. Swan, and E. H. Wood. 1957. *Br R Heart* 19:70. [7] Schafer, H., R. C. Blain, and R. J. Bing. 1956. *Ann Int M* 44:505. [8] Chapman, D. W., J. P. Abbott, and J. L. Latham. 1957. *Circulation* N Y 15:35.

126. CEREBRAL FUNCTION IN COR PULMONALE WITH CO₂ RETENTION: MAN

Condition	Blood Flow, ml/100 g/min	Vascular Resistance, mm Hg/100 g/min	O ₂ Consumption, ml/100 g/min	Mean Arterial Pressure, mm Hg	Arterial CO ₂ Pressure, mm Hg	Arterial 100% Saturation	Arterial pH	Reference
1 Breathing air	50	6.8	3.0	2.7	91	96	7.40	(P)
2 Breathing O ₂	44	7.9	3.2	2.7	91	96	7.40	1
3 Breathing O ₂	55	7.82	3.64	97	49	96	7.38	2
4 Exercising	53	1.85	2.93	103	54	96	7.34	2

1/ 8 normal controls, 9 patients. [1] 2/ Unstated number of controls, 9 patients [2]

Contributor: Samet, Philip

References: [1] Patterson, J. L., Jr., A. Heyman, and T. W. Duke. 1952. *Am J Med* 12:382. [2] Scheinberg, P., I. Blackburn, M. Saslaw, M. Rich, and G. Baum. 1953. *J Clin Invest* 32:750.

Part I: CARDIAC INDEX AND PULMONARY ARTERY PRESSURE VS ARTERIAL O₂ SATURATION

Measurements made under resting conditions. Cardiac output calculated from O₂ content of mixed venous blood and arterial blood, and from O₂ uptake by the lungs (sample of mixed venous blood obtained from pulmonary artery or right ventricle through a catheter). Cardiac index then calculated by dividing cardiac output by body surface area in square meters. Pressure measured, through a catheter in the pulmonary artery, with a high frequency manometer. Values in parentheses are ranges, estimate "c" (cf. Introduction).

Arterial O ₂ Saturation %	Cardiac Index ¹ L/min/sq m	Pulmonary Artery Pressure ² mm Hg	Reference
(A)	(B)	(C)	(D)
1 Above 90	3.7(2.2-6.6)	18(9-28)	B, 1-5; C, 1-4
2 85-90	3.2(2.3-5.4)	26(15-38)	B, 1-6; C, 1-4, 6
3 80-85	3.6(1.9-5.8)	28(13-43)	B, 1-6; C, 1, 2, 4, 6
4 75-80	3.6(2.7-4.9)		1-3, 5, 6
5 70-80		34(16-57)	1, 2, 6
6 70-75	3.8(2.6-6.4)		2, 3, 5, 6
7 70 and below	3.7(1.7-6.1)	53(44-74)	B, 1, 3-6; C, 1, 4, 6


/1/ No demonstrable relationship in any one patient between cardiac index and arterial O₂ saturation. /2/ Integrated mean pressure.

Contributors: (a) Ebert, Richard V., (b) Stone, Daniel J.

References: [1] Harvey, R. M., M. I. Ferrer, D. W. Richards, Jr., and A. Cournand. 1951. Am. J. Med. 10:719. [2] Yu, P. N. G., F. W. Lovejoy, Jr., H. A. Joos, R. E. Nye, Jr., and W. S. McCann. 1953. J. Clin. Invest. 32:130. [3] Dexter, L., J. L. Whittenberger, R. Gorlin, B. M. Lewis, F. W. Haynes, and R. J. Siegel. 1951. Tr. Ass. Am. Physicians 64:226. [4] Sancetta, S. M. 1955. Am. Heart J. 49:501. [5] Mounsey, J. P. D., L. W. Ritzman, N. J. Selverstone, W. A. Briscoe, and G. A. McLemore. 1952. Brit. Heart J. 14:153. [6] Fowler, N. O., R. N. Westcott, R. C. Scott, and E. Hess. 1952. Circulation, N. Y. 6:888.

Part II. MEAN PULMONARY ARTERY PRESSURE VS ARTERIAL O₂ SATURATION

Graph constructed from data in column C, Part I.



Contributor: Stone, Daniel J.

129. CEREBRAL BLOOD FLOW AND O₂ CONSUMPTION IN ENDOCRINE DISORDERS: MAN

Method X = intermittent sampling method of Kety [13], S = continuous sampling method of Scheinberg [14]. Values in parentheses are ranges, estimate "b" unless otherwise indicated (cf. Introduction).

	No. of		Brain		
16	8	K	45(15-55)	2.7	11
17	6	K	65(38-92)	1.7(1.3-2.1)	11
18	7	K	62(51-73)	2.6(1.9-3.3) ¹	12

¹/ 1.9(0.9-2.9) in coma

Contributor: Gordan, Gilbert S.

References. [1] Kennedy, C. 1956. Progress in neurobiology. Harper and Brothers, New York. v. 1, p. 230. [2] Gordan, G. S. 1956. Hormones and the aging process. Academic Press, New York. p. 299. [3] Fazekas, J. F. 1955. Am. J. Med. 18:477. [4] Fazekas, J. F. 1953. J. Am. Geriatr. Soc. 1:836. [5] Gordan, G. S. 1956. Recent Progr. Hormone Res. 12:153. [6] Gordan, G. S. 1951. Ann. N. York Acad. Sc. 54:575. [7] Sensenbach, W. 1954. J. Clin. Invest. 33:1434. [8] Scheinberg, P. 1950. Ibid. 29:1010. [9] Sokoloff, L. 1953. Ibid. 32:202. [10] Schieve, J. F. 1951. Ibid. 30:1527. [11] Kety, S. S. 1949. Ibid. 27:500. [12] Kety, S. S. 1948. Am. J. Psychiat. 104:765. [13] Kety, S. S., and C. F. Schmidt. 1948. J. Clin. Invest. 27:476. [14] Scheinberg, P., and E. A. Stead, Jr. 1949. Ibid. 28:1163.

127. RENAL FUNCTION IN CONDITIONS OF NORMAL, LOW, AND HIGH CARDIAC OUTPUT: MAN

Values in parentheses are ranges, estimate "c" (cf. Introduction).

Variable	Normal		Chronic Rheumatic Heart Disease (Low Output)		Chronic Cor Pulmonale (High Output)		Reference
	No. of Subjects	Value	No. of Subjects	Value	No. of Subjects	Value	
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)
1 Glomerular filtration rate, ml/min	14	145(119-174)	8	68(50-94)	10	76(28-134)	1
2							
3							
4							
5							
6							
7 output, %							
7 Renal arterio-venous O ₂ difference, vol %	3	2.1(1.9-2.5)	5	3.8(3.2-4.4)	2	2.1(2.0-2.25)	1

Contributor: Helm, Robert A.

References: [1] Davies, C. E., and J. A. Kilpatrick. 1951. Clin. Sc., Lond. 10:53. [2] McMichael, J., and E. P. Sharpey-Schafer, 1944. Brit. Heart J. 6:33.

128. RIGHT HEART CATHETERIZATION VALUES IN ACQUIRED CARDIAC DISEASES, AT REST AND DURING EXERCISE: MAN

Exercise: pedalling (bicycle ergometer) at approximately 40 revolutions per minute for a period of approximately 5 minutes. Values in parentheses are ranges, estimate "c" (cf. Introduction).

Variable	Condition	Normal	Mitral Stenosis	Mitral Insufficiency	Aortic Stenosis	Left Ventricular Failure ¹
(A)	(B)	(C)	(D)	(E)	(F)	(G)
1 Number of subjects		25	24	20	8	8
2 Right atrial pressure (mean), mm Hg	Rest	6(2-10)	6(2-17)	8(2-16)	5(1-15)	5(2 to +8)
3 Pulmonary artery pressure (mean), mm Hg	Rest	17(10-22)	40(18-72)	44(19-78)	35(10-61)	32(15-46)
4 Pulmonary artery "wedge" pressure (mean), mm Hg	Exercise	24(17-33)	65(35-112)	54(28-80)		49(29-63)
5 Pulmonary artery "wedge" pressure (mean), mm Hg	Rest	12(8-15)	24(12-46)	23(14-39)	23(8-37)	
6	Exercise		34(23-47)	32(21-42)		
7 Cardiac index, L/min/sq m	Rest	3.5(2.8-6.3)	2.6(1.5-3.9)	2.6(1.6-3.9)	2.7(2.2-4.2)	2.9(2.1-3.7)
8	Exercise	5.7(4.8-7.7)	3.5(1.8-5.5)	3.2(2.0-4.1)		2.7(1.9-4.3)
9 Pulmonary arteriolar resistance, dynes sec cm ⁻⁵	Rest	67(44-110)	350(90-1110)	200(40-490)		
10	Exercise		550(40-2000)	300(70-590)		
11 Total pulmonary resistance, dynes sec cm ⁻⁵	Rest	189(90-290)	750(320-1870)	600(440-990)		514(260-1070)
12	Exercise	162(110-230)	1050(420-3200)	750(390-1180)		839(362-1420)

¹/ Excluded are diseases (such as hyperthyroidism, arterio-venous fistula, beriberi) resulting in left ventricular failure, with so-called "high output" failure.

Contributor: Witham, A. Calhoun

References: [1] Anderson, M. W., and E. H. Wood. 1957. Clinical cardiopulmonary physiology. Grune and Stratton, New York. [2] Ball, D., H. Kopelman, and A. C. Witham. 1952. Brit. Heart J. 14:363. [3] Hickam, J. E., and W. H. Cargill. 1948. J. Clin. Invest. 27:10.

Specification (A)	Animal (B)	Graves' Disease ¹ (C)	Simmonds' Disease ² (D)	Myxedema ³ (E)	Addison's Disease ⁴ (F)	Diabetes ⁵ (G)	Reference (H)
Blood (concluded)							
pH	Man					-6	G,7
	Dog				-		F,60,61
Total blood count	Man			-	+		E,75,78
	Cat				+		F,62
	Dog		-		+		D,52,57,63,F,53,54,64
	Rabbit				+		F,62
	Rat		-	-	+		D,55,57,65,E,66,F,62
	Rat				-		F,66
RBC count	Man					+	G,6
RBC life span	Man			-			E,67
	Rat		0				D,68
O ₂ capacity	Man				+		F,4
	Dog				+		F,64
CO ₂ capacity	Cat				-		F,65
	Dog				-		F,60,64
Arterial O ₂ saturation	Man					0	F,32
Venous O ₂ saturation	Man					+	F,32
Glucose (fasting)	Man	+	-	0	-	+	C,17,43,70,D,2,71;E,1,72;F,4,17,49,G,5-7,28
	Cat				+		F,73,74,G,6
	Dog		-		-	+	D,9,75,F,76,G,6
	Monkey		-		+		D,75,G,6
	Pig				+		G,6
	Rabbit		-			+	D,75
	Rat		-				D,75
Insulin	Man		-		-		D,c,F,77,78
	Dog		-				D,79
	Rat		-			-	G,80
Cholesterol	Man	-	0	+		+	D,79
	Man					+	C,91,92,D,63,E,25,81,82,G,6,7,84,85
	Dog		-			0	G,5,26,86
	Dog				+	+	D,87,F,88,G,89-91
	Monkey			0	0		F,92
	Rabbit		-				E,93
	Rat						D,11
Phospholipids	Man	-		+	0		E,94,F,88
	Dog					+	C,81,82,G,6,84,85
Total fatty acids	Man		-			+	D,87,G,c
	Dog					+	G,c
β -Hydroxybutyric acid	Man		-			+	D,87,G,c
	Dog					+	G,6
Creatinine	Man					+	G,91
Creatinine	Cat			-			E,25
Glycogen	Man				-		F,73
Thiocyanate	Man					-	G,28
Insulin sensitivity	Man	-	+	+	+	+	G,95
	Man				+	-	C,E,3,D,2,71;F,4,17,G,6
	Cat			0		-6	E,96,G,7
	Dog			+			E,97-99
	Monkey		+	+	+	-	D,100,101;E,97-99
	Rabbit		+				F,76,G,c
	Rat		+				D,101-103
Glucose tolerance	Man	+	+	+	+	+	D,101
	Man					+	D,101,F,7
	Dog			0			C,3,17,104,D,2,105;E,3,25,F,4,G,5-7,26
			+		+	+	E,72
					+	+	D,106,F,c,G,107

1/1 Hyperthyroidism. 1/2 Hypopituitarism. 1/3 Hypothyroidism. 1/4 Hypoadrenalism. 1/5 Hypoparathyroidism.

130. EFFECT OF ENDOCRINE DISORDERS: MAMMALS

Symbols: (+) = Increase, (-) = decrease, (0) = no change.

Specification	Animal	Graves' Disease ¹	Simmonds' Disease ²	Myxedema ³	Addison's Disease ⁴	Diabetes ⁵	Reference
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)
1 Body temperature	Man	+	-	-	-	-	C,1,D,2;E,3,F,4,G,5
2	Man					-6	G,5,7
3	Cat				-		F,8
4	Dog		-		-		D,9,10,F,8
5	Rabbit		0				D,11
6	Rat		-	-	-		D,12;E,13;F,14-16
7 Activity	Man	-	-	-	-		C,E,17;D,2;F,4
8	Cat		-		-		F,8
9	Dog		-		-		D,9,F,8
10	Rat					-	G,18
11 Spontaneous running	Rat	-	-	-	-		C,19,20;D,21;E,20,22;F,23
12	Rat			0			E,19
13 Forced activity	Rat				-		F,24
14 Heart							
Size	Man	+	-	+	-	+	C,4,17;D,1,E,17,25,26;F,4,27;G,28
15	Rat				-		F,29
16 Rate	Man	+	-	-		+	C,30,D,8,E,25,31;G,5,6,32
17	Dog		-				D,b
18	Monkey		-				D,b
19 Cardiac output	Man	+	-	-	-		C,33-35,D,36;E,25,33,34,F,c
20	Dog		-		-		D,36,F,37
21	Monkey		-				D,b
22 ECG							
Amplitude ⁷	Man			-			E,17,25,38
23 Height	Man				-		F,17
24 ST segment	Man				-		G,39
25 PR interval	Man				+		F,17,27
26 QT interval	Man				+	+	F,17,27;G,39
27 Pressure							
Blood	Man	+8	-	+	-9	-	C,4;D,2;E,17,25,26;F,4,17;G,5,6,32
28	Man					-6	G,7
29	Dog		-		-		D,9,40,41;F,37
30	Rat				-		F,42
31 Pulse	Man	+	-	-			C,d,D,b,E,38
32	Dog		-				D,b
33 Capillary permeability	Man			+			E,43
34	Dog				+		F,44
35	Rat				+		F,44
36 Circulation time	Man	-	+	+			C,3;D,1,E,3,38
37 Prothrombin time	Dog					-	G,45
38 Blood							
Volume	Man	+	-	-	-	-6	C,46,D,f,E,47,48;F,49;G,6,7
39	Cat				-		E,48
40	Dog				-		F,29
41	Rat		-				D,50
42	Rat		0				D,51
43 Hemoglobin	Man		-	-	-	+	D,2;E,3,F,17,G,32
44	Man					+6	G,6
45	Cat				+		F,8
46	Dog		-		+		D,52,F,53
47	Dog		-		0		F,54
48	Rat		-		-		D,50,55,56,F,57
49 Hematocrit	Man		-	-	+	+	D,7,E,3,F,4,G,32
50	Cat				-		F,8
51	Dog				+		F,53,58
52	Rat		-	0	-		D,50,55,56,E,59,F,57

/1/ Hyperthyroidism. /2/ Hypopituitarism. /3/ Hypothyroidism. /4/ Hypoadrenalism. /5/ Hypoinsulinism.
 /6/ In ketosis. /7/ T wave inverted. /8/ Systolic and mean, right ventricular. /9/ Systolic and diastolic.

130. EFFECT OF ENDOCRINE DISORDERS: MAMMALS (Continued)

Specification	Animal	Graves' Disease ¹	Simmonds' Disease ²	Myxedema ³	Addison's Disease ⁴	Diabetes ⁵	Reference
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)
Blood electrolytes (concluded)							
Urea	Man		0			+6	D,2,G,6
	Cat				+		F,73,74
	Dog				+		F,53,58
	Rat			+	+		E,139,F,8
Uric acid	Man			+		0	G,118
	Cat				-		F,73
	Dog				0		F,53,54
Blood flow							
Brain	Man	+		-			C,f,E,34
	Rat				-		F,140
Lung	Man	+					C,33
Muscle	Man	+					C,141,142
Peripheral artery	Man					-6	G,5
	Man						G,7,32
Skin	Man	+		-			C,141,142,E,142
Kidney	Man			-	-		E,25,143,F,49
Plasma Volume	Man	+	-	-	-	-6	C,17,D,f,E,17,38,F,4,G,144
	Dog				-		F,29
	Rat		-				D,56
CO ₂ combining power	Man					-6	G,6,7
	Dog					-6	G,112
Proteins	Man	-		+	+	+	C,3,17,E,3,25,26,F,4,G,32
	Cat				+		F,74,145
	Dog		-		+		D,146,F,145
	Dog				0		F,54
	Rat		-				D,59
Total proteins	Man					0	G,147
	Rat		-				D,56
Total lipids	Man					+	G,6
	Dog					+	G,89,91
	Dog					-	G,148
Choline	Dog					-	G,149
Cholinesterase activity	Man					+	G,150
2,3-Diphosphoglyceric acid	Man					0	G,151
Albumin	Man					-	G,147
β-Globulin	Man					+	G,147
Amino acid nitrogen	Dog					+	G,91
Acetoacetic acid	Man					+	G,91
Plasma flow, renal	Dog					+6	G,91
Serum							
Albumin	Man						D,40
	Dog						E,25
	Rat						D,59,F,145
Total globulin	Dog			0			D,E,59
	Rat				+		D,59,F,145
γ-Globulin	Man			+			D,E,59
α-Globulin	Man			+			E,25
Lipoprotein	Dog			-			E,25
	Dog					0	G,90
Phospholipids	Man					+6	G,90
Glucuronic acid	Man				+		E,25,81,82
Free amino acids							G,152
		+				+	G,153
					+		C,17,F,154
						+	G,5,28,32
		+				+	G,28,155
				+		+11	C,f,E,17,G,f

sm. /3/ Hypothyroidism. /4/ Hypoadrenalism. /5/ Hypoparathyroidism.

130. EFFECT OF ENDOCRINE DISORDERS: MAMMALS (Continued)

Specification	Animal	Graves' Disease ¹	Summonds' Disease ²	Myxedema ³	Addison's Disease ⁴	Diabetes ⁵	Reference
(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)
107 Glucose tolerance (concluded)	Monkey		+				D,102
108 Blood electrolytes Bicarbonate	Rat						F,78
109	Man				0		G,d
110	Cat					-6	F,69
111	Dog				-		F,61,108,G,f
112	Man	0	0	0	+	-6	C,4;D,2;E,4,38;F,73
113	Cat				-		D,109-111;F,53,9
114	Dog		0		+		G,112
115	Rat					-	F,113
116	Man	0			-		C,F,4;E,4,25,G,11
117	Man			0	-	-6	E,38;G,6
118	Man						F,115
119	Cat			+			D,110,111,F,53,10
120	Dog		0		-		G,112
121	Rat						C,116;F,117
122	Man	0			-		G,118
123	Man				+		G,119
124	Man					0	D,120
125	Dog		0			-6	F,121
126	Man						F,117
127	Man				0		D,122;F,123;G,124,125
128	Dog		0		+	-	D,124
129	Dog	+	+				C,E,126,127,G,114,128,129
130	Man			-			F,115
131	Cat		0		+		D,110,G,112
132	Dog		+				D,111
133	Man		0			+	D,2;F,17;G,6
134	Cat				+	+	F,130
135	Dog				+		F,53
136	Man	0	+	0	+		D,E,59;F,8
137	Cat		0	0	+		C,E,4,D,2;F,49
138	Dog		0		+		D,131,F,73,74
139	Man				+	+	D,109,110;F,60,61;G,124,132
140	Dog		-				D,133;F,54
141	Rat		-		0		D,134
142	Man	0	+	0	+	0	C,F,4,D,2,E,4,25,G,135
143	Man						G,114
144	Man					+	G,7
145	Cat					-6,10	F,115,130
146	Dog		0		+		D,110,F,108
147	Monkey		+		+		D,102
148	Rat						F,42
149	Man	+	-	-	+		C,E,3,D,f
150	Man	0	-	0	-	-	C,4;D,2;E,4,25;F,4,136,G,114
151	Man					-6	E,137,G,6
152	Cat						F,115
153	Dog		0		-		D,110,111;F,108,G,112
154	Monkey						D,102
155	Rat		-				F,8
156	Man				-		F,49
157	Dog				+		F,61,138
158	Dog				+		F,138
159	Man				+	-6	G,6

1/1 Hyperthyroidism. /2/ Hypopituitarism. /3/ Hypothyroidism. /4/ Hypoadrenalism. /5/ Hypoinsulinism.
 /6/ In ketosis. /10/ During insulin treatment.


- [36] Grollman, A. 1930. Cardiac output of man in health and disease. C. C. Thomas, Springfield, Ill.
- [37] Bialock, A., and J. W. Beard. 1933. *J. Laborat. Clin. M.* 18:941. [38] Thompson, W. O. 1942. Glandular physiology and therapy. J. B. Lippincott, Philadelphia. [39] Martin, H. E., and M. Wertman. 1947. *Am. Heart J.* 34:646. [40] Earle, D. P., S. J. Farber, R. C. deBodo, M. Kurtz, and M. W. Sikoff. 1953. *Am. J. Physiol.* 173:189. [41] Braun-Menendez, E. 1932. *C. rend. Soc. biol.* 111:477. [42] Freed, S. C., R. H. Rosenman, and M. K. Smith. 1954. *Circul. Res.* N. Y. 2:494. [43] Lange, K. 1944. *Am. J. M. Sc.* 208:5. [44] Gaunt, R., J. H. Birnie, and W. J. Eversole. 1949. *Physiol. Rev.* 29:281. [45] Kuhlman, F. H. 1941. *Klin. Wochr.* 20:889.
- [46] Blumgart, R. L., S. L. Cowgill, and D. R. Gillman. 1930. *J. Clin. Invest.* 9:69. [47] Thompson, W. O. 1946. *Ibid.* 2:477. [48] Friedlander, M., N. Laskey, and S. Silbert. 1935. *Endocrinology* 19:342. [49] Loeb, R. F. 1942. Glandular physiology and therapy. J. B. Lippincott, Philadelphia. [50] Gemzell, C. A., and T. Sjostrand. 1954. *Acta endocr., Kbh.* 16:6. [51] Wyman, L. C., and Caroline von Suden. 1930. *Am. J. Physiol.* 94:579. [52] Houssay, B. A., M. Royer, and O. Orías. 1931. *Rev. Soc. Argent. biol.* 7:314. [53] Rogoff, J. M., and G. N. Stewart. 1926. *Am. J. Physiol.* 78:711. [54] Viale, G., and A. A. Bruno. 1927. *Rev. Soc. Argent. biol.* 3:166. [55] Meineke, H. A., and R. C. Crafts. 1956. *Anat. Rec.* 124:47. [56] Bond, C. F., and S. L. Leonard. 1957. *Am. J. Physiol.* 191:296. [57] Gordon, A. S. 1954. *Recent Progr. Hormone Res.* 10:339. [58] Lucas, G. H. W. 1926. *Am. J. Physiol.* 77:14. [59] Levin, L., and J. H. Leatham. 1942. *Ibid.* 136:306. [60] Yonkman, F. F. 1928. *Ibid.* 86:471. [61] Swingle, W. W., and W. F. Werner. 1927. *Proc. Soc. Exp. Biol., N. Y.* 25:169.
- [62] Corey, E. L., and S. W. Britton. 1932. *Am. J. Physiol.* 102:699. [63] Houssay, B. A., M. Royer, and O. Orías. 1931. *C. rend. Soc. biol.* 108:496. [64] Pico, O. M., and V. Deulofeu. 1926. *Ibid.* 95:1505. [65] Garcia, J. F., D. C. Van Dyke, and N. L. Berlin. 1952. *Proc. Soc. Exp. Biol., N. Y.* 80:472. [66] Crafts, R. C. 1941. *Endocrinology* 29:596. [67] Basó, R., S. T. Olavarrieta, and R. M. Suarez. 1959. *J. Clin. Endocr.* 18:501.
- [68] Berlin, N. L., D. C. Van Dyke, and C. Lotz. 1953. *Proc. Soc. Exp. Biol., N. Y.* 82:287. [69] Swingle, W. W., and A. J. Eisenman. 1927. *Am. J. Physiol.* 79:679. [70] Geyelin, H. R. 1915. *Arch. Int. M.* 16:975.
- [71] Sodeman, W. A. 1956. *Pathologic physiology.* W. B. Saunders, Philadelphia. [72] Gilligan, D. R., M. I. Abrams, and B. Stern. 1934. *Am. J. M. Sc.* 188:790. [73] Hartman, F. A., C. G. MacArthur, F. D. Gunn, W. E. Hartman, and J. J. MacDonald. 1921. *Am. J. Physiol.* 81:244. [74] Swingle, W. W. 1927. *Ibid.* 79:666.
- [75] Smith, P. E., L. Dotti, H. H. Tyndale, and E. T. Engle. 1936. *Proc. Soc. Exp. Biol., N. Y.* 34:247. [76] Zucker, T. F., and B. N. Berg. 1937. *Am. J. Physiol.* 119:539. [77] Wyman, L. C., and B. S. Walker. 1929. *Ibid.* 89:215. [78] Banerji, H., and C. Reid. 1934. *Ibid.* 81:93. [79] Randle, P. J. 1953. The influence of growth hormone on blood insulin and glucagon activity. In R. W. Smith, Jr., O. H. Gaebler, and C. N. H. Long, ed. *The hypophyseal growth hormone, nature and actions.* McGraw-Hill, New York. [80] Gellhorn, E., J. Feldman, and A. Allen. 1941. *Endocrinology* 29:849. [81] Peters, J. P., and E. B. Man. 1943. *J. Clin. Invest.* 22:715.
- [82] Peters, J. P., and E. B. Man. 1950. *Ibid.* 29:1. [83] Van Aradell, P. P., Jr., and R. H. Williams. 1956. *Am. J. Med.* 20:4. [84] Peterson, V. P. 1943. *Acta med. scand.* 146:375. [85] Felch, W. C., and L. B. Dotti. 1949. *Proc. Soc. Exp. Biol., N. Y.* 72:376. [86] Katz, L. B., G. J. Rhodes, R. S. George, and C. Moosa. 1953. *Am. J. M. Sc.* 225:120. [87] Munoz, J. M. 1932. *Rev. Soc. Argent. biol.* 8:587. [88] Shlossow, S. 1926. *Sc. Rep. Inst. Infec. Dis., Tokyo* 5:541. [89] Morimoto, M. 1928. *Phügera Arch.* 219:733. [90] Fasoli, A., E. B. Magid, M. D. Glasman, and P. P. Foa. 1954. *Proc. Soc. Exp. Biol., N. Y.* 85:609. [91] Bodansky, M., and O. Bodansky. 1952. *Biochemistry of disease.* Ed. Z. Macmillan, New York. p. 474. [92] Rogoff, J. M., and G. N. Stewart. 1928. *Am. J. Physiol.* 86:25. [93] Fleischmann, W. 1947. *Q. Rev. Biol.* 22:119. [94] Enterman, C., I. L. Chazkoff, and F. L. Reichert. 1942. *Endocrinology* 30:194. [95] Becher, E. 1942. *Klin. Wochr.* 21:2.
- [96] Abrams, M. I., and D. R. Gilligan. 1934. *Am. J. M. Sc.* 188:796. [97] Britton, S. W., and W. K. Myers. 1928. *Am. J. Physiol.* 84:132. [98] Burn, J. H., and H. P. Marks. 1925. *J. Physiol., Lond.* 60:133. [99] Assadi, Z., and A. Ernst. 1927. *Arch. ges. Physiol.* 215:431. [100] Houssay, B. A., and M. A. Magenia. 1924. *Rev. As. arg. argent.* 37:389. [101] Russell, J. A. 1938. *Physiol. Rev.* 18:1. [102] Knobil, E. 1955. *Am. J. Physiol.*

	Specification	Animal	Graves' Disease ¹	Simmonds' Disease ²	Myxedema ³	Addison's Disease ⁴	Diabetes ⁵	Reference
	(A)	(B)	(C)	(D)	(E)	(F)	(G)	(H)
214	Respiratory rate	Man	+	-	-	-	+	C,156;D,4;E,31;G,5,6
215		Man					+6	G,7
216	Respiratory quotient	Cat					-	G,157
217		Dog					-	G,91,124
218		Rat					-	F,16
219	O ₂ consumption (BMR)	Man		-	-	-	+6	D,2,71;E,4;F,4,27;G,158
220		Cat				0	+	F,159;G,160
221		Dog		-				D,9,161,162
222		Guinea pig	+					C,163,164
223		Rat		-	-	-		D,12,165,166;E,165,166,165,167
224	Vital capacity	Man	-					C,168
225	Ventilatory response to exercise	Man	+					C,169
226	Ventilatory equivalent							
227	Resting	Rat			-			E,170
228	Active	Rat			-			E,170
229	Coefficient of pulmonary elastic resistance	Man	0					C,169
229	Mean inspiratory non-elastic resistance	Man	0					C,169

/1/ Hyperthyroidism. /2/ Hypopituitarism. /3/ Hypothyroidism. /4/ Hypoadrenalism. /5/ Hypoinsulinism.
/6/ In ketosis.

Contributors: (a) Fregly, Melvin J., (b) Grollman, Arthur, (c) Knobil, Ernest, (d) Ganong, William F.,
(e) Leatham, James H., (f) Engel, Frank L.

References: [1] Grollman, A. 1947. *Essentials of endocrinology*. J. B. Lippincott, Philadelphia. [2] Soffer, L. J. 1956. *Diseases of the endocrine glands*. Lea and Febiger, Philadelphia. p. 142. [3] Houssay, B. A. 1955. *Human physiology*. Ed. 2. McGraw-Hill, New York. [4] Duncan, G. G. 1953. *Diseases of metabolism*. Ed. 3. W. B. Saunders, Philadelphia. [5] Forsham, P. H., and G. W. Thorn. 1955. In R. H. Williams, ed. *Textbook of endocrinology*. Ed. 2. W. B. Saunders, Philadelphia. [6] Duncan, G. G. 1947. *Diseases of metabolism*. Ed. 2. W. B. Saunders, Philadelphia. [7] Schneider, R. W. 1954. *Glandular physiology and therapy*. Ed. 5. J. B. Lippincott, Philadelphia. p. 386. [8] Grollman, A. 1936. *The adrenals*. Williams and Wilkins, Baltimore. [9] Koester, S., and A. Geesink. 1929. *Pflügers Arch.* 222:293. [10] Solari, L. A. 1931. *Rev. Soc. argent. biol.* 7:216. [11] Ikeda, M. 1933. *Jap. J. Obst.* 15:213. [12] Smith, P. E. 1930. *Am. J. Anat.* 45:205. [13] Fregly, M. J. Unpublished. [14] Iampietro, P. F., M. J. Fregly, and E. R. Buskirk. 1956. *Canad. J. Biochem. Physiol.* 34:721. [15] Hermanson, V., and F. A. Hartman. 1945. *Am. J. Physiol.* 144:108. [16] Carr, C. J., and F. F. Beck. 1937. *Ibid.* 119:589. [17] Williams, R. H., ed. 1955. *Textbook of endocrinology*. Ed. 2. W. B. Saunders, Philadelphia. [18] Souto-Mator, M. G., and V. G. Foglia. 1944. *Rev. Soc. argent. biol.* 20:55. [19] Hoskins, R. G. 1927. *Endocrinology* 11:97. [20] Richter, C. P. 1933. *Ibid.* 17:73. [21] Richter, C. P., and G. B. Wislocki. 1930. *Am. J. Physiol.* 95:481. [22] Hall, V. E., and M. Lindsay. 1938. *Endocrinology* 22:66. [23] Richter, C. P. 1936. *Ibid.* 20:657. [24] Gans, H. M., and H. H. Miley. 1927. *Am. J. Physiol.* 82:1. [25] Asper, S. P., and J. G. Wiswell. 1956. *Am. J. Med.* 20:732. [26] Lerman, J., R. J. Clark, and J. H. Means. 1934. *Ann. Int. M.* 8:82. [27] Soffer, L. J. 1956. *Diseases of the endocrine glands*. Lea and Febiger, Philadelphia. [28] Joslin, E. P., H. F. Root, P. White, A. Marble, and C. C. Bailey. 1946. *Treatment of diabetes mellitus*. Ed. 8. Lea and Febiger, Philadelphia. [29] Estrada, O. P. 1926. *C. rend. Soc. biol.* 95:1506. [30] Leblond, C. P., and H. E. Hoff. 1944. *Am. J. Physiol.* 141:32. [31] Turner, C. D. 1955. *General endocrinology*. W. B. Saunders, Philadelphia. [32] Schectter, A. E., G. H. Wiesel, and C. Cohn. 1941. *Am. J. M. Sc.* 202:364. [33] Myers, J. D., E. S. Brannon, and B. C. Holland. 1950. *J. Clin. Invest.* 29:1069. [34] Scheinberg, P., E. A. Stead, Jr., E. S. Brannon, and J. V. Warren. 1950. *Ibid.* 29:1139. [35] Gladstone, S. A. 1936. *Proc. Soc. Exp. Biol., N. Y.* 34:587.

131. PHYSIOLOGIC ALTERATIONS AND COMPENSATORY MECHANISMS IN CHRONIC ANEMIA (O₂ transport constant)

- 183:635. [103] Smith, P. E., H. H. Tyndale, L. Dotti, and E. T. Engle. 1936. *Proc. Soc. Exp. Biol.*, N. Y. 34:250.
- [104] Popper, H. L., and S. Hirschhorn. 1932. *Arch. klin. Chir.* 169:355. [105] Hart, J. F., and M. Magiday. 1941. *Arch. Int. M.* 68:893. [106] Blasotti, A. 1934. *Rev. Soc. argent. biol.* 10:124. [107] Galehr, O., P. Ladurner, and L. Unterrichter. 1927. *Pflügers Arch.* 218:477. [108] Loeb, R. F., D. W. Atchley, E. M. Benedict, and J. Leland. 1933. *J. Exp. M.* 57:775. [109] Gerschman, R. 1931. *C. rend. Soc. biol.* 108:494. [110] Mazzocco, P. 1927. *Rev. Soc. argent. biol.* 3:213. [111] Marenzi, A. D., and R. Gerschman. 1934. *Ibid.* 10:131. [112] Gerschman, R., and A. D. Marenzi. 1935. *Ibid.* 11:500. [113] Blumenfeld, C. M., and F. W. Clausen. 1940. *Am. J. Physiol.* 128:577. [114] Corte-Real, E. 1943. *Med. contemp.*, Lisbon. 60:273. [115] Baumann, E. J., and S. Kurland. 1927. *J. Biol. Chem.* 71:281. [116] Kasanen, A., and I. Vutanen. 1956. *Acta med. scand.* 153:467. [117] Sachs, A., V. E. Levine, and W. O. Griffith. 1937. *Proc. Soc. Exp. Biol.*, N. Y. 37:46. [118] Stock, R., and J. D. Currence. 1950. *J. Clin. Endocr.* 10:313. [119] Illing, E. K. B., C. H. Gray, and R. D. Lawrence. 1951. *Biochem. J.*, Lond. 48:637. [120] Maveroff de Lissner, H. 1933. *Rev. Soc. argent. biol.* 9:227. [121] Larsen, V., and N. A. Lassen. 1956. *Acta med. scand.* 154:65. [122] Marenzi, A. D. 1934. *Rev. Soc. argent. biol.* 10:121. [123] Rabboni, F. 1934. *Arch. Ital. chir.* 37:687. [124] Binet, L., and J. Klukowski. 1933. *J. Physiol. path. gén.* 31:372. [125] Callazo, J. A., and E. Morelli. 1926. *Ibid.* 24:508. [126] Dine, R. F., and P. H. Lavietes. 1942. *J. Clin. Invest.* 21:781. [127] Soffer, L. J., C. Cohn, E. B. Grossman, M. Jacobs, and H. Sobotka. 1941. *Ibid.* 20:429. [128] Hauray, V. G., and A. Cantarow. 1942. *J. Laborat. Clin. M.* 27:616. [129] Stutzman, F. L., and D. S. Amatuzio. 1953. *Ibid.* 41:215. [130] Zwemer, R. L., and R. C. Sullivan. 1934. *Endocrinology* 18:97. [131] Anderson, A. B., and E. C. Oestler. 1938. *J. Physiol.*, Lond. 92:124. [132] Gerschman, R., and A. D. Marenzi. 1935. *Rev. Soc. argent. biol.* 11:391. [133] Jones, L. M., and G. Y. Shunowara. 1942. *J. Biol. Chem.* 142:935. [134] Li, C. H., I. Gerschwind, and H. M. Evans. 1949. *Endocrinology* 44:67. [135] Shinn, M. B. 1943. *Univ. Pittsburgh Bull.* 39:405. [136] Loeb, R. F. 1932. *Science* 76:420. [137] Curtis, R. H. 1956. *Ann. Int. M.* 44:376. [138] DeMeio, R. H. 1933. *Rev. Soc. argent. biol.* 9:324. [139] Peralke, E. C. 1948. *Endocrinology* 42:356. [140] Bergen, J. R., C. A. Hunt, and H. Hoaglund. 1953. *Am. J. Physiol.* 175:327. [141] Stewart, H. J., and W. F. Evans. 1940. *Am. Heart J.* 20:715. [142] Stewart, H. J., and W. F. Evans. 1942. *Arch. Int. M.* 69:808. [143] Corcoran, A. C., and I. H. Page. 1948. *J. Clin. Endocr.* 7:801. [144] Jacobson, S. D., and R. H. Lyons. 1942. *J. Laborat. Clin. M.* 27:1169. [145] Hartman, F. A., L. A. Lewis, J. S. Thatcher, and H. R. Street. 1942. *Endocrinology* 31:287. [146] Baker, R., and G. H. Miller, Jr. 1951. *Ibid.* 49:484. [147] Lewis, L. A., R. W. Schneider, and E. P. McCullagh. 1944. *J. Clin. Endocr.* 4:535. [148] Rubin, S. H., and E. P. Ralli. 1940. *Am. J. Physiol.* 129:578. [149] Neurath, H., and F. W. Putnam. 1945. *J. Biol. Chem.* 160:397. [150] Strehler, E., and H. Meyer. 1952. *Helvet. med. Acta* 19:555. [151] Lennerstrand, A. 1944. *Acta med. scand.* 116:225. [152] Saltzman, A., W. T. Caraway, and I. A. Beck. 1954. *Metabolism* 3:11. [153] Suvarnakich, K., G. V. Mann, and F. J. Stare. 1952. *J. Nutr.* 47:105. [154] Nicholson, W. M., and L. J. Soffer. 1935. *Bull. Johns Hopkins Hosp.* 55:236. [155] Herzstein, J., and L. A. Weinroth. 1945. *Arch. Int. M.* 76:34. [156] Turner, C. D. 1955. *General endocrinology*. W. B. Saunders, Philadelphia. p. 32. [157] Ring, G. C., and C. W. Hampel. 1932. *Am. J. Physiol.* 102:460. [158] Fisher, P., and J. I. Kleinerman. 1952. *J. Clin. Invest.* 31:126. [159] Kojima, T. 1929. *Tohoku J. Exp. M.* 13:379. [160] Ring, G. C., and C. W. Hampel. 1933. *Am. J. Physiol.* 104:298. [161] Artundo, A. 1930. *Rev. Soc. argent. biol.* 6:497. [162] Houssay, B. A., and A. Artundo. 1933. *Ibid.* 9:66. [163] Friedgood, H. B. 1934. *Bull. Johns Hopkins Hosp.* 54:48. [164] Dobyns, B. M. 1946. *Surg. Gyn. Obst.* 82:290. [165] Evans, H. S., A. N. Contopoulos, and M. E. Simpson. 1957. *Endocrinology* 60:403. [166] Foster, G. L., and P. E. Smith. 1926. *J. Am. M. Ass.* 87:2151. [167] Horvath, S. M. 1938. *Endocrinology* 23:223. [168] Rabinowitch, I. M. 1926. *Boston M. & S. J.* 194:199. [169] McIlroy, M. B., F. L. Eldridge, and R. W. Stone. 1956. *Clin. Sc.*, Lond. 15:353. [170] Kao, F. F., and B. B. Schlögl. 1958. *Am. J. Physiol.* 195:229.

133. CARDIOVASCULAR AND PULMONARY SYMPTOMS OF ANEMIA: MAN AND OTHER ANIMALS

Variable	Chronic Anemia ^{1,2}				Acute Anemia	
	Uncomplicated	Reference	With Congestive Heart Failure	Reference	Large Hemorrhage ³	Reference
(A)	(B)	(C)	(D)	(E)	(F)	(G)
1 Vessels	Hyperkinemia, pulsating neck vessels, collapsing pulse, capillary pulsation, Duroziez' and Traube's phenomena, venous hum (bruit de diable). Dilated and tortuous retinal veins. ⁴ Capillary dilatation and stasis leading to ischemic necrosis of tissues. ⁴ Thickening of the small and medium arteries by intimal proliferation, thrombosis. ⁴	1 2 3 4	Engorged neck veins.	5	Pallor, cyanosis, coldness of extremities, absence of pulsation over large arteries. Ischemic optic neuritis with pale edematous discs, attenuated retinal arterioles and mildly distended and tortuous veins, hemorrhages and exudates near the discs, edema of retina in macular areas, and subsequent optic atrophy. ⁵ Visceral ischemia; visceral congestion, edema, and petechial hemorrhages if death occurs despite blood transfusions.	6 7 8
2 Heart	Dilatation, hypertrophy and increased weight, fatty degeneration with yellow subendocardial streaks at venous end of capillaries perpendicular to the fibers of the trabeculae and papillary muscles (tiger lily heart). Interstitial edema, polymorphonuclear infiltration, myocardial degeneration, vacuolated sarcoplasm, Zenker's degeneration. ⁴ Increased interarterial coronary anastomosis Pericardial lemon-yellow or hemorrhagic effusion. Flabby musculature Cor pulmonale ⁴	9-11 4 12 13,14 11 15	Flabby myocardium, obliterative endarteritis of coronary and pericardial vessels, vacuolization of sarcoplasm, interstitial edema, increased size and weight, dilatation, hypertrophy ⁴	4	Firm musculature. Diffuse focal or confluent stress of myocardial necrosis, chiefly in the subendocardium of the left ventricle and especially the papillary muscles. Scattered areas of fatty infiltration and acute necrosis in both the ventricles and inter-ventricular septum, most frequently beneath the endocardial surface and in the papillary muscles. ⁶	27 28 29
3 Impulse	Forceful apical thrust. Diffuse and wavy. Pulmonary diastolic tap.	16 2,17,18 2,17,18	Forceful apical thrust, may be diffuse and wavy.	17		
4 Borders	Enlarged. ⁴	19-21	Enlarged.	17		
5 Sounds	M ₁ accentuated, P ₂ accentuated or split Gallop rhythm.	22 2,19	M ₁ accentuated, P ₂ accentuated or split. Gallop rhythm.	17 26	Disappearance of sounds. Tic-tac sounds if there is tachycardia.	6 1
6 Murmurs	Systolic pulmonary, mitral, aortic. Diastolic or presystolic mitral, pulmonary, aortic.	5,9,10, 19,22 5,9,23- 25	Systolic pulmonary, mitral. Diastolic, mitral	17,26 17,26		
7 Thrills	Systolic or diastolic	5	Systolic	17		

- /1/ Since effects have no linear relationship to the severity of anemia, no quantitative values are given [5]
 /2/ Moderate abnormalities appear when hemoglobin level is between 8 and 9 grams per cent, serious abnormalities do not occur until hemoglobin is less than 7 grams. [31] /3/ A liter or more, data include observations during hemorrhagic shock. /4/ Observations made in sickle cell anemia, many of the effects are peculiar to sickle cell disease only, and the possibility exists that associated histologic and pathologic findings are unrelated to anemia per se [b]. /5/ Usually observed after repeated hemorrhage [2]. /6/ Experimental observations in dogs.
 /7/ Enlargement usually reversible, but may persist in cases of long duration [20].

Chronic Anemia ^{1, 2}				Acute Anemia	
Uncomplicated	Reference	With Congestive Heart Failure	Reference	Large Hemorrhage ^{3, 4}	Reference
(A)	(B)	(C)	(D)	(E)	(F)
Palpitation, dyspnea, edema. Intermittent claudication Gross hematuria. Angina. Circulatory death.	1-3 2, 4 5 2, 4, 5-9 10, 11	Palpitation, edema, dyspnea (at rest), orthopnea. Cardiac asthma.	3 10, 11	Angina. Oliguria, anuria, acute tubular failure. Increased respiratory rate and amplitude, sweating, fainting, twitching of extremities, unconscious- ness. Dimness or loss of vision. Yawning, desire to defecate or urinate, rotation of eyes. Waves of tingling over body, restlessness, carpal spasms, hyperventilation. Manifestations of circulatory and respiratory failure.	12-14 15 16, 17 18, 19 17 16 20

/1/ Since effects have no linear relationship to the severity of anemia, no quantitative values are given [3].
 /2/ Moderate abnormalities appear when hemoglobin level is between 8 and 9 grams per cent; serious abnormalities do not occur until hemoglobin is less than 7 grams. [21] /3/ A liter or more; data include observations during hemorrhagic shock. /4/ May precipitate congestive failure [12, 22], or myocardial infarction [23].

Contributors: (a) Sanghvi, L. M., (b) Wintrobe, M. M.

References: [1] Blumgart, H. L., and M. D. Altschule. 1948. *Blood*, N. Y. 3:329. [2] Hunter, A. 1946. *Q. J. Med.*, Oxf. 15:107. [3] Sanghvi, L. M., R. Sharma, and S. N. Misra. 1957. *Circulation*, N. Y. 15:373.
 [4] Pickering, G. W., and E. J. Wayne. 1934. *Clin. Sc.*, Lond. 1:305. [5] Crone, R. L., S. C. Jefferson, V. J. Pileggi, and E. C. Lowry. 1957. *Arch. Int. M.* 100:597. [6] Elliot, A. H. 1934. *Am. J. M. Sc.* 187:185.
 [7] Herrick, J. B. 1927. *Am. Heart J.* 2:35. [8] Williams, F. A., and H. Z. Giffin. 1927. *Am. J. M. Sc.* 174:30.
 [9] Wilson, J. L., and S. A. Levine. 1952. *Ann. Int. M.* 36:897. [10] Friedberg, C. K. 1956. *Diseases of the heart*. W. B. Saunders, Philadelphia. pp. 1046-1053. [11] Sanghvi, L. M. Unpublished. [12] Dack, S. E., Corday, and A. M. Master. 1951. *Am. Heart J.* 42:161. [13] Master, A. M., S. Dack, H. Horn, B. Freedman, and L. E. Field. 1950. *Circulation*, N. Y. 1:1302. [14] Zimmerman, S. L., and R. Barnett. 1944. *Ann. Int. M.* 21:1045. [15] Van Slyke, D. D. 1948. *Ibid.* 28:701. [16] Shenkin, H. A., R. H. Cheney, S. R. Govans, J. D. Hardy, A. G. Fletcher, and I. Starr. 1944. *Am. J. M. Sc.* 208:421. [17] Wallace, J., and E. P. Sharpey-Schafer. 1941. *Lancet*, Lond. 2:393. [18] Ebert, R. V., E. A. Stead, Jr., and J. G. Gibson. 1941. *Arch. Int. M.* 68:578.
 [19] Hollenhorst, R. W., and H. P. Wagener. 1950. *Am. J. M. Sc.* 219:209. [20] McDowall, R. J. S. 1956. *The control of the circulation of the blood*. Dawson and Sons, London. pp. 449-455. [21] Brannon, E. S., A. J. Merrill, J. V. Warren, and E. A. Stead, Jr. 1945. *J. Clin. Invest.* 24:332. [22] Kinney, T. D., and G. K. Mallory. 1945. *N. England J. M.* 232:215. [23] Friedberg, C. K., and H. Horn. 1939. *J. Am. M. Ass.* 112:1675.

**134. ELECTROCARDIOGRAPHIC, BALLISTOCARDIOGRAPHIC, AND
ROENTGENOGRAPHIC FINDINGS IN ANEMIA: MAN AND OTHER ANIMALS**

Method	Chronic Anemia ^{1,2}				Acute Anemia	
	Uncomplicated	Reference	With Congestive Heart Failure	Reference	Large Hemorrhage ³	Reference
(A)	(B)	(C)	(D)	(E)	(F)	(G)
1 Electrocardiographic ⁴	Low, flat, or inverted T. Notched T. ⁵ Depressed RT junction and ST segment. Low, flat, or inverted P. Tall, peaked, notched P. ⁵ Decreased QRS amplitude. Large, diphasic QRS. ⁵ Increased QT interval. Increased PR interval. Incomplete right bundle branch block. ⁵ Left ventricular hypertrophy pattern. Left ventricular strain. Right axis deviation. ⁵ Nodal tachycardia, nodal escape, and arteriovenous dissociation. ⁵ Premature beats. Atrial fibrillation. ⁵ Parasytostole. Atrial bigeminy. Isolated T inversion in precordial leads.	1-4 3 1,3,4 1,5,6 3 1,4 3 1-3 2,3,7-10 11 1,7,11 3,12 2 3 1-3,8 5 12 12 12	Low, flat, or inverted T. Depressed ST segment. Low, flat, or inverted P. Decreased QRS amplitude. QT interval more often increased. Increased PR interval. Left ventricular hypertrophy pattern. Premature beats. Atrial tachycardia.	12 12 12 12 1 12 12 12 12	Low, flat, or inverted T. Depressed ST segment. Low, wide, or slurred QRS.	13-16 13,14,16
2 Ballistocardiographic ⁴	Minimal respiratory variation of I and J, increased amplitude of systolic complexes I, J, K, abnormal pattern. ⁶ Large systolic complexes with normal or abnormal pattern, tall or bifid R, plateau-like I valley, slurred or notched I-J and J-K segments, broad or bifid J, fused H-J, short, absent, or notched K, tall N, L, totally bizarre complexes of low amplitude, grade 1-4 abnormal patterns.	17 18	Pattern may be normal. Increased systolic complexes, tall L and N. Large systolic complexes with normal or abnormal pattern, tall H, L, N, slurred or notched I-J and J-K segments, notched or bifid J, short or absent K, totally bizarre complexes of low amplitude, grade 1-4 abnormal patterns.	17 18	Minimal respiratory variation, normal or reduced I and J, J slopes off gradually into the small L wave instead of returning quickly to the base line, absent K, large H.	17
3 Roentgenographic Vessels	Increased aortic and pulmonary artery pulsations. Dilated pulmonary artery. Prominent vascular shadows.	19 6,19 20	Prominent vascular shadows. Dilated pulmonary artery.	19 19		

[1] Since effects have no linear relationship to severity of anemia, no quantitative values are given [1].
 [2] Moderate abnormalities appear when hemoglobin level is between 8 and 9 grams per cent, serious abnormalities do not occur until hemoglobin is less than 7 grams [28]. [3] A liter or more, data include observations during hemorrhagic shock. [4] In some patients with chronic anemia (either uncomplicated or with congestive failure), and in the absence of any other cardiovascular disease, an increase in pattern abnormality may occur after improvement of anemia, or abnormality may persist after cure of anemia. [5] Observations made in sickle cell anemia; many of the effects are peculiar to sickle cell disease only, and the possibility exists that associated histologic and pathologic findings are unrelated to anemia per se [6]. [6] Pattern may be normal, in anemia with cardiac damage, abnormal pattern often apparent only during normal expiration. [7] [7] Marked abnormal pattern may appear when anemia is corrected in some subjects known to have heart disease, transient abnormal curves may appear when hemoglobin is rapidly raised in severe anemia and the heart is severely damaged by anoxia [17].

Variable	Chronic Anemia ^{1,2}				Acute Anemia	
	Uncomplicated	Refer- ence	With Congestive Heart Failure	Refer- ence	Large Hemorrhage ³	Refer- ence
(A)	(B)	(C)	(D)	(E)	(F)	(G)
8 Lungs	Obliterative endarteritis and thrombosis of small and medium vessels, polymorphonuclear in- filtration, infarction, pleural effusion. ⁴ Capillary dilatation and stasis, bronchopneumonia, atelectasia. ⁴ Congestion. Edema. ⁴ Pneumonia. ⁴	4 3 4,5 2,4,25, 27 30	Edema, obliterative endarteritis and thrombosis of pul- monary arterioles, bilateral pleural effusion. ⁴ Signs of congestion, pleural effusion. Pulmonary edema.	4 5 4,17		

/1/ Since effects have no linear relationship to the severity of anemia, no quantitative values are given [5].
 /2/ Moderate abnormalities appear when hemoglobin level is between 8 and 9 grams per cent; serious abnormalities do not occur until hemoglobin is less than 7 grams. [31] /3/ A liter or more; data include observations during hemorrhagic shock. /4/ Observations made in sickle cell anemia; many of the effects are peculiar to sickle cell disease only, and the possibility exists that associated histologic and pathologic findings are unrelated to anemia per se [b].

Contributors: (a) Sanghvi, L. M., (b) Wintrobe, M. M.

References: [1] Ruskin, A. 1953. Physiological cardiology. C. C. Thomas, Springfield, Ill. [2] Klinefelter, H. F. 1942. Am. J. M. Sc. 203:34. [3] Kimmelstiel, P. 1948. Ibid. 216:11. [4] Winsor, T., and G. E. Burch. 1945. Am. Heart J. 29:685. [5] Sanghvi, L. M., R. Sharma, and S. N. Misra. 1957. Circulation, N. Y. 15:373. [6] Wallace, J., and E. P. Sharpey-Schafer. 1941. Lancet, Lond. 2:393. [7] Hollenhorst, R. W., and H. P. Wagener. 1950. Am. J. M. Sc. 219:209. [8] Moon, V. H. 1942. Shock, Lea and Febiger, Philadelphia. p. 107. [9] Cabot, R. C., and O. Richardson. 1919. J. Am. M. Ass. 72:991. [10] Ellis, L. B., and J. M. Faulkner. 1939. N. England J. M. 220:943. [11] Gould, S. E. 1953. Pathology of the heart. C. C. Thomas, Springfield, Ill. p. 532. [12] Zoll, P. M., S. Wessler, and M. J. Schlesinger. 1951. Circulation, N. Y. 4:797. [13] Soloff, L. A., and C. T. Bello. 1950. Ibid. 2:298. [14] Whipple, G. H., and W. L. Bradford. 1936. J. Pediat., S. Louis 9:279. [15] Yater, W. M., and G. H. Hansmann. 1936. Am. J. M. Sc. 191:474. [16] Eddleman, E. E., Jr., L. Hefner, T. J. Reeves, and T. R. Harrison. 1957. Arch. Int. M. 99:400. [17] Sanghvi, L. M. Unpublished. [18] Reich, N. E. 1954. Uncommon heart diseases. C. C. Thomas, Springfield, Ill. p. 320. [19] Hunter, A. 1946. Q. J. Med., Oxf. 15:107. [20] Porter, W. B. 1937. Am. Heart J. 13:550. [21] Wintrobe, M. M. 1946. Blood, N. Y. 1:121. [22] Mukerji, S. K., M. P. Kahirasagar, and C. R. Bhandari. 1955. J. Ind. M. Ass. 24:369. [23] Goldstein, B., and E. P. Boas. 1927. Arch. Int. M. 39:226. [24] Gunewardene, H. O. 1933. J. Trop. M. Hyg., Lond. 36:49. [25] Tung, C. L., W. N. Bien, Y. C. Chu, and S. H. Wang. 1937. Chin. M. J. 52:479. [26] Massias, C. 1947. Progr. méd., Par. 75:454. [27] Friedberg, C. K. 1956. Diseases of the heart. W. B. Saunders, Philadelphia. pp. 1046-1053. [28] Friedberg, C. K., and H. Horn. 1939. J. Am. M. Ass. 112:1675. [29] Melcher, G. W., Jr., and W. W. Walcott. 1951. Am. J. Physiol. 164:832. [30] Henderson, A. B. 1950. Am. J. Med. 9:757. [31] Brannon, E. S., A. J. Merrill, J. V. Warren, and E. A. Stead, Jr. 1945. J. Clin. Invest. 24:332.

115. HEMODYNAMIC AND PULMONARY EFFECTS OF ANEMIA: MAN AND OTHER ANIMALS

Variable	Chronic Anemia ^{1,2}			Acute Anemia		Reference
	Uncomplicated	With Congestive Heart Failure		Small Hemorrhage ³ (D)	Large Hemorrhage ⁴ (E)	
(A)	(B)	(C)		(D)	(E)	(F)
1 Cardiac function						
2 Output	Increased	Increased or normal		Slightly decreased ⁵	Decreased	B,1-9,C,6,8,10, D,11-13,E, 14-19
3	May be normal	Increased or decreased		May be normal		B,6,20,C,6,D,13, 17,19
4 Stroke volume	Increased	Increased		Normal ⁵	Decreased	B,4,9,D,19,E,18
5 Rate	Increased	Increased		Slightly increased ⁶	Increased ⁷	B,1,3,5,C,6,D, 12,13,17,E,11, 19,21
6	Normal or slow	Increased or decreased				B,6,20,22,C,6
7 Work	Normal				Decreased	B,4,E,18
8	May increase ⁸					B,7
9 Utilization co-efficient					Decreased	E,18
10 Pyruvate, lactate, glucose						
11 Utilization, absolute					Decreased	E,18
12 Pyruvate					Increased	E,18
13 Lactate					Decreased or normal	E,18
14 Glucose						
15 Blood					Decreased	B,5,8,23,C,10, E,17,21,24
16 Total volume	Decreased	May increase			Normal	B,5,25,E,26
17 Viscosity	Decreased				Decreased, may increase terminally	E,27
18					Immediately decreased ⁹	B,28,C,10,E,24, 29,30
19 Proteins	May decrease	Often decreased			Decreased	B,23,C,10,E,17, 24,29
20 Plasma volume	Increased	Increased			Increased, may increase terminally	B,C,28,E,27
21 Capillary permeability	Increased in severe cases	Increased				
22 Blood pressure					Decreased	B,5,9,31,C,32, D,17,19,E,17, 19,33
23 Systolic	Decreased ¹⁰	Decreased		Decreased ⁵	Decreased	B,34,E,35
24	May be normal				Normal or increased	B,36
25 Diastolic	Decreased	Decreased		Decreased ¹¹	Decreased	B,5,9,31,C,32, D,17,E,17,19
26 Mean	Decreased	Decreased			Decreased	B,9,C,32,E,17, 18
27 Pulse	Increased	Increased		Decreased	Decreased	B,1,6,9,C,32,D, E,36
28 Right atrium	Normal	Increased		Decreased	Decreased	B,1,C,8,D,12, 13,17,E,11,17
29 Pulmonary artery	Normal, may increase	May increase			Decreased	B,7,C,37,E,38
30 Pulmonary arterio-capillary gradient	Normal, may increase	May increase				B,7,C,37

11/ Effects have no linear relationship to the severity of anemia, and therefore no numerical values can be given [6]
 12/ Moderate abnormalities appear when hemoglobin level is between 8 and 9 grams per cent, serious abnormalities do not occur until hemoglobin is less than 7 grams [3]. 13/ Effects of small hemorrhage are increased respiratory rate and occasionally orthostatic syncope [61]. 14/ A liter or more, data include observations during hemorrhagic rate and occasionally orthostatic syncope [61]. 15/ May be normal in supine position, but increased when standing [19]. 16/ Bradycardia, particularly during fainting [11,19,33]. 17/ Observations made in sickle cell anemia, many of the effects are peculiar to sickle cell disease only, and the possibility exists that associated histological and pathological findings are unrelated to anemia per se [b]. 18/ May be little change [33], increased later [27]. 19/ Hypertension may occur after recovery [4,9]. 20/ May be normal in supine position, but decreased when standing [19].

134. ELECTROCARDIOGRAPHIC, BALLISTOCARDIOGRAPHIC, AND
ROENTGENOGRAPHIC FINDINGS IN ANEMIA: MAN AND OTHER ANIMALS (Concluded)

Method	Chronic Anemia ^{1,2}				Acute Anemia	
	Uncomplicated	Reference	With Congestive Heart Failure	Reference	Large Hemorrhage ³	Reference
(A)	(B)	(C)	(D)	(E)	(F)	(G)
4 Roentgenographic (concluded) Heart	Hyperactive cardiac pulsations. Enlarged silhouette. Generalized, globular, left ventricular or right ventricular enlargement. Straightening of left border. Prominent conus. Mitralized configuration. Enlargement of left atrium accompanying general cardiac enlargement. Pericardial effusion. ⁸	19 21 1 22 1 1,5,23,24 6 25	Enlargement.	1	Decreased size and oscillations.	27
5 Lungs	Increased bronchovascular markings. Irregular patchy infiltrations, symmetrical nodular densities (hemosiderosis), infarction. ⁵ Massive consolidation. ⁵	19,20 20 20	Congestion, unilateral or bilateral pleural effusion. Encysted pleural effusion. Pericardial effusion	1 26 19		

/1/ Since effects have no linear relationship to severity of anemia, no quantitative values are given [1].
/2/ Moderate abnormalities appear when hemoglobin level is between 8 and 9 grams per cent; serious abnormalities do not occur until hemoglobin is less than 7 grams [28]. /3/ A liter or more, data include observations during hemorrhagic shock. /5/ Observations made in sickle cell anemia; many of the effects are peculiar to sickle cell disease only, and the possibility exists that associated histologic and pathologic findings are unrelated to anemia per se [b]. /8/ Only with pericardial inflammation, or in heart failure [b].

Contributors: (a) Sanghvi, L. M., (b) Wintrobe, M. M.

References: [1] Sanghvi, L. M., R. Sharma, and S. N. Misra. 1957. *Circulation*, N. Y. 15:373. [2] Winsor, T., and G. E. Burch. 1945. *Am. Heart J.* 29:685. [3] Lindo, C. L., and L. R. Doctor. 1955. *Ibid.* 50:218.
[4] Szekely, P. 1940. *Brit. Heart J.* 2:1. [5] Henderson, A. B. 1950. *Am. J. Med.* 9:757. [6] Pietrafesa, E. R. 1942. *Am. Heart J.* 23:736. [7] Pickering, G. W., and E. J. Wayne. 1934. *Clin. Sc., Lond.* 1:305. [8] Klinefelter, H. F. 1942. *Am. J. M. Sc.* 203:34. [9] Grover, V. 1947. *Ann. Int. M.* 26:843. [10] King, J. T., and C. A. Janeway. 1937. *Internat. Clin.* 3:41. [11] Leight, L., T. H. Snider, G. O. Clifford, and H. K. Hellems. 1954. *Circulation*, N. Y. 10:653. [12] Sanghvi, L. M., S. N. Misra, K. Banerjee, and K. D. Gupta. 1958. *Am. Heart J.* 56:79. [13] Dack, S., E. Corday, and A. M. Master. 1951. *Ibid.* 42:161. [14] Master, A. M., S. Dack, H. Horn, B. Freedman, and L. E. Field. 1950. *Circulation*, N. Y. 1:1302. [15] Zimmerman, S. L., and R. Barnett. 1944. *Ann. Int. M.* 21:1045. [16] Scherf, D., and S. D. Klotz. 1944. *Ibid.* 20:438. [17] Dock, W., H. Mandelbaum, and R. A. Mandelbaum. 1953. *Ballistocardiography*. Mosby Company, St. Louis. pp. 65, 99, 106, 118, 120, 126.
[18] Sanghvi, L. M., and K. Banerjee. 1959. *Circulation*, N. Y. 19:41. [19] Sanghvi, L. M. Unpublished.
[20] Rubin, E. H. 1956. *The lung as a mirror of systemic disease*. C. C. Thomas, Springfield, Ill. p. 55.
[21] Ball, D. 1931. *Am. Heart J.* 6:517. [22] Hunter, A. 1946. *Q. J. Med., Oxf.* 15:107. [23] Goldstein, B., and E. P. Boas. 1927. *Arch. Int. M.* 39:226. [24] Heilig, R. 1942. *Ind. M. Gaz.* 77:257. [25] Soloff, L. A., and C. T. Bello. 1950. *Circulation*, N. Y. 2:298. [26] Sanghvi, L. M., and S. N. Misra. 1958. *Am. Heart J.* 55:421.
[27] Zdansky, E. 1953. *Roentgen diagnosis of the heart and great vessels*. Grune and Stratton, New York. p. 214.
[28] Brannon, E. S., A. J. Merrill, J. V. Warren, and E. A. Stead, Jr. 1945. *J. Clin. Invest.* 24:332.

Variable	Congestive Anemia		Acute Anemia		Reference
	Uncomplicated	With Congestive Heart Failure	Small Hemorrhage	Large Hemorrhage	
(A)	(B)	(C)	(D)	(E)	(F)
Pulmonary function (continued)					
O ₂ consumption	Normal	Increased	Normal	Decreased	B, 1, 2, 3, 5, 17, 18
Cardiac output	Increased	Increased	Decreased	Decreased	B, 4, 5, 16, 18
Blood lactate and acid	Increased	Increased	Increased	Increased	17, 18, 19
Renal function	Normal	Decreased	Increased	Increased	C, D, E
Respiratory volume	Normal	Decreased	Increased	Increased	B, C
Respiratory pressure gradient	Normal	Decreased	Increased	Increased	C, 1, 2, 3, 5, 17, 18
Respiratory volume	Normal	Decreased	Increased	Increased	C, 1
Arterial O ₂ pressure gradient	Normal	Decreased	Increased	Increased	C, 1
Arterial O ₂ pressure	Normal	Decreased	Increased	Increased	C, 1
Pulmonary arterial O ₂ pressure	Normal	Decreased	Increased	Increased	C, 1
Peripheral venous O ₂ pressure	Normal	Decreased	Increased	Increased	C, 1
Pulmonary elastic resistance coefficient	Normal	Decreased	Increased	Increased	C, 1
Mean arterial pressure	Normal	Decreased	Increased	Increased	C, 1
Renal function					
Effective plasma flow	Decreased	Decreased	Decreased	Decreased	B, 9, 49, 53, C, 9
Glomerular filtration rate	Decreased	Usually no decrease	Decreased	Decreased	B, 9
Renal excretion	Decreased	Decreased	Decreased	Decreased	B, 9, 49, 53, C, 9
Renal excretion	Decreased	Decreased	Decreased	Decreased	B, 9
Renal excretion	Decreased	Decreased	Decreased	Decreased	B, 4, 5, 16
Renal excretion	Decreased	Decreased	Decreased	Decreased	B, 4, 5, 16

11/ Effects have no linear relationship to the severity of anemia, and therefore no numerical values can be given [5].
 12/ Moderate abnormalities appear when hemoglobin level is between 8 and 9 grams per cent; serious abnormalities do not occur until hemoglobin is less than 7 grams [3]. 13/ Effects of small hemorrhage are increased respiratory rate and occasionally orthostatic syncope [61]. 14/ A liter or more, data include observations during hemorrhagic shock. 15/ Observations made in sickle cell anemia, many of the effects are peculiar to sickle cell disease only, and the possibility exists that associated histological and pathological findings are unrelated to anemia per se [6].
 16/ Hyperventilation may cause alkalemia [62]. 17/ In minimal or early congestive heart failure [6]. 18/ May be normal in children, but decreases with age [53].

Contributors: (a) Sanghvi, L. M., (b) Wintrobe, M. M.

References: [1] Shumgart, H. L., and M. D. Anschule 1943. Blood, N. Y. 3:329. [2] Shumgart, H. L.,

135. HEMODYNAMIC AND PULMONARY EFFECTS OF ANEMIA: MAN AND OTHER ANIMALS (Continued)

Variable (A)	Chronic Anemia ^{1,2}		Acute Anemia		Reference (F)
	Uncomplicated (B)	With Congestive Heart Failure (C)	Small Hemorrhage ³ (D)	Large Hemorrhage ⁴ (E)	
27 Blood pressure (concluded)					
28 Mesenteric artery					
29 Hepatic artery				Decreased	E,39
30 Venous	Almost always normal	Increased	Slightly decreased	Decreased	E,39
31 Inferior vena cava	May increase			Decreased or normal	B,1,6;C,6;D,4 E,19
32 Portal vein				Decreased	B,6,8 E,39
33 Blood flow				Immediately decreased, then increased	E,39
34 Coronary	Increased				
35 Pulmonary	Increased				
36 Cutaneous	Decreased			Decreased	B,41,42;E,18,4
37 Skeletal	Increased in forearm		Decreased	Decreased	B,1;E,44
38 Cerebral	Increased		Decreased	Decreased	B,10,D,E,40
39 Mesenteric				Increased during fainting	B,45,D,40,E,11
40 Hepatic	Increased			Decreased	B,46;E,47
41 Renal	Decreased	Decreased		Decreased	E,39
42 Vascular resistance				Increased	B,36,E,48
43 Total	Decreased	Decreased ¹²	Normal	Decreased	B,9,49,50;C,9, 50,E,16,48
44				Normal or increased	B,1,3,5,7,9,42, C,10,D,E,17
45			May increase	May decrease in shock	D,12,E,17
46 Coronary	Decreased		Decreased during fainting		D,11
47 Pulmonary	Normal			Decreased	B,42,E,18,43
48 Cutaneous	Increased			Inconsistent change	B,7;E,42
49 Cerebral	Decreased		Increased	Increased	B,51;D,E,40
50 Mesenteric				Decreased	B,46,E,47
51 Hepatic				Immediately increased, then decreased	E,39
52 Renal	Increased	Increased		Increased	E,39
53 Pulmonary function					B,C,9
54 O ₂ utilization coefficient ¹³					
55 Systemic	Increased				
56 Cardiac	Increased in severe cases			Increased	B,5,7-9,22,E, 18
Arterio-venous O ₂ difference				Normal	B,41,E,18,42
53 Systemic	Decreased				
54 Coronary	Decreased			Increased	B,3,5,9,E,17,18
55 O ₂ dissociation curve	Displaced to right			Normal	B,41,E,18
56				Decreased	B,42
					B,52

/1/ Effects have no linear relationship to the severity of anemia.

/2/ Moderate anemia.

do not occur in severe anemia and in shock. /

/13/ Relative to the rate of flow.

135. HEMODYNAMIC AND PULMONARY EFFECTS OF ANEMIA: MAN AND OTHER ANIMALS

Variable	Chronic Anemia ^{1,2}		Acute Anemia		Reference
	Uncomplicated	With Congestive Heart Failure	Small Hemorrhage ³	Large Hemorrhage ⁴	
(A)	(B)	(C)	(D)	(E)	(F)
Pulmonary function (concluded)				Normal	B, 1, 2-9, E, 17, 18
O ₂ consumption	Normal			Decreased	B, 41, E, 18, 42
Cardiac	Increased in severe cases			Increased ¹⁴	B, 1, E, 18
Blood lactic acid	Increased		Increased		C, 1; D, 53
Vital capacity		Decreased ¹⁵			B, 54
0.5 sec expiratory capacity/vital capacity x 100	Normal				B, 54
Physiologic dead space	Normal				C, 1
Reserve air		Decreased ¹⁵		Increased	C, 1, 7; E, 55
Respiratory minute volume		Increased ¹⁵			C, 1
Residual air		Increased ¹⁵			B, 56
Alveolar-arterial O ₂ pressure gradient	Increased			Normal	B, 1, E, 55
Arterial O ₂ pressure	Normal				B, 56
Pulmonary arterial O ₂ pressure	May increase				B, 54
Pulmonary arterial O ₂ pressure	Increased				B, 54
Peripheral venous O ₂ pressure	Increased				B, 54
Pulmonary elastic resistance coefficient	Normal				B, 57
Mean inspiratory non-elastic resistance	Normal				B, 57
Renal function					
Effective plasma flow	Decreased ^{8, 16}	Decreased		Decreased	B, 9, 49, 58, C, 9, E, 59
Glomerular filtration rate	Decreased ^{8, 16}	Usually no decrease		Decreased	B, 9, 49, 58, C, 9, E, 59
Tubular excretory capacity	Decreased				B, 49, 58
Salt and water excretion	Decreased				B, 49, 60

- 11) Effects have no linear relationship to the severity of anemia, and therefore no numerical values can be given [6].
 12) Moderate abnormalities appear when hemoglobin level is between 8 and 9 grams per cent, serious abnormalities do not occur until hemoglobin is less than 7 grams [3]. 13) Effects of small hemorrhage are increased respiratory rate and occasionally orthostatic syncope [61]. 14) A liter or more, data include observations during hypovolemic shock. 18) Observations made in sickle cell anemia, many of the effects are similar to those in iron deficiency anemia and the possibility exists that associated histological and pathologic changes are similar. 19) Hyperventilation may cause alkalemia [62]. 115) In man, the effects are similar to those in man, but are normal in children, but decreases with age [58].

Contributors (a) Sangster, L. M., (b) Wintrobe, M. M.

References [1] Blumgart, H. L. and M. D. Altchule 1948 Blood, N. Y. 3:329. [2] Blumgart, H. L.,

- S. L. Gargill, and D. R. Gilligan. 1931. *J. Clin. Invest.* 9:679. [3] Brannon, E. S., A. J. Merrill, J. V. Warren, and E. A. Stead, Jr. 1945. *Ibid.* 24:332. [4] Stewart, H. J., N. F. Crane, and J. E. Deltick. 1937. *Ibid.* 16:431. [5] Wintrobe, M. M. 1946. *Blood*, N. Y. 1:121. [6] Sanghvi, L. M., R. Sharma, and S. N. Misra. 1957. *Circulation*, N. Y. 15:373. [7] Leight, L., T. H. Snider, G. O. Clifford, and H. K. Hellems. 1954. *Ibid.* 10:653. [8] Sharpey-Schafer, E. P. 1944. *Clin. Sc.*, Lond. 5:125. [9] Whitaker, W. 1956. *Q. J. Med.*, Oxf. 25:175. [10] Friedberg, C. K. 1956. *Diseases of the heart*. W. B. Saunders, Philadelphia. pp. 1046-1053. [11] Barcroft, H., O. G. Edholm, J. McMichael, and E. P. Sharpey-Schafer. 1944. *Lancet*, Lond. 1:489. [12] McMichael, J. 1947. *Advances in internal medicine*. Interscience, New York. v. 2, p. 92. [13] Warren, J. V., E. S. Brannon, E. A. Stead, Jr., and A. J. Merrill. 1945. *J. Clin. Invest.* 24:337. [14] Courmand, A., and R. L. Riley, et al. 1943. *Surgery* 13:964. [15] Mukherjee, S. R., and S. Rowlands. 1951. *Lancet*, Lond. 1:1041. [16] Richards, D. W., Jr. 1944. *Bull. N. York Acad. M.* 20:363. [17] Brannon, E. S., E. A. Stead, Jr., J. V. Warren, and A. J. Merrill. 1946. *Am. Heart J.* 31:407. [18] Edwards, W. S., A. Siegel, and R. J. Bing. 1954. *J. Clin. Invest.* 33:1646. [19] Shenkin, H. A., R. H. Cheney, S. R. Govons, J. D. Hardy, A. G. Fletcher, and I. Starr. 1944. *Am. J. M. Sc.* 208:421. [20] Porter, W. B., and D. W. James. 1953. *Circulation*, N. Y. 8:111. [21] Walcott, W. W. 1945. *Am. J. Physiol.* 143:247. [22] Bishop, J. M., K. W. Donald, and O. L. Wade. 1955. *Clin. Sc.*, Lond. 14:329. [23] Gibson, J. G., A. W. Harris, and V. W. Swigert. 1939. *J. Clin. Invest.* 18:621. [24] Noble, R. P., and M. I. Gregersen. 1946. *Ibid.* 25:172. [25] Mendiowitz, M. 1943. *Ibid.* 27:565. [26] Fine, J., H. A. Frank, and A. M. Seligman. 1944. *Ibid.* 23:731. [27] Moon, V. H. 1942. *Shock*. Lea and Febiger, Philadelphia. p. 105. [28] Fishberg, A. M. 1940. *Heart failure*. Lea and Febiger, Philadelphia. pp. 197-199. [29] Ebert, R. V., E. A. Stead, Jr., and J. G. Gibson. 1941. *Arch. Int. M.* 68:578. [30] Stead, E. A., Jr. 1940. *J. Clin. Invest.* 19:776. [31] Hunter, A. 1946. *Q. J. Med.*, Oxf. 15:107. [32] Sanghvi, L. M. Unpublished. [33] Wallace, J., and E. P. Sharpey-Schafer. 1941. *Lancet*, Lond. 2:393. [34] Klinefelter, H. F. 1942. *Am. J. M. Sc.* 203:34. [35] Dack, S. E. Corday, and A. M. Master. 1951. *Am. Heart J.* 42:161. [36] Ruskin, A. 1953. *Physiological cardiology*. C. C. Thomas, Springfield, Ill. p. 266. [37] Warren, J. V., J. T. Doyle, E. H. Estes, and J. S. Wilson. 1951. *J. Clin. Invest.* 30:680. [38] Hild, R., K. Mechelke, and E. Nusser. 1956. *Pflügers Arch.* 263:401. [39] Cull, T. E., M. P. Scibetta, and E. E. Selkurt. 1956. *Am. J. Physiol.* 185:365. [40] McDowall, R. J. S. 1956. *The control of the circulation of the blood*. Dawson and Sons, London. pp. 449-455. [41] Bing, R. J. 1951. *Bull. N. York Acad. M.* 27:407. [42] Case, R. B., E. Berglund, and S. J. Sarnoff. 1955. *Am. J. Med.* 18:397. [43] Opdyke, D. F., and R. C. Foreman. 1947. *Am. J. Physiol.* 148:726. [44] Richards, D. W., Jr. 1945. *Fed. Proc.*, Balt. 4:215. [45] Abramson, D. L., S. M. Fierst, and K. Flachs. 1943. *Am. Heart J.* 25:609. [46] Heyman, A., J. L. Patterson, Jr., and T. W. Duke. 1952. *J. Clin. Invest.* 31:824. [47] Fazekas, J. F., J. Kien, and A. E. Parrish. 1955. *Am. J. M. Sc.* 229:41. [48] Heinemann, H. O., C. M. Smythe, and P. A. Marks. 1953. *Am. J. Physiol.* 174:352. [49] Bradley, S. E., and G. P. Bradley. 1947. *Blood*, N. Y. 2:192. [50] Merrill, A. J. 1946. *J. Clin. Invest.* 25:389. [51] Fahr, G. E., and E. Ronzone. 1922. *Arch. Int. M.* 29:331. [52] Kennedy, A. C., and D. J. Valtis. 1954. *J. Clin. Invest.* 33:1372. [53] Glaser, E. M., and J. McMichael. 1940. *Lancet*, Lond. 2:230. [54] Sproule, B. J., E. R. Halden, and W. F. Miller. 1958. *J. Clin. Invest.* 37:486. [55] Frank, H. A. 1953. *N. England J. M.* 249:445. [56] Ryan, J. M., and J. B. Hickam. 1952. *J. Clin. Invest.* 31:186. [57] McIlroy, M. B., F. L. Eldridge, and R. W. Stone. 1956. *Clin. Sc.*, Lond. 15:353. [58] Eiteldorf, J. N., J. D. Smith, A. H. Tuttle, and L. W. Diggs. 1955. *Am. J. Med.* 18:243. [59] Van Slyke, D. D. 1948. *Ann. Int. M.* 28:701. [60] Sirausa, M. B., and H. J. Fox. 1940. *Am. J. M. Sc.* 200:454. [61] Barcroft, H., O. G. Edholm, J. McMichael, and E. P. Sharpey-Schafer. 1944. *Lancet*, Lond. 1:489. [62] Bald, F. W. 1927. *Am. J. Physiol.* 81:222

APPENDIXES

4. Mitral valve resistance = $\frac{(LA_{dm} - LV_{dm}) \times 1332}{CO \times \frac{\text{duration of total cardiac cycles (measured)}}{\text{duration of diastoles of same cardiac cycles (measured)}}}$ = dynes sec cm⁻⁵
5. Tricuspid valve resistance = $\frac{(RA_{dm} - RV_{dm}) \times 1332}{CO \times \frac{\text{duration of total cardiac cycles (measured)}}{\text{duration of diastoles of same cardiac cycles (measured)}}}$ = dynes sec cm⁻⁵
6. Pulmonary valve resistance = $\frac{(RV_{sm} - PA_{sm}) \times 1332}{PBF \times \frac{\text{duration of total cardiac cycles (measured)}}{\text{duration of systolic ejections of same cardiac cycles (measured)}}}$ = dynes sec cm⁻⁵
7. Mitral valve flow (ml/sec) = $\frac{\text{cardiac output in ml/min}}{\text{duration of the sum of diastolic periods in sec/min}}$
8. Mitral valve area¹ = $\frac{MVF}{31 \sqrt{PC_m - 5}} = \frac{MVF}{31 \sqrt{LA_{dm} - LV_{dm}}}$
9. Right ventricular work² = $\frac{13.6 (PF \times 1.055) (PA_{sm} - RA_{sm})}{1000}$ = kg m/min/sq m body surface area

/1/ Formulas for calculating area of pulmonary, tricuspid, and aortic valves, and for calculating size of an atrial or ventricular septal defect, or for measuring diameter of a patent ductus arteriosus, may be found in references 7 and 12. /2/ Left ventricular work may be estimated by substituting brachial or femoral for pulmonary, arterial systolic mean pressure, and left for right, atrial mean pressure.

Contributors: (a) Witham, A. Calhoun, (b) Chapman, Don W., (c) Luisada, Aldo A.

References: [1] Cournand, A., R. L. Riley, E. S. Breed, E. deF. Baldwin, and D. W. Richards, Jr. 1945. J. Clin. Invest. 24:106. [2] Aperia, A. 1940. Skand. Arch. Physiol., Berl., Suppl. 16, v. 83. [3] Bing, R. T., L. D. Vandam, and F. D. Gray, Jr. 1947. Bull. Johns Hopkins Hosp. 80:107. [4] Bradford, J. R., and H. P. Dean. 1894. J. Physiol., Lond. 16:34. [5] Fick, A. 1870. Sitzber. Phys. med. Ges. Würzburg 2.16. [6] Evans, C. L. 1936. Starling's Principles of human physiology. Ed. 7. Lea and Febiger, Philadelphia. [7] Luisada, A. A., and C. K. Liu. 1958. Intracardiac phenomena. Grune and Stratton, New York. [8] Burton, A. C. 1953. Annual Rev. Physiol. 15:223. [9] Cournand, A., J. S. Baldwin, and A. Himmelstein. 1949. Cardiac catheterization in congenital heart disease. Commonwealth Fund, New York. [10] Cournand, A., H. L. Molley, A. Himmelstein, D. Dresdale, and J. S. Baldwin. 1947. Am. J. Physiol. 150:267. [11] Facquet, J., J. M. Lemoine, P. M. Alhomme, and J. Lefebvre. 1952. Arch. mal. coeur 45:741. [12] Gorlin, R., and S. G. Gorlin. 1951. Am. Heart J. 41:1. [13] Gorlin, R., and L. Dexter. 1952. Ibid. 43:188. [14] Hellem, H. K., F. W. Haynes, L. Dexter, and T. D. Kinney. 1948. Am. J. Physiol. 155:98. [15] Lagerlöf, H., and L. Werkö. 1949. Scand J. Clin. & Invest., Oslo 1:147. [16] Wiggers, C. J. 1949. Physiology in health and disease. Ed. 5. Lea and Febiger, Philadelphia.

APPENDIX II. REGRESSION EQUATIONS

I. Relations Between Stroke Volume, Left Ventricular Work, and Various Aspects of Blood Pressure, Determined by Blood Perfusion Method Simulating Systole in Cadavers. [1-4]

Data for regression equations were secured from cadavers perfused by blood at necropsy. Stroke volume was measured in the syringe, work was estimated by integration of the product of pressure and flow values at each instant, and blood pressures were obtained from a capacitance manometer attached to a needle in the femoral artery.

Abbreviations: SV = stroke volume, ml; W = left ventricular work, g-m; FSP = femoral systolic pressure, mm Hg; FDP = femoral diastolic pressure, mm Hg; FPP = femoral pulse pressure, mm Hg; FMP = femoral mean pressure, mm Hg; ASP = aortic systolic pressure, mm Hg; AMP = aortic mean pressure, mm Hg; A = age, yr; SA = body surface area, sq m; PWV = pulse wave velocity, m/sec; Q = cross section area of the root of the aorta, sq cm; ml = milliliter, g-m = gram meters.

Regression Equation (A)	Standard Deviation (B)
1 SV = 48.1 - 0.38 PWV	14.3 ml
2 SV = 50.5 - 0.09 A	14.3 ml
3 SV = 19.3 + 15.66 SA	14.0 ml
4 SV = 37.3 - 0.18 FDP	13.9 ml
5 SV = 28.0 + 0.30 FPP	13.9 ml
6 SV = 51.6 - 0.07 A - 0.28 PWV	14.3 ml
7 SV = 35.9 + 0.03 A + 16.49 SA	14.0 ml
8 SV = 16.3 + 21.13 SA - 1.14 PWV	13.8 ml
9 SV = 55.6 - 0.20 FDP + 0.43 PWV	13.8 ml
10 SV = 74.2 - 0.22 FDP - 0.22 A	13.6 ml
11 SV = 26.4 - 0.22 FDP + 20.24 SA	13.2 ml
12 SV = 14.8 + 0.29 FPP + 8.41 SA	11.8 ml
13 SV = 38.6 + 0.32 FPP - 0.19 A	11.7 ml
14 SV = 42.6 + 0.49 FPP - 3.55 PWV	9.6 ml
15 SV = 47.4 + 0.43 FPP - 0.40 FDP	8.9 ml
16 SV = 26.4 + 0.42 FPP - 0.42 FDP + 13.95 SA	8.5 ml
17 SV = 54.3 + 0.55 FPP - 0.32 FDP - 2.60 PWV	7.5 ml
18 SV = 91.0 + 0.34 FPP - 0.57 FDP - 0.61 A	5.9 ml
19 SV = 94.4 + 0.54 FPP - 0.58 FDP - 0.52 A - 1.60 SA	5.9 ml
20 SV = 54.7 + 0.57 FPP - 0.44 FDP - 0.31 A + 13.80 SA - 2.30 PWV	5.2 ml
21 SV = 90.97 + 0.73 FPP - 0.57 (FDP + 1/3 FPP) - 0.61 A	5.9 ml
22 SV = 90.97 + 0.82 FPP - 0.57 (FDP + 1/2 FPP) - 0.61 A	5.9 ml
23 SV = 90.96 + 0.92 FPP - 0.57 (FDP + 2/3 FPP) - 0.61 A	5.9 ml
When $Z_1 = FMP = 1/2 (FSP + FDP)$ then	5.9 ml
24 SV = 20 + $\frac{45}{Z_1}$ (FPP)	11.5 ml
25 SV = 47 + $\frac{56}{Z_1}$ (FPP) - 0.59 A	9.3 ml
When $Z_2 = 1/3 (2 FSP + FDP)$ then	
26 SV = 18 + $\frac{53}{Z_2}$ (FPP)	11.6 ml
27 SV = 43 + $\frac{71}{Z_2}$ (FPP) - 0.59 A	9.4 ml

APPENDIX II. REGRESSION EQUATIONS (Continued)

Regression Equation (A)		Standard Deviation (B)
28	$SV = 50 - 0.7 (Q \times 10)$	14.2 ml
29	$SV = 45 - 0.003 \left(\frac{Q}{PWV} \times 10^2 \right)$	14.4 ml
30	$SV = 40 + 0.14 (FPP) (Q) (10^{-1})$	14.0 ml
31	$SV = 40 + 0.10 \frac{(FPP) (Q)}{PWV}$	14.0 ml
32	$SV = -13.4 + 20.8 \sqrt{\frac{FPP}{PWV}}$	9.3 ml
33	$SV = -11 + 0.05 FPP + 1.08 \left(\frac{1}{PWV} \right) (10^2)$	8.9 ml
Standard deviation of the values for stroke volume, found in the various experiments, about their own mean.		14.4 ml
34	$W = -8.83 + 0.93 SV (AMP) (13.6) (10^{-3})$	10.9 g-m
35	$W = -6.37 + 0.89 SV (FMP) (13.6) (10^{-3})$	12.0 g-m
Let $X_4 = SV$ estimated by equation 18		
36	$\therefore W = -1.67 + 0.83 X_4 (FMP) (13.6) (10^{-3})$	14.9 g-m
Let $X_5 = SV$ estimated by equation 25		
37	$\therefore W = -7.96 + 1.22 X_5 (FMP) (13.6) (10^{-3})$	16.5 g-m
38	$W = 1.1 + 1.13 FPP$	18.3 g-m
39	$W = 0.54 + 68.4 \frac{FPP}{A}$	16.6 g-m
40	$W = -9 + 0.78 FMP$	26.2 g-m
41	$W = 44.9 + 1.21 FPP - 0.79 A$	15.3 g-m
42	$W = 51.5 + 1.13 FPP + 31.48 SA$	16.6 g-m
43	$W = 15.5 + 1.20 FPP - 0.65 A + 12.9 SA$	15.1 g-m
44	$W = 13.6 + 1.30 FPP - 3.12 PWV$	17.2 g-m
45	$W = 15.16 + 1.13 FPP - 2.22 Q$	16.7 g-m
46	$W = 29.57 + 1.31 FPP - 3.38 PWV - 2.35 Q$	15.4 g-m
Standard deviation of the values for work, found in the various experiments, about their own mean		32.2 g-m

II. Relations Between Stroke Volume, Left Ventricular Work, and Various Aspects of Blood Pressure, Determined by Arm Auscultatory Method [1-6]

Abbreviations: SV = stroke volume, ml, W = left ventricular work, g-m; SP (aus) = auscultatory systolic pressure, mm Hg; DP (muff) = auscultatory diastolic pressure estimated as the level of muffling of sounds; DP (dis) = auscultatory diastolic pressure estimated as the level of disappearance of sounds, BSP, BDP, BPP and RSP, RDP, RPP = brachial systolic, brachial diastolic, brachial pulse, radial systolic, radial diastolic, and radial pulse pressures, respectively, mm Hg, FMP = femoral mean pressure, mm Hg, A = age, yr.

Regression Equation (A)		Standard Deviation (B)
1	$BSP = -6.08 + 1.04 SP (aus)$	10.3 mm Hg
2	$BDP = 4.86 + 0.83 DP (muff)$	7.7 mm Hg
3	$BPP = 8.56 + 1.01 [SP (aus) - DP (muff)]$	10.9 mm Hg
4	$SV = 93 + 0.54 [SP (aus) - DP (muff)] - 0.47 DP (muff) - 0.61 A$	
5	$RPP = 14.54 + 0.94 [SP (aus) - DP (dis)]$	3.9 mm Hg
6	$RSP = 16.62 + 1.00 SP (aus)$	12.5 mm Hg
7	$RDP = -3.2 + 1.03 DP (dis)$	12.5 mm Hg

APPENDIX II. REGRESSION EQUATIONS (Continued)

Regression Equation (A)		Standard Deviation (B)
8	$SV = 101 + 0.5 [SP (aus) - DP (dis)] - 0.59 DP (dis) - 0.61 A$	
9	$SV = 100 + 0.5 [SP (aus) - DP (dis)] - 0.6 DP (dis) - 0.6 A$ Let X_6 = an estimate of SV by equation 4 and Let Y_6 = an estimate of FMP as follows.	
10	$FMP = -0.61 + 0.52 [SP (aus) + 0.42 DP (muff)]$	
11	$W = -1.67 + 0.0113 X_6 Y_6$ Let X_4 = an estimate of SV by equation 8 and Let Y_4 = an estimate of FMP as follows:	
12	$FMP = 3.21 + 0.5 [SP (aus) + 0.52 DP (dis)]$	
13	$W = -1.67 + 0.0113 X_4 Y_4$	

III. Relation Between Stroke Volume and Amplitude of Certain Waves of the High-Frequency Ballistocardiogram. [7]

Data for regression equations were secured from experiments on cadavers perfused with blood while lying on a high-frequency ballistocardiograph. Ballistocardiograms show influence of size and age of subjects; abnormal tracings were excluded.

Abbreviations: SV = stroke volume, ml, Ht = height of subject, cm, Wt = weight of subject, kg; A = age of subject, yr, SA = body surface area, sq m, C = duration of cardiac cycle, sec, I and J = areas of I and J waves from base line of record, mm sec, when calibration of record is such that 280 grams displaces the base line 1 centimeter

Regression Equation (A)		Standard Deviation (B)
1	$SV = 11 + 0.27 Ht \sqrt{(21 + J)} \sqrt{C}$	9 ml
2	$SV = 14 + 0.64 Wt \sqrt{(21 + J)} \sqrt{C}$	7.9 ml
3	$SV = 11 + \frac{2700}{A} \sqrt{(21 + J)} \sqrt{C}$	7.8 ml
4	$SV = 10 + 27 SA \sqrt{(21 + J)} \sqrt{C}$	7.7 ml
5	$SV = 15 + 38.5 \sqrt{(21 + J)} \sqrt{C}$	9.9 ml
6	$SV = -27 + 42 \sqrt{(21 + J)} \sqrt{C} + 24 SA$	8.0 ml
7	$SV = 31 + 45 \sqrt{(21 + J)} \sqrt{C} - 0.49 A$	8.0 ml
8	$SV = 4 + 45 \sqrt{(21 + J)} \sqrt{C} + 15 SA - 0.29 A$	7.5 ml
Standard deviation of the values for stroke volume, in 51 experiments, about their own mean.		14.6 ml

IV. Empirical Normal Standards for High-Frequency Ballistocardiograms. [8]

Normal subjects tested after 15-minute rest on the high-frequency ballistocardiograph, and never within 2 hours of a meal. The instrument was calibrated statically so that a force of 280 grams displaced the light spot 1 centimeter.

Abbreviations: I and J = the altitude or areas of waves typical of the largest wave of the respiratory cycle, I₂ and J₂ = the altitude or areas of waves typical of the smallest waves of the respiratory cycle, PR = pulse rate, beats/min; MBP = mean blood pressure, mm Hg; mm sec = millimeter second.

Aspect Studied	Measurement Analyzed	Subjects		Results		
		Age, yr	No. and Sex	Mean	Standard Deviation	S. D. % ¹
(A)	(B)	(C)	(D)	(E)	(F)	(G)
1 Wave altitudes	$\sqrt{I + J + I_2 + J_2}$ mm	20-39	56♂	5.4 mm	0.58 mm	10.7
2			48♀	4.4 mm	0.49 mm	11.2
3 Wave altitudes x PR	$\sqrt{I + J + I_2 + J_2}$ mm x PR	20-39	56♂	385	52.5	13.7
4			48♀	318	55.5	17.4
5 Wave altitudes	$\sqrt{I + J + I_2 + J_2}$ mm	20-49	106♂	5.3 mm	0.66 mm	12.5
6			88♀	4.5 mm	0.62 mm	13.8
7 Wave areas	$\sqrt{I + J + I_2 + J_2}$ mm sec	20-49	103♂	0.83 mm sec	0.11 mm sec	13.3
8			81♀	0.67 mm sec	0.07 mm sec	10.4
9 Wave areas x PR	$\sqrt{I + J + I_2 + J_2}$ mm sec x PR	20-49	101♂	81	12.2	15.0
10			81♀	68	12.4	18.3
11 Wave areas x MBP	$\sqrt{I + J + I_2 + J_2}$ mm sec x MBP	20-49	98♂	122	16.0	14.1
12			81♀	89	12.6	14.2

/1/ Standard deviation as % of mean.

V. Normal Standards for Amplitude of Ballistocardiograms, Allowing for the Effect of Age. [8]

Normal subjects, ranging from 20-49 years of age, tested after 15-minute rest on the high-frequency ballistocardiograph, and never within 2 hours of a meal. The instrument was calibrated statically so that a force of 280 grams displaced the light spot 1 centimeter.

Abbreviations: I and J = the altitude or areas of waves typical of the largest waves of the respiratory cycle; I₂ and J₂ = the altitude or areas of waves typical of the smallest waves of the respiratory cycle; A = age, yr; PR = pulse rate, beats/min, mm sec = millimeter second

Aspect Studied	Subjects, No. and Sex	Regression Equation	Standard Deviation about the Regression	S. D. % ¹
(A)	(B)	(C)	(D)	(E)
1 Wave altitudes	105♂	$\sqrt{I + J + I_2 + J_2}$ mm = -0.04 A + 6.7	0.60 mm	11.3
2	87♀	$\sqrt{I + J + I_2 + J_2}$ mm = -0.03 A + 5.5	0.53 mm	11.8
3 Wave areas	101♂	$\sqrt{I + J + I_2 + J_2}$ mm sec = -0.0071 A + 1.4	0.142 mm sec	12.1
4	81♀	$\sqrt{I + J + I_2 + J_2}$ mm sec = -0.0042 A - 1.07	0.085 mm sec	9.0
5 Wave area x PR	101♂	$\sqrt{I + J + I_2 + J_2}$ mm sec x PR = -0.608 A + 102.3	11.0	13.6
6	81♀	$\sqrt{I + J + I_2 + J_2}$ mm sec x PR = -0.495 A + 82.4	11.6	17.0

/1/ Standard deviation about the regression as % of mean.

APPENDIX II. REGRESSION EQUATIONS (Concluded)

VL. Relation Between Various Aspects of the Femoral Pulse Wave and Aspects of Left Ventricular Function. [9]

Data for regression equations were secured from blood perfusion experiments in which systole was simulated in cadavers. Femoral pulse wave secured by a needle in the femoral artery.

Abbreviations MEV = maximum ejection velocity, ml/sec (the integral of the preceding accelerations), AMP = aortic mean blood pressure, mm Hg; FPP = femoral pulse pressure, mm Hg; Z = maximum slope of femoral pulse wave front, mm Hg/sec, PWV = pulse wave velocity, m/sec; SV = stroke volume, ml; FASA = adjusted systolic area of femoral pulse wave, mm Hg/sec (the area above a straight line intersecting the pressure record at the beginning and end of ejection). FMP = femoral mean blood pressure, mm Hg.

Regression Equation	Standard Deviation about the Regression	Standard Deviation of MEV x AMP(10 ⁻²), MEV, and SV about its own mean
(A)	(B)	(C)
1 MEV x AMP(10 ⁻²) = 63.8 + 0.29 Z	65	166
2 MEV x AMP(10 ⁻²) = -11.0 + 5.43 FPP	67	160
3 MEV = 158 + 1.36 $\frac{Z}{PWV}$	59	109 ml/sec
4 MEV = 10 + 3.16 $\sqrt{\frac{Z}{PWV} \times 100}$	53	109 ml/sec
5 SV = 22 + 350 $\frac{FASA}{FMP}$	9.3 ml	14.7 ml
6 SV = -2.7 + 190 $\sqrt{\frac{FASA}{FMP}}$	8.6 ml	14.7 ml
7 SV = 18.6 + 30 $\frac{FASA}{PWV}$	7.7 ml	14.7 ml
8 SV = -5.4 + 55 $\sqrt{\frac{FASA}{PWV}}$	7.4 ml	14.7 ml

Contributor Starr, Isaac

References [1] Starr, L. T. G. Schnabel, Jr., and R. L. Mayock 1953. Circulation, N. Y. 8 44. [2] Starr, L., T. G. Schnabel, Jr., S. I. Askovitz, and A. Schild. 1954. Ibid. 9, 648. [3] Starr, L., S. I. Askovitz, W. Feder, and A. Schild. 1955. Ibid. 12 1005. [4] Starr, L. Unpublished. [5] Hagan, C., and J. Bordley. 1941. Bull. Johns Hopkins Hosp. 69 526. [6] Steele, J. M. 1941. J. Mount Sinai Hosp. N. York 8, 1042. [7] Starr, L. 1955. J. Appl. Physiol. 8: 115. [8] Starr, L. 1955. Circulation, N. Y. 11: 914. [9] Starr, L. 1958. Ibid. 17: 589.

**APPENDIX III. EMPIRICAL NORMAL STANDARDS FOR ULTRA-LOW
FREQUENCY BALLISTOCARDIOGRAMS**

Recorded are normal displacement, velocity, and acceleration of the human body during the cardiac cycle, when the body is free to move in space. Mean values are given, with the standard deviation in parentheses.

Data on Subjects

Subjects, No. and Sex	Age yr	Height in.	Body Weight kg	Body and Platform Weight kg	Body Surface Area sq m	Pulse Rate beats/min
(A)	(B)	(C)	(D)	(E)	(F)	(G)
1 50♂	29.6 (4.8)	70.72 (2.32)	76.25 (10.15)	83.05 (10.15)	1.929 (0.137)	65.3 (8.8)
2 50♀	28.8 (7.1)	65.12 (2.41)	58.98 (8.36)	65.78 (8.36)	1.635 (0.132)	71.1 (9.4)

I Standards for Amplitude.

1. Displacement.

Abbreviations: "IJ = vertical distance from tip of I wave to tip of J wave, "JM = vertical distance from tip of J wave to tip of M wave; μ = microns measuring actual movement of subject in space.

Subjects	"IJ		"JM		"IJ"/"JM
	μ	g-cm	μ	g-cm	%
1 50♂	37.73	309.4	86.80	722.0	44.4
20-39 yr	(10.33)	(88.51)	(22.50)	(225.4)	(12.8)
2 50♀	26.55	174.0	68.84	453.3	39.8
20-39 yr	(8.70)	(59.4)	(18.80)	(130.8)	(12.2)

2. Velocity.

Abbreviations: 'I and 'J = velocity equivalent to deflection of these waves from the base line; 'HI and 'IJ = velocity equivalent to the vertical distance between wave tips.

Subjects	'I		'J		'HI		'IJ	
	mm/sec	g-m/sec	mm/sec	g-m/sec	mm/sec	g-m/sec	mm/sec	g-m/sec
1 50♂	0.66	54.4	0.66	54.6	0.78	64.8	1.31	108.9
20-39 yr	(0.15)	(13.2)	(0.13)	(12.9)	(0.16)	(14.2)	(0.22)	(20.8)
2 50♀	0.48	32.1	0.57	37.7	0.57	37.5	1.07	69.9
20-39 yr	(0.13)	(8.3)	(0.10)	(8.2)	(0.12)	(8.6)	(0.19)	(14.5)

3. Acceleration.

Abbreviations: H, I, and J = force equivalent to displacement of the tip of these waves from the base line, g = force in grams.

Subjects	H	I	J
	g	g	g
1 50♂	94.0	226.2	239.2
20-39 yr	(33)	(56.3)	(49.7)
2 50♀	52.6	146.0	177.3
20-39 yr	(24)	(42.5)	(51.0)

APPENDIX III. EMPIRICAL NORMAL STANDARDS FOR ULTRA-LOW FREQUENCY BALLISTOCARDIOGRAMS (Concluded)

Standards for Time.

1. Displacement.

Abbreviations: Q-"G₀, Q-"H, Q-"I, Q-"J, and Q-"M = time in seconds between the Q wave of the electrocardiogram and the peak of the corresponding wave of the displacement ballistocardiogram when recorded simultaneously. "G₀ = onset of G wave.

Subjects	Q-"G ₀	Q-"H	Q-"I	Q-"J	Q-"M
1 50σ	0.064 ¹	0.087 ²	0.134	0.228	0.440
20-39 yr	{0.018}	{0.022}	{0.013}	{0.020}	{0.025}
2 50σ	0.075 ³	0.086 ⁴	0.128	0.212	0.433
20-39 yr	{0.024}	{0.016}	{0.012}	{0.016}	{0.029}

1/ Value for 32 subjects. 2/ Value for 49 subjects. 3/ Value for 25 subjects. 4/ Value for 46 subjects.

2. Velocity.

Abbreviations: Q-"H, Q-"I, Q-"J, Q-"L, and Q-"M = time in seconds between the Q wave of the electrocardiogram and the peaks of the corresponding waves of the velocity ballistocardiogram when recorded simultaneously; 'I₁ and 'J₁ = duration of the 'I and 'J waves on the base line.

Subjects	Q-"H	Q-"I	Q-"J	Q-"L	Q-"M	'I ₁	'J ₁
1 50σ	0.115	0.182	0.305	0.426 ¹	0.533 ²	0.098	0.219
20-39 yr	{0.013}	{0.013}	{0.021}	{0.021}	{0.029}	{0.019}	{0.029}
2 50σ	0.109	0.170	0.282	0.422	0.490	0.086	0.229
20-39 yr	{0.013}	{0.008}	{0.026}	{0.021}	{0.028}	{0.017}	{0.020}

1/ Value for 49 subjects. 2/ Value for 44 subjects.

3. Acceleration.

Abbreviations: Q-"H, Q-"I, Q-"J, Q-"L₀, Q-"L, Q-"M, Q-"N = time in seconds between the Q wave of the electrocardiogram and the peaks of the corresponding wave of the acceleration ballistocardiogram when recorded simultaneously. L₀ = small negative wave usually preceding the L wave, I₁ and J₁ = duration in seconds of the I and J waves on the base line.

Subjects	Q-"H	Q-"I	Q-"J	Q-"L ₀	Q-"L	Q-"M	Q-"N	I ₁	J ₁
1 50σ	0.101	0.154	0.235	0.390	0.418	0.451	0.557 ¹	0.064	0.127
20-39 yr	{0.014}	{0.014}	{0.020}	{0.021}	{0.023}	{0.024}	{0.031}	{0.012}	{0.011}
2 50σ	0.099	0.150	0.216	0.387	0.415	0.447	0.544 ¹	0.057	0.116
20-39 yr	{0.013}	{0.012}	{0.017}	{0.023}	{0.021}	{0.024}	{0.032}	{0.011}	{0.016}

1/ Value for 48 subjects

Contributor: Starr, Isaac

Reference: Scarborough, W R, E E Folk, III, P M Smith, and J H Condon. 1958. Am J. Cardiology 2 613.

INDEX

AMMONIUM SULFAMATE, translocation (plants), 301
AMPLITUDE (See also ELECTROCARDIOGRAM; BALLISTOCARDIOGRAM; specific waves)
ballistocardiogram, normal standards, 360, 362, 363
electrocardiogram (man), 142-145; (farm animals), 146
at altitude, 165
during hypoxia, 165
in myxedema, 336
AMYLOIDOSIS, pulse configuration, 156*
ANABASINE SULFATE, effect on heart (insects), 278
ANACROTIC PULSE, configuration, 157*
ANALEPTICS
cardiovascular action, 251
hemodynamic effects of specific drugs (mammals), 257, 258, 270
site of action, 246
ANALGESICS
cardiovascular action, 250
hemodynamic effects of specific drugs (mammals), 255, 262, 269
ANATOMY, COMPARATIVE (vertebrates)
blood vessels, 12-15
heart, 10, 11
lymphatics, 16-18
ANEMIA, 343*, 344-352
arterio-venous O₂ difference (man), 330; (man and other animals), 350
blood pressures (man), 330; (man and other animals), 349, 350
cardiac index, 78, 330
cardiac output (man), 78, 330, (man and other animals), 349
circulation time during, 121
heart rate, 330
hemoglobin, 330
from ionizing radiation (vertebrates), 217-219, 223-227
O₂ consumption, 330
physiologic alterations and compensatory mechanisms, 343*
ANESTHESIA, effect on
blood flow
cerebral, 125
digital, 133, 134
hepatic (mammals), 128, 129
limb, 133
cardiac output (mammals other than man), 80
ANESTHETICS
cardiovascular action, 250
hemodynamic effects of specific drugs (mammals), 255, 257, 259, 262, 264, 268, 269, 271
sites of action, 246
ANIONS, cardiovascular action, 252
ANIONS (BLOOD), effect of shock (dog), 320
ANOXIA, effect on cardiac output, 78
ANTERIOR CARDINAL VEINS, comparative anatomy (vertebrates), 13
ANTHRACENEACETIC ACID, translocation (plants), 301
ANTIARRHYTHMIC DRUGS
general effects, 250
hemodynamic effects of specific drugs (mammals), 255, 264, 269
ANTIBIOTICS
cardiovascular action, 251
hemodynamic effects of penicillin, 260

ANTICOAGULANTS (concluded)
dosages in vivo (man), 235, 240, 241, 259, (vertebrates), 236-238
effect on
circulatory function (vertebrates), 242-245
clotting time (man), 241; (vertebrates), 236-238
natural-occurring: preparation and assay, 233, 234
oral, 235
ANTICONVULSANTS (See also ANTIARRHYTHMIC DRUGS)
cardiovascular action, 250
hemodynamic effects of diphenylhydantoin sodium, 241
ANTIFIBRILLATORY DRUGS (See also ANTIARRHYTHMIC DRUGS)
hemodynamic effects of quinidine sulfate (mammals), 255, 269
site of action, 246
ANTIHEMOPHILIC FACTOR
in coagulation stages, 231*
preparation and assay, 233
ANTIHEMOPHILIC GLOBULIN (See ANTIHEMOPHILIC FACTOR)
ANTIHISTAMINES (See also ANTIARRHYTHMIC DRUGS)
cardiovascular action, 251
hemodynamic effects of diphenhydramine HCl, 269
site of action, 246
ANTIHYPERTENSIVES (See also VERATRUM, RAUWOLFIA ALKALOIDS)
general effects, 247
hemodynamic effects of specific drugs (mammals), 253, 257, 258, 261, 265, 267, 268, 271
sites of action, 246
ANTIMALARIALS
cardiovascular action, 251
hemodynamic effects of specific drugs, 269
ANTIMYOCIN A, effect on heart (insect), 278
ANTIPLATELET SERUM, effect on circulatory function (dog), 242
ANTITHROMBIN
in coagulation stages, 231*, 232*
preparation and assay, 233
ANXIETY, effect on cardiac index, 78
AORTA, 7*, 8* (See also AORTIC ARCH(ES), ARTERY(IES))
comparative anatomy (vertebrates), 12
development (man), 4, 5, (chick), 2, (frog), 1; (pig), dimensions, 21
elasticity (man, dog), 60, 61
lumina, growth at various ages, 20
O₂ saturation in congenital heart malformations, 152, 153
thickness of wall, 20
thoracic (See THORACIC AORTA)
AORTIC ARCH(ES), 7*, 8*
blood flow, during hypercapnia (dog), 171
comparative anatomy (vertebrates), 12
development (man), 4, 5; (frog), 1, (pig), 3
pulse waves, 113
stretch receptors: drug stimulants and depressants, 246
AORTIC INSUFFICIENCY
effect on
blood flow (man), 133, 325, (man, dog), 130

AORTIC INSUFFICIENCY (concluded)

effect on (concluded)
 blood pressures (man), 322; (man, dog), 130
 vascular resistance (man), 325; (man, dog), 130
 murmur in: phonocardiographic configuration, 154*
 pulse configuration, 157*

AORTIC LESIONS, effect on circulation time, 123

AORTIC SAC (See TRUNCUS ARTERIOSUS)

AORTIC STENOSIS

effect on
 blood flow (man), 325; (man, dog), 130
 blood pressures (man), 322; (man, dog), 130
 vascular resistance (man), 325; (man, dog), 130
 pulse configuration, 157*
 right heart catheterization values, 334

AORTIC VALVE, 6*

ostial breadth at various ages, 20

AORTIC VALVULAR DISEASE (See AORTIC INSUFFICIENCY, AORTIC STENOSIS)

APICAL SYSTOLIC MURMUR, 150

ARGININE, translocation (plant), 314

ARM (See FOREARM, LIMB BLOOD)

ARRHYTHMIA(S)

cardiac index in, 78
 drugs for control of, 250, 255, 264, 269
 in endocrine disorders, 339
 during gravitational acceleration, 191

ARTERIAL BLOOD (See also specific organ or body area)

CO₂ content, 176, 177
 effect of shock, 317
 during hypercapnia, 176, 177

CO₂ pressure, 176, 177
 in cor pulmonale, 333
 effect of shock, 317
 during hypercapnia, 176, 177

flow (See also specific organ or body region)
 calculation formula, 355

during hypercapnia (mammals), 171

O₂ concentration, in cor pulmonale, 333

O₂ content, effect of shock, 317

O₂ pressure, in anemia (man and other animals), 351

O₂ saturation

in cardiac tamponade, 329

in cor pulmonale, 333

in diabetes, 337

effect of shock, 317

in pericarditis, 328

in pulmonary emphysema, 331, 332*

reduced: hemodynamic effects, 166, 167

pH (man), 176, 177; (vertebrates), 178, 179 (See also

specific organ or body region)

effect of shock (man), 317; (dog, rat), 321

during hypercapnia (man), 172, 176, 177; (dog),

173-175; (vertebrates), 178, 179

vascular resistance (See also specific organ or body

region)

in pathologic conditions, 325, 326

ARTERIAL BLOOD PRESSURE (man), 100, 101, 105, 176,

177, (birds), 108, 109; (fishes), 110; (inverte-

brates), 110; (mammals other than man), 106,

107; (reptiles, amphibians), 109; (vertebrates),

176, 179 (See also specific organ or body region)

vs age and sex; 102*-104*, 105

at altitude, 161, 162

in anemia (man), 330; (man and other animals), 349

in arterio-venous communication, 330

basal values, 100, 101

in beriberi, 330

in cardiac tamponade, 329

ARTERIAL BLOOD PRESSURE (concluded)

diastolic (See DIASTOLIC BLOOD PRESSURE)

effect of

albumin infusion, 324

drugs (man, dog), 252-255

exercise and recovery, 100, 101

explosive decompression (mammals), 186, 189

hypoxia, 161

ionizing radiation (vertebrates), 196-205, 228

reduced arterial O₂ saturation, 166

saline infusion, 324

shock, 317

during gravitational acceleration (mammals), 190, 192

during hypercapnia (man), 172, 176, 177; (dog), 184,

185; (mammals), 170, 171; (vertebrates), 178, 179

during hyperoxia (mammals), 168, 169

mean (See MEAN ARTERIAL PRESSURE)

during O₂ inhalation at 3-4 atmospheres, 186

in Paget's disease, 330

in pericarditis, 328

during pregnancy and postpartum, 158

during recompression (dog), 187

regression equations, 357, 358

syntolic (See SYSTOLIC BLOOD PRESSURE)

in thyrotoxicosis, 330

ARTERIAL BRACHIAL PULSE PRESSURE, regression

equation, 358

..... 55

..... : circula-

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

ASCITES, incidence in pericarditis, 328
ASPARAGINE, translocation (plants), 314, 315
ASPARTIC ACID, translocation (plant), 314
ASP VENOM, effect on circulatory function (rabbit), 242
ASSIMILATES, translocation (plants), 283-289, 290, 313
ASTHMA, circulation time during, 122
ATRIAL BLOOD PRESSURE (man), 112, 114; (man, dog), 147*, 148 (See also MEAN RIGHT ATRIAL PRESSURE)
 in anemia (man and other animals), 349
 during gravitational acceleration (mammals), 190
 during pregnancy, 158
ATRIAL FIBRILLATION, incidence in pericarditis, 329
ATRIAL MUSCLE, effect of drugs, 269-271
ATRIAL SEPTAL DEFECT (See SEPTAL DEFECTS)
ATRIOVENTRICULAR CONDUCTION SYSTEM, effect of drugs, 269-271
ATRIOVENTRICULAR JUNCTION, comparative anatomy (vertebrates), 10, 11
ATRIOVENTRICULAR NODE, 6*
ATRIOVENTRICULAR VALVE, development, 4, 5
ATRIUM (IA), 6* (See also LEFT ATRIUM; RIGHT ATRIUM)
 action potential of single fibers (vertebrates), 56-58
 atrio-ventricular weight ratio, during first year, 22*
 blood flow, during cardiac cycle (man, dog), 148
 capillary vascularity (cat), 37
 chemical composition (man), 38, (cattle), 45; (dog), 40, 41; (rabbit), 42
 comparative anatomy (vertebrates), 10, 11
 development, 4, 5
 elasticity (dog), 62, 63
 electrical properties (vertebrates), 56-58
 fiber diameter (vertebrates), 56, 57
 O₂ saturation, in congenital heart malformations, 152, 153
 pressure (See ATRIAL BLOOD PRESSURE; MEAN RIGHT ATRIAL PRESSURE)
 pulse waves, 113
 resting potential of single fibers (vertebrates), 56, 57
ATROPINE
 effect on heart (insects), 278
 translocation (plants), 316
AURICLE, AURICULAR (See ATRIUM, ATRIAL)
AUTOMATICITY, CARDIAC: effect of inorganic ions, 276
AUTOMATICITY, ECTOPIC: effect of drugs, 269-271
AUTOMATICITY, VENTRICULAR: effect of drugs, 269-271
AUTOPROTHROMBIN-1 (See FACTOR VII)
AUTOPROTHROMBIN-2 (See PLASMA THROMBOPLASTIN COMPONENT)
AUXIN, NATIVE: translocation (plants), 292, 299-301, 304-306
AXILLARY ARTERY, 7*
AXILLARY VEIN, 8*
AZURE A, effect on circulatory function (vertebrates), 244

BALISTOCARDIOGRAM
 amplitude
 normal standards, allowing for age, 360
 in relation to stroke volume. regression equations, 359
 determinations for
 cardiac index, 72, 78, 79
 cardiac output, 168, 176
 findings in anemia (man and other animals), 347

BALISTOCARDIOGRAM (concluded)
 normal standards
 for high-frequency table, 360
 for ultra-low frequency table, 362, 363
BASAL VALUES
 blood pressure, 100, 101
 cardiac index, 72
 cardiac output (mammals other than man), 80
 heart rate, 95, 99-101
BASILIC VEIN, MEDIAN. blood pressure, 111
BAZETT'S K (See ELECTROCARDIOGRAM, QT index)
BENZENE, effect on heart (insect), 282
BENZENE HEXACHLORIDE, effect on heart (insects), 278
BENZOIC ACID, translocation (plant), 296
BERBERINE SULFATE, translocation (plants), 290
BERIBERI, 78, 330
BHC, effect on heart (insects), 278
BICARBONATE (BLOOD)
 effect of shock (dog), 320
 in endocrine disorders (mammals), 338
BILIRUBIN (PLASMA), effect of shock, 318
BLOOD (See also CAPILLARY(IES); PLASMA; PLATELETS; RBC, SERUM; WHOLE BLOOD, specific organ or body region)
 coagulation (See also COAGULANTS, ANTICOAGULANTS)
 stages, 231*, 232*
 tests, 240
 concentration in tissues (vertebrates), 70, 71
 constituents (See specific constituents)
 effect of
 anemia (See ANEMIA)
 ionizing radiation (vertebrates), 196-205
 shock (man), 317, 318, (laboratory animals), 319-321
 in endocrine disorders (mammals), 336-339
 fractional extraction of diffusible substances by tissues (man, dog), 47
 during gravitational acceleration (mammals), 190, 195
 hematocrit values (See HEMATOCRIT)
 hemoglobin (See HEMOGLOBIN)
 O₂ saturation (See OXYGEN SATURATION)
 pH (See pH, specific organ or body region)
 pressure (See specific organ or body region)
BLOOD CELL(S) (See also RBC; WHITE BLOOD CELLS)
 count, in endocrine disorders (mammals), 337
 development (chick), 2; (frog), 1
BLOOD FLOW (See also specific organ or body region)
 in anemia (man and other animals), 350
 in aortic insufficiency, (man), 133, 325, (man, dog), 130
 atrial, during cardiac cycle (man, dog), 148
 -cardiac output ratio, in congenital heart malformations, 152, 153
 effect of
 albumin infusion, 326
 anticoagulants (vertebrates), 242-244
 coagulants (vertebrates), 241, 242
 norepinephrine, 326
 saline infusion, 326
 in endocrine disorders (mammals), 339
 during gravitational acceleration (mammals), 191-194
 during hypercapnia (mammals), 171
 methods of quantitating flow rates, 355
 in pathologic conditions, 325, 326, 339, 350
 during pregnancy and postpartum, 159
 in tissue (man, dog), 47
 ventricular, during cardiac cycle (man, dog), 148

BLOOD PRESSURE(S) (See **HYPERTENSION**, **HYPOTENSION**, specific pressure, specific organ or body region)

BLOOD VESSELS (See also specific vessel)

in anemia (man and other animals), 345

cardiovascular and pulmonary responses (mammals), 55

comparative anatomy (vertebrates), 12-15

development (man), 4, 5; (chick), 2; (frog), 1; (pig), 3

effect of

anticoagulants (vertebrates), 242-244

coagulants (vertebrates) 241, 242

ionizing radiation (vertebrates), 196-204

during hypoxia (mammals), 169

BLOOD VOLUME (crustacean, mollusk), 68, (insects), 69, 70, (mammals), 64-67; (vertebrates other than mammals), 67, 68 (See also specific organ or body region)

in anemia (man and other animals), 349

effect of

ionizing radiation (vertebrates), 229, 230

shock, 517

in endocrine disorders (mammals), 336

in pericarditis, 328

during pregnancy and parturition, 158

in pulmonary emphysema, 331

in tissue (vertebrates), 70, 71

BODY TEMPERATURE (See **TEMPERATURE**, **BODY**)

BONE, blood volume (mouse, rat), 70, 71

BONE MARROW

development, 4, 5

effect of ionizing radiation (vertebrates), 207-213

BORON, as heart constituent (rabbit), 42

BRACHIAL ARTERY

blood flow, during hypercapnia (dog), 171

blood pressure, during hypercapnia (dog), 171

causing hypercapnia, 182

blood volume (vertebrates), 70, 71

capillary filtration coefficient (vertebrates), 46

effect of

drugs, 258-261

gravitational acceleration (mammals), 191, 193

O₂ consumption in endocrine disorders, 335

resistance to arrested circulation (mammals), 135

white matter capillary density, surface, and diffusion distance, 34

BROEMER-RANKE METHOD, determination for cardiac output (rabbit), 80

BROMIDE IONS, cell surface permeability to (squid), 51; (plants), 51

BROMINE, as heart constituent (man), 38, (cattle), 45, (dog), 40, (rabbit, rat), 42, 43

2-BROMO-3,5-DICHLOROBENZOIC ACID, translocation (plant), 293

BRONCHIAL ARTERIES

blood pressures, 112

pulse waves, 113

BRONCHIAL ASTHMA, circulation time during, 322

CAFFEINE

effect on heart (insects), 278

translocation (plant), 314

CALCIFICATION, incidence in pericarditis, 329

CALCIUM, as heart constituent (man), 38, (cattle), 45; (dog), 40, (rodents), 42, 43

CALCIUM (BLOOD)

effect of shock (dog), 319

in endocrine disorders (mammals), 338

CALCIUM ION(S)

in blood coagulation stages, 231*, 232*

effect on

cardiac properties, 276

heart (man), 273, 274; (insects), 277

-potassium ion ratio required to maintain beat of isolated heart (insects), 276

CALP (LEC) (See also **LAMB BLOOD**)

blood flow, 132, 133

CAPILLARY(IES)

blood pressure, various tissues (vertebrates), 111

density, surface, and diffusion distance in tissue (vertebrates), 34

effect of

hypoxia, 163

ionizing radiation (vertebrates), 197-204

exchange of large molecules (mammals), 48

filtration

coefficients, various tissues (vertebrates), 46

formula (vertebrates), 46

fractional extraction of diffusible substances by tissue (man, dog), 47

permeability

in anemia (man and other animals), 349

coefficients, various substances (cat), 48

in endocrine disorders (mammals), 336

vascularity, muscle (vertebrates), 35-37

CARBOHYDRATES (See also specific carbohydrate)

translocation (plants), 283-288, 313

CARBON DIOXIDE

arterial blood. effect of shock on content (man), 317, (cat), 320

effect of exposure to high concentration (See **HYPERCAPNIA**)

effect on heart (insects), 282

retention with cor pulmonale: effect on cerebral function, 333

CARBON DIOXIDE COMBINING POWER (PLASMA)

in diabetes (man, dog), 339

3,3'-CARBOXYMETHYLENE-BIS-4-HYDROXY-COUMARIN

effect on clotting time (vertebrates), 237

therapeutic effect (man), 235, (vertebrates) 247

CARDIAC

CARDIAC CYCLE (concluded)
 correlation of events during (man, dog), 147*, 148
 normal and pathologic heart sounds in, 155*

CARDIAC DISEASE (See also specific disease)
 circulation time during, 122, 123

CARDIAC FAILURE (See also HEART FAILURE, CONGESTIVE)
 circulation time during, 122, 123

CARDIAC FIBERS (See also specific fiber)
 action potential (vertebrates), 56-58
 diameter (vertebrates), 56, 57
 resting potential (vertebrates), 56, 57

CARDIAC FORCE, MAXIMUM. effect of hypoxia, 167

CARDIAC FUNCTION. in anemia (man and other animals), 349

CARDIAC GLYCOSIDES (See GLYCOSIDES, CARDIAC)

CARDIAC IMPULSE. in anemia (man and other animals), 345

CARDIAC INDEX (See also CARDIAC OUTPUT)
 in acute hemorrhage, 326
 vs age, 73
 in anoxia, 78
 in chest injury, 326
 in disease, 78, 79, 326-334
 effect of
 albumin infusion, 326
 hypoxia, 167
 norepinephrine, 326
 reduced arterial O₂ saturation, 167
 saline infusion, 326
 shock, 317
 during hyperoxia, 79
 methods
 acetylene, 79
 ballistocardiogram, 78, 79
 dye, 73, 74
 dye (Hamilton), 72, 78, 79
 ethyl iodide, 72, 78
 Fick, 72-74, 78, 79, 328, 329
 vs O₂ uptake, during exercise, 74, 75*
 during pregnancy, 79, 158

CARDIAC INFARCTION. effect on circulation time, 123

CARDIAC MUSCLE (See also HEART MUSCLE)
 capillary vascularity (vertebrates), 37

CARDIAC OUTPUT (See also CARDIAC INDEX)
 vs age, 72
 basal values (man), 77, (mammals other than man), 80
 -blood flow ratio, in congenital heart malformation, 152, 153
 vs body temperature (man, dog), 77
 in disease, 78, 79, 330, 334, 336, 349
 effect of
 drugs (man, dog), 252-255
 exercise (man), 77; (mammals other than man), 80
 posture, 77, 79
 during gravitational acceleration (mammals), 191, 193
 during hypercapnia (man), 170, 176; (vertebrates), 178
 during hyperoxia, 168
 methods
 acetylene, 77, 79, 158, 168, 176
 ballistocardiogram, 168, 176
 Broemser-Ranke, 80
 dye-dilution, 72, 158, 176
 dye (Hamilton), 78, 79
 ethyl iodide, 77
 Fick, 72, 77-80, 158
 rotameter, 80
 Stewart's sodium chloride, 80

CARDIAC OUTPUT (concluded)
 vs O₂ consumption, basal and during exercise (man), 77; (mammals), 80
 during postpartum, 158
 during pregnancy, 79, 158

CARDIAC OXYGEN CONSUMPTION. in anemia (man), 330; (man and other animals), 351

CARDIAC OXYGEN UPTAKE. effect of drugs (man, dog), 257

CARDIAC PROPERTIES. effect of
 drugs, 269-271
 inorganic ions, 276

CARDIAC PULSATIONS. in pericarditis, 329

CARDIAC SINUS. electrical properties (dog), 56, 58

CARDIAC STROKE VOLUME (See STROKE VOLUME, CARDIAC)

CARDIAC TAMPONADE. ACUTE, 329

CARDIAC TISSUES (See also specific tissue)
 electrical properties (vertebrates), 56-59

CARDIAC VEIN. 6*, 8*

CARDINAL VEINS. comparative anatomy (vertebrates), 12, 13

CARDIO-ACCELERATOR CENTER. drug stimulants and depressants, 246

CARDIO-INHIBITORY CENTER. drug stimulants and depressants, 246

CARDIOTONIC DRUGS (See GLYCOSIDES, CARDIAC)

CARDIOVASCULAR DISEASE (See also specific disease)
 effect on circulation time, 123

CARDIOVASCULAR RESPONSES (mammals), 55

CAROTID ARTERY(IES). 6*, 7* (See also specific artery)
 blood pressures, 112
 comparative anatomy (vertebrates), 12
 pulse waves, 113

CAROTID SINUSES. stretch receptors: drug stimulants and depressants, 246

CASTRATION. effect on cerebral blood flow, 335

CATHETERIZATION (See CARDIAC CATHETERIZATION)

CATIONS. cardiovascular action, 252

CATIONS (BLOOD). effect of shock (dog), 320

CAUDAL VEIN. comparative anatomy (vertebrates), 12

CELIAC ARTERY. 7*
 comparative anatomy (vertebrates), 14
 dimensions, 21

CELLS (See specific cells)

CELL SURFACE. permeability to
 ions (animals), 50, 51; (plants), 51
 organic substances (animals), 52; (plants), 53, 54
 water (animals, plants), 49

CELLULOSE SULFURIC ACID ESTER. effect on
 circulatory function (rabbit), 243

CEPHALIN. effect on circulatory function (dog, frog), 242

CEREBELLUM TISSUE. capillary density, surface,
 and diffusion distance (mouse), 34

CEREBRAL ARTERIOSCLEROSIS. 125

CEREBRAL BLOOD
 arterial CO₂ pressure, 182, 183
 arterial pH, 182, 183
 arterio-venous O₂ difference, 182, 183
 flow (man), 182, (man, monkey), 125
 in anemia (man and other animals), 350
 in cor pulmonale, 333
 effect of drugs, 258-261
 in endocrine disorders, 335
 during hyperoxia (mammals), 169
 during pregnancy, 159

CEREBRAL BLOOD [concluded]
during gravitational acceleration, 191, 193
during hypercapnia, 182, 183
mean arterial blood pressure, 182, 183
O₂ consumption (man), 335, (man, monkey), 125
in cor pulmonale, 333
in endocrine disorders, 335
O₂ uptake, effect of drugs, 258-261
vascular resistance (man), 182, (man, monkey), 125
in anemia (man and other animals), 350
in cor pulmonale, 333
effect of drugs, 258-261

CEREBRAL CORTEX TISSUE
capillary density, surface, and diffusion distance
(man, mouse), 34
resistance to arrested circulation (mammals), 135

CEREBRAL VASCULAR DISEASE and hypercapnia
effect on cerebral blood, 182

CERVICAL LYMPHATICS, lymph flow and lymphocyte
content (mammals), 136

CERVICAL VEINS, incidence of distention in pericarditis, 328

CHEST INJURY, effect on
blood flow, 326
blood pressures, 323
vascular resistance, 326

CHLORAZOL FAST PINK, effect on circulatory function
(vertebrates), 243

CHLORDANE, effect on heart (insect), 27b

CHLORIDE IONS, cell permeability to (frog), 50;
(plant), 51

CHRISTMAS FACTOR (See PLASMA THROMBO-
PLASTIN COMPONENT)

CIRCULATION, ARRESTED resistance of tissue to
(mammary). 135

CIRCULATORY SYSTEM, development (man), 4, 5;
(chick), 2; (frog), 1; (pig), 3

CISTERNA CHYLII, 9*
comparative anatomy (vertebrates), 17
CISTERNA CHYLII REGION, development of lymph
vessels, 4, 5

CLOTTING TIME
effect of
anticoagulants (man), 241; (vertebrates), 236-238
heparin, 241
tests, 240

COAGULANTS (See also specific coagulant)
in coagulation stages, 231^a, 232^a
effect on circulatory function (vertebrates), 241, 242
natural-occurring, preparation and assay, 233, 234
vitamin K, 239

COBRA VENOM, effect on circulatory function (rabbit).
242

COLIC VEINS. *see* COLIC VEINS.

[illegible]

CONDUCTIVITY, heart muscle. effect of inorganic ions. 276

CONSTRUCTIVE PERICARDITIS (See PERICARDITIS, CONSTRUCTIVE)
CONTINUOUS MURMURS
 in congenital heart malformations, 152, 153
 phonocardiographic configurations, 154*
CONTRACTILITY, heart muscle
 effect of
 drugs, 269-271
 inorganic ions, 276
CONUS ARTERIOSUS, comparative anatomy (vertebrates), 10, 11
COPPER, as heart constituent (man), 38; (artio-, perissodactyla), 45; (cat), 40; (rodents), 42, 43
COPPER (BLOOD), in endocrine disorders, 338
CORONARY ARTERIAL DISEASE, effect on cardiac index, 78
CORONARY ARTERIES, 6*
CORONARY BLOOD
 arterial CO₂ content (dog), 180, 181
 arterial pH (cat, dog), 180, 181
 arterio-venous O₂ difference, in anemia (man and other animals), 350
 flow (cat, dog), 180
 in anemia (man and other animals), 350
 effect of drugs (mammals), 257
 during hypercapnia (cat, dog), 180, 181
 vascular resistance (dog), 180, 181
 in anemia (man and other animals), 350
 effect of drugs (mammals), 257
CORONARY OCCLUSION, effect on digital blood flow and pressure, 134
CORONARY SCLEROSIS, in diabetes, 339
CORONARY THROMBOSIS (See MYOCARDIAL INFARCTION)
CORONARY VEIN, 8*
COR PULMONALE, 333
 blood flow, 130, 325
 blood pressures, 130, 322, 333
 cardiac index, 325, 333
 with CO₂ retention effect on cerebral blood, 333
 effect on renal function, 334
 and emphysema, 331
 and hypoxia, 130
 vascular resistance, 130, 325
CORRIGAN PULSE, configuration, 157*
COTHROMBOPLASTIN (See FACTOR VII)
COUGH, incidence in pericarditis, 328
CREATINE, as heart constituent (man), 39; (birds), 46; (cat, dog), 40, 41; (cattle, sheep), 45; (rabbit), 42, 43
CREATINE (BLOOD), in myxedema, 337
CREATINE NITROGEN (BLOOD), effect of shock (cat, rat), 319
CREATININE (BLOOD)
 in Addison's disease (cat), 337
 effect of shock (dog), 320
 fractional extraction by tissue (dog), 47
CREATININE (PLASMA), effect of shock, 318
CUMOPYRAN, series I and II: therapeutic effect, 235
CURARE, effect on heart (insects), 278
CUSHING'S SYNDROME, effect on digital blood flow and pressure, 134
CUTANEOUS BLOOD, in anemia (man and other animals), 350
CUTANEOUS VEIN, comparative anatomy (vertebrates), 13
CYANIDE, effect on heart (insects), 282
CYANOL, translocation (plants), 290
CYANOSIS, incidence in pericarditis, 329
CYSTIC ARTERIES, dimensions, 21

2, 4-D, translocation (plants), 292-296, 298-307
 3, 4-D, translocation (plants), 303
DDT, effect on heart (insects), 278, 279
DEAD SPACE, PHYSIOLOGIC, in anemia (man and other animals), 351
DECOMPRESSION, EXPLOSIVE
 effect on
 arterial blood pressures (dog), 189; (mammals), 188
 heart rate (mammals), 188
DEXTRAN
 cardiovascular action, 252
 effect on pulmonary blood, 265
 transcapillary exchange (dog), 48
DEXTRAN SULFATE
 clotting time (rabbit), 236
 duration and side effects, 240
DIABETES, (mammals), 336-340
DIABETES MELLITUS, effect on cerebral blood flow, 335
 cerebral O₂ consumption, 335
 digital blood flow, 134
DIACETIN, cell surface permeability to (animals), 52; (plant), 53
DIAPHRAGM
 capillary vascularity (guinea pig), 35
 cell surface permeability to ions (rat), 50
DIASTOLIC BLOOD PRESSURE (man), 105, 176, 177; (birds), 108, 109; (fishes), 110; (mammals other than man), 106, 107; (reptiles, amphibians), 109
 vs age and sex, 102*, 104*, 105
 at altitude, 161, 162
 atrial, in newborn, 114
 basal values, 100, 101
 brachial: regression equation, 158
 effect of
 exercise and recovery, 100, 101
 hypoxia, 161, 162, 166
 norepinephrine, 130
 reduced arterial O₂ saturation, 166
 shock, 317
 femoral, in newborn, 114
 during hypercapnia, 170-172, 176, 177
 during hyperoxia, 168
 in pathologic conditions, 102*, 103*, 134, 322, 323, 328-330, 349
 pulmonary, 112, 114
 effect of exercise, 130
 effect of hypoxia (man), 130; (man, dog), 164
 effect of norepinephrine, 130
 effect of posture, 130
 in newborn, 114
 in pathologic conditions, 130, 322, 323, 328-331, 333
 radial: regression equation, 358
 right ventricular, 112, 114
 in pathologic conditions, 322, 323, 328, 329, 333
 ventricular, 114
 various chambers and vessels, 112
DIASTOLIC INTERVAL, effect of inorganic ions, 273
DIASTOLIC MURMURS
 in heart malformations, 152, 153
 in heart failure, 154
 in heart disease, 155
 in heart failure, 156
 in heart failure, 157
 in heart failure, 158
 in heart failure, 159
 in heart failure, 160
 in heart failure, 161
 in heart failure, 162
 in heart failure, 163
 in heart failure, 164
 in heart failure, 165
 in heart failure, 166
 in heart failure, 167
 in heart failure, 168
 in heart failure, 169
 in heart failure, 170
 in heart failure, 171
 in heart failure, 172
 in heart failure, 173
 in heart failure, 174
 in heart failure, 175
 in heart failure, 176
 in heart failure, 177
 in heart failure, 178
 in heart failure, 179
 in heart failure, 180
 in heart failure, 181
 in heart failure, 182
 in heart failure, 183
 in heart failure, 184
 in heart failure, 185
 in heart failure, 186
 in heart failure, 187
 in heart failure, 188
 in heart failure, 189
 in heart failure, 190
 in heart failure, 191
 in heart failure, 192
 in heart failure, 193
 in heart failure, 194
 in heart failure, 195
 in heart failure, 196
 in heart failure, 197
 in heart failure, 198
 in heart failure, 199
 in heart failure, 200
 in heart failure, 201
 in heart failure, 202
 in heart failure, 203
 in heart failure, 204
 in heart failure, 205
 in heart failure, 206
 in heart failure, 207
 in heart failure, 208
 in heart failure, 209
 in heart failure, 210
 in heart failure, 211
 in heart failure, 212
 in heart failure, 213
 in heart failure, 214
 in heart failure, 215
 in heart failure, 216
 in heart failure, 217
 in heart failure, 218
 in heart failure, 219
 in heart failure, 220
 in heart failure, 221
 in heart failure, 222
 in heart failure, 223
 in heart failure, 224
 in heart failure, 225
 in heart failure, 226
 in heart failure, 227
 in heart failure, 228
 in heart failure, 229
 in heart failure, 230
 in heart failure, 231
 in heart failure, 232
 in heart failure, 233
 in heart failure, 234
 in heart failure, 235
 in heart failure, 236
 in heart failure, 237
 in heart failure, 238
 in heart failure, 239
 in heart failure, 240
 in heart failure, 241
 in heart failure, 242
 in heart failure, 243
 in heart failure, 244
 in heart failure, 245
 in heart failure, 246
 in heart failure, 247
 in heart failure, 248
 in heart failure, 249
 in heart failure, 250
 in heart failure, 251
 in heart failure, 252
 in heart failure, 253
 in heart failure, 254
 in heart failure, 255
 in heart failure, 256
 in heart failure, 257
 in heart failure, 258
 in heart failure, 259
 in heart failure, 260
 in heart failure, 261
 in heart failure, 262
 in heart failure, 263
 in heart failure, 264
 in heart failure, 265
 in heart failure, 266
 in heart failure, 267
 in heart failure, 268
 in heart failure, 269
 in heart failure, 270
 in heart failure, 271
 in heart failure, 272
 in heart failure, 273
 in heart failure, 274
 in heart failure, 275
 in heart failure, 276
 in heart failure, 277
 in heart failure, 278
 in heart failure, 279
 in heart failure, 280
 in heart failure, 281
 in heart failure, 282
 in heart failure, 283
 in heart failure, 284
 in heart failure, 285
 in heart failure, 286
 in heart failure, 287
 in heart failure, 288
 in heart failure, 289
 in heart failure, 290
 in heart failure, 291
 in heart failure, 292
 in heart failure, 293
 in heart failure, 294
 in heart failure, 295
 in heart failure, 296
 in heart failure, 297
 in heart failure, 298
 in heart failure, 299
 in heart failure, 300
 in heart failure, 301
 in heart failure, 302
 in heart failure, 303
 in heart failure, 304
 in heart failure, 305
 in heart failure, 306
 in heart failure, 307
 in heart failure, 308
 in heart failure, 309
 in heart failure, 310
 in heart failure, 311
 in heart failure, 312
 in heart failure, 313
 in heart failure, 314
 in heart failure, 315
 in heart failure, 316
 in heart failure, 317
 in heart failure, 318
 in heart failure, 319
 in heart failure, 320
 in heart failure, 321
 in heart failure, 322
 in heart failure, 323
 in heart failure, 324
 in heart failure, 325
 in heart failure, 326
 in heart failure, 327
 in heart failure, 328
 in heart failure, 329
 in heart failure, 330
 in heart failure, 331
 in heart failure, 332
 in heart failure, 333
 in heart failure, 334
 in heart failure, 335
 in heart failure, 336
 in heart failure, 337
 in heart failure, 338
 in heart failure, 339
 in heart failure, 340
 in heart failure, 341
 in heart failure, 342
 in heart failure, 343
 in heart failure, 344
 in heart failure, 345
 in heart failure, 346
 in heart failure, 347
 in heart failure, 348
 in heart failure, 349
 in heart failure, 350
 in heart failure, 351
 in heart failure, 352
 in heart failure, 353
 in heart failure, 354
 in heart failure, 355
 in heart failure, 356
 in heart failure, 357
 in heart failure, 358
 in heart failure, 359
 in heart failure, 360
 in heart failure, 361
 in heart failure, 362
 in heart failure, 363
 in heart failure, 364
 in heart failure, 365
 in heart failure, 366
 in heart failure, 367
 in heart failure, 368
 in heart failure, 369
 in heart failure, 370
 in heart failure, 371
 in heart failure, 372
 in heart failure, 373
 in heart failure, 374
 in heart failure, 375
 in heart failure, 376
 in heart failure, 377
 in heart failure, 378
 in heart failure, 379
 in heart failure, 380
 in heart failure, 381
 in heart failure, 382
 in heart failure, 383
 in heart failure, 384
 in heart failure, 385
 in heart failure, 386
 in heart failure, 387
 in heart failure, 388
 in heart failure, 389
 in heart failure, 390
 in heart failure, 391
 in heart failure, 392
 in heart failure, 393
 in heart failure, 394
 in heart failure, 395
 in heart failure, 396
 in heart failure, 397
 in heart failure, 398
 in heart failure, 399
 in heart failure, 400
 in heart failure, 401
 in heart failure, 402
 in heart failure, 403
 in heart failure, 404
 in heart failure, 405
 in heart failure, 406
 in heart failure, 407
 in heart failure, 408
 in heart failure, 409
 in heart failure, 410
 in heart failure, 411
 in heart failure, 412
 in heart failure, 413
 in heart failure, 414
 in heart failure, 415
 in heart failure, 416
 in heart failure, 417
 in heart failure, 418
 in heart failure, 419
 in heart failure, 420
 in heart failure, 421
 in heart failure, 422
 in heart failure, 423
 in heart failure, 424
 in heart failure, 425
 in heart failure, 426
 in heart failure, 427
 in heart failure, 428
 in heart failure, 429
 in heart failure, 430
 in heart failure, 431
 in heart failure, 432
 in heart failure, 433
 in heart failure, 434
 in heart failure, 435
 in heart failure, 436
 in heart failure, 437
 in heart failure, 438
 in heart failure, 439
 in heart failure, 440
 in heart failure, 441
 in heart failure, 442
 in heart failure, 443
 in heart failure, 444
 in heart failure, 445
 in heart failure, 446
 in heart failure, 447
 in heart failure, 448
 in heart failure, 449
 in heart failure, 450
 in heart failure, 451
 in heart failure, 452
 in heart failure, 453
 in heart failure, 454
 in heart failure, 455
 in heart failure, 456
 in heart failure, 457
 in heart failure, 458
 in heart failure, 459
 in heart failure, 460
 in heart failure, 461
 in heart failure, 462
 in heart failure, 463
 in heart failure, 464
 in heart failure, 465
 in heart failure, 466
 in heart failure, 467
 in heart failure, 468
 in heart failure, 469
 in heart failure, 470
 in heart failure, 471
 in heart failure, 472
 in heart failure, 473
 in heart failure, 474
 in heart failure, 475
 in heart failure, 476
 in heart failure, 477
 in heart failure, 478
 in heart failure, 479
 in heart failure, 480
 in heart failure, 481
 in heart failure, 482
 in heart failure, 483
 in heart failure, 484
 in heart failure, 485
 in heart failure, 486
 in heart failure, 487
 in heart failure, 488
 in heart failure, 489
 in heart failure, 490
 in heart failure, 491
 in heart failure, 492
 in heart failure, 493
 in heart failure, 494
 in heart failure, 495
 in heart failure, 496
 in heart failure, 497
 in heart failure, 498
 in heart failure, 499
 in heart failure, 500
 in heart failure, 501
 in heart failure, 502
 in heart failure, 503
 in heart failure, 504
 in heart failure, 505
 in heart failure, 506
 in heart failure, 507
 in heart failure, 508
 in heart failure, 509
 in heart failure, 510
 in heart failure, 511
 in heart failure, 512
 in heart failure, 513
 in heart failure, 514
 in heart failure, 515
 in heart failure, 516
 in heart failure, 517
 in heart failure, 518
 in heart failure, 519
 in heart failure, 520
 in heart failure, 521
 in heart failure, 522
 in heart failure, 523
 in heart failure, 524
 in heart failure, 525
 in heart failure, 526
 in heart failure, 527
 in heart failure, 528
 in heart failure, 529
 in heart failure, 530
 in heart failure, 531
 in heart failure, 532
 in heart failure, 533
 in heart failure, 534
 in heart failure, 535
 in heart failure, 536
 in heart failure, 537
 in heart failure, 538
 in heart failure, 539
 in heart failure, 540
 in heart failure, 541
 in heart failure, 542
 in heart failure, 543
 in heart failure, 544
 in heart failure, 545
 in heart failure, 546
 in heart failure, 547
 in heart failure, 548
 in heart failure, 549
 in heart failure, 550
 in heart failure, 551
 in heart failure, 552
 in heart failure, 553
 in heart failure, 554
 in heart failure, 555
 in heart failure, 556
 in heart failure, 557
 in heart failure, 558
 in heart failure, 559
 in heart failure, 560
 in heart failure, 561
 in heart failure, 562
 in heart failure, 563
 in heart failure, 564
 in heart failure, 565
 in heart failure, 566
 in heart failure, 567
 in heart failure, 568
 in heart failure, 569
 in heart failure, 570
 in heart failure, 571
 in heart failure, 572
 in heart failure, 573
 in heart failure, 574
 in heart failure, 575
 in heart failure, 576
 in heart failure, 577
 in heart failure, 578
 in heart failure, 579
 in heart failure, 580
 in heart failure, 581
 in heart failure, 582
 in heart failure, 583
 in heart failure, 584
 in heart failure, 585
 in heart failure, 586
 in heart failure, 587
 in heart failure, 588
 in heart failure, 589
 in heart failure, 590
 in heart failure, 591
 in heart failure, 592
 in heart failure, 593
 in heart failure, 594
 in heart failure, 595
 in heart failure, 596
 in heart failure, 597
 in heart failure, 598
 in heart failure, 599
 in heart failure, 600
 in heart failure, 601
 in heart failure, 602
 in heart failure, 603
 in heart failure, 604
 in heart failure, 605
 in heart failure, 606
 in heart failure, 607
 in heart failure, 608
 in heart failure, 609
 in heart failure, 610
 in heart failure, 611
 in heart failure, 612
 in heart failure, 613
 in heart failure, 614
 in heart failure, 615
 in heart failure, 616
 in heart failure, 617
 in heart failure, 618
 in heart failure, 619
 in heart failure, 620
 in heart failure, 621
 in heart failure, 622
 in heart failure, 623
 in heart failure, 624
 in heart failure, 625
 in heart failure, 626
 in heart failure, 627
 in heart failure, 628
 in heart failure, 629
 in heart failure, 630
 in heart failure, 631
 in heart failure, 632
 in heart failure, 633
 in heart failure, 634
 in heart failure, 635
 in heart failure, 636
 in heart failure, 637
 in heart failure, 638
 in heart failure, 639
 in heart failure, 640
 in heart failure, 641
 in heart failure, 642
 in heart failure, 643
 in heart failure, 644
 in heart failure, 645
 in heart failure, 646
 in heart failure, 647
 in heart failure, 648
 in heart failure, 649
 in heart failure, 650
 in heart failure, 651
 in heart failure, 652
 in heart failure, 653
 in heart failure, 654
 in heart failure, 655
 in heart failure, 656
 in heart failure, 657
 in heart failure, 658
 in heart failure, 659
 in heart failure, 660
 in heart failure, 661
 in heart failure, 662
 in heart failure, 663
 in heart failure, 664
 in heart failure, 665
 in heart failure, 666
 in heart failure, 667
 in heart failure, 668
 in heart failure, 669
 in heart failure, 670
 in heart failure, 671
 in heart failure, 672
 in heart failure, 673
 in heart failure, 674
 in heart failure, 675
 in heart failure, 676
 in heart failure, 677
 in heart failure, 678
 in heart failure, 679
 in heart failure, 680
 in heart failure, 681
 in heart failure, 682
 in heart failure, 683
 in heart failure, 684
 in heart failure, 685
 in heart failure, 686
 in heart failure, 687
 in heart failure, 688
 in heart failure, 689
 in heart failure, 690
 in heart failure, 691
 in heart failure, 692
 in heart failure, 693
 in heart failure, 694
 in heart failure, 695
 in heart failure, 696
 in heart failure, 697
 in heart failure, 698
 in heart failure, 699
 in heart failure, 700
 in heart failure, 701
 in heart failure, 702
 in heart failure, 703
 in heart failure, 704
 in heart failure, 705
 in heart failure, 706
 in heart failure, 707
 in heart failure, 708
 in heart failure, 709
 in heart failure, 710
 in heart failure, 711
 in heart failure, 712
 in heart failure, 713
 in heart failure, 714
 in heart failure, 715
 in heart failure, 716
 in heart failure, 717
 in heart failure, 718
 in heart failure, 719
 in heart failure, 720
 in heart failure, 721
 in heart failure, 722
 in heart failure, 723
 in heart failure, 724
 in heart failure, 725
 in heart failure, 726
 in heart failure, 727
 in heart failure, 728
 in heart failure, 729
 in heart failure, 730
 in heart failure, 731
 in heart failure, 732
 in heart failure, 733
 in heart failure, 734
 in heart failure, 735
 in heart failure, 736
 in heart failure, 737
 in heart failure, 738
 in heart failure, 739
 in heart failure, 740
 in heart failure, 741
 in heart failure, 742
 in heart failure, 743
 in heart failure, 744
 in heart failure, 745
 in heart failure, 746
 in heart failure, 747
 in heart failure, 748
 in heart failure, 749
 in heart failure, 750
 in heart failure, 751
 in heart failure, 752
 in heart failure, 753
 in heart failure, 754
 in heart failure, 755
 in heart failure, 756
 in heart failure, 757
 in heart failure, 758
 in heart failure, 759
 in heart failure, 760
 in heart failure, 761
 in heart failure, 762
 in heart failure, 763
 in heart failure, 764
 in heart failure, 765
 in heart failure, 766
 in heart failure, 767
 in heart failure, 768
 in heart failure, 769
 in heart failure, 770
 in heart failure, 771
 in heart failure, 772
 in heart failure, 773
 in heart failure, 774
 in heart failure, 775
 in heart failure, 776
 in heart failure, 777
 in heart failure, 778
 in heart failure, 779
 in heart failure, 780
 in heart failure, 781
 in heart failure, 782
 in heart failure, 783
 in heart failure, 784
 in heart failure, 785
 in heart failure, 786
 in heart failure, 787
 in heart failure, 788
 in heart failure, 789
 in heart failure, 790
 in heart failure, 791
 in heart failure, 792
 in heart failure, 793
 in heart failure, 794
 in heart failure, 795
 in heart failure, 796
 in heart failure, 797
 in heart failure, 798
 in heart failure, 799
 in heart failure, 800
 in heart failure, 801
 in heart failure, 802
 in heart failure, 803
 in heart failure, 804
 in heart failure, 805
 in heart failure, 806
 in heart failure, 807
 in heart failure, 808
 in heart failure, 809
 in heart failure, 810
 in heart failure, 811
 in heart failure, 812
 in heart failure, 813
 in heart failure, 814
 in heart failure, 815
 in heart failure, 816
 in heart failure, 817
 in heart failure, 818
 in heart failure, 819
 in heart failure, 820
 in heart failure, 821
 in heart failure, 822
 in heart failure, 823
 in heart failure, 824
 in heart failure, 825
 in heart failure, 826
 in heart failure, 827
 in heart failure, 828
 in heart failure, 829
 in heart failure, 830
 in heart failure, 831
 in heart failure, 832
 in heart failure, 833
 in heart failure, 834
 in heart failure, 835
 in heart failure, 836
 in heart failure, 837
 in heart failure, 838
 in heart failure, 839
 in heart failure, 840
 in heart failure, 841
 in heart failure, 842
 in heart failure, 843
 in heart failure, 844
 in heart failure, 845
 in heart failure, 846
 in heart failure, 847
 in heart failure, 848
 in heart failure, 849
 in heart failure, 850
 in heart failure, 851
 in heart failure, 852
 in heart failure, 853
 in heart failure, 854
 in heart failure, 855
 in heart failure, 856
 in heart failure, 857
 in heart failure, 858
 in heart failure, 859
 in heart failure, 860
 in heart failure, 861
 in heart failure, 862
 in heart failure, 863
 in heart failure,

(plant), 303
DICROTIC PULSE, configuration, 157*
DICUMAROL
 in coagulation stages, 231*, 232*
 effect on
 clotting time (vertebrates), 237
 circulatory function (mammals), 244
 therapeutic effects (man), 235; (vertebrates), 237
DICUMAROL-INDUCED HYPOFIBRINOGENEMIA,
 response to vitamin K, 239
DIELDRIEN, effect on heart (insect), 279
DIETHYLENE GLYCOL, cell surface permeability to
 (animals), 52
DIFFUSION AREA, various substances in muscle
 tissue (cat), 48
DIFFUSION, CAPILLARY various substances (man,
 dog), 47
DIFFUSION DISTANCE, CAPILLARY (vertebrates), 34
DIGITAL BLOOD, 132-134 (See also **LDIB BLOOD**)
DIGITALIS (See also **GLYCOSIDES, CARDIAC**)
 effect on heart (insects), 279
 general effects, 249
 hemodynamic effects (man, dog), 254, 259, 264, 270
 1,2-DIHYDROXYRIBIDINE-3,6-DIONE, translocation
 (plants), 293, 299-301, 305, 306
 4,6-DINITRO-o-CRESOL, effect on heart (insect), 279
DINITRO-o-CYCLOHEXYLPHENOL, effect on heart
 (insect), 279
DINONYLPHTHALATE, effect on heart (insect), 279
DIPAXIN
 effect on clotting time (vertebrates), 238
 therapeutic effects (man), 235, (vertebrates), 238
 2,3-DIPHOSPHOGLYCERIC ACID (PLASMA), in
 diabetes, 339
DIPHOTHERIA TOXIN, effect on heart (insect), 279
DISEASE (See also specific disease)
 cardiac index in, 78, 79, 328-334
 cardiac output in (man), 78, 79, 330, 334, (man and
 other animals), 349, (mammals), 336
 circulation time in, 121-125
 effect on
 digital blood flow, 134
 hepatic blood flow (mammals), 128, 129
 limb blood flow, 133
 pulmonary circulation, 130
 renal function, 334
 heart (See **HEART DISEASE, specific disease**)
 hemodynamic effects of various, 130, 322, 323, 330,
 349-351
DISEASE, RHEUMATIC HEART effect on renal
 function, 334
DIURETICS
 cardiovascular action, 251
 hemodynamic effects of mercurial sodium, 255
DIAC, effect on heart (insect), 279
DNOCHF, effect on heart (insect), 279
DNP effect on heart (insect), 279
D 111
DORNA
DOER

DRUGS

anticoagulants (See **ANTICOAGULANTS**)
 with cardiovascular effects: sites of action, 246
 coagulants (See **COAGULANTS**)
 for disorders of the circulatory system, 247-250
 for disorders not limited to the circulatory system,
 250-252
 effect on
 cardiac properties, 269-271
 coronary circulation (mammals), 257, 258
 heart (insects), 278-282
 hepatic circulation (man, dog), 261-263
 limb circulation (man, dog), 266-268
 pulmonary circulation (man), 130, (man, dog), 263-266
 renal circulation (man, dog), 261-263
 systemic circulation (man, dog), 252-255
DUCTS (See specific duct)
DUCTUS ARTERIOSUS, development, 4, 5
DUODENUM
 capillary density, surface, and diffusion distance
 (mouse), 34
 effect of ionizing radiation (chicken), 230
DYE METHOD (See also **HAMILTON DYE METHOD**)
 determinations for
 cardiac index, 72-74
 cardiac output, 72, 158
DYES, translocation (plants), 290, 291
DYSPEA, incidence in pericarditis, 328
EAR
 effect of hypoxia on circulation, 163
 pulse effect of gravitational acceleration, 191
 129
muscles
 effect of drugs, 269-271
 heart drug stimulants and depressants, 246
EDC, effect on clotting time (man, rabbit), 238
EDEMA
 incidence in pericarditis, 328
 peripheral, in endocrine disorders, 339
EEL SERUM, effect on circulatory function (rabbit),
 242
EGG, cell surface permeability to
 inorganic ions (invertebrates), 50
 organic substances (invertebrates), 52
 water (animals), 49
EISENMENGER'S COMPLEX (Syndrome)
 effect on pulmonary circulation, 130
 murmurs in, 152
ELASTICITY, arterio-venous system (man, dog), 60-63
ELASTIC RESISTANCE COEFFICIENT. PULMONARY
 STANCE

 in anemia (man and other animals), 347
 correlation with heart sounds, 155*
 effect of
 anticoagulants (vertebrates), 242, 244
 coagulants (rabbit), 242
 gravitational acceleration (mammals), 191, 193
 hypoxia, 164-166
 inorganic ions (man, dog), 273, 274
 ionizing radiation (vertebrates), 197-200, 203
 pH (dog), 59
 reduced arterial O₂ saturation, 166

ELECTROCARDIOGRAM (concluded)
 in endocrine disorders, 336
 and events in cardiac cycle (man, dog), 147*
 findings, in congenital heart malformations, 152, 153
 during hypercapnia (dog), 173-175
 incidence of low voltage: in pericarditis, 329
 intervals (See also specific interval)
 at altitude, 164
 duration, 140*; (vertebrates), 141
 effect of hypoxia, 164
 effect of inorganic ions (man, dog), 273, 274
 effect of pH (dog), 59
 intracardiac pulse, 156*
 leads, 137*-139* (See also LEADS ELECTRO-
 CARDIOGRAPHIC; specific lead)
 amplitudes of waves, in various (man), 142-145;
 (farm animals), 146
 R wave as percentage of RS deflection, in various,
 145
 QRS complex (See QRS (ELECTROCARDIOGRAM))
 QT index (QT_c; Bazett's K), (vertebrates), 141
 definition, 140
 effect of pH (dog), 59
 ST segment (See ST SEGMENT (ELECTRO-
 CARDIOGRAM))
 waves (See also specific wave)
 amplitudes in various leads (man), 142-145; (farm
 animals), 146
 duration, 140*; (vertebrates), 141
 effect of altitude, 165
 effect of hypoxia, 165
 effect of inorganic ions (man, dog), 273, 274
 effect of pH (dog), 59
ELECTROLYTES (See ION(S), INORGANIC; specific
 electrolyte)
ELECTROLYTES (BLOOD), in endocrine disorders
 (mammals), 338, 339
EMBRYO (See also FETUS)
 development of circulatory system (man), 4, 5;
 (chick), 2; (frog), 1; (pig), 3
 heart rate (mammals), 94
EMPHYSEMA, PULMONARY (See PULMONARY
 EMPHYSEMA)
ENDOCARDITIS, pulse configuration, 156*
ENDOCRINE DISORDERS (mammals), 336-342
 cerebral blood flow and O₂ consumption, 335
EPIGASTRIC ARTERIES, comparative anatomy
 (vertebrates), 13
EPIGASTRIC VEINS, comparative anatomy (verte-
 brates), 13
EPINEPHRINE
 effect on limb blood flow, 133
 general effects, 249
 hemodynamic effects (mammals), 254, 257, 259,
 262, 264, 267, 270
EPN, effect on heart (insect), 279
EPONYCHUM TISSUE, capillary blood pressure, 111
EQUATIONS, REGRESSION, 357-361
ERGOTAMINE
 and dicumarol: effect on circulatory function (rat),
 244
ERYTHROCYTES (See also RBC;
 surface permeability to
 ions (vertebrates), 50
 organic substances (animals), 52
 water (vertebrates), 49

ESERINE, effect on heart (insects), 279
ETHER
 cardiovascular action, 250
 effect on heart (insects), 282
 hemodynamic effects (man, dog), 255, 271
ETHYL ACETATE, effect on heart (insect), 282
ETHYLENE-BIS-4-HYDROXYCOUMARIN, effect on
 clotting time (man, rabbit), 238
ETHYLENE GLYCOL, cell surface permeability to
 (animals), 52; (plants), 53
ETHYL-IODIDE METHOD, determinations for
 cardiac index, 72
 in pathologic conditions, 78
 cardiac output at various temperatures, 77
**ETHYL p-NITROPHENYL THIONOBENZENE PHOS-
 PHATE**, effect on heart (insect), 279
EXCITABILITY, heart muscle: effect of inorganic ions,
 276
EXERCISE
 and age: effect on
 blood pressures, 100, 101
 heart rate, 100, 101
 and anemia: effect on cardiac index, 78
 cardiac index vs O₂ uptake, during, 74, 75*
 cardiac output vs O₂ consumption, during, 77
 during compression, 186
 and cor pulmonale: effect on cerebral function, 333
 effect on circulation time, 115, 116
 in pathologic conditions, 122, 123
 effect on
 limb blood flow, 133
 pulmonary circulation, 130
 and endocrine disorders (mammals), 336
 and heart disease: right heart catheterization values,
 334
 and hyperthyroidism: effect on cardiac index, 78
 O₂ uptake vs arterio-venous O₂ difference, during, 76*
 and tricuspid abnormalities: effect on
 blood flow, 325
 blood pressures, 322
EXTENSOR ILEOTIBIALIS MUSCLE, capillary
 vascularity (salamander), 36
EXTERNAL CAROTID ARTERY, 7*
EXTERNAL ILIAC ARTERY, 7*
EXTERNAL ILIAC VEIN, 8*
EXTERNAL JUGULAR VEINS, 8*
EXTREMITIES (See DIGITAL BLOOD; LIMB BLOOD)
EYE MUSCLE, capillary vascularity (fishes), 36
FACTOR V (See also Ac-GLOBULIN)
 coagulation test, 240
FACTOR VII
 in coagulation stages, 231*
 coagulation test, 240
 preparation and assay, 233
FACTORS, CLOTTING (See CLOTTING FACTORS)
FASTING
 effect on hepatic blood flow (mammals), 129
 and hypercapnia: effect on arterial blood and heart
 rate, 176, 177
FAT, as heart constituent (cattle), 45
FATTY ACIDS, in endocrine disorders (man, dog), 337
FEMORAL ARTERY(IES), 7*
 blood flow, during hypercapnia (dog), 171
 blood pressures, 112, 114
 O₂ content of blood: effect of shock (dog), 321
 pulse waves, 113
 regression equation for mean pressure, 359

FEMORAL VEINS, 8*
 blood pressure, 111
 O₂ content of blood, effect of shock (dog), 321
FEMORAL VENOUS SYSTEM, elasticity (dog), 63
FERRIC CYANIDE (BLOOD), fractional extraction by tissue (dog), 47
FERRIC β_2 -GLOBULIN (BLOOD), transcapillary exchange (guinea pig), 48
FETUS (See also EMBRYO)
 cardiac muscle: capillary vascularity, 37
 circulation time (sheep), 119
FEVER, circulation time during, 123
FIBERS (See specific fiber or tissue)
FIBRILLATION
 atrial incidence in pericarditis, 329
 pulse configuration, 156*

FIBRIN
 in coagulation stages, 231*, 232*
 effect on circulatory function (dog, rabbit), 241

FIBRINOGEN
 clotting test, 240
 in coagulation stages, 231*, 232*
 effect on circulatory function (cat), 241
 effect of shock (dog), 320
 preparation and assay, 233

FIBRINOLYSIS (See also PLASMIN)
 effect on circulatory function (vertebrates), 242

FIBRINOPEPTIDE
 in coagulation stages, 232*
 effect on circulatory function (frog), 241

FIBROBLASTS, surface permeability to water (vertebrates), 49

FICK METHOD, determinations for
 cardiac index, 72-74
 in cardiac tamponade, 329
 in pericarditis, 328
 cardiac output, 72, 77
 in pathologic and other conditions, 78, 79, 158
 left ventricular work, during pregnancy, 158
 peripheral resistance, during pregnancy, 158

FINGER (See also DIGITAL BLOOD, LIMB BLOOD)
 arterial blood pressure, 134
 blood flow, 132-134
 effect of hypoxia, 163
 during pregnancy, 159

FLORIGEN, translocation (plants), 312, 313

FLUORESCENCE, translocation (plants), 286, 290, 291

FLUORINE, as heart constituent, 38

FOOT (See also LIMB BLOOD)
 blood flow, 132, 133

FOREARM (See also LIMB BLOOD)
 blood flow, 132, 133, 180
 effect of hypoxia, 163
 during hypercapnia, 180
 during pregnancy and postpartum, 159
 capillary filtration coefficient, 48
 fractional extraction of diffusible substances from capillaries, 47
 vascular resistance, 180, 181
 during hypercapnia, 180, 181

FORELIMB ARTERIES, comparative anatomy (vertebrates), 14, 15

FORELIMB VEINS, comparative anatomy (vertebrates), 15

FORMALDEHYDE, effect on heart (insect), 282

FORMAMIDE, cell surface permeability to (animals), 52

FORMULAS AND CALCULATIONS proposed for clinical use, 355, 356

FRUCTOSE, translocation (plants), 283, 284, 287

FUNGAL METABOLITES, translocation (plants), 298

GANGLION BLOCKING DRUGS

effect on circulation
 cerebral, 258
 coronary (man, dog), 257
 hepatic (man, dog), 261
 limb, 266, 267
 pulmonary, 263
 renal (man, dog), 261
 systemic (man, dog), 252, 253
 general effects, 247
 site of action, 246

GANGLION STIMULANTS, site of action, 246

GASES, effect on heart (insects), 282

GASTRIC ARTERIES, dimensions, 21

GASTROCNEMIUS MUSCLE
 capillary vascularity (vertebrates), 35, 36
 cell surface permeability to ions (vertebrates), 50

GASTRODUODENAL ARTERY, dimensions, 21

GIBBERELLIC ACID, translocation (plants), 297, 299, 304, 307

GLOBULIN (SERUM)
 in endocrine disorders (dog, rat), 339
 in pericarditis, 328

GLOMERULAR FILTRATION RATE (See KIDNEY)

GLUCOSE, translocation (plants), 283, 284, 286, 287

GLUCOSE (BLOOD)
 capillary permeability coefficient (cat), 48
 effect of shock (dog, rat), 319
 in endocrine disorders (mammals), 337, 338
 utilization in anemia (man and other animals), 349

GLUCOSE (PLASMA), effect of shock, 318

GLUCURONIC ACID (SERUM), in diabetes, 339

GLUTAMIC ACID, translocation (plant), 314

GLUTAMINE, translocation (plants), 314

GLUTATHIONE (BLOOD), in endocrine disorders (man, dog), 338

GLUTEAL MUSCLE, capillary vascularity (rabbit), 35

GLYCEROL, cell surface permeability to (animals), 52; (plants), 53

GLYCEROL (SERUM), effect of shock (rabbit), 320

GLYCINE, translocation (plants), 314

GLYCOCOLL, translocation (plants), 314

GLYCONE, as heart constituent (cat, dog), 40, 41; (horse), 45; (pigeon), 46; (rodents), 42-44

GLYCONE (BLOOD), in diabetes, 337

GLYCOSIDES, CARDIAC
 general effects, 249, 250
 hemodynamic effects of specific drugs (mammals), 254, 257, 259, 262, 264, 270

GLYCYL-GLYCYL-GLYCINE (PLASMA), effect of shock (rat), 320

GRACILIS MUSCLE
 capillary vascularity (dog, frog), 35, 36
 fractional extraction of diffusible substances from capillaries (dog), 47

GRAVES' DISEASE, (mammals), 336-340 (See also HYPERTHYROIDISM)

GRAVITY, ACCELERATION OF
 definition, 190
 effect on circulatory and respiratory function (mammals), 190-195

GRAVITY-INDUCED SHOCK, effect on blood composition (laboratory animals), 319-321

GREAT SAPHENOUS VEIN, 8*
 blood pressure, 111

GROWTH REGULATORS, translocation (plants), 292-308

HEPATIC BLOOD (concluded)
 flow (mammals), 128, 129
 effect of cirrhosis, 128
 effect of hepatitis, 128
 during pregnancy, 128, 159
 vascular resistance (dog), 184, 185
 during hypercapnia (dog), 184, 185

HEPATIC VEIN, 7*

HEPATITIS, effect on hepatic blood flow, 128

HEPATOMEGALY, incidence in pericarditis, 328

HERBICIDES (See also specific herbicide)
 translocation (plants), 292-308

1,2,3,4,5,6-HEXACHLOROCYCLOHEXANE, effect on heart (insect), 279

HEXAMETHONIUM
 effect on heart (insect), 279
 hemodynamic effects (man, dog), 252, 253, 257, 258, 261, 263, 266

HEXOSE PHOSPHATES, translocation (plants), 284

HISTAMINE
 cardiovascular action, 252
 hemodynamic effects (man, dog), 257, 258, 262, 265, 267
 site of action, 246

HPO₄ (BLOOD), effect of shock (dog), 319

H₂PO₄ (BLOOD), effect of shock (dog), 319

HYDROGEN, effect on heart (insect), 282

β -HYDROXYBUTYRIC ACID (BLOOD), in diabetes (man, dog), 337

β -HYDROXYETHYL-4-CHLOROPHENOXYACETATE, translocation (plant), 293

HYOSCINE, translocation (plants), 316

HYOSCYAMINE, translocation (plants), 316

HYPERADRENOCORTICISM, effect on cerebral blood flow and O₂ consumption, 335

HYPERCAPNIA, effect on
 arterial blood pressure (man), 172, 176-179, (dog), 173-175, (mammals), 170, 171
 blood flow (mammals), 171
 cerebral (man), 182
 coronary (cat, dog), 180
 limb (man, dog), 180
 mesenteric (dog), 184
 pulmonary (mammals), 184, 185
 renal (man, dog), 182, 183
 cardiac output (man), 170, 176, (vertebrates), 178

CO₂ pressure
 arterial blood, 176, 177
 cerebral blood, 182, 183
 limb blood, 180, 181
 electrocardiogram (dog), 173-175
 heart rate (man), 176, 177, (vertebrates), 178, 179
 mean pulmonary pressure (mammals), 184, 185
 pH (vertebrates), 172-183
 pulse rate (man), 172, (man, hedgehog), 170, (dog), 173-175
 vascular resistance
 cerebral, 182
 coronary (dog), 180, 181
 hepatic (dog), 184, 185
 limb (man, dog), 180, 181
 mesenteric (dog), 184, 185
 renal (man, dog), 182, 183

HYPEREMIA, effect on capillary blood pressure (man, frog), 111

HYPEROXIA, effect on
 blood flow, 326
 blood pressures (mammals), 168, 169, 323
 blood vessels (mammals), 168, 169
 cardiac index, 79

HYPEROXIA, effect on (concluded)
 cardiac output, 168
 pulse rate (mammals), 168
 vascular resistance, 326

HYPERTENSION (See also PULMONARY HYPERTENSION)
 and age, 102*, 103*
 drugs to relieve, 247
 effect on
 capillary blood pressure, 111
 cardiac index, 78
 circulation time, 123
 digital blood flow and pressure, 134
 limb blood flow, 133
 and hypercapnia
 effect on cerebral blood, 182
 effect on renal blood, 182, 183
 from ionizing radiation (mammals), 197, 199, 203
 systolic and diastolic pressures, 102*, 103*
 right ventricular pulse configuration, 156*

HYPERTENSION, PULMONARY (See PULMONARY HYPERTENSION)

HYPERTENSIVE DRUGS (See SYMPATHOMIMETIC AMINES)

HYPERTHYROIDISM (See also GRAVES' DISEASE)
 circulation time in, 123, 124
 effect on
 cardiac index, 78
 cerebral blood flow and O₂ consumption, 335
 limb blood flow, 133

HYPNOTICS
 cardiovascular action, 250
 hemodynamic effects of specific drugs, 255, 267, 271

HYPOADRENALISM (See ADDISON'S DISEASE)

HYPOGLYCEMIA, effect on cerebral blood flow and O₂ consumption, 335

HYPOGONADISM, effect on cerebral blood flow and O₂ consumption, 335

HYPOINSULINISM (See DIABETES)

HYPOPIUITARISM (See SIMMONDS' DISEASE)

HYPOPROTHROMBINEMIA, dicumarol-induced response to vitamin K, 239

HYPOTENSION
 and age, 102*, 103*
 drugs to relieve, 248, 249
 from ionizing radiation (vertebrates), 196, 201, 205
 systolic and diastolic pressures, 102*, 103*

HYPOTENSIVE DRUGS, effect on
 cerebral circulation, 258
 pulmonary circulation (man, dog), 264, 265

HYPOTHYROIDISM (See also MYXEDEMA)
 effect on
 cerebral blood flow and O₂ consumption, 335
 circulation time, 124

HYPOXIA (See also ALTITUDE, ANOXIA)
 effect on
 blood flow, 326
 blood pressures, 161, 162, 166, 167, 323
 cardiac index, 167
 electrocardiogram, 164-166
 heart rate, 166
 heart size (mammals), 160
 left ventricular work, 167
 maximum cardiac force, 167
 peripheral blood (man, dog), 163
 pulmonary circulation (man, dog), 130, 164
 pulse rate, 162, 167
 stroke volume, 167
 vascular resistance, 326

ILEOCOLIC VEIN, 8*
ILIAC ARTERY(IES), 7* (See also specific artery)
ILIAC GLANDS, 9*
ILIAC REGION, POSTERIOR: development of lymph vessels and nodes, 4, 5
3-INDOLEACETIC ACID, translocation (plants), 292, 293, 296, 297, 299-301, 303-306
INDOLEBUTYRIC ACID, translocation (plants), 301, 306
INFANTS (See NEWBORN)
INFARCTION, MYOCARDIAL: effect on cardiac index, 79
INFERIOR MESENTERIC ARTERY, 7*
 dimensions, 21
INFERIOR MESENTERIC VEINS, 8*
INFERIOR VENA CAVA, 6*-8*
 elasticity (dog), 62, 63
INFERIOR VENA CAVAL PRESSURE, (dog), 184, 185
 in anemia (man and other animals), 350
 during hypercapnia (dog), 184, 185
INJURY (See CHEST INJURY; COLD INJURY)
INNOMINATE ARTERY, 6*
INNOMINATE VEIN, 8*
INORGANIC IONS (See ION(S), INORGANIC; specific ion)
INSECTICIDES (See also specific insecticide)
 translocation (plants), 316
INSUFFICIENCY (See specific)
INSULIN
 cardiovascular action, 251
 effect on cerebral blood, 259
INSULIN (BLOOD), in endocrine disorders (mammals), 337
INTERNAL CAROTID ARTERY, 7*
 effect of hypoxia on peripheral blood (dog), 163
INTERNAL JUGULAR VEIN, 7*, 8*
 blood pressure: effect of breathing O₂ at 3-4 atmospheres, 186
INTERVALS (See ELECTROCARDIOGRAM; specific interval)
INTERVENTRICULAR SEPTAL DEFECT (See SEPTAL DEFECTS)
INTESTINAL LACTEAL LYMPHATICS, lymph flow and lymphocyte content (mouse, rat), 136
INTESTINE(S) (See also SMOOTH MUSCLE)
 blood volume (vertebrates), 70, 71
 resistance to arrested circulation (dog), 135
INTRACARDIAC PULSE, configuration in normal and pathologic conditions, 156*
INTRAPLEURAL PRESSURE, effect of gravitational acceleration (cat), 192
INULIN (BLOOD)
 capillary permeability coefficient (cat), 48
 fractional extraction by tissue (dog), 47
IODIDE ION, cell surface permeability to (plant), 51
2-iodo-4-CHLOROPHENOXACETIC ACID, translocation (plants), 299
2-iodo-3-NITROBENZOIC ACID, translocation (plants), 299
ION(S), INORGANIC (See also specific ion)
 cell surface permeability to (animals), 50, 51;
 (plants), 51
 effect on
 cardiac properties, 276
 heart (man, dog), 273-275, (insects), 277
 ratio required to maintain beat of isolated heart (insects), 276
IRON, as heart constituent (man), 38, (cattle), 45; (dog), 40; (rabbit, rat), 42, 43
IRON (BLOOD)
 in Addison's disease, 338
 protein-bound: in endocrine disorders, 338

IRON (PLASMA), effect of shock (dog), 320
ISOPROPYL N-PHENYLCARBAMATE, translocation (plants), 301
JUGULAR LYMPHATIC TRUNK, comparative anatomy (vertebrates), 17
JUGULAR REGION, development of lymph vessels and nodes, 4, 5
JUGULAR VEIN(S), 7*, 8* (See also specific vein)
 O₂ content of blood: effect of shock (dog), 321
 pulse waves, 113
K₃, effect on circulatory function (dog), 242
KAOLIN, effect on circulatory function (dog), 242
KIDNEY (See also RENAL BLOOD, RENAL PLASMA FLOW)
 blood flow, in endocrine disorders, 339
 blood volume (vertebrates), 70, 71
 effect of ionizing radiation (chicken), 230; (salmon), 213
 glomerular filtration rate, 334
 in anemia, 351
 in cor pulmonale, 334
 in pericarditis, 328
 in rheumatic heart disease, 334
 peripheral blood: effect of hypoxia (dog), 163
KINASE (BLOOD), in coagulation stages, 232*
LABILE FACTOR (See Ac-GLOBULIN)
LACTATE (BLOOD)
 effect of shock (dog), 319
 utilization, in anemia (man and other animals), 349
LACTEAL LYMPHATICS, INTESTINAL: lymph flow and lymphocyte content (mouse, rat), 136
LACTIC ACID (BLOOD)
 in anemia (man and other animals), 351
 effect of shock (dog, rat), 319
 in endocrine disorders (dog), 338
LACTIC ACID N-3-CHLOROPHENYLCARBAMATE, translocation (plants), 292
LACTIC ACID N-PHENYLCARBAMATE, translocation (plants), 292
LAENNEC'S CIRRHOSIS (See also CIRRHOSIS)
 effect on
 cardiac index, 78
 hepatic blood flow, 128
LATERAL CUTANEOUS VEIN, comparative anatomy (vertebrates), 13
LATERAL SEGMENTAL VESSELS, comparative anatomy (vertebrates), 14
LEAD, as heart constituent (man), 38; (cat), 40
LEADS, ELECTROCARDIOGRAPHIC
 amplitudes of waves, in various (man), 142-145;
 (farm animals), 146
 R wave as percentage of RS deflection, in various, 145
 standard limb (man), 137*, 142, (farm animals), 146
 unipolar limb, 138*, 142, 143
 unipolar precordial, 139*, 143-145
LEAF, cell surface permeability to water, 49
LEFT ATRIUM, 6* (See also ATRIUM(IA); ATRIAL BLOOD PRESSURE; MEAN LEFT ATRIAL PRESSURE)
 blood pressures, 112, 114
 in newborn, 114

LEFT ATRIUM (concluded)
 elasticity (dog), 62, 63
 pulse waves, 113

LEFT CAROTID ARTERY, 64

LEFT COLIC VEIN, 8*

LEFT CORONARY ARTERY, 6*

LEFT HEART (See also **LEFT ATRIUM**, **LEFT VENTRICLE**)
 blood pressure, 114

LEFT VENTRICLE, 6* (See also **VENTRICLE(S)**, **VENTRICULAR PRESSURE**, **LEFT VENTRICULAR WORK**)
 blood pressures, 112, 114
 capillary density, surface, and diffusion distance, 34
 capillary vascularity (cat, rabbit), 37
 chemical composition (man), 38, 39; (artio-, perisodactyla), 45; (cat, dog), 39-41; (pigeon), 46, (rabbit), 42
 pulse waves, 113
 weight at various ages, 22

LEFT VENTRICULAR FAILURE
 effect on pulmonary circulation, 130
 right heart catheterization values, 334

LEFT VENTRICULAR WORK
 effect of
 hypoxia, 167
 reduced arterial O₂ saturation, 167
 regression equations, 358, 359

LEG (See also **CALF**, **LEMB BLOOD**)
 blood flow (man), 132, 133, (dog), 180
 effect of temperature, 132
 during hypercapnia (dog), 180
 during pregnancy, 159
 capillary filtration coefficient (cat, dog), 46
 peripheral blood effect of hypoxia, 163

LESION(S) (See specific)

LETHANE 66, effect on heart (insect), 279

LETHANE 384, effect on heart (insect), 279

LEUKOCYTES, surface permeability to water (vertebrates), 49

LEMB BLOOD (See also **CALF**; **LEG**; **FOREARM**, **HAND**; **DIGITAL BLOOD**)
 CO₂ pressure (man, dog), 180, 181
 flow (man), 132, (man, dog), 180
 effect of drugs (man, dog), 266-268
 effect of exercise, 133
 effect of temperature, 132
 in pathologic conditions, 133
 during hypercapnia (man, dog), 180, 181
 vascular resistance (man, dog), 180, 181
 vessels effect of drugs (man, dog), 266-268

LEMB LEADS, **ELECTROCARDIOGRAPHIC** (See **LEADS**, **ELECTROCARDIOGRAPHIC**)

LINDANE, effect on heart (insect), 279

LINGUAL TONSIL, development, 4, 5

LIPID ANTICOAGULANT, preparation and assay, 233

LIPIDS (PLASMA), in diabetes (man, dog), 319

LIPOPROTEIN (SERUM), in diabetes (dog), 319

LIPOMAR
 effect on clotting time, 235
 therapeutic effects (man), 235, (vertebrates), 237

LIQUOID, effect on circulatory function (cat, rabbit), 243

LIVER (See also **HEPATIC BLOOD**)
 blood volume (vertebrates), 70, 71
 effect of ionizing radiation (chicken), 230
 capillary density, surface, and diffusion distance (mouse), 34
 cirrhosis (See **CIRRHOSIS**)
 development, 4, 5

LIVER (concluded)
 fatty effect on cardiac index, 78
 fractional extraction of diffusible substances from capillaries (dog), 47
 function in blood coagulation, 231v
 resistance to arrested circulation (dog), 135

LOBELINE, effect on heart (insect), 279

LONGITUDINAL BLOOD VESSELS, comparative anatomy (vertebrates), 12, 13

LONGITUDINAL LYMPHATIC TRUNK, comparative anatomy (vertebrates), 17

LUMBAR SYMPATHETOMY, effect on circulation time, 124

LUMINA, ARTERIAL: growth at various ages, 20

LUNG(S) (See also **PULMONARY** entries)
 blood flow, in Graves' disease, 359
 blood volume (vertebrates), 70, 71
 effect of ionizing radiation (chicken), 230
 diseases (See also specific disease)
 circulation time in, 124
 effect of anemia (man and other animals), 346, 348
 fractional extraction of diffusible substances from capillaries (man, dog), 47
 reflexes (mammals), 55
 resistance to arrested circulation (dog), 135
 vessels: effect of drugs (man, dog), 263-265

LUNG VOLUMES, during gravitational acceleration, 192, 194

LYMPHATICS, 9*
 comparative anatomy (vertebrates), 16-18
 cervical lymph flow and lymphocyte content (mammals), 136
 intestinal lacteal: lymph flow and lymphocyte content (mouse, rat), 136

LYMPHATIC TISSUE, development (man), 4, 5; (chick), 2; (frog), 1; (pig), 3

LYMPHATIC TRUNKS (See also specific trunk)
 comparative anatomy (vertebrates), 16, 17

LYMPH DUCT, RIGHT: lymph flow and lymphocyte content (mammals), 136

LYMPHEDEMA, effect on digital blood flow and pressure, 134

LYMPH FLOW (mammals), 136

LYMPH HEARTS
 comparative anatomy (vertebrates), 16
 development (frog), 1

LYMPH NODES
 comparative anatomy (vertebrates), 16
 development, 4, 5
 effect of ionizing radiation (vertebrates), 208-212

LYMPHOCYTE(S)
 content of lymph (mammals), 136
 development (chick), 2; (frog), 1
 effect of
 ionizing radiation (vertebrates), 207-227
 shock (rat), 321

LYMPHOID TISSUE, effect of ionizing radiation (mammals), 207-209, 213

LYMPH SACS
 comparative anatomy (vertebrates), 16
 development (chick), 2

LYMPH SINUSES, comparative anatomy (vertebrates), 16

LYMPH VESSELS, development, 4, 5

MAGNESIUM, as heart constituent (man), 38; (cattle), 45, (dog), 40, (pigeon), 46; (rodents), 42, 43

MAGNESIUM (BLOOD)

in endocrine disorders (mammals), 338

effect of shock (dog), 319

MAGNESIUM (PLASMA), effect of shock, 318**MAGNESIUM IONS**, effect on
cardiac properties, 276

heart (man, dog), 273, 274; (insects), 277

MALEIC HYDRAZIDE, translocation (plants), 293,
299-301, 305, 306**MALONAMIDE**, cell surface permeability to (invertebrates), 52; (plants), 53**MAMMARY VEIN, INTERNAL**, elasticity, 62**MANGANESE**, as heart constituent (man), 38; (artio-,
perissodactyls), 45; (dog), 40; (rodents), 42, 43**MANNITOL**cell surface permeability to (mammals), 52;
(plant), 53

translocation (plant), 283

MASSETER MUSCLE, capillary vascularity (mouse,
guinea pig), 35**MEAN ARTERIAL PRESSURE** (man), 176, 177, 184, 185;
(birds), 108, 109; (fishes), 110; (invertebrates),
110; (mammals other than man), 106, 107;
(reptiles, amphibians), 109; (vertebrates),
178, 179 (See also specific organ or body region)

at altitude, 162

in anemia (man and other animals), 349

effect of

albumin infusion, 324

drugs (man, dog), 252-255

explosive decompression (dog), 189

hypoxia, 167

norepinephrine, 323

reduced arterial O₂ saturation, 167

saline infusion, 324

shock, 317

during hypercapnia (man), 176; (dog), 184; (vertebrates), 178, 179

during hyperoxia, 323

in pathologic conditions, 322, 323, 349

MEAN BLOOD PRESSURE, various chambers and
vessels, 112**MEAN FEMORAL PRESSURE**, 114

regression equation, 359

MEAN INSPIRATORY NON-ELASTIC RESISTANCE, in
anemia (man and other animals), 351**MEAN LEFT ATRIAL PRESSURE**, 112, 114

in newborn, 114

MEAN PULMONARY ARTERIAL PRESSURE (man),

112, 114, (mammals), 184, 185

in acquired cardiac disease, 334

in beriberi, 330

effect of

hypoxia, 164

norepinephrine, 323

saline infusion, 324

during hypercapnia (mammals), 184, 185

during hyperoxia, 323

in newborn, 114

during pregnancy, 158

in pulmonary emphysema, 332

URE, in

130

effect of

drugs, 130

norepinephrine, 323

posture, 130

saline infusion, 324

MEAN PULMONARY CAPILLARY PRESSURE (concluded)

during hyperoxia, 323

in pathologic conditions, 130, 322, 323

MEAN RIGHT ATRIAL PRESSURE, 114

in acquired cardiac disease, 334

in cardiac tamponade, 329

effect of

norepinephrine, 323

saline infusion, 324

in newborn, 114

in pathologic conditions, 322, 323, 328-330

in pericarditis, 328

in thyrotoxicosis, 330

MECHOLYL (METHACHOLINE CHLORIDE)

effect on heart (insect), 279

general effects, 248

hemodynamic effects (man, dog), 254, 267, 271

MENADIONE, effect on circulatory function (dog), 242**MESENTERIC ARTERY(IES)**, 7* (See also specific
artery)

comparative anatomy (vertebrates), 14

pressure, in anemia (man and other animals), 350

MESENTERIC BLOOD

in anemia (man and other animals), 350

flow (dog), 184

during hypercapnia (dog), 184, 185

vascular resistance (dog), 184, 185

MESENTERIC CAPILLARIES

blood pressure (vertebrates), 111

filtration coefficient (frog), 46

MESENTERIC VEINS, 8* (See also specific vein)**MESENTERIC VENOUS SYSTEM**, elasticity (dog), 62,
63**METHOXYCHLOR**, effect on heart (insect), 279**a-METHOXYPHENYLACETIC ACID**, translocation
(plants), 293, 297**3,3'-METHYLENE-BIS-4-HYDROXYCOUMARIN**

54

MITRAL INSUFFICIENCY

effect on cardiac index, 78

right heart catheterization values, 334

MITRAL LESIONS, effect on

cardiac index, 78

circulation time, 123

MITRAL MURMUR(S)

characteristics, 149

regurgitant: phonocardiographic configuration, 154*

MITRAL STENOSIS

effect on

blood flow, 325

blood pressures, 322

cardiac index, 78

circulation time, 123

limb blood flow, 133

pulmonary circulation, 130

vascular resistance, 325

murmur in phonocardiographic configuration, 154*

pulse configuration, 156*

right heart catheterization values, 334

MITRAL VALVE, 6*

area: formula, 356

blood flow: formula, 356

ostial breadth, 20

resistance: formula, 356

MOLECULES, transcappillary exchange (mammals), 48**MONOACETIN**, cell surface permeability to (animals),

52, (plant), 53

OXYGEN UPTAKE (concluded)

vs cardiac index

during exercise, 74, 75*

at rest, 74

cerebral: effect of drugs, 258-260

PAGET'S DISEASE, 330

effect on

cardiac index, 79

cardiac output, 79

PALATINE TONSIL, development, 4, 5**PANCREAS**capillary density, surface, and diffusion distance
(guinea pig), 34

residual blood volume, 71

PANTOTHENIC ACID, translocation (plant), 315**PAPILLARY MUSCLE, 6***

capillary density, surface, and diffusion distance, 34

capillary vascularity (cat, rabbit), 37

electrical properties (cat, dog), 56, 58

fiber diameter (cat, dog), 56

PARA-OXON, effect on heart (insect), 280**PARASTERNAL MURMURS, characteristics, 149****PARASYMPATHETIC BLOCKING DRUGS,**

cardiovascular action, 251

hemodynamic effects of atropine sulfate, 267, 270

site of action, 246

PARASYMPATHOMIMETICS

general effects, 248

hemodynamic effects of specific drugs (mammals),

253, 254, 257, 265, 267, 270, 271

site of action, 246

PARATHION, effect on heart (insects), 280**PARIETAL MUSCLE, capillary vascularity (trout), 36****PARITOL, duration and side effects, 240****PATENT DUCTUS ARTERIOSUS**

effect on

blood pressures, 323

cardiac index, 325

O₂ saturation, 152, 153

pulmonary circulation, 130

electrocardiographic findings, 152, 153

murmurs in phonocardiographic configuration, 154*

roentgenographic findings, 152, 153

PENTOSE (PLASMA), effect of shock (rat), 320**PERICARDIAL CALCIFICATION, incidence in pericarditis, 329****PERICARDIAL CAVITY, development, 4, 5****PERICARDIAL TAMPONADE, effect on blood flow, 326****PERICARDITIS, CONSTRUCTIVE, 328**

accompanying conditions, 328, 329

effect on

blood flow, 326

blood pressure, 323

pulmonary circulation, 130

from ionizing radiation (man, dog), 196, 198

pulse configuration, 156*

PERIPHERAL BLOOD

at altitude (man, dog), 163

in diabetes, 339

effect of

ionizing radiation (vertebrates), 214-227

hypoxia (man, dog), 163

shock, 317

during gravitational acceleration (mammals), 191, 192, 194

low resistance: pulse configuration, 157*

PERIPHERAL BLOOD (concluded)O₂ pressure, in anemia (man and other animals), 31

total resistance, during pregnancy, 158

PERIPHERAL EDEMA, in endocrine disorders, 339**PERMEABILITY**

capillary

in anemia (man and other animals), 349

coefficients, various substances (cat), 48

in endocrine disorders (mammals), 336

cell surface

to ions (animals), 50, 51; (plants), 51

to organic substances (animals), 52; (plants), 53, 54

to water (animals, plants), 49

coefficients, capillary: various substances (cat), 48

PEYER'S PATCHES, development, 4, 5

pH

arterial (vertebrates), 178, 179

effect of shock (man), 317; (dog, rat), 321

during hypercapnia (man), 172; (dog), 173-175;

(vertebrates), 178, 179

cerebral arterial, 182, 183, 333

in cor pulmonale, 333

during hypercapnia, 182, 183

coronary arterial (cat, dog), 180, 181

during hypercapnia, (cat, dog), 180, 181

effect on heart (dog), 59

in endocrine disorders (man, dog), 337

PHARYNGEAL TONSIL, development, 4, 5**PHENINDIONE(2-PHENYL-1,3-INDANEDIONE)**

effect on clotting time (dog, rabbit), 238

therapeutic effects (man), 235; (dog, rabbit), 238

PHENOTHIAZINE, effect on heart (insects), 280**PHENYLACETIC ACID, translocation (plant), 306****PHENYLINDANEDIONE, in blood coagulation stages, 231*****2-PHENYL-1,3-INDANEDIONE**

effect on clotting time (dog, rabbit), 238

therapeutic effects (man), 235; (dog, rabbit), 238

PHENYLPROPIONIC ACID, translocation (plant), 306**3-(1'-PHENYLPROPYL)-4-HYDROXYCOUMARIN**

effect on clotting time (vertebrates), 237

therapeutic effects (man), 235; (vertebrates), 237

PHONOCARDIOGRAM, normal and pathologic heart

sounds in the cardiac cycle, 155*

PHONOCARDIOGRAPHIC CONFIGURATION, heart

murmurs, 154*

PHOSPHATE IONS, cell surface permeability to (frog),

50; (plants), 51

PHOSPHATES, HEXOSE, translocation (plants), 284**PHOSPHOLIPIDS, as heart constituents (man), 39;**

(cat, dog), 40, 41; (rodents), 42, 44

PHOSPHOLIPIDS (BLOOD)

in coagulation stages, 232*

in endocrine disorders (man, dog), 337

PHOSPHOLIPIDS (SERUM)

effect of shock (rabbit), 320

in myxedema, 339

PHOSPHORUS

as heart constituent (man), 38, (cat, dog), 40;

(cattle), 45, (rabbit, rat), 42, 43

translocation (plants), 286

PHOSPHORUS (BLOOD), in endocrine disorders

(mammals), 338

PHOSPHORUS (PLASMA), effect of shock (man), 318;

(rabbit, rat), 319, 320

PHOSPHORUS (RBC), effect of shock (rabbit, rat), 319**PHOSPHORUS (WHOLE BLOOD), effect of shock (rat), 320****PHOTOSYNTHATE, translocation (plants), 284-288****PILOCARPINE, effect on heart (insects), 280**

PITUITARY GLAND, residual blood volume (cattle, sheep), 71

PLANTS, cell surface permeability to ions, 51
organic substances, 53, 54
water, 49

PLASMA
Ac-globulin (See Ac-GLOBULIN)
amino acid nitrogen (See AMINO ACID NITROGEN (PLASMA))
antithrombin factor (See ANTITHROMBIN FACTOR)
antithrombin (See ANTITHROMBIN)
CO₂ combining power, in diabetes (man, dog), 339
in endocrine disorders (mammals), 339
factor VII clotting test, 240
natural-occurring coagulants and anticoagulants in, 231*, 232*, 233
renal flow (See RENAL PLASMA FLOW)
volume (mammals), 64-67, (vertebrates other than mammals), 67, 68
in anemia (man and other animals), 349
effect of ionizing radiation (vertebrates), 198-203, 229, 230
in endocrine disorders (mammals), 339
during pregnancy and postpartum, 138

PLASMA EXPANDERS
cardiovascular action, 252
hemodynamic effects of specific drugs, 260, 262, 265

PLASMA THROMBOPLASTIN, formation in blood coagulation, 231*

PLASMA THROMBOPLASTIN COMPONENT
in coagulation stages, 231*
preparation and assay, 233

PLASMIN
in coagulation stages, 231*, 232*
effect on circulatory function (vertebrates), 242

PLASMINOGEN, preparation and assay, 233

PLATEAU PULSE, configuration, 157*

PLATELET FACTORS AND COFACTORS, in coagulation stages, 231*, 232*

PLATELETS
in coagulation stages, 231*, 232*
effect on circulatory function (dog), 242
preparation and assay, 233

PLEURAL EFFUSION, incidence in pericarditis, 329

PNEUMONICOSIS, effect on circulation time, 124

POISEUILLE'S LAW, 355

POLYCYTHEMIA, effect on circulation time, 124

PORTAL VEIN, 74, 8*

O₂ content of blood effect of shock (dog), 321
pressure, in anemia (man and other animals), 350

PORTAL VENOUS PRESSURE (dog), 184
during hypercapnia (dog), 184

POSTERIOR CARDINAL VEINS, comparative anatomy (vertebrates), 12

POSTERIOR ILIAC REGION, development of lymph vessels and nodes, 4, 5

POSTERIOR VENA CAVA, comparative anatomy (vertebrates), 12

POSTPARTUM
circulation time during, 117, 118
circulatory function during, 138, 139

POSTURE
and CO₂ inhalation effect on
blood pressure, 176, 177
cardiac output, 176
heart rate, 176, 177
and compression effect on blood pressures and pulse rate, 166

POSTURE (concluded)

effect on
cardiac output (man), 79, (mammals other than man), 80
circulation time, 115-118
heart murmur intensity, 149
heart rate, 99
O₂ consumption (mammals other than man), 80
pulmonary circulation (man, dog), 130

POTASSIUM, as heart constituent (man), 36, (cat, dog), 40, 41; (cattle), 45; (rodents), 42, 43

POTASSIUM (BLOOD)
effect of shock (dog, rabbit), 119
in endocrine disorders (mammals), 338
fractional extraction by tissue (dog), 47

POTASSIUM CYANIDE, effect on heart (insect), 280

POTASSIUM FLUORESCIN, translocation (plants), 290, 291

POTASSIUM ION(S)
-calcium, -sodium ion ratios required to maintain beat of isolated heart (insect), 276
cell surface permeability to (animals), 50, (plants), 51
effect on
cardiac properties, 276
heart (man), 275, 276, (insects), 277

PQ INTERVAL (ELECTROCARDIOGRAM), effect of hypoxia, 164

PRECARDIAL LEADS, UNIPOLAR, 139*
amplitudes of waves, in various, 143-145
wave as, 143-145

129

PREGNANCY, 158, 159
effect on
cardiac index, 79
cardiac output, 79
circulation time, 115, 117
digital blood flow and pressure, 134
hepatic blood flow, 128

PRESSURE (See specific pressure, specific organ or body region)

PRESSURE EQUIVALENTS, 186

PRETRACHEAL MUSCLE, capillary vascularity (guinea pig), 35

PRIMULONE, translocation (plants), 290

PR INTERVAL (ELECTROCARDIOGRAM)
in Addison's disease, 136
at altitude, 164
in anemia (man and other animals), 347
definition, 140
duration (vertebrates), 141
effect of
hypoxia, 164
inorganic ions (man, dog), 273
ionizing radiation (dog), 199
pH (dog), 59
during hypercapnia (dog), 173-175

231*

hemodynamic effects of specific drugs, 255, 271
site of action, 246

PROFIBRINOLYSIS, in coagulation stages, 231*

PROPIONAMIDE, cell surface permeability to (animals), 52; (plants), 53

PROPYL ALCOHOL, cell surface permeability to (ox), 52

PROPYLENE GLYCOL, cell surface permeability to (animals), 52, (plant), 53

PROTAMINES, effect on circulatory function (vertebrates), 243

PROTEINS (BLOOD)
in anemia (man and other animals), 349
effect of shock (cat), 320

PROTEINS (PLASMA)
effect of shock (dog), 320
in endocrine disorders (mammals), 329

PROTEINS (SERUM)
in pericarditis, 328

PROTHROMBIN (See also **BLOOD COAGULATION**)
in coagulation stages, 231*, 232*
effect of shock (dog), 320
preparation and assay, 233
tests, 240

PROTHROMBIN TIME
in diabetes (dog), 336
effect of
anticoagulants (vertebrates), 236-238
vitamin K therapy, 239

PROTODIASTOLIC SOUND, incidence in pericarditis, 329

PROVENTRICULUS
blood volume (chicken), 71
effect of ionizing radiation (chicken), 230

PTC (See **PLASMA THROMBOPLASTIN COMPONENT**)

PT INTERVAL (ELECTROCARDIOGRAM), effect of hypoxia, 164

PULMONARY ARTERIAL PRESSURE (man), 112, 114; (mammals), 184, 185
in anemia (man and other animals), 349
in aortic insufficiency, 130, 322
in aortic stenosis (man, dog), 130
in beriberi, 330
in cardiac tamponade, 329
in congenital heart malformations, 152, 153
in cor pulmonale, 333
diastolic (See **DIASTOLIC BLOOD PRESSURE**, pulmonary)
effect of
breathing air at 4 atmospheres (rat), 187
drugs (man, dog), 130, 263-265
exercise, 130
hypoxia (man), 161, (man, dog), 130
posture (man, dog), 130
mean (See **MEAN PULMONARY ARTERIAL PRESSURE**)
in newborn, 114
in pathologic conditions (man), 130, 322, 323, 328-331, (man and other animals), 349
in pericarditis, 328
in pulmonary emphysema, 331
systolic (See **SYSTOLIC BLOOD PRESSURE**, pulmonary)
in thyrotoxicosis, 330

PULMONARY ARTERIAL PRESSURE RECEPTORS (mammals), 55

PULMONARY ARTERIOLAR RESISTANCE
in acquired cardiac disease, 334
formula, 355

PULMONARY ARTERIOLES
blood pressures, 112
pulse waves, 113

PULMONARY ARTERY(IES), 6*, 8*
development, 4, 5
effect of ionizing radiation, 197
O₂ saturation of blood, in congenital heart malformations, 152, 153

PULMONARY ARTERY(IES) (concluded)
pulse pressure, 112
pulse waves, 113
thickness of wall, 20
ventricular pressure, 112

PULMONARY BLOOD (man, dog), 130, 184, 185 (See also **LUNG(S)**)
effect of
drugs (man), 130; (man, dog), 263-265
exercise, 130
hypoxia (man, dog), 130
flow
in anemia (man and other animals), 350
calculation, 355
during hypercapnia (mammals), 184, 185
in pathologic conditions (man), 130, (man and other animals), 350
volume, effect of hypoxia, 164

PULMONARY CAPILLARY PRESSURE, effect of hypoxia, 164

PULMONARY DEFLATION RECEPTORS (mammals), 55

PULMONARY DISEASE (See also specific disease)
effect on
cardiac index, 79
circulation time, 123

PULMONARY ELASTIC RESISTANCE COEFFICIENT
in anemia (man and other animals), 351
in Graves' disease, 340

PULMONARY EMPHYSEMA, 331-333
effect on
blood flow, 325
blood pressures, 322, 331-333
pulmonary circulation, 130
vascular resistance, 325
pulmonary arterial O₂ saturation vs blood pressure, 332*
and silicosis effect on circulation time, 123

PULMONARY HYPERTENSION (See also **HYPERTENSION**)
effect on
blood flow, 325
blood pressures, 323
pulmonary circulation, 130, 333
vascular resistance, 325

PULMONARY HYPOTENSIVE DRUGS (man, dog), 264, 265

PULMONARY INSUFFICIENCY, murmur in phonocardiographic configuration, 154*

PULMONARY MURMURS, characteristics, 149

PULMONARY REFLEXES (mammals), 55

PULMONARY RESISTANCE (See also **PULMONARY VASCULAR RESISTANCE**)
in acquired cardiac disease, 334
effect of hypoxia, 164
formula, 355
during hyperoxia, 326
in pathologic conditions, 325, 326, 334

PULMONARY STENOSIS, 152
effect on
blood flow, 325
blood pressures, 323
pulmonary circulation, 130
vascular resistance, 325

PULMONARY STRETCH RECEPTORS (mammals), 55

PULMONARY SYSTOLIC MURMUR, 151

PULMONARY VALVE, 6*
ostial breadth, 20
resistance: formula, 356

PULMONARY VASCULAR BED, elasticity (dog), 62, 63

PULMONARY VASCULARITY, effect of congenital heart malformations, 152, 153

RED BLOOD CELLS (See RBC, ERYTHROCYTES)

REFLEXES, CARDIOVASCULAR AND PULMONARY (mammals), 55

REFLEX HEATING, effect on limb blood flow, 133

REFRACTORY PERIOD, atrial and ventricular muscles: effect of drugs, 269-271

REGRESSION EQUATIONS, 357-361

RELAXATION, heart muscle: effect of inorganic ions, 276

RENAL BLOOD (See also KIDNEY)
arterio-venous O₂ difference, 334
effect of drugs (man, dog), 261-263
flow (man), 126, 334; (dog), 182, 183; (mammals), 127
in anemia (man and other animals), 350
in cor pulmonale, 334
during pregnancy and postpartum, 159
in rheumatic heart disease, 334
during gravitational acceleration (goat), 190, 194
during hypercapnia (man, dog), 182, 183
mean pressure (dog), 182, 183
vascular resistance (dog), 182, 183
in anemia (man and other animals), 350

RENAL CORTEX, capillary density, surface, and diffusion distance (mouse), 34

RENAL FUNCTION
in anemia (man and other animals), 351
in conditions of normal, low, and high cardiac output, 334

RENAL MEDULLA, capillary density, surface, and diffusion distance (mouse), 34

RENAL PLASMA FLOW (man), 126, 334, (mammals), 127 (See also KIDNEY)
in anemia (man and other animals), 351
in cor pulmonale, 334
in pericarditis, 328
in rheumatic heart disease, 334
in Simmonds' disease (dog), 339

RENAL PORTAL SYSTEM, comparative anatomy (vertebrates), 12

RENAL VEIN, 8*
O₂ content of blood: effect of shock (dog), 321

RESIDUAL BLOOD VOLUME, concentration in tissue (vertebrates), 71

RESISTANCE (See specific organ or body region)

RESISTANCE COEFFICIENT, ELASTIC (See PULMONARY ELASTIC RESISTANCE COEFFICIENT)

RESPIRATION
effect of
anticoagulants (vertebrates), 242-244
coagulants (vertebrates), 241, 242
during gravitational acceleration (mammals), 192, 194
heart murmur intensity, during, 149

RESPIRATORY RATE
in cardiac tamponade, 329
effect of O₂ inhalation at 3-4 atmospheres, 186
in endocrine disorders, 340

REST
and anemia: effect on cardiac index, 78
and cor pulmonale: effect on cerebral function, 333
effect on cardiac index and O₂ uptake, 74
vs exercise: effect on circulation time, 115
and heart disease: right heart catheterization values, 334
and hyperthyroidism: effect on cardiac index, 78
and tricuspid abnormalities: effect on blood flow, 325
blood pressures, 322
vascular resistance, 325

RESTING POTENTIAL, cardiac fibers (vertebrates), 56, 57

RETINAL CIRCULATION, during gravitational acceleration (man, monkey), 191

RETRO-PERITONEAL REGION, development of lymph vessels and nodes, 4, 5

RHEUMATIC CARDITIS, effect on circulation time, 123

RHEUMATIC HEART DISEASE, effect on renal function, 334

RHODAMINE B, translocation (plants), 290, 291

RHODAMINE 6G, translocation (plants), 290

RHYTHM, HEART (See also HEART RATE; ARRHYTHMIA(S))
effect of inorganic ions, 276

RIBOFLAVIN, translocation (plant), 315

RIBOFLAVIN (SERUM), in diabetes, 339

RIGHT ATRIAL PRESSURE, 112, 114 (See also ATRIAL BLOOD PRESSURE; MEAN RIGHT ATRIAL PRESSURE)

RIGHT ATRIUM, 6* (See also ATRIUM(IA); MEAN RIGHT ATRIAL PRESSURE)
chemical composition (man), 38; (cattle), 45; (dog), 40, 41; (rabbit), 42
elasticity (dog), 62
O₂ saturation, in congenital heart malformations, 152, 153
pulse waves, 113

RIGHT COLIC VEIN, 8*

RIGHT CORONARY ARTERY, 6*

RIGHT HEART (See also RIGHT ATRIUM; RIGHT VENTRICLE; MEAN RIGHT ATRIAL PRESSURE; RIGHT VENTRICULAR PRESSURE)
blood pressures, 114
O₂ content of blood, during shock (dog), 320

RIGHT VENTRICLE, 6* (See also VENTRICLE(S); RIGHT VENTRICULAR PRESSURE)
capillary density, surface, and diffusion distance, 34
capillary vascularity (cat, rabbit), 37
chemical composition (man), 38, 39, (artio-, perissodactyls), 45; (cat, dog), 39-41; (rabbit, rat), 42, 43
O₂ saturation, in congenital heart malformations, 152, 153
pulse waves, 113
weight at various ages, 22

RIGHT VENTRICULAR HYPERTENSION, pulse configuration, 156*

RIGHT VENTRICULAR PRESSURE, 112, 114
at altitude, 161
in cardiac tamponade, 329
in congenital heart malformations, 152, 153
in cor pulmonale, 333
diastolic (See DIASTOLIC BLOOD PRESSURE, ventricular)
effect of hypoxia, 161
in newborn, 114
during pregnancy, 158
systolic (See SYSTOLIC BLOOD PRESSURE, ventricular)

RIGHT VENTRICULAR PULSE, in constrictive pericarditis: configuration, 156*

RIGHT VENTRICULAR WORK, 347, 348
ultra, 356
347, 348

RUBIDIUM IONS, effect on heart (insects), 283
cell surface permeability to (invertebrates), 50, (plants), 51

RUSSEL VIPER VENOM, effect on circulatory function (vertebrates), 242
R WAVE (ELECTROCARDIOGRAM)
amplitude in various leads (man), 142-145; (farm animals), 146
as percentage of RS deflection, 145

SHOCK, effect on (concluded)
cardiac index, 79
cardiac output, 79
SHUNTS, calculation of, 355
SILICON, as heart constituent (man), 38; (dog), 41
SILICOSIS
effect on
blood pressures, 322
cardiac index, 325
and emphysema: effect on circulation time, 123
SIMMONDS' DISEASE (mammals), 336-340
SINO-ATRIAL JUNCTION, comparative anatomy
(vertebrates), 10, 11
SINO-ATRIAL NODE, 6*
comparative anatomy (vertebrates), 10, 11
effect of drugs, 269-271
electrical properties (rabbit), 56, 58
SINO-ATRIAL NODE, 6* (See SINO-ATRIAL NODE)
SINUS, therapeutic effects, 235
SINUS, CARDIAC: electrical properties (dog), 56, 58
SINUS VENOSUS
comparative anatomy (vertebrates), 10, 11
electrical properties (frog), 57, 58
SKELETAL BLOOD FLOW, in anemia (man and other
animals), 359
SKELETAL MUSCLE
blood volumes (vertebrates), 70, 71
effect of ionizing radiation (chicken), 230
capillary vascularity (vertebrates), 35, 36
resistance to arrested circulation (mammals), 135
SKIN
blood flow, in endocrine disorders, 339
blood volume (vertebrates), 71
circulation, during hypoxia (dog), 163
SLEEP, heart rate during, 99
SMALL INTESTINE, resistance to arrested circulation
(dog), 135

cell surface permeability to (animals), 50; (plants), 51
effect on
 cardiac properties, 276
 heart (insects), 277
-potassium ion ratio required to maintain beat of
 isolated heart (insects), 276
SODIUM AND POTASSIUM

SPLEEN
 blood volume (vertebrates), 70, 71
 development (man), 4, 5; (pig), 3
 effect of ionizing radiation (vertebrates), 209-213, 230

SPLenic ARTERY, dimensions, 21

SPLenic VEIN, 8*

STABLE FACTOR (See **FACTOR VII**)

STACHYSE, translocation (plant), 283

STANDARD LIMB LEADS, electrocardiographic, 137*

amplitudes of waves, in (man), 142, (farm animals), 146

STARCH, translocation (plants), 283, 284, 286

STENOSIS (See specific)

STEWART'S METHOD (sodium chloride), determination for cardiac output (dog), 80

STOMACH, blood volume (rat), 71

STREETER'S HORIZONS, in development of the circulatory system, 4, 5

STREPTOKINASE, in coagulation stages, 231*, 232*

STROKE INDEX, CARDIAC (See also **STROKE VOLUME**, CARDIAC)
 in pathologic conditions, 325, 326

STROKE VOLUME, CARDIAC (See also **STROKE INDEX**, CARDIAC)
 vs age, 72
 in anemia (man and other animals), 349
 in cor pulmonale, 333
 effect of
 body temperature (man, dog), 77
 hypoxia, 167
 reduced arterial O₂ saturation, 167
 shock, 317
 regression equations, 357-359, 361

STRYCHNINE, effect on heart (insects), 281

ST SEGMENT (ELECTROCARDIOGRAM)
 at altitude, 165
 in anemia (man and other animals), 347
 in diabetes, 336
 effect of
 hypoxia, 165
 inorganic ions (man, dog), 273, 274
 pH (dog), 59
 during hypercapnia (dog), 173-175

SUBCLAVIAN ARTERY (IES), 6*, 7*

comparative anatomy (vertebrates), 14, 15
 growth of lumen at various ages, 20

SUBCLAVIAN LYMPH DUCT, lymph flow and lymphocyte content (monkey, rabbit), 136

SUBCLAVIAN VEIN, 8*

SUBMAXILLARY MUSCLE, capillary vascularity (frog), 36

SUBMAXILLARY GLAND, blood volume (mouse, rat), 70, 71

SUBVERTEBRAL LYMPHATIC TRUNKS, comparative anatomy (vertebrates), 16, 17

SUCROSE
 cell surface permeability to, 52
 translocation (plants), 315, 283-288

SUCROSE (BLOOD), capillary permeability coefficient (cat), 48

SUGARS, translocation (plants), 283-287, 314 (see also **CARBOHYDRATES**)

SULFATE (BLOOD), in Addison's disease (man, dog), 338

SULFUR (BLOOD), in Addison's disease (dog), 338

.. .. . dog, 171

.. .. .

circumferential growth, 18*

SUPERIOR VENA CAVAL PRESSURE
 in cardiac tamponade, 329
 in pericarditis, 328

S WAVE (ELECTROCARDIOGRAM), amplitudes in various leads (man), 142-145; (farm animals), 146

SYMPATHETOMY, effect on limb blood flow, 133

SYMPATHOMIMETIC AMINES
 general effects, 248, 249
 hemodynamic effects of specific drugs, 254, 257, 259, 262, 264, 267, 270, 271
 site of action, 246

SYMPATHOMIMETIC BRONCHODILATORS
 cardiovascular action, 251
 hemodynamic effects of isoproterenol HCl, 254, 264, 267, 270

SYSTOLIC BLOOD PRESSURE (man), 105, 176, 177, (birds), 108, 109; (fishes), 110; (invertebrates), 110; (mammals other than man), 106, 107; (reptiles, amphibians), 109
 vs age and sex, 102*-104*, 105
 at altitude, 161, 162
 atrial, in newborn, 114
 basal values, 100
 brachial: regression equation, 358
 during compression, 186
 effect of
 exercise and recovery, 100, 101
 hypoxia, 161, 162, 166
 norepinephrine, 130
 reduced arterial O₂ saturation, 166
 shock, 317
 femoral, in newborn, 114
 during hypercapnia, 170-172, 176, 177
 during hyperoxia, 168
 in pathologic conditions, 102*, 103*, 134, 328-331, 349

pulmonary, 112, 114
 effect of exercise, 130
 effect of hypoxia (man), 130, (man, dog), 164
 effect of norepinephrine, 130
 effect of posture, 130
 in newborn, 114
 in pathologic conditions, 130, 328-331, 333
 radial: regression equation, 358
 right ventricular, 112, 114
 in pathologic conditions, 328-330, 333
 ventricular: various chambers and vessels, 112

SYSTOLIC CLICK, 150

SYSTOLIC INTERVAL, effect of inorganic ions, 273

SYSTOLIC MURMURS, 150
 in congenital heart malformations, 152, 153
 phonocardiographic configurations, 154*

2, 4, 5-T, translocation (plants), 293, 294, 296, 301, 302

TAMPONADE, ACUTE CARDIAC, 329

TEMPERATURE, BODY
 vs cardiac output (man, dog), 77
 effect on
 circulation time (chicken), 119
 digital blood flow and pressure, 134
 limb blood flow, 132
 heart rate (vertebrates), 96-98
 in endocrine disorders (mammals), 336

TEPP, effect on heart (insect), 281

TESTIS
 blood volume (vertebrates), 70, 71
 resistance to arrested circulation (rat), 135

TETRAETHYLENE GLYCOL, cell surface permeability to (invertebrates), 52

TETRAETHYLENE PYROPHOSPHATE, effect on heart (insect), 281

TETRALOXY OF FALLOT, 152
circulation time during, 123
effect of pulmonary circulation, 130

THIAMINE, translocation (plants), 315

THIGH MUSCLE, capillary vascularity (guinea pig), 35

THIOCYANATE, effect on heart (insect), 281

THIOCYANATE (BLOOD)
in diabetes, 137
fractional extraction by tissue (man, dog), 47

THIOUREA
cell surface permeability to (animals), 52, (plants), 53
translocation (plant), 314

THORACIC AORTA
elasticity (man, dog), 60, 61
pulse waves, 113

THORACIC DUCT
comparative anatomy (vertebrates), 16, 17
development (chick), 2; (frog), 1

THORACIC DUCT, ABDOMINAL, lymph flow and lymphocyte content (mouse, rat), 136

THORACIC LYMPH DUCT, lymph flow and lymphocyte content (mammals), 136

THRILLS, HEART in anemia (man and other animals), 345

THROMBIN
in coagulation stages, 231*, 232*
effect on circulatory function (dog, rabbit), 241
preparation and assay, 234

THROMBOANGIITIS OBLITERANS
circulation time in, 124
effect on digital blood flow, 134

THROMBOCID, effect on clotting time (rabbit), 237

THROMBOPHLEBITIS, effect on blood flow and pressure, 134

THROMBOPHLEBITIS MIGRANS, effect on circulation time, 124

THROMBOPLASTIN
in coagulation stages, 231*
effect on circulatory function (dog, frog), 241, 242
generation test, 240

THROMBOPLASTINOGEN (See **ANTHEMOPHILIC FACTOR**)

THYMUS
development (man), 4, 5, (pig), 3
effect of ionizing radiation (mouse, rat), 209, 212

THYROID GLAND, blood volume (rat), 71

THYROTOXICOSIS, 330
effect on circulation time, 124

TIBIAL ARTERIES
blood pressures, 112
pulse waves, 113

TIBIALIS ANTERIOR MUSCLE, capillary vascularity (vertebrates), 35, 36

TIDAL VOLUME, during gravitational acceleration, 194

TIN, as heart constituent (artio-, perissodactyla), 45

TISSUE(S) (See also specific tissue)
adipose capillary density, surface, and diffusion distance (rat), 34
blood flow (man, dog), 47
capillary blood pressure (vertebrates), 111
capillary density, surface, and diffusion distance (vertebrates), 34
capillary filtration coefficients (vertebrates), 46
capillary permeability, various substances (cat), 48
concentration of blood (vertebrates), 70, 71
fractional extraction of diffusible substances (man, dog), 47
natural-occurring coagulants and anticoagulants in, 231*, 132*, 233, 234
resistance to arrested circulation (mammals), 135

TOE, blood flow and pressure, 133, 134

TOLAZOLINE HCL, effect on limb blood flow, 133

TOLUIDINE BLUE, effect on circulatory function (vertebrates), 244

TONSILS, development (man), 4, 5; (pig), 3

TOURNQUET-INDUCED SHOCK, effect on blood composition (laboratory animals), 319-321

TOXAPHENE, effect on heart (insect), 281

TRACHEAL RECEPTORS (mammals), 55

TRANQUILIZERS
cardiovascular action, 251
hemodynamic effects of chlorpromazine, 259, 271

TRANSLLOCATION, organic substances (plants), 283-316

TRAUMA (See **SHOCK**)

TREBURON, duration and alde effects, 240

TRICHLOROACETIC ACID, translocation (plant), 301

2, 3, 5, 6-TRICHLOROBENZOIC ACID, translocation (plant), 297

2, 3, 6-TRICHLOROBENZOIC ACID, translocation (plant), 297

2, 4, 5-TRICHLOROPHENOXACETIC ACID, translocation (plants), 293, 294, 296, 301, 302

TRICUSPID ATRESIA, 152

TRICUSPID INSUFFICIENCY
effect on
blood flow, 325
circulation time, 123
vascular resistance, 325
pulse configuration, 156*

TRICUSPID REGURGITANT MURMUR, phonocardiographic configuration, 154*

TRICUSPID STENOSIS
effect on
blood flow, 325
blood pressures, 322
vascular resistance, 325
pulse configuration, 156*

TRICUSPID VALVE, 6*
ostial breadth, 20
murmurs phonocardiographic configurations, 154*
resistance formula, 156

TRIETHYLENE GLYCOL, cell surface permeability to (animals), 52

TRIMETHYLENE GLYCOL, cell surface permeability to (animals), 52

TROMEXAN
effect on clotting time (vertebrates), 237
therapeutic effects (man), 235, (vertebrates), 237

TRUNCUS ARTERIOSUS
comparative anatomy (vertebrates), 10, 11
development (man), 4, 5, (frog), 1; (pig), 3
electrical properties (frog), 57, 58

TRUNCUS ARTERIOSUS, PERSISTENT, 152

TRYPSIN
in coagulation stages, 231*, 232*
effect on circulatory function (dog, rabbit), 242

T WAVE (ELECTROCARDIOGRAM)
at altitude, 165
amplitudes in various leads (man), 142-145, (farm animals), 146
effect of
hypoxia, 165, 166
inorganic ions, 273, 274
ionizing radiation (vertebrates), 197-200, 203
reduced arterial O₂ saturation, 166
pH (dog), 59
and heart sounds, 155
during hypercapnia (dog), 173-175
low inverted incidence in chronic pericarditis, 329

TWO-TUBE CLOTTING TESTS, 240

UNIPOLAR LIMB LEADS, electrocardiographic, 138*
 amplitudes of waves in, 142, 143
 UNIPOLAR PRECORDIAL LEADS, electrocardiographic, 139*
 amplitudes of waves in, 143-145
 R wave as percentage of RS deflection, in, 145
 UREA
 cell surface permeability to (animals), 52; (plants), 54
 translocation (plants), 293, 300, 303, 314, 315
 UREA (BLOOD)
 capillary permeability coefficient (cat), 48
 in endocrine disorders (mammals), 339
 fractional extraction by tissue (dog), 47
 UREA NITROGEN (BLOOD)
 effect of shock (cat, rat), 319
 in pericarditis, 328
 URIC ACID (BLOOD)
 effect of shock (rat), 319
 in endocrine disorders (mammals), 339
 URIC ACID (PLASMA), effect of shock (man), 318;
 (dog, rat), 319
 UTERINE BLOOD FLOW, during pregnancy and postpartum, 159
 U WAVE (ELECTROCARDIOGRAM), effect of inorganic ions, 274
 VALVE(S) (See also specific valve)
 blood flow: formula, 355
 ostial breadth, 20
 VALVULAR DISEASE (See also specific disease)
 circulation time in, 123
 VALVULAR RESISTANCE, formulas, 356
 VAPORS, effect on heart (insects), 282
 VARICOSE VEINS
 circulation time with, 124
 effect on limb blood flow, 133
 VASCULAR BED, PULMONARY. elasticity (dog), 62, 63
 VASCULAR RESISTANCE (See specific organ or body region)
 VASOCONSTRICTION, effect on limb blood flow, 132
 VASOCONSTRICTOR CENTER, drug stimulants and depressants, 246
 VASOCONSTRICTORS (See SYMPATHOMIMETIC AMINES)
 VASODILATATION, effect on
 digital blood flow, 132, 134
 digital blood pressure, 134
 limb blood flow, 132
 VASODILATORS (See GANGLION BLOCKING DRUGS; ADRENERGIC BLOCKING DRUGS; CHOLINERGIC, MUSCULOTROPIC VASODILATORS)
 VASTUS LATERALIS MUSCLE, capillary vascularity (guinea pig), 35
 VEIN(S), 8* (See also specific vein)
 blood pressure, 111, 112
 comparative anatomy (vertebrates), 12-15
 development (chick), 2, (frog), 1, (pig), 3
 elasticity (man, dog), 62, 63
 pulse waves, 113
 VENA CAVA (See also INFERIOR, POSTERIOR, SUPERIOR VENA CAVA)
 mean pressure, 112
 pulse waves, 113
 VENOM, effect on circulatory function (vertebrates), 242

VENOUS BLOOD

CO₂ pressure, during shock, 317
 O₂ pressure, in anemia (man and other animals), 351
 O₂ saturation
 in diabetes, 337
 effect of shock, 317
 VENOUS BLOOD PRESSURE, 111 (See also PORTAL VENOUS PRESSURE)
 at altitude, 161
 in anemia (man and other animals), 350
 in beriberi, 330
 in cardiac tamponade, 329
 effect of hypoxia, 161
 during gravitational acceleration (mammals), 190, 192, 193
 in pericarditis, 328
 during pregnancy and postpartum, 158
 VENOUS HEMATOCRIT (mammals), 64-67
 VENOUS PRESSURE RECEPTORS (mammals), 55
 VENOUS SYSTEM, elasticity (man, dog), 62, 63
 VENTRICLE(S), 6* (See also LEFT VENTRICLE; RIGHT VENTRICLE)
 action potential of single fibers (vertebrates), 56-58
 atrio-ventricular weight ratio during first year, 22*
 blood flow, during cardiac cycle (man, dog), 148
 chemical composition (man), 38, 39, (artio-, perisodactyls), 45, (birds), 46; (cat, dog), 39-41; (rodents), 42-44
 comparative anatomy (vertebrates), 10, 11
 development, 4, 5
 electrical properties (vertebrates), 56-58
 fiber diameter (vertebrates), 56, 57
 resting potential of single fibers (vertebrates), 56, 57
 weight at various ages, 22
 VENTRICULAR FAILURE, LEFT (See LEFT VENTRICULAR FAILURE)
 VENTRICULAR HYPERTENSION, RIGHT: pulse configuration, 156*
 VENTRICULAR MUSCLE, effect of drugs, 269-271
 VENTRICULAR PRESSURE, 112, 114 (See also RIGHT VENTRICULAR PRESSURE)
 during cardiac cycle (man, dog), 147*, 148
 diastolic (See DIASTOLIC BLOOD PRESSURE, ventricular, right ventricular)
 in newborn, 114
 systolic (See SYSTOLIC BLOOD PRESSURE, ventricular, right ventricular)
 VENTRICULAR RECEPTORS (mammals), 55
 VENTRICULAR SEPTAL DEFECT (See SEPTAL DEFECTS, ventricular)
 VENTRICULAR SEPTUM (See SEPTUM(TA), ventricular)
 VENTRICULAR WORK (See also LEFT VENTRICULAR WORK; RIGHT VENTRICULAR WORK)
 against blood pressure
 effect of hypoxia, 323
 in pathologic conditions, 322
 VERATRINE, effect on heart (insects), 281
 VERATRUM ALKALOIDS
 general effects, 247
 hemodynamic effects of specific drugs, 253, 258, 261, 264, 268
 site of action, 246
 VERBASCOS, translocation (plant), 283
 VESSELS (See also ARTERY(IES), VEIN(S); BLOOD VESSELS, specific vessel)
 effect of anemia (man and other animals), 347
 "VIBRATORY" MURMUR, PRECORDIAL, 151
 VIRUSES, translocation (plants), 309-312
 VISCOSITY (WHOLE BLOOD), during pregnancy and postpartum, 159

VITAL CAPACITY, respiratory
 in disease, 328, 340, 351
 during gravitational acceleration, 194

VITAMIN K
 in coagulation stages, 231*, 232*
 as dietary deficiency: effect on circulatory function
 (mouse), 242
 effect on dicumarol-induced hypoprote thrombinemia,
 239

VITAMINS
 translocation (plant), 315
 hemodynamic effect of specific vitamins (man, dog)
 254, 259, 267

WARE PROTHROMBIN TEST, normal values, 240

WARFARIN
 effect on clotting time (mammals), 237
 therapeutic effects (man), 235, (mammals), 237

WATER
 capillary permeability coefficient (cat), 48
 cell surface permeability to (animals, plants), 49
 as heart constituent (man), 38, (cat, dog), 39, 40,
 (cattle, horse), 45, (rabbit, rat), 42, 43
 translocation (plants), 294

WATER (PLASMA), effect of shock (dog), 320

WATER (RADIOACTIVE), fractional extraction from
 blood by tissue (man, dog), 47

WAVES (BALLISTOCARDIOGRAM), normal standards
 for amplitude, 380

WAVES (ELECTROCARDIOGRAM) (See also specific
 wave)
 amplitudes in various leads (man), 142-145, (farm
 animals), 146
 effect of
 altitude, 165
 hypoxia, 165
 inorganic ions (man, dog), 273, 274
 pH (dog), 59

WEB TISSUE, capillary blood pressure (frog), 111

WEIGHT, atrio-ventricular ratio during first year, 22*

WEIGHT, HEART (man), 24, (birds), 30, 31; (fishes),
 33, 34; (mammals other than man), 25-30,
 (reptiles, amphibians), 32
 vs age and sex, 23
 vs body weight (man), 24; (birds), 30, 31; (fishes),
 33, 34, (mammals other than man), 25-30;
 (reptiles, amphibians), 32
 during first year, 23

WEIGHT, VENTRICULAR, at various ages, 22

WETTERER METHOD, determination for cardiac
 output (mammals), 80

WHITE BLOOD CELLS (See also LEUKOCYTES)
 count, in pericarditis, 328
 effect of ionizing radiation (vertebrates), 214-217,
 219-227

WHOLE BLOOD
 amino acid nitrogen: effect of shock (rat), 319
 phosphorus: effect of shock (rat), 320
 viscosity, during pregnancy and postpartum, 159
 volume: effect of shock, 317

WORK (See VENTRICULAR WORK, EXERCISE)

XYLAN POLYSULFURIC ACID
 effect on clotting time (rabbit), 237
 ester: effect on circulatory function (rabbit), 243

XYLOSE, cell surface permeability to (ground hog, ox),
 52

XYLOSE (BLOOD), fractional extraction by tissue (dog),
 47

YOLK SAC, development, 4, 5

ZINC, as heart constituent (man), 38, (dog), 41,
 (horse), 45; (rabbit), 42

